

Genes Interacting with Occupational Exposures to Low Molecular Weight Agents and Irritants on Adult-Onset Asthma in Three European Studies

Marta Rava, Ismail Ahmed, Manolis Kogevinas,
Nicole Le Moual, Emmanuelle Bouzigon, Ivan Curjuric,
Marie-Hélène Dizier, Orianne Dumas, Juan R. Gonzalez,
Medea Imboden, Amar J. Mehta, Pascale Tubert-Bitter,
Jan-Paul Zock, Deborah Jarvis, Nicole M. Probst-Hensch,
Florence Demenais, and Rachel Nadif

<http://dx.doi.org/10.1289/EHP376>

Received: 25 January 2016

Revised: 18 April 2016

Accepted: 13 June 2016

Published: 9 August 2016

Note to readers with disabilities: *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



National Institute of
Environmental Health Sciences

Genes Interacting with Occupational Exposures to Low Molecular Weight Agents and Irritants on Adult-Onset Asthma in Three European Studies

Marta Rava^{1,2*}, Ismail Ahmed^{3,4,5*}, Manolis Kogevinas^{6,7#}, Nicole Le Moual^{1,8#}, Emmanuelle Bouzigon^{9,10}, Ivan Curjuric^{11,12}, Marie-Hélène Dizier^{9,10}, Orianne Dumas^{1,8}, Juan R. Gonzalez^{6,7}, Medea Imboden^{11,12}, Amar J. Mehta^{11,12,13}, Pascale Tubert-Bitter^{3,4,5}, Jan-Paul Zock^{6,7,14}, Deborah Jarvis¹⁵, Nicole M. Probst-Hensch^{11,12}, Florence Demenais^{9,10†}, and Rachel Nadif^{1,8†}

*These authors are joint first authors, #these authors are joint second authors, and †these authors are joint last authors

¹INSERM, U1168, VIMA: Aging and chronic diseases. Epidemiological and public health approaches, F-94807, Villejuif, France

²Spanish National Cancer Research Centre (CNIO), Genetic & Molecular Epidemiology Group, Human Cancer Genetics Program, 28029 Madrid, Spain

³Inserm UMR 1181 « Biostatistics, Biomathematics, Pharmacoepidemiology and Infectious Diseases » (B2PHI), F-94807 Villejuif, France

⁴Institut Pasteur, UMR 1181, B2PHI, F-75015 Paris, France

⁵Univ Versailles St Quentin-en-Yvelines, UMR 1181, B2PHI, F-78180, Montigny le Bretonneux, France

⁶ISGlobal, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain

⁷CIBER Epidemiología y Salud Pública, Madrid, Spain

⁸Univ Versailles St-Quentin-en-Yvelines, UMR-S 1168, F-78180, Montigny le Bretonneux, France

⁹Inserm, UMR-946, Genetic Variation and Human Diseases unit, F-75010, Paris, France

¹⁰Univ Paris Diderot, Sorbonne Paris Cité, Institut Universitaire d'Hématologie, F-75007, Paris, France

¹¹Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel, Switzerland

¹²University of Basel, Switzerland

¹³Department of Environmental Health, Harvard School of Public Health, Boston, United States

¹⁴Universitat Pompeu Fabra (UPF), Barcelona, Spain

¹⁵Respiratory Epidemiology and Public Health, Imperial College, and MRC-HPA Centre for Environment and Health, London, United Kingdom

Correspondence:

Rachel NADIF, PhD

INSERM UMR-S 1168, VIMA: Aging and chronic diseases. Epidemiological and public health approaches. 16, avenue Paul Vaillant Couturier, F-94807, Villejuif, France. Phone number: 33 (0) 145 59 51 89, Fax number: 33 (0) 145 59 51 69,

E-mail: rachel.nadif@inserm.fr

Running title: candidate genes and occupational exposures in asthma

Competing Financial Interest: Glaxo Wellcome and Glaxo France, producers of asthma medication, partially funded the local studies (respectively Bergen and Paris) in ECRHS II.

Acknowledgments

We also thank all study members and staff involved in data collections in each cohort:

EGEA: We thank the Epidemiological Study on Genetics and Environment of Asthma (EGEA) cooperative group members as follows. **Coordination:** V Siroux (epidemiology, PI since 2013); F Demenais (genetics); I Pin (clinical aspects); R Nadif (biology); F Kauffmann (PI 1992-2012).

Respiratory epidemiology: Inserm U 700, Paris: M Korobaeff (Egea1), F Neukirch (Egea1); Inserm U 707, Paris: I Annesi-Maesano (Egea1-2); Inserm U1168 (ex-CESP/U 1018), Villejuif: F Kauffmann, N Le Moual, R Nadif, MP Oryszczyn (Egea1-2), R Varraso; Inserm U 823, Grenoble: V Siroux. **Genetics:** Inserm U 393, Paris: J Feingold; Inserm U 946, Paris: E Bouzigon, F Demenais, MH Dizier; CNG, Evry: I Gut (now CNAG, Barcelona, Spain), M Lathrop (now Univ McGill, Montreal, Canada). **Clinical centers:** Grenoble: I Pin, C Pison; Lyon: D Ecochard (Egea1), F Gormand, Y Pacheco; Marseille: D Charpin (Egea1), D Vervloet (Egea1-2); Montpellier: J Bousquet; Paris Cochin: A Lockhart (Egea1), R Matran (now in Lille); Paris Necker: E Paty (Egea1-2), P Scheinmann (Egea1-2); Paris Troussseau: A Grimfeld (Egea1-2), J Just. **Data and quality management:** Inserm ex-U155 (Egea1): J Hochez; Inserm U1168 (ex-CESP/U 1018), Villejuif: N Le Moual; Inserm ex-U780: C Ravault (Egea1-2); Inserm ex-U794: N Chateigner (Egea1-2); Grenoble: J Quentin-Ferran (Egea1-2).

SAPALDIA: We thank the team of the Swiss study on Air Pollution and Lung and Heart Diseases in Adults (SAPALDIA).

Study directorate: NM Probst-Hensch (PI; e/g); T Rochat (p), C Schindler (s), N Künzli (e/exp), JM Gaspoz (c)

Scientific team: JC Barthélémy (c), W Berger (g), R Bettschart (p), A Bircher (a), C Brombach (n), PO Bridevaux (p), L Burdet (p), Felber Dietrich D (e), M Frey (p), U Frey (pd), MW Gerbase (p), D Gold (e), E de Groot (c), W Karrer (p), F Kronenberg (g), B Martin (pa), A Mehta (e), D Miedinger (o), M Pons (p), F Roche (c), T Rothe (p), P Schmid-Grendelmeyer (a), D Stolz (p), A Schmidt-Trucksäss (pa), J Schwartz (e), A Turk (p), A von Eckardstein (cc), E Zemp Stutz (e).

Scientific team at coordinating centers: M Adam (e), I Aguilera (exp), S Brunner (s), D Carballo (c), S Caviezel (pa), I Curjuric (e), A Di Pascale (s), J Dratva (e), R Ducret (s), E Dupuis Lozeron (s), M Eeftens (exp), I Eze (e), E Fischer (g), M Foraster (e), M Germond (s), L Grize (s), S Hansen (e), A Hensel (s), M Imboden (g), A Ineichen (exp), A Jeong (g), D Keidel (s), A Kumar (g), N Maire (s), A Mehta (e), R Meier (exp), E Schaffner (s), T Schikowski (e), M Tsai (exp)

(a) allergology, (c) cardiology, (cc) clinical chemistry, (e) epidemiology, (exp) exposure, (g) genetic and molecular biology, (m) meteorology, (n) nutrition, (o) occupational health, (p) pneumology, (pa) physical activity, (pd) pediatrics, (s) statistics

The study could not have been done without the help of the study participants, technical and administrative support and the medical teams and field workers at the local study sites.

Local fieldworkers : Aarau: S Brun, G Giger, M Sperisen, M Stahel, Basel: C Bürl, C Dahler, N Oertli, I Harreh, F Karrer, G Novicic, N Wyttenbacher, Davos: A Saner, P Senn, R Winzeler, Geneva: F Bonfils, B Blicharz, C Landolt, J Rochat, Lugano: S Boccia, E Gehrig, MT Mandia, G

Solari, B Viscardi, Montana: AP Bieri, C Darioly, M Maire, Payerne: F Ding, P Danieli A Vonnez, Wald: D Bodmer, E Hochstrasser, R Kunz, C Meier, J Rakic, U Schafroth, A Walder.

Administrative staff: N Bauer Ott, C Gabriel, R Gutknecht.

ECRHS: The ECRHS data incorporated in this analysis would not have been available without the collaboration of the following individuals and their research teams.

ECRHS Co-ordinating centre. P Burney, D Jarvis, S Chinn, J Knox (ECRHS II), C Luczynska†, J Potts.

Steering Committee for ECRHS II. P Burney, D Jarvis, S Chinn, U. Ackermann-Liebrich, J.M Anto, I.Cerveri, R.deMarco†, T.Gislason, J.Heinrich, C. Janson, N. Kunzli, B. Leynaert, F. Neukirch, J. Schouten, J. Sunyer; C. Svanes, P. Vermeire†, M. Wjst.

Principal Investigators and Senior Scientific Teams for ECRHS II centres within this analysis: Estonia: Tartu (R Jogi, A Soon), France: Paris (F Neukirch, B Leynaert, R Liard, M Zureik), Grenoble (I Pin, J Ferran-Quentin), Germany: Erfurt (J Heinrich, M Wjst, C Frye, I Meyer) Hamburg (K Richter, D Nowak), Norway: Bergen (A Gulsvik, E Omenaa, C Svanes, B Laerum), Spain: Barcelona (JM Anto, J Sunyer, M Kogevinas, JP Zock, X Basagana, A Jaen, F Burgos), Huelva (J Maldonado, A Pereira, JL Sanchez), Albacete (J Martinez-Moratalla Rovira, E Almar), Galdakao (N Muniozguren, I Urritia), Oviedo (F Payo), Sweden: Uppsala (C Janson, G Boman, D Norback, M Gunnbjornsdottir), Umea (E Norrman, M Soderberg, K Franklin, B Lundback, B Forsberg, L Nystrom), Switzerland: Basel (N Kunzli, B Dibbert, M Hazenkamp, M Brutsche, U Ackermann-Liebrich); UK: Norwich (D Jarvis, B Harrison), Ipswich (D Jarvis, R Hall, D Seaton).

†Deceased.

Grant information

The genotyping of all three studies was funded by the French National Agency of Research (ANR-PRSP 2009: IAGO), and by the European Commission (contract n° 018996) (GABRIEL) and the Wellcome Trust grant (WT 084703MA), both awarded to the GABRIEL consortium (a multidisciplinary study to identify the genetic and environmental causes of asthma in the European Community).

EGEA: Research funded by the French Agency of health safety, environment and work (AFSSET, EST-09-15).

SAPALDIA: The Swiss National Science Foundation (grants no 33CS30-148470/1, 33CSCO-134276/1, 33CSCO-108796, , 324730_135673, 3247BO-104283, 3247BO-104288, 3247BO-104284, 3247-065896, 3100-059302, 3200-052720, 3200-042532, 4026-028099, PMPDP3_129021/1, PMPDP3_141671/1), the Federal Office for the Environment, the Federal Office of Public Health, the Federal Office of Roads and Transport, the canton's government of Aargau, Basel-Stadt, Basel-Land, Geneva, Luzern, Ticino, Valais, and Zürich, the Swiss Lung League, the canton's Lung League of Basel Stadt/ Basel Landschaft, Geneva, Ticino, Valais, Graubünden and Zurich, Stiftung ehemals Bündner Heilstätten, SUVA, Freiwillige Akademische Gesellschaft, UBS Wealth Foundation, Talecris Biotherapeutics GmbH, Abbott Diagnostics,.

ECRHS: The co-ordination of ECRHS II was supported by the European Commission, as part of their Quality of Life program. This work was also funded by the US National Institutes of Health (NIH grant 1R01HL062633) and the Carlos III Health Institute of the Spanish Ministry of Health and Consumption (FIS grant 01/3058).

The following bodies funded the local studies in ECRHS II included in this paper: **Albacete:** Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01 and 99/0034-02),

Hospital Universitario de Albacete, Consejeria de Sanidad; **Barcelona:** SEPAR, Public Health Service (grant code: R01 HL62633-01), Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01 and 99/0034-02) CIRIT (grant code: 1999SGR 00241) Red Respira ISCII; **Basel:** Swiss National Science Foundation, Swiss Federal Office for Education & Science, Swiss National Accident Insurance Fund (SUVA), USC NIEHS Center grant 5P30 ES07048; **Bergen:** Norwegian Research Council, Norwegian Asthma & Allergy Association (NAAF), Glaxo Wellcome AS, Norway Research Fund; **Erfurt:** GSF-National Research Centre for Environment & Health, Deutsche Forschungsgemeinschaft (DFG) (grant code FR 1526/1-1); **Galdakao:** Basque Health Dept; **Grenoble:** Programme Hospitalier de Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health, Direction de la Recherche Clinique, Ministere de l'Emploi et de la Solidarite, Direction Generale de la Sante, CHU de Grenoble, Comite des Maladies Respiratoires de l'Isere; **Hamburg:** GSF-National Research Centre for Environment & Health, Deutsche Forschungsgemeinschaft (DFG) (grant code MA 711/4-1); **Ipswich and Norwich:** Asthma UK (formerly known as National Asthma Campaign); **Huelva:** Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01 and 99/0034-02); **Oviedo:** Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01 and 99/0034-02); **Paris:** Ministere de l'Emploi et de la Solidarite, Direction Generale de la Sante, UCB-Pharma (France), Aventis (France), Glaxo France, Programme Hospitalier de Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health, Direction de la Recherche Clinique, CHU de Grenoble; **Tartu:** Estonian Science Foundation; **Umeå:** Swedish Heart Lung Foundation, Swedish Foundation for Health Care Sciences & Allergy Research, Swedish Asthma & Allergy Foundation, Swedish Cancer & Allergy Foundation; **Uppsala:** Swedish Heart Lung Foundation,

Swedish Foundation for Health Care Sciences & Allergy Research, Swedish Asthma & Allergy Foundation, Swedish Cancer & Allergy Foundation.

ABSTRACT

Background: The biological mechanisms by which cleaning products and disinfectants - an emerging risk factor - affect respiratory health remain incompletely evaluated. Studying genes by environment interactions (GxE) may help identify new genes related to adult-onset asthma.

Objectives: To identify interactions between genetic polymorphisms of a large set of genes involved in the response to oxidative stress, and occupational exposures to low molecular weight (LMW) agents or irritants on adult-onset asthma.

Methods: Data came from three large European cohorts: EGEA, SAPALDIA, and ECRHS. A candidate pathway-based strategy identified 163 genes involved in response to oxidative stress and potentially related with exposures to LMW agents/irritants. Occupational exposures were evaluated using an asthma job-exposure matrix and job-specific questionnaires for cleaners and healthcare workers. Logistic regression models were used to detect GxE interactions, adjusted for age, sex and population ancestry in 2599 adults (Mean age: 47 years, 60% women, 36% exposed, 18% asthmatics). P-values were corrected for multiple comparisons.

Results: Ever exposure to LMW agents/irritants was associated with current adult-onset asthma ($OR(95\%CI)=1.28(1.04,1.58)$). Eight SNP by exposure interactions at five loci were found at $p<0.005$: *PLA2G4A* (rs932476, chromosome 1), near *PLA2R1* (rs2667026, chromosome 2), near *RELA* (rs931127, rs7949980, chromosome 11), *PRKD1* (rs1958980, rs11847351, rs1958987, chromosome 14), and *PRKCA* (rs6504453, chromosome 17). Results were consistent across the three studies and after accounting for smoking.

Conclusions: Using a pathway-based selection process, we identified novel genes potentially involved in the adult asthma in relation with occupational exposure. These genes play a role in the NF- κ B pathway involved in inflammation.

INTRODUCTION

Recent reviews regarding the role of environmental risk factors in adult-onset asthma showed that occupational exposures are important causes of asthma in adults (Le Moual et al. 2013; Beasley et al. 2015). Approximately 15% of adult asthma is likely to be attributable to occupational exposures (Toren and Blanc 2009), and occupational asthma is known to be a good model to study the pathophysiology of asthma in general (Malo et al. 2015). Exposure to cleaning agents is an emerging risk factor for adult-onset asthma. Evidence of an adverse effect of cleaning products or disinfectants in asthma mostly comes from studies on occupational risk factors (Siracusa et al. 2013), but a deleterious role of domestic cleaning exposure has also been observed (Quinn et al. 2015; Le Moual et al. 2013; Dumas et al. 2013). Some of the numerous agents contained in cleaning products and disinfectants are chemical sensitizers, but most are hypothesized to act as respiratory irritants (Siracusa et al. 2013). The biological mechanisms by which cleaning products and disinfectants affect respiratory health remain incompletely evaluated (Tarlo and Lemiere 2014; Le Moual et al. 2013; Tarlo 2014). However, inhalation of low molecular weight (LMW) agents and irritants is likely to induce the release of reactive oxygen species through the epithelium, and oxidative stress is known as one of the potential mechanisms causing epithelium injury (Mittal et al. 2014). Furthermore, there is strong evidence that an imbalance between the reducing and oxidizing systems favoring the oxidative state is present in asthma. Reactive oxygen and nitrogen species from endogenous and exogenous sources play a major role in the airway inflammation, and oxidative stress is an important pathophysiological component of asthma (Chung and Marwick 2010; Aldakheel et al. 2016). Thus, to better understand the mechanism of LMW chemical sensitizers and irritants in asthma, it

may be particularly relevant to focus on the oxidative pathway (Tarlo and Lemiere 2014; Tarlo, 2014).

Asthma is a heterogeneous disease, and it is now well established that it is due to a complex interplay of environmental and genetic factors (Kauffmann and Demenais 2012). There have been considerable efforts to characterize the genetic determinants of asthma (Holloway et al. 2010), however, the identified genetic factors explain only a small part of the genetic component of asthma. One of the reasons is that many genetic factors are likely to be involved in the development, the activity and the severity of asthma, and that they act primarily through complex mechanisms that involve interactions with environmental factors (GxE) and with other genes (GxG), notably through pathways and networks. Furthermore, the effect of such genetic factors may be missed if genes are considered individually, regardless of the biological functions they share with other genes or the pathways they are involved in (Liu et al. 2012). Candidate GxE interaction studies conducted on genes involved in the response to oxidative/nitrosative stress and their interaction with environmental exposures in asthma focused more on children than in adults and mostly on outdoor air pollution and smoking (Romieu et al. 2010; Minelli et al. 2011). Furthermore, they have explored a limited number of genes (Kauffmann and Demenais 2012; Kogevinas 2014; Rava et al. 2015). In order to widen the number of genes to be investigated, we recently proposed a candidate pathway-based strategy to select an enriched gene-set for GxE interaction studies (Rava et al. 2013). This gene selection process integrates information on the biological processes shared by genes, the canonical pathways to which genes belong and the biological knowledge related to the environmental exposure under study. This approach represents a powerful alternative strategy between genome wide and candidate approaches to detect relevant associations of environmental exposures with biological markers as well as GxE

interactions.

In the present paper, we hypothesized that genes involved in the response to oxidative stress modify the associations of exposure to LMW agents and irritants with current asthma. We first applied the candidate pathway-based strategy to select oxidative stress related genes that may interact with occupational exposures to LMW agents and irritants in current adult-onset asthma. We then tested for interactive effect of single nucleotide polymorphisms (SNPs) of these genes and LMW agents and irritants on current adult-onset asthma in 2599 participants from the French Epidemiological family-based study of the Genetics and Environment of Asthma (EGEA), the Swiss Cohort Study on Air Pollution and Lung and Heart Disease in Adults (SAPALDIA), and the European Community Respiratory Health Survey (ECRHS).

METHODS

Study population

Data came from three multicentre epidemiological European studies: the French Epidemiological family-based study of the Genetics and Environment of Asthma ([EGEA], Kauffmann et al. 1997; Kennedy et al. 2000) (see Figure S1A), and two population-based studies: the Swiss Cohort Study on Air Pollution and Lung and Heart Disease in Adults ([SAPALDIA], Downs et al. 2007; Mehta et al. 2012; Ackermann-Liebrich et al. 2005) (see Figure S1B), and the European Community Respiratory Health Survey ([ECRHS], ECRHS 2002; Kogevinas et al. 2007) (see Figure S1C). All three cohorts applied comparable study design and highly comparable questionnaires. Participants included in the analyses were derived from the entire study population for EGEA and from the nested case-control samples within ECRHS (Smit et al. 2014) and SAPALDIA cohorts (Curjuric et al. 2012). Participants had genome-wide SNP data, occupational history regarding LMW agents and irritants, especially cleaning/disinfecting

products, and data on adult-onset asthma and relevant covariates such as age, sex and smoking status.

Ethical approval was obtained in each study from the appropriate institutional ethics committees, and written informed consent was obtained from each participant. Detailed cohort descriptions are given in the online supplemental material.

Current adult-onset asthma

In all cohorts, current asthma was defined as ever diagnosis of asthma (Moffatt et al. 2010; Smit et al. 2014) and presence of respiratory symptoms (wheeze; nocturnal chest tightness; attacks of breathlessness after activity, at rest, or at night; asthma attacks) or using asthma medications in the last 12 months. Participants without asthma were those without asthma at baseline and at follow-up. Participants with ever asthma, but without symptoms or treatment in the last 12 months were excluded. Since we were interested in participants who may have developed asthma due to occupational exposure, we restricted the current adult-onset asthma definition to asthmatics with an age of onset \geq age 16.

Occupational exposures to LMW agents and irritants

In all cohorts, occupational history was recorded by interview and job codes were linked to an asthma-specific job-exposure matrix (JEM) evaluating exposure to 22 agents, and including a local expert re-evaluation step (Kennedy et al. 2000). Healthcare workers and cleaners were further asked to answer a job-specific questionnaire regarding exposure to cleaning/disinfecting products.

In the present study, we considered only exposures to substances hypothesized to cause irritant-asthma, or to cause asthma through mechanisms induced by LMW agents. Exposure to LMW agents was evaluated by the JEM, and included products typically classified as LMW agents (e.g., highly reactive chemicals, metals), but also mixed environments with potential exposure to high molecular weight (HMW) and LMW agents (e.g., agriculture, textile). Exposure to irritants was evaluated 1) by the JEM, for high peaks irritant exposure, and 2) using self-reported exposure to cleaning/disinfecting products, with a focus on those that are more likely to be respiratory irritants (see Supplemental Material Table S1 for more details). Participants who had ever been exposed to any of the LMW agents, mixed environments, irritants or any specific cleaning/disinfecting products were classified as “exposed”. Unexposed participants were those who were never exposed to any of the 22 agents of the asthma JEM (including HMW agents) or to other agents potentially at risk for respiratory health (vapors, general dusts, gases and fumes) evaluated by another JEM (ALOHA JEM, Matheson et al. 2005; de Jong et al. 2015). The three cohorts used same definitions.

Genotyping

The three cohorts (EGEA, SAPALDIA and ECRHS) were part of the European Gabriel consortium (<http://www.gabriel-fp6.org/>) for asthma genetics (Moffatt et al. 2010), and constitutes the ESE consortium. Participants were genotyped using Illumina 610 Quad array (Illumina, San Diego, CA) at the Centre National de Génotypage (CNG, Evry, France). Stringent quality criteria, as detailed by Imboden et al. (2012), were used to select both individuals and SNPs for analysis. The quality control (QC) criteria were call rate $\geq 97\%$, minor allele frequency $\geq 5\%$, and Hardy-Weinberg (HW) P-value $>10^{-4}$.

Gene coverage, which indicates the fraction of common HapMap markers successfully tagged by the set of selected SNPs, was obtained with Haploview 4.2 (Barrett et al. 2005). We specified that all HapMap markers being captured by the set of tags should be correlated at $r^2 \geq 0.8$ with at least one marker in the set.

Gene selection through a candidate pathway-based strategy

For this study, a large set of genes was selected according to the candidate pathway-based strategy previously published (Rava et al. 2013). Briefly, the selection process followed three steps: **Step1- Gene selection:** we used the Gene Ontology (GO) database (Gene Ontology Consortium (Ashburner et al. 2000, <http://amigo2.berkeleybop.org/amigo>, version 1.8) to select genes involved in the "response to oxidative stress" (GO:0006979). This list was further enlarged by literature reviews of asthma related genome-wide association studies, and biological studies on response to oxidative stress related to environmental exposures of interest; **Step 2 - Pathway enrichment:** using Ingenuity Pathway Analysis (IPA, <http://www.ingenuity.com/>) we identified the canonical pathways that contained at least 5 genes out of the set of the genes selected in step 1 and which were significantly enriched in these genes ($p < 0.05$); **Step 3 - Environment integration:** we selected the subset of pathways identified at step 2 that contained genes selected at step 1 expected to be involved in the response to oxidative stress potentially caused by occupational exposure to LMW agents or and irritants. This strategy is fully detailed in Rava et al. (2013).

For each of the genes belonging to the selected pathways, we examined all SNPs passing the quality control QC process and lying from 20 kb upstream to 20 kb downstream of the gene (UCSC genome browser hg18 assembly; Build 37.1).

Statistical analysis strategy

The three ESE cohorts were pooled to increase statistical power as done before (Smit et al. 2012, 2014); this also allowed assessing consistency of results across cohorts. SNP-occupational exposure interactions were investigated using a logistic regression model that included the SNP effect assumed to be additive, a binary exposure (E) variable (1=exposed, 0=unexposed) and a multiplicative term for SNPxE interaction. All models were adjusted for age, sex and the four first principal components (PCs) to account for population stratification as previously done (Smit et al. 2014). No additional adjustment for study was done since PCs are capturing any possible variability caused by geographical location. Smoking status was further included as a potential confounder.

Test of SNPxE interaction was based on a Wald test. To account for multiple testing, the Benjamini and Hochberg procedure (1995) was implemented. For interactions belonging to the top 1% of P-values distribution, consistency of interaction effect estimates across studies was assessed by use of the Cochran Q test statistic and the extent of heterogeneity was measured by I^2 , which ranges from 0% to 100%. The I^2 statistic describes the percentage of variation across studies that is due to heterogeneity rather than chance (Higgins and Thompson 2002; Higgins et al. 2003) and $I^2 = 100\% \times (Q\text{-degree of freedom})/Q$. I^2 values of 0%-24% suggest little heterogeneity, of 25%-49% reflect moderate heterogeneity, of 50%-74% reflect large heterogeneity, and of >75% reflect very large heterogeneity (Viechtbauer and cheung 2010). As smoking may also induce oxidative stress, a sensitivity analysis excluding current smokers was performed. The robustness of the results to the family dependency existing in the EGEA study

was investigated by using generalized estimating equations (GEE) with an exchangeable working correlation matrix to take into account potential clustering within families.

For each of the genes belonging to the selected pathways, interactions with occupational exposure for current adult-onset asthma were also investigated at the gene level by using the versatile gene-based test (VEGAS, Liu et al. 2010). This gene-based statistic sums up the chi-square test statistics of SNPxE interaction (square of the Wald test statistics) for all SNPs of a gene. The correlation (r^2) between these statistics is taken into account by computing an empirical P-value through Monte-Carlo simulations using the linkage-disequilibrium pattern of HapMap Utah residents with ancestry from northern and western Europe (CEU) reference sample; this empirical P-value is estimated by the proportion of simulated test statistics that exceeds the observed gene-based test statistic. The empirical P-values were then adjusted for multiple testing using the method of Benjamini and Hochberg.

eQTL analysis, functional annotation and chemical–gene/protein interactions

We investigated whether the SNPs (or their proxies, $r^2 \geq 0.8$) found to interact with occupational exposures to LMW agents or irritants were cis-expression quantitative trait loci (cis-eQTLs). We used the eQTL browser (<http://www.gtexportal.org/home/>) that includes e-QTL data from many tissues; the Genotype-Tissue Expression project (GTEx, Gibson 2015). Furthermore, functional annotations of these SNPs (or proxies) were done using the HaploReg tool (<http://www.broadinstitute.org/mammals/haploreg/haploreg.php>). HaploReg annotates SNPs in terms of predicted ROADMAP and ENcyclopedia Of DNA Elements (ENCODE), chromatin states (promoter and enhancer histone modification signals), DNase I hypersensitivity sites, and transcription factor (TF) and protein binding sites.

Furthermore, curated [chemical–gene interactions|chemical–disease|gene–disease] data were retrieved from the Comparative Toxicogenomics Database (CTD, Davis et al. 2014, MDI Biological Laboratory, Salisbury Cove, Maine, and NC State University, Raleigh, North Carolina. World Wide Web, URL: <http://ctdbase.org/>). [April, 2016]. CTD is a robust, publicly available database that aims to advance understanding about how environmental exposures affect human health. It provides manually curated information about chemical–gene/protein interactions, chemical–disease and gene–disease relationships.

RESULTS

Data description

The study population included 2599 participants with a mean age of 46.7 years and 60% of women (Table 1). ECRHS participants were younger, and the proportion of women was lower in SAPALDIA. Almost half of the participants were never smokers. The proportion of current smokers varied from 18.6% (EGEA) to 31.4% (ECRHS), and 463 had current adult-onset asthma. Among the 927 exposed participants, 25.4% were exposed to LMW agents only, 4.4% were exposed to irritants only, 23.7% were health care workers or cleaners (exposure to cleaning products), 12.6% were exposed to mixed environment only, and 33.9% had combined exposures (i.e. two or more of the aforementioned exposures).

A positive and significant association was found between lifetime occupational exposure to LMW agents or irritants and current adult-onset asthma: age and sex adjusted pooled Odds-Ratio (OR_a)=1.28; 95% Confidence Interval (95%CI) 1.04-1.58). Across the three cohorts, the associations between exposure and asthma were: age and sex adjusted OR_a=1.09 (95%CI: 0.72-1.65; n=122/689, cases/all) in EGEA, 0.89 (95%CI: 0.56-1.42; n=107/574) in SAPALDIA, and 1.55 (95%CI: 1.15-2.08; n=234/1336) in ECRHS.

Genes selected with the candidate pathway-based strategy

Step1- Gene selection: 387 genes were selected through GO and further enriched by literature reviews and biological studies to get a list of 411 genes; **Step 2 - Pathway enrichment:** we identified 277 pathways that contained at least 5 genes out of the 411 genes selected at step 1 and which were enriched in these genes ($p<0.05$); **Step 3 - Environment integration:** 17 of the 277 pathways were further selected because they included genes involved in response to oxidative stress and potentially related with exposures to LMW agents or irritants. These pathways had pathway enrichment P-values ranging from 0.03 to 1.58×10^{-31} (Excel File Table S1) and included from 5 up to 47 genes (15-20 genes on average); more than 50% of the genes were involved in more than one pathway. The final analyzed set included a total of 163 unique genes (Excel File Tables S2) and 3297 SNPs.

Analysis of SNPs x occupational exposure interactions

At the SNP level, none of the interactions with LMW/irritants on current adult-onset asthma reached the significance level after correction for multiple testing ($P=0.05/3297=1.5 \cdot 10^{-5}$). However, we selected 14 interactions belonging to the top 1% of P-values distribution ranked from lowest (top) to highest (bottom) (Supplemental Material Table S2). Among these 14 interactions, 8 interactions at five loci showed little heterogeneity ($I^2 < 24\%$) between the three studies (Table 2 and Supplemental Table S3): rs932476 in *PLA2G4A* (phospholipase A2, group IVA (cytosolic, calcium-dependent) gene, chromosome 11, $P=0.005$), rs2667026 near *PLA2R1* (phospholipase A2 Receptor 1, chromosome 2, $P=0.005$), rs931127 and rs7949980 near *RELA* (V-Rel Avian Reticuloendotheliosis Viral Oncogene Homolog A gene, chromosome 11, $P=0.001$

and $P=0.003$ respectively), rs1958980, rs11847351, and rs1958987 in *PRKD1* (protein kinase D1, chromosome 14, P-values ranging from 0.004 to 0.005), and rs6504453 in *PRKCA* (protein kinase C, alpha, chromosome 17, $P=0.003$). The two SNPs near *RELA* were in moderate linkage disequilibrium (LD, $r^2=0.65$), whereas the three SNPs in *PRKD1* were in strong LD ($r^2>0.8$, see Figures S2A to S2E). Further, rs932476 in *PLA2G4A* and rs931127 in *RELA* were also marginally associated with asthma ($P=0.0036$ and $P=0.035$ respectively, Table 2). Similar interactive and marginal estimates were obtained by taking into account family dependency (Supplemental Material Table S4) or by adjusting for study/centre (data not shown). Excluding current smokers from the analysis showed consistent results except for the *PLA2G4A* gene (Supplemental Material Table S4). Finally, adjusting for smoking gave similar estimates (Supplemental Material Table S4).

Associations between SNPs and current adult-onset asthma in unexposed and exposed participants are reported in Figure 1. "Flip-Flop" interactions were observed. Near *RELA*, the risk of current adult-onset asthma was increased in G carriers of rs931127 and in T carriers of rs7949980 among exposed participants ($OR=1.54$, $P=2\times 10^{-4}$, and $OR=1.40$, $P=0.005$ respectively), whereas inverse but not significant effects were observed among unexposed participants. The risk was also increased - although not statistically significant - among exposed participants in G carriers of rs2667042 near *PLA2RI*, whereas inverse and significant effects ($OR=0.74$, $P=0.009$) were observed among unexposed participants. On the contrary, the risk of current adult-onset asthma was decreased but not significantly among exposed participants in G carriers in *PLA2G4A*, in G carriers of rs1958980 or G carriers of rs11847351 or T carriers of rs1958987 in *PRKD1*, and in T carriers of rs6504453 in *PRKCA*, and significantly in T carriers

of rs6504453 in *PRKCA* ($OR=0.79$, $P=0.05$), whereas inverse and significant effects were observed among unexposed participants ($OR=1.25$ to 1.50 , $P=0.01$ to 3×10^{-4}).

PRKD1 and *PRKCA* are involved together in the "NRF2-mediated Oxidative Stress Response" pathway, in association with *RELA* in three other pathways: "Xenobiotic Metabolism Signaling", "Production of Nitric Oxide and Reactive Oxygen Species in Macrophages", and "N-formyl-methionyl-leucyl-phenylalanine (fMLP) signaling in neutrophils", or in association with *PLA2G4A* in the "CCR3 signaling in eosinophils" pathway (Excel File Table S3). Furthermore, *RELA* and *PRKCA* are involved together in the "Apoptosis signaling" pathway, and *RELA*, *PLA2G4A* and *PLA2R1* are involved together in the "Antioxidant Action of Vitamin C" pathway. Gene coverage for the SNPs in *PLA2G4A*, *PLA2R1*, *PRKD1* and *PRKCA* were quite high: 55% ($r^2=0.98$), 76% ($r^2=0.97$), 74% ($r^2=0.96$) and 68% ($r^2=0.96$) respectively. A low coverage was observed for *RELA* (<10%).

Analysis of Gene x occupational exposure interactions

At the gene level, *RELA* and *PRKD1* were among the top gene interaction with occupational exposures to LMW/irritants that were detected by the gene-based test among all 163 studied genes ($P\text{-value}=0.009$ and $P=0.04$ respectively, Supplemental Material Table S5), but none of them reached the significance level after correction for multiple testing.

eQTL, functional annotations and chemical–gene/protein interactions

Using the eQTL browser GTEx, we found that the T allele at rs6504453 in *PRKCA* was associated with increased gene expression in lung tissue (see Figure S3, $P=0.017$). No eQTL was

found among the SNPs (or proxies) interacting with exposures at *PLA2G4A*, *PLA2R1*, *RELA* and *PRKD1* loci.

Using the functional annotation HaploReg tool v3, we found that the SNPs rs932476 in *PLA2G4A*, rs2667026 near *PLA2R1*, rs931127 and rs7949980 near *RELA*, and rs1958980, rs11847351 and rs1958987 in *PRKD1* mapped to marks of active regulatory elements notably in B cells, small airways epithelial cells, and lymphoblastoid cell lines. These marks included enhancer histone marks, DNase hypersensitivity sites, and binding protein sites for NFKB, Histone Deacetylase 2 (HDAC2), and Nuclear factor erythroid 2-related factor 2 (Nrf2) (Excel File Table S4).

Further, from the Comparative Toxicogenomics Database, we found that chlorine, formaldehyde and hydrogen peroxide have been reported to modify the localization of *PRKCA* protein, the expression of *PLA2R1* or *PLA2G4A* mRNA, and the expression and the activity of *RELA* protein (Excel File Table S5). We also found that exposures known to contain compounds with irritant properties (air pollutants and vehicle emissions) modified the expression of *PRKD1* mRNA and methylation of *PLA2R1* (Excel File Table S5).

DISCUSSION

This study identified interactions between genetic variants near or within five genes, *PLA2G4A*, *PLA2R1*, *RELA*, *PRKD1* and *PRKCA*, and occupational exposures to LMW agents or irritants for current adult-onset asthma. The evidence rests on the results obtained in pooled data of three large European epidemiological studies and the consistency of results across these studies. Functional annotations of the interacting SNPs at these loci and functional supports specific for the GxE interactions detected suggest that a few of these SNPs might be involved in regulatory mechanisms.

Up to now, a limited number of genes were explored in GxE interaction studies conducted with candidate gene approaches. The most commonly studied genes were those coding for the enzymes NAD(P)H dehydrogenase [quinine] 1 (*NQO1*), the glutathione S-transferases (*GSTs*), the heme oxygenase 1 (*HMOX-1*), the catalase (*CAT*) and the superoxide dismutase (*SOD*) (Minelli et al. 2011). Our study relies on an original strategy to select and enlarge the list of candidate genes. Supported by biological knowledge, we think this approach allows a good tradeoff between GEWIS and candidate gene approaches. It is interesting to note that our set of 163 genes included the few genes mentioned previously and studied in interaction with other exposures related to oxidative stress (smoking, outdoor air pollution) in asthma following a candidate gene approach. We cannot exclude that our selection may have overrepresented the anti-oxidative defense, and may lose a number of relevant genes that are not targeted by our analysis. However, we were able to highlight that genes modulating exposure to LMW agents and irritants have all a prominent role in the NF-kappa-B pathway and our strategy had also the capacity to generate new hypotheses. One of the difficulties in GxE studies is the need of large studies or consortia to detect significant interaction, which in turn might be affected by heterogeneity in both outcome and exposure definition of the participating studies. To overcome these limitations, definition of adult-onset asthma as well as those of occupational exposures to LMW agents or irritants were harmonized across the three epidemiological studies, and genotyping was performed identically in the three studies in the framework of the European Gabriel consortium asthma GWAS (Moffatt et al. 2010). Despite the fact that the three studies were pooled, we obtained 463 exposed participants with adult onset asthma to detect GxE interactions. This small number of exposed cases may have hampered our findings, and we acknowledge that the lack of replication is a limitation. However, replication is very difficult

because EGEA, SAPALDIA and ECRHS are, to the best of our knowledge, the only three cohorts having such specific information on occupational exposures (the asthma-specific JEM with the expertise step that increases the precision of exposure assessment, and the specific questionnaires in cleaners and health care workers). By adding other studies using only the asthma-specific JEM, we would lose part of the specificity of our analysis. None of our GxE interaction tests reached the significance level after correction for multiple testing, so we focused on SNPs with P-values for SNPsxE in top 1% of the distribution, and reduced false positives by only selecting consistent interactions across the three studies. As regards the method used, various study designs and statistical methods have been proposed to investigate GxE interactions (Liu et al. 2012). We used the classical GxE interaction test based on a case-control design, which may not be the most powerful approach. Indeed, when one can assume independence between exposure and SNPs, it has been shown that case-only based approaches (Mukherjee et al. 2008) are better alternatives. However, these approaches could not be applied to our study because our gene-selection process aimed at selecting genes potentially associated with the environmental exposure due to their biological function. We further repeated the analyses using a joint test of gene and gene-environment interaction (Dai et al. 2012) but similar results were obtained (data not shown).

Irritant-induced asthma is usually described as a separate, “nonsensitizing”, type of occupational asthma (Maestrelli et al. 2009; Tarlo and Lemiere 2014). On the other side, low molecular weight agents are generally classified as sensitizers, although most of them are not associated with the production of specific IgE (Tarlo and Lemiere 2014). The precise mechanisms linking irritants and LMW chemicals to asthma are poorly known, and it is therefore challenging to classify most asthmogenic chemicals (*e.g.*, cleaning products) into definite categories. However,

both occupational exposures to LMW chemicals and irritants may result in oxidative stress (Dumas et al. 2015). We could thus investigate a relatively broad spectrum of exposures by carefully selecting genes through our pathway-based strategy integrating hypotheses about the environment. Smoking is also known to be related to oxidative stress. Our results remained almost consistent after running analysis without current smokers or after accounting for smoking, suggesting that the detected interactions were not due to the effect of smoking.

To our knowledge, none of our findings have been reported previously by published GWASs of asthma (GWAS-Catalog of Published Genome-Wide Association Studies, <http://genome.gov/gwastudies>, Hindorff et al.), or by GEWIS studies in asthma. Differences in length of microsatellite sequences in the promoter region of *PLA2G4A* were reported between patients with severe asthma and healthy controls, with a direct impact on mRNA and protein expression, suggesting a role in asthma pathogenesis (Sokolowska et al. 2010). Scarce candidate G x occupational exposure interaction studies have been published for asthma (Kauffmann et al. 2010; Kogevinas 2014; Smit et al. 2014; Cherry et al. 2015). Focusing on occupational exposures, *CTNNA3* (catenin alpha 3, alpha-T catenin) was reported by GWAS as the strongest candidate gene for toluene diisocyanate (TDI)-induced asthma in Korean patients (Kim et al. 2009), and only one genome-wide study of interaction (GEWIS) was published that identified novel susceptibility loci for occupational exposure to biological dust, mineral dust, and gases and fumes in relation to FEV₁ levels (de Jong et al. 2015).

Interestingly, all the genes we detected play a role in the NF-kappa-B pathway. NF-kappa-B is an ubiquitous transcription factor involved into the mechanism whereby oxidants affect the pathophysiology of asthma (Schuliga 2015). The genetic variants interacting with exposure do not belong to protein-coding regions, but are more likely to have a regulatory function, as

indicated by the functional annotations of a few of these SNPs. *RELA* encodes the RelA protein that is complexed with NFKB1, the most abundant form of NF- κ B. *PRKD1* encodes a serine/threonine kinase, called PKD1 that activates NF- κ B in response to oxidative stress conditions (Sundram et al. 2011; Storz 2007). Exposure to photochemically altered air pollutant mixture, was associated with a decrease in expression of *PRKD1* mRNA in human lung epithelial cells (Rager et al. 2011). On the contrary, exposure to Zinc Oxide nanoparticles, that is associated with acute pulmonary oxidative stress and inflammation (Vandebriel and de Jong 2012), was reported to activate NF- κ B in human bronchial epithelial cells through a mechanism that involves RelA-NF- κ B phosphorylation (Wu et al. 2010). Interestingly, in a similar opposite manner, we found negative associations between genetic variants in *PRKD1* and adult-onset asthma (decreased risk), and positive associations between genetic variants near *RELA* and adult-onset asthma (increased risk) in participants exposed to LMW or irritant agents. All these effects are "Flip-flop effects", and we can only speculate on the mechanism behind an opposite effect among the exposed and unexposed subjects. Finally, the protein encoded by *PRKCA* was suggested as a regulator of NF- κ B-induced expression of genes involved in inflammatory responses (Nakashima 2002), and was associated with generation of reactive oxygen species through a biological interaction with other genes including member of the mammalian PLA2 family (Chi et al. 2014). A role of the secretory phospholipase A2 receptor in the development of asthma was recently reported in animal models of asthma and in human lung cells (Murakami et al. 2014; Leslie 2015). It is noteworthy that the SNPs interacting with exposure identified by this study mapped to protein binding sites that included NFKB, Histone Deacetylase 2 (HDAC2) whose activity is regulated by oxidative stress and Nuclear factor erythroid 2-related factor 2 (Nrf2) which plays a crucial role in the cellular defense against oxidative stress. Lastly, chlorine,

formaldehyde and hydrogen peroxide were reported to affect the localization of the *PRKCA* protein or to modify the expression of *PLA2G4A* and *PLA2R1* mRNA, or the activity or expression of *RELA* protein (CTD, <http://ctdbase.org/>, Davis et al. 2014). Overall, all these data suggest that *PLA2G4A*, *PLA2R1*, *RELA*, *PRKD1* or *PRKCA* may play a role in risk of asthma in adults in relationship with exposure to LMW agents or irritants.

CONCLUSIONS

In conclusion, the present study identified new promising candidate genes interacting with occupational exposures to LMW agents or irritants in current adult-onset asthma. More generally, this study highlights the interest to perform GxE interaction analysis to identify new genes and mechanisms of asthma occurrence related to specific environmental exposures.

REFERENCES

- Ackermann-Liebrich U, Kuna-Dibbert B, Probst-Hensch NM, Schindler C, Felber Dietrich D, Stutz EZ, et al. 2005. Follow-up of the Swiss Cohort Study on Air Pollution and Lung Diseases in Adults (SAPALDIA 2) 1991-2003: methods and characterization of participants. *Soz Praventivmed* 50:245-263.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, et al. 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nature genetics* 25:25-29.
- Barrett JC, Fry B, Maller J, Daly MJ. 2005. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics*. 21:263-265.
- Beasley R, Semprini A, Mitchell EA. 2015. Risk factors for asthma: is prevention possible? *Lancet* 386:1075-1085.
- Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B*. 57:289-300.
- Cherry N, Beach J, Burstyn I, Parboosinh J, Schouchen J, Senthilselvan A, et al. 2015. Genetic susceptibility to beryllium: a case-referent study of men and women of working age with sarcoidosis or other chronic lung disease. *Occup Environ Med* 72:21-27.
- Chi PL, Liu CJ, Lee IT, Chen YW, Hsiao LD, Yang CM. 2014. HO-1 induction by CO-RM2 attenuates TNF- α -induced cytosolic phospholipase A2 expression via inhibition of PKC α -dependent NADPH oxidase/ROS and NF- κ B. *Mediators Inflamm* 2014:279171.
- Curjuric I, Imboden M, Nadif R, Kumar A, Schindler C, Haun M, et al. 2012. Different genes interact with particulate matter and tobacco smoke exposure in affecting lung function decline in the general population. *PLoS One* 7(7): e40175.

- Dai JY, Logsdon BA, Huang Y, Hsu L, Reiner AP, Prentice RL, et al. 2012. Simultaneously testing for marginal genetic association and gene-environment interaction. *Am J Epidemiol* 176:164-173.
- Davis AP, Grondin CJ, Lennon-Hopkins K, Saraceni-Richards C, Sciaky D, King BL, et al. 2015. The Comparative Toxicogenomics Database's 10th year anniversary: update 2015. *Nucleic Acids Res* 43(Database issue):D914-20.
- de Jong K, Vonk JM, Timens W, Bossé Y, Sin DD, Hao K, et al. 2015. Genome-wide interaction study of gene-by-occupational exposure and effects on FEV(1) levels. *J Allergy Clin Immunol* 136:1664-1672.
- Downs SH, Schindler C, Liu LJ, Keidel D, Bayer-Oglesby L, Brutsche MH, et al. 2007. Reduced exposure to PM10 and attenuated age-related decline in lung function. *N Engl J Med* 357:2338–2347.
- Dumas O, Kauffmann F, Le Moual N. 2013. Asthma and exposures to cleaning products [Asthme et expositions aux produits de nettoyage]. *Arch Mal Prof* 74:117-129.
- Dumas O, Matran R, Zerimech F, Decoster B, Huyvaert H, Ahmed I, et al. 2015. Occupational exposures and fluorescent oxidation products in 723 adults of the EGEA study. *Eur Respir J* 46:258-261.
- European Community Respiratory Health Survey II Steering Committee. 2002. The European Community Respiratory Health Survey II. *Eur Respir J* 20:1071–1079.
- Gibson G. 2015. Human genetics. GTEx detects genetic effects. *Science* 348:640-641.
- Higgins JPT, Thompson SG. 2002. Quantifying heterogeneity in a meta-analysis. *Stat Med* 21:1539-58

- Higgins JPT, Thompson SG, Deeks JJ, Altman DG. 2003. Measuring inconsistency in meta-analyses. *British Medical Journal* 327:557-560.
- Hindorff LA, MacArthur J (European Bioinformatics Institute), Morales J (European Bioinformatics Institute), Junkins HA, Hall PN, Klemm AK, and Manolio TA. A Catalog of Published Genome-Wide Association Studies. Available at: www.genome.gov/gwastudies. Accessed 7/16/2015.
- Holloway JW, Yang IA, Holgate ST. 2010. Genetics of allergic disease. *J Allergy Clin Immunol* 125(2 Suppl 2):S81-S94.
- Imboden M, Bouzigon E, Curjuric I, Ramasamy A, Kumar A, Hancock DB,, et al. 2012. Genome-wide association study of lung function decline in adults with and without asthma. *J Allergy Clin Immunol* 129:1218–1228.
- Kauffmann F, Castro-Giner F, Smit LAM, Nadif R, Kogevinas M. 2010. Gene-environment interactions in occupational asthma. In: Occupational Asthma, T. Sigsgaard/D. Heederik (Editors). *Progress in Inflammation Research*, Birkhäuser Verlag AG. pp 205-228.
- Kauffmann F, Demenais F. 2012. Gene-environment interactions in asthma and allergic diseases: challenges and perspectives. *J Allergy Clin Immunol* 130:1229-1240.
- Kauffmann F, Dizier MH, Pin I, Paty E, Gormand F, Vervloet D, et al. 1997. Epidemiological study of the genetics and environment of asthma, bronchial hyperresponsiveness, and atopy: phenotype issues. *Am J Respir Crit Care Med* 156:S123-S129.
- Kennedy SM, Le Moual N, Choudat D, Kauffmann F. 2000. Development of an asthma specific job exposure matrix and its application in the epidemiological study of genetics and environment in asthma (EGEA). *Occup Environ Med* 57:635–641.

- Kim SH, Cho BY, Park CS, Shin ES, Cho EY, Yang EM, et al. 2009. Alpha-T-catenin (CTNNA3) gene was identified as a risk variant for toluene diisocyanate-induced asthma by genome-wide association analysis. *Clin Exp Allergy* 39:203-212.
- Kogevinas M, Zock JP, Jarvis D, Kromhout H, Lillienberg L, Plana E, Radon K, et al. 2007. Exposure to substances in the workplace and new-onset asthma: an international prospective population-based study (ECRHS-II). *Lancet* 370:336–341.
- Kogevinas M. 2014. Individual variability, from candidate G*E to GEWIS. *Occup Environ Med* 71 Suppl 1:A123-A124.
- Le Moual N, Jacquemin B, Varraso R, Dumas O, Kauffmann F, Nadif R. 2013. Environment and asthma in adults. *Presse Med* 42:e317-e333.
- Leslie CC. 2015. Cytosolic phospholipase A2: Physiological function and role in disease. *J Lipid Res* 56:1386-1402.
- Liu C, Maity A, Lin X, Wright RO, Christiani DC. 2012. Design and analysis issues in gene and environment studies. *Environ Health global access scie source* 11:93.
- Liu JZ, McRae AF, Nyholt DR, Medland SE, Wray NR, Brown KM, et al. 2010. A versatile gene-based test for genome-wide association studies. *Am J Hum Genet* 87:139-45.
- Maestrelli P, Boschetto P, Fabbri LM, Mapp CE. 2009. Mechanisms of occupational asthma. *J Allergy Clin Immunol* 123:531-542.
- Malo JL, Tarlo SM, Sastre J, Martin J, Jeebhay MF, Le Moual N, et al; on behalf of the ATS ad hoc committee on Asthma in the Workplace. 2015. An Official American Thoracic Society Workshop Report: Presentations and Discussion of the Fifth Jack Pepys Workshop on Asthma in the Workplace Comparisons between Asthma in the Workplace and Non-Work-related Asthma. *Ann Am Thorac Soc* Vol 12, No 7, pp S99–S110.

- Matheson MC, Benke G, Raven J, Sim MR, Kromhout H, Vermeulen R, et al. 2005. Biological dust exposure in the workplace is a risk factor for chronic obstructive pulmonary disease. *Thorax* 60:645-651.
- Mehta AJ, Miedinger D, Keidel D, Bettschart R, Bircher A, Bridevaux PO, et al. 2012. Occupational exposure to dusts, gases, and fumes and incidence of chronic obstructive pulmonary disease in the Swiss Cohort Study on Air Pollution and Lung and Heart Diseases in Adults. *Am J Respir Crit Care Med* 185: 1292–1300.
- Minelli C, Wei I, Sagoo G, Jarvis D, Shaheen S, Burney P. 2011. Interactive effects of antioxidant genes and air pollution on respiratory function and airway disease: a HuGE review. *Am J Epidemiol.* 173:603-620.
- Mittal M, Siddiqui MR, Tran K, Reddy SP, Malik AB. 2014. Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal* 20:1126-1167.
- Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, et al. 2010. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med* 363:1211–1221.
- Mukherjee B, Ahn J, Gruber SB, Rennert G, Moreno V, Chatterjee N. 2008. Tests for gene-environment interaction from case-control data: a novel study of type I error, power and designs. *Genet Epidemiol* 32:615-26.
- Murakami M, Taketomi Y, Miki Y, Sato H, Yamamoto K, Lambeau G. 2014. Emerging roles of secreted phospholipase A2 enzymes: the 3rd edition. *Biochimie* 107 PtA:105-113.
- Nakashima S. 2002. Protein kinase C alpha (PKC alpha): regulation and biological function. *J Biochem* 132:669-675.
- Quinn MM, Henneberger PK, and members of the National Institute for Occupational Safety and Health (NIOSH), National Occupational Research Agenda (NORA) Cleaning and

- Disinfecting in Healthcare Working Group. 2015. Cleaning and disinfecting environmental surfaces in health care: Toward an integrated framework for infection and occupational illness prevention. *American Journal of Infection Control* 43:424-434.
- Rager JE, Lichtveld K, Ebersviller S, Smeester L, Jaspers I, Sexton KG, et al. 2011. A toxicogenomic comparison of primary and photochemically altered air pollutant mixtures. *Environ Health Perspect* 119:1583-1589.
- Rava M, Ahmed I, Demenais F, Sanchez M, Tubert-Bitter P, Nadif R. 2013. Selection of genes for gene-environment interaction studies: a candidate pathway-based strategy using asthma as an example. *Environmental Health* 12:56.
- Rava M, Smit LA, Nadif R. 2015. Gene-environment interactions in the study of asthma in the postgenomewide association studies era. *Curr Opin Allergy Clin Immunol* 15:70-78.
- Romieu I, Moreno-Macias H, London SJ. 2010. Gene by environment interaction and ambient air pollution. *Proc Am Thorac Soc* 7:116-122.
- Schuliga M. 2015. NF-kappaB Signaling in Chronic Inflammatory Airway Disease. *Biomolecules* 5:1266-1283.
- Siracusa A, De Blay F, Folletti I, Moscato G, Olivieri M, Quirce S, et al. 2013. Asthma and exposure to cleaning products - a European Academy of Allergy and Clinical Immunology task force consensus statement. *Allergy* 68:1532-1545.
- Smit LA, Kogevinas M, Antó JM, Bouzigon E, González JR, Le Moual N, et al. 2012. Transient receptor potential genes, smoking, occupational exposures and cough in adults. *Respir Res* 13:26.

- Smit LA, Strachan DP, Vermeulen R, de Bakker PI, Demenais F, Dumas O, et al. 2014. Human leukocyte antigen class II variants and adult-onset asthma: does occupational allergen exposure play a role? *Eur Respir J* 44:1234-1242.
- Sokolowska M, Stefanska J, Wodz-Naskiewicz K, Cieslak M, Pawliczak R. 2010. Cytosolic phospholipase A2 group IVA is overexpressed in patients with persistent asthma and regulated by the promoter microsatellites. *J Allergy Clin Immunol* 125:1393-1395.
- Storz P. 2007. Mitochondrial ROS--radical detoxification, mediated by protein kinase D. *Trends Cell Biol* 17:13-18.
- Tarlo SM, Lemiere C. 2014. Occupational asthma. *N Engl J Med* 370:640–649.
- Tarlo SM. 2014. Irritant-induced asthma in the workplace. *Curr Allergy Asthma Rep* 14:406.
- Torén K, Blanc PD. 2009. Asthma caused by occupational exposures is common - a systematic analysis of estimates of the population-attributable fraction. *BMC Pulm Med* 9:7.
- Vandebriel RJ, De Jong WH. 2012. A review of mammalian toxicity of ZnO nanoparticles. *Nanotechnol Sci Appl* 5:61-71.
- Sundram V, Chauhan SC, Jaggi M. 2011. Emerging Roles of Protein Kinase D1 in Cancer Mol Cancer Res 9:985–996.
- Viechtbauer W, Cheung MW. 2010. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods* 1:112-25.
- Wu W, Samet JM, Peden DB, Bromberg PA. 2010. Phosphorylation of p65 is required for zinc oxide nanoparticle-induced interleukin 8 expression in human bronchial epithelial cells. *Environ Health Perspect* 118:982-987.

Table 1. Characteristics of adult participants in the three studies

	All (N=2599)	ECRHS (N=1336)	SAPALDIA (N=574)	EGEA (N=689)
Age, year, mean (SD)	46.7 (11.3)	43.1 (7.1)	53.4 (10.9)	48.0 (14.9)
Sex, women, n (%)	1563 (60.1)	822 (61.5)	311 (54.2)	430 (62.4)
Smoking habits, n (%)				
Never smokers	1167 (44.9)	569 (42.6)	248 (43.2)	350 (50.8)
Ex-smokers	735 (28.3)	337 (25.2)	191 (33.3)	207 (30.0)
Current smokers	682 (26.2)	419 (31.4)	135 (23.5)	128 (18.6)
Missing	15 (0.6)	11 (0.8)	0 (0.0)	4 (0.6)
Occupational exposure, n (%) ^a	927 (35.7)	440 (32.9)	175 (30.5)	312 (45.3)
Current adult onset asthma, n (%)	463 (17.8)	234 (17.5)	107 (18.6)	122 (17.7)

^a% ever exposed to Low Molecular Weight (LMW) agents or to mixed environments or to high pick irritants, or to specific cleaning products or disinfectants in the population selected for the analyses, *i.e.*; after exclusion of adults with occupational exposures to other potentially asthmagenic agents (High Molecular Weight (HMW) agents).

Table 2. Interactive effects of SNPs by occupational exposure to LMW agents or irritants on current adult-onset asthma

Chr	Gene	SNP	Reference /Effect Allele	EAF ^a	Cases/Control s	Marginal effect		Interaction - CC	
						N/N	OR/P-value	OR/P-value	OR/P-value
1	<i>PLA2G4A</i>	rs932476	A/G	0.35	463/2136	1.25	0.0036	0.64	0.0050
2	<i>PLA2R1</i>	rs2667026	A/G	0.83	463/2136	0.89	0.2354	1.77	0.0050
11	<i>RELA</i> ^b	rs931127	A/G	0.43	462/2135	1.17	0.0350	1.61	0.0014
11	<i>RELA</i>	rs7949980	C/T	0.51	463/2133	1.07	0.3421	1.56	0.0030
14	<i>PRKD1</i> ^b	rs1958980	A/G	0.67	463/2136	1.08	0.3344	0.64	0.0042
14	<i>PRKD1</i>	rs1184735	A/G	0.67	463/2133	1.08	0.3429	0.64	0.0043
14	<i>PRKD1</i>	rs1958987	C/T	0.68	459/2127	1.07	0.3609	0.64	0.0050
17	<i>PRKCA</i>	rs6504453	C/T	0.35	462/2134	1.04	0.6086	0.63	0.0032

Chr: chromosome, CC: case-control. ^aEffect Allele Frequency (EAF) calculated in controls.

^bThe two SNPs near *RELA* were in moderate Linkage Disequilibrium (LD) with $r^2=0.65$, whereas the three SNPs in *PRKD1* are in strong LD ($r^2>0.8$).

Figure legends

Figure 1. Associations between SNPs that showed an interactive effect with occupational exposure to LMW agents or irritants on current adult-onset asthma in unexposed (grey) and exposed (black) participants.

Figure 1.

