

AUTHOR MANUSCRIPT

Published in final edited form as: Cell Death and Differentiation, 1, October 2017 doi: 10.1038/cdd.2017.155

A role for caspase-8 and TRAIL-R2/DR5 in ER-stress induced apoptosis

Authors: Cristina Muñoz-Pinedo1* and Abelardo López-Rivas2*

*corresponding authors

- 1. Cell Death Regulation Group. Oncobell Program. Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet, Spain
- 2. Cell Death Signalling Group. Andalusian Center for Molecular Biology and Regenerative Medicine –CABIMER. CSIC-Universidad de Sevilla-Universidad Pablo de Olavide

*Corresponding author: Cristina Muñoz Pinedo IDIBELL - Hospital Duran i Reynals 3a planta Gran Via de l'Hospitalet, 199 08908 L'Hospitalet de Llobregat Barcelona – Spain phone +34 93 260 7130 fax +34 93 260 7426 cmunoz@idibell.cat

*Corresponding author: Abelardo López Rivas CABIMER Avda Américo Vespucio, 24 41092 Sevilla-Spain phone + 34 954 46 7997 fax + 34 954 46 1664 abelardo.lopez@cabimer.es

Publisher Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain. © Journal editor. All rights reserved.

Correspondence:

A role for caspase-8 and TRAIL-R2/DR5 in ER-stress induced apoptosis

Running title: DR5 and caspase-8 in ER-stress induced apoptosis

Authors: Cristina Muñoz-Pinedo1* and Abelardo López-Rivas2*

*corresponding authors

- 3. Cell Death Regulation Group. Oncobell Program. Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet, Spain
- 4. Cell Death Signalling Group. Andalusian Center for Molecular Biology and Regenerative Medicine –CABIMER. CSIC-Universidad de Sevilla-Universidad Pablo de Olavide

*Corresponding author: Cristina Muñoz Pinedo IDIBELL - Hospital Duran i Reynals 3a planta Gran Via de l'Hospitalet, 199 08908 L'Hospitalet de Llobregat Barcelona – Spain phone +34 93 260 7130 fax +34 93 260 7426 cmunoz@idibell.cat

*Corresponding author: Abelardo López Rivas CABIMER Avda Américo Vespucio, 24 41092 Sevilla-Spain phone + 34 954 46 7997 fax + 34 954 46 1664 abelardo.lopez@cabimer.es

Word count: 407

The authors declare no conflict of interests

Text

Glab and colleagues¹ examine in a recent paper apoptosis induced by some driverss of endoplasmic reticulum (ER) stress. They conclude that in contrast to a previously published report², DR5/TRAIL-R2 and caspase-8 are universally dispensable in ER stress-induced apoptosis. We argue here that their own data and other published reports indicate that in many models, DR5 and/or caspase-8 are essential players in apoptosis mediated by the unfolded protein response (UPR), upon chronic ER stress.

The authors analyze the effects of knocking down caspase-8, DR5, Bim and Bid. The authors describe that Bim is not essential at least in one of their three models. In line with these results, we and other authors have shown that Bim is dispensable for apoptosis mediated by the UPR in some systems even though the protein is induced; in these cases the apoptotic cell death can be attributed to Noxa or to Caspase-8 (please see table 1 in Ref. 3, and Refs. 4-6). We find inconsistent that the title and abstract mention that caspase-8 and DR5 are dispensable in ER stress-induced apoptosis but not that Bim is not required in their system, in contrast to earlier findings (Ref. 7).

The authors' own data (Ref. 1, Fig. 3c) indicates that Bid is crucial in one of their models: HCT116. Moreover, although the authors report a drastic reduction of apoptosis with some treatments, they also show that over 20% of Bax, Bak deficient cells undergo apoptosis (Ref.1, Fig. 2E,F). We suggest that this apoptosis is mediated by caspase-8 as described in Bax, Bak deficient HCT116 and MEFs treated with UPR-inducing stimuli^{4,6}. Indeed, a certain role for caspase-8 is shown by the authors in Supplementary fig. 2 but this is not mentioned in the title or abstract.

Regarding the role of DR5, authors do observe reduction of cell death in one knock-out clone. Importantly, the authors seem focused to disprove one study, while ignoring numerous studies that have shown a role of DR5 and caspase-8 in apoptosis activated by different UPR-inducing stimuli such as thapsigargin or glucose deprivation^{3,8,9}, table 1). Moreover, it is possible that in cells knocked out for DR5, it is DR4 or other death receptors (reviewed in Ref. 6) that may be participating in caspase-8 activation and Bid cleavage, as we have recently described in ATF4-mediated cell death⁶.

Altogether, these data indicate that DR5, caspase-8, Bim, Bid and Noxa are required for ER stress and UPR-mediated death in a cell type-dependent manner.

References

- 1. Glab, JA et al., (2017). Cell Death Differ 24: 944.
- 2. Lu, M et al., (2014). Science (New York, N.Y.) 345: 98.
- 3. Iurlaro, R et al, (2016). The FEBS Journal 283: 2640.
- 4. Caro-Maldonado, A et al., (2010). Cell Death Differ 17: 1335-44.
- 5. Ramírez-Peinado, S et al., (2011). Cancer Res 71: 6796-806.
- 6. Iurlaro, R et al., (2016). Molecular and Cellular Biology 37: e00479-16.
- 7. Puthalakath H et al (2007). Cell. 2007 Jun 29;129(7):1337-49.
- 8. Yamaguchi, H et al (2004). Journal of Biological Chemistry 279: 45495-45502.
- 9. Martín-Pérez, R et al., (2014). Cancer Research 74: 1766-1777.

Table 1. Examples of reports showing a role for caspase-8 and/or DR5 in apoptosismediated by components of the UPR or the Integrated Stress Response (ISR)

Mediator	Stimulus	Cell type	Reference
Caspase-8	Tunicamycin	HEK293, MCF7	Tomar <i>et al</i> 2013, PMID <u>24021263</u>
		<i>bax^{-/-}bak^{-/-}</i> BMK	Ullman <i>et al</i> 2011, PMID <u>21576355</u>
	Bortezomib/MG132	HEK293, MDAMB231, MCF7	Pan <i>et al</i> 2001, PMID <u>21628531</u>
Caspase-8, FADD	Bortezomib/MG132	Atg5+/+ and Atg5-/- MEFs,FADD+/+ and FADD-/- MEFs, KG-1	Young <i>et al</i> 2012, PMID <u>22362782</u>
		HeLa, H460, MEFs	Laussmann <i>et al</i> 2011, PMID <u>21455219</u>
	Tunicamycin, Thapsigargin	CASP9-/- MEFs, bax/Bak-/- MEFs	Deegan <i>et al</i> 2014, PMID <u>25470234</u>
Caspase-8, RIPK1	Tunicamycin	MEFs and HEK293T	Estornes <i>et al</i> 2014, PMID <u>25476903</u>
DR5	Thapsigargin	HCT116, LNCaP, A2780S and DU145	Yamaguchi <i>et al</i> 2004, PMID <u>15322075</u>
DR5, Caspase-8	Thapsigargin, Tunicamycin	MCF10A	Martín-Pérez <i>et al</i> 2014, PMID <u>24453000</u>
	Thapsigargin	HCT116, SK-MES-1, KSM11, RPMI-8226	Lu <i>et al</i> 2014, PMID <u>24994655</u>
	CB-5083	HCT116	Anderson <i>et al</i> 2015, PMID <u>26555175</u>
DR5, DR4	Glucose deprivation	HCT116, HCT116 Bax, Bak (-/-), HeLa	lurlaro <i>et al</i> 2017, PMID <u>28242652</u>
	Tunicamycin, thapsigargin, brefeldin A	HCT116, MDA-MB-231, H1703	Dufour et al 2017, PMID <u>28039489</u>
DR4/TRAIL-R1	Tunicamycin, Thapsigargin	A549	Li <i>et al</i> 2015, PMID 25770212