

Cite as: Lomas, T., Medina, J. C., Ivtzan, I., Rupperecht, S., Hart, R., & Eiroa-Orosa, F. J. (2017). A systematic review of the impact of mindfulness on the wellbeing of healthcare professionals. *Journal of Clinical Psychology*. <https://doi.org/10.1002/jclp.22515>

## **A systematic review of the impact of mindfulness on the wellbeing of healthcare professionals.**

Tim Lomas<sup>1†</sup>, Juan Carlos Medina<sup>2</sup>, Itai Ivtzan<sup>1</sup>, Silke Rupperecht<sup>3</sup>, Francisco José Eiroa-Orosa<sup>1</sup>

<sup>1</sup> School of Psychology, University of East London, Arthur Edwards Building, Water Lane,  
London, E15 4LZ, United Kingdom

<sup>2</sup> Faculty of Psychology, University of Barcelona, Passeig de la Vall d'Hebron, 08035  
Barcelona, Spain

<sup>3</sup> Leuphana University, Scharnhorststraße 1, 21335 Lüneburg, Germany

† Author responsible for correspondence:

Email: [t.lomas@uel.ac.uk](mailto:t.lomas@uel.ac.uk)

## **Abstract**

**Objectives:** Among efforts to improve the wellbeing of healthcare professionals are initiatives based around mindfulness meditation. To understand the value of such initiatives, we conducted a systematic review of empirical studies pertaining to mindfulness in healthcare professionals.

**Design:** Databases were reviewed from the start of records to January 2016. Eligibility criteria included empirical analyses of mindfulness and wellbeing outcomes acquired in relation to practice. 81 papers met the eligibility criteria, consisting of a total 3,805 participants. Studies were principally examined for outcomes such as burnout, distress, anxiety, depression and stress.

**Results:** Mindfulness was generally associated with positive outcomes in relation to most measures (although results were more equivocal with respect to some outcomes, most notably burnout).

**Conclusion:** Overall, mindfulness does appear to improve the wellbeing of healthcare professionals. However, the quality of the studies was inconsistent, so further research is needed, particularly high-quality randomised control trials.

**Keywords:** mindfulness; meditation; healthcare professionals; wellbeing; systematic review.

## **A systematic review of the impact of mindfulness on the wellbeing of healthcare professionals.**

Healthcare professionals (HCPs) can face particular challenges that can be detrimental to their physical and mental health. A wealth of research has accumulated indicating that HCPs are liable to experience a range of mental health issues, including anxiety (Gao et al., 2012), burnout (Khamisa, Oldenburg, Peltzer, & Ilic, 2015), depression (Givens & Tjia, 2002), and stress (Bidwal, Ip, Shah, & Serino, 2015). Moreover, these problems may be particularly acute among HCPs relative to people in other professions (Brooks, Gerada, & Chalder, 2011). A recent survey of over 3,700 public sector workers in the UK found that staff working for the National Health Service were the most stressed, with 61% reporting feeling stress all or most of the time, and 59% stating that stress is worse this year than last year (Dudman, Isaac, & Johnson, 2015).

Analyses of these problems include attempts to understand why HCPs are especially vulnerable to mental health issues. Some scholars explain outcomes like burnout according to the model of effort-reward imbalance, finding that HCPs face a particularly disadvantageous imbalance due to the considerable effort required by their work, emotionally and physically (Rasmussen et al., 2015). Such efforts include factors such as emotional demands (Tyssen, Vaglum, Grønvold, & Ekeberg, 2000), exacerbated by often limited resources, such as time allocation per patient (Mossialos, Wenzl, Osborn, & Anderson, 2015). Another factor is adverse events in healthcare settings, which can mean that HCPs may be ‘second victims’ (Draper, Kőlves, De Leo, & Snowdon, 2014). Particular HCP populations can be especially vulnerable, such as younger and/or less experienced workers; Bidwal et al. (2015) found that levels of stress among trainees in the healthcare professions were roughly twice as high as in the general adult population. Professionals may also fare worse than others owing to their specific occupational context, such as work demands in their particular national healthcare

system. For instance, a survey of general practitioners in 11 developed countries found that workers in the UK reported the highest levels of stress, with 29% saying they intended to quit general practice within five years.

These issues represent a significant problem: obviously for the wellbeing of the HCPs themselves, but also for patients (e.g., the ability of HCPs to treat them skilfully), and for the healthcare system (e.g., the economic cost of staff burnout) (Toppinen-Tanner, Ojajärvi, Väänaänen, Kalimo, & Jäppinen, 2005). As such, efforts are underway to protect against or ameliorate work-related mental health issues in HCPs. Among the most prominent of these types of initiatives are programmes based around mindfulness meditation – mindfulness-based interventions (MBIs) – which is the focus of this review.

## **Mindfulness**

The past few decades have seen a burgeoning interest in mindfulness in the West, spanning clinical practice, academia, and society more broadly. Originating in the context of Buddhism around the 5<sup>th</sup> century B.C.E, mindfulness came to prominence in the West through Kabat-Zinn (1982), who created a pioneering ‘mindfulness-based stress reduction’ (MBSR) programme for chronic pain. ‘Mindfulness’ can refer to: (1) a state/quality of mind; and (2) a meditation practice that enables one to cultivate this. The most prominent operationalisation of mindfulness as a mental state/quality is Kabat-Zinn’s (2003, p.145) definition of it as ‘the awareness that arises through paying attention on purpose, in the present moment, and nonjudgmentally to the unfolding of experience moment by moment.’ Expanding on this, Shapiro, Carlson, Astin, and Freedman (2006) deconstruct it into three components: intention (motivation for paying attention thus); attention (cognitive processes through which attention is enacted); and attitude (emotional qualities with which one imbues one’s attention).

‘Mindfulness’ is also deployed for meditation practices which facilitate this state. Meditation broadly refers to mental activities which share a common focus on training the self-

regulation of attention and awareness, thereby enhancing control of mental processes, and consequently increasing wellbeing (Walsh & Shapiro, 2006). According to Lutz, Slagter, Dunne, and Davidson (2008), most practices feature either ‘focused attention’ or ‘open-monitoring’ processes. Focused attention can be operationalised in terms of the co-ordination of various attention modalities (Posner & Petersen, 1990), including sustained, executive, and selective attention. By contrast, open-monitoring delineates a broader receptive capacity to detect events within an open ‘field’ of awareness (Raffone & Srinivasan, 2010). Mindfulness – as a practice, and a state of mind – is commonly presented as a case of open-monitoring (Kabat-Zinn, 2003). However, in practice, mindfulness meditation usually involves both focused attention and open-monitoring, e.g., beginning with a period of focused attention on the breath, in order to stabilise one’s awareness, followed by the more receptive state of open-monitoring (Chiesa, Calati, & Serretti, 2011).

According to Shapiro et al. (2006), the main significance of mindfulness – as a quality/state, and a practice – is that it involves a meta-mechanism known as *reperceiving*. The three components of mindfulness (intention, attention and attitude) combine to generate a ‘fundamental shift in perspective,’ in which ‘rather than being immersed in the personal drama or narrative of our life story, we are able to stand back and witness it’ (p.377). This process, also known as ‘*decentring*,’ is defined as ‘the ability to observe one’s thoughts and feelings as temporary, objective events in the mind, as opposed to reflections of the self that are necessarily true’ (Fresco et al., 2007, p.234). This ability is theorised as having a positive impact upon wellbeing. In MBIs, the aim is not to change participants’ thoughts/feelings per se, as cognitive therapy might seek to, but to help people ‘become more aware of, and relate differently to’ this content (Shapiro, Astin, Bishop, & Cordova, 2005, p.165). For example, in Mindfulness-Based Cognitive Therapy (MBCT), designed to prevent depressive relapse, people are taught to *decentre* from their cognitions, thus helping prevent a ‘downward spiral’ of negative thoughts

and worsening negative affect which could otherwise trigger relapse (Segal, Williams, & Teasdale, 2002). Thus MBCT, and MBIs generally, involve ‘retraining awareness’ so that people have greater choice in how they relate and respond to their subjective experience, rather than habitually responding in maladaptive ways (Chambers, Gullone, & Allen, 2009, p.659). For instance, the development of decentring can help people tolerate distressing qualia, which is important given that *inability* to tolerate such qualia is a transdiagnostic factor underlying diverse psychopathologies (Aldao, Nolen-Hoeksema, & Schweizer, 2010).

Mindfulness interventions were initially limited to clinical settings, such as Kabat-Zinn’s (1982) MBSR program and subsequent adaptations like MBCT (Segal et al., 2002). However, since the late 1990s, there has been increasing use of mindfulness in occupational contexts, not only for staff who may be suffering with stress and mental health issues, but for workers ‘in general’ (e.g., as a protective measure against future issues). To assess the state of this literature with regard to HCPs, we conducted a systematic review of relevant research. Although a number of reviews have already been conducted in this area, these have tended to have fairly narrow remits in terms of population and/or outcome. These include reviews focused only on certain healthcare professions, such as General Practitioners (Murray, Murray, & Donnelly, 2016), social workers (Trowbridge & Lawson, 2016), and nurses (Botha, Gwin, & Purpora, 2015), all of which featured small numbers of studies (e.g., Murray et al. selected just four papers, not all of which were even exclusively concerned with mindfulness per se). Or, such reviews have concentrated on HCPs more generally, but have only been concerned with specific outcomes, such as stress in the case of Burton, Burgess, Dean, Koutsopoulou, and Hugh-Jones (2017), who only included nine studies, or empathy and emotional competencies in the case of Lamothe, Rondeau, Malboeuf-Hurtubise, Duval, and Sultan (2016), which focused just on MBSR, and identified 14 such studies. By contrast, the current paper aims for greater inclusivity, reporting the results of a far broader systematic review, encompassing: (a)

workers across all HCP contexts; (b) a wide range of wellbeing outcomes; and (c) the impact of mindfulness generally (not limited to any one intervention).

## **Methods**

The literature search was conducted using the MEDLINE and Scopus electronic databases. The search was conducted as part of a broader systematic review on mindfulness in all occupations (which is still ongoing). The criteria for the broader review were: mindfulness AND work OR occupation OR profession OR staff (in all fields in MEDLINE and limited to article title, abstract, and keywords in Scopus). The dates selected were from the start of the database records to 10<sup>th</sup> January 2016. For this current review into HCPs, in terms of PICOS (participants, interventions, comparisons, outcomes and study design) the key inclusion criteria were: participants – currently employed in a healthcare context; outcomes – any pertaining to mindfulness, wellbeing, and job performance; and study design – any empirical study featuring data collection. Exclusion criteria were theoretical articles or commentaries without statistical or qualitative analyses. Although we were principally interested in studies of MBIs in healthcare workplaces, as a secondary concern we were also interested in non-intervention studies (e.g., regression analyses of the association between trait mindfulness and wellbeing outcomes). Studies were required to be published (or in press) in English in a peer-reviewed academic journal. The review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Moher, Liberati, Tetzlaff, & Altman, 2009). The review protocol for the broader systematic review was registered with the International Prospective Register of Systematic Reviews (PROSPERO) database on 5<sup>th</sup> January 2016 (registration number: CRD42016032899).

Papers were divided into intervention studies and non-intervention studies. For intervention studies, the following variables were extracted from each paper: type of design (e.g., RCT versus convenience sample); occupation of participants; number of experimental

participants; number of control participants (if applicable); type of MBI; length of MBI; nature of control; principle wellbeing and performance outcomes; and the significance level and effect size of principle outcomes. For non-intervention studies, the following variables were extracted: type of analysis (e.g., regression versus qualitative); occupation of participants; number of experimental participants; principle wellbeing and performance outcomes; and the significance level of principle outcomes.

The primary summary measures were mindfulness and wellbeing outcomes. These were principally psychometric scales pertaining to mindfulness, mental health (e.g., anger, anxiety, burnout, depression, distress, stress, satisfaction, wellbeing), and physical health (e.g., illness, diet, exercise, and sleep). Secondary summary measures of interest were outcomes that *pertain* to wellbeing (e.g., compassion, empathy, emotional intelligence and regulation, relationships, resilience, and spirituality). Tertiary summary measures of interest were outcomes relating to job performance. The Quality Assessment Tool for Quantitative Studies (QATQS; National Collaborating Centre for Methods and Tools, (Tools, 2008) was used to assess the quality of the studies. QATQS assesses methodological rigor in six areas: (a) selection bias; (b) design; (c) confounders; (d) blinding; (e) data collection method; and (f) withdrawals and drop-outs. Each area is assessed on a quality score of 1 to 3 (1 = strong; 2 = moderate; 3 = weak). Scores for each area were collated, and a global score assigned to each study. If there are no weak ratings, the study is scored 1 (strong); one weak rating leads to a 2 (moderate); and two or more weak ratings generates a 3 (weak). QATQS scoring was conducted by the fourth author, and checked independently by the first author. Any discrepancies were resolved by discussion with agreement reached in all cases.

## **Results**

For the broader systematic review (i.e., mindfulness across all occupations), following removal of duplicate citations, 721 potentially relevant papers were identified. In the current



specific systematic review (focusing specifically on HCPs), from reviewing the abstract, 543 papers were excluded. From the full text reviews of 178 papers, 97 further papers were excluded. Thus, a total of 81 papers were included in the systematic analysis (66 intervention studies, and 15 non-intervention studies). Two of these papers pertained to the same trial (Cohen-Katz, Wiley, Capuano, Baker, Kimmel, et al., 2005; Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005), and so the 81 papers included in the analysis represented results from 80 independent participant samples. The studies comprised a total of 3,805 participants (discounting participants not including in analyses due to attrition). There were 2,645 participants in the intervention studies, as below in table 1, including 1,869 undertaking MBIs, 663 separate control participants (excluding Singh et al., 2006, in which participants acted their own controls), plus one study (Grepmaier, Mitterlehner, Loew, & Nickel, 2007) in which participants were not HCPs per se, but rather patients being treated by them. There were 1,160 participants in non-intervention studies, as detailed in table 2. The studies covered a range of occupations, including physicians ( $n = 9$ ), nurses ( $n = 16$ ), disability professionals ( $n = 4$ ), therapists, psychologists and counsellors ( $n = 24$ , mixed (non-specific) mental health professionals ( $n = 8$ ), and mixed (non-specific) healthcare professionals ( $n = 20$ ).

**Table 1.** Overview of studies

Authors	Occupation	Design	Expt. group	Control group	Intervention	Length	Control	Primary outcome(s)
(Aggs & Bambling, 2010)	Psychotherapists	Convenience sample	47	-	Mindful therapy	8 weeks	N/A	PI < stress & strain ( $p < .01$ ). PI > mindfulness & awareness ( $p < .01$ ).
(Barbosa et al., 2013)	Healthcare graduates	Convenience sample	13	15	MBSR	8 weeks	Nothing	PI < anxiety ( $d = -.09$ , $p < .001$ ), burnout (emotional exhaustion, $d = -.41$ ; depersonalisation, $d = -.26$ ; and personal accomplishment, $d = .29$ ; $p < .001$ ). PI > compassion & empathy (physician empathy, $d = .77$ , $p < .01$ ).
(Bazarko, Cate, Azocar, & Kreitzer, 2013)	Nurses (corporate)	Convenience sample	36	-	MBSR adaptation (6 sessions by telephone)	8 weeks	N/A	PI < burnout (personal burnout, $d = -.97$ ; work-related burnout, $d = -.67$ ; and client-related burnout, $d = -.30$ ; $p < .001$ ), and stress & strain (perceived stress, $d = -1.21$ , $p < .001$ ). PI > compassion & empathy (physician empathy, $d = .76$ ; and self-compassion, $d = 1.25$ ; $p < .001$ ), health (physical health, $d = -.38$ , $p < .001$ ; and mental health, $d = 1.40$ $p < .05$ ), and wellbeing, satisfaction & flourishing (serenity, $d = 1.48$ $p < .001$ ).
(Beckman et al., 2012)	Primary care physicians	Convenience sample	20	-	Program in mindful communication (Krasner et al., 2009).	52 hours	N/A	Qualitative interviews ( $n = 20$ ): PI > mindfulness & awareness, and relationships.
(Beddoe & Murphy, 2004)	Trainee nurses	Convenience sample	16 (23)*	-	MBSR	8 weeks	N/A	PI < stress & strain ( $p < .05$ ). PI >> compassion & empathy.
(Bond et al., 2013)	Trainee doctors	Convenience sample	27	-	Mind-body course**	11 weeks	N/A	PI > compassion & empathy (self-compassion, $d = .17$ , $p = .04$ ), emotional intelligence & regulation (self-regulation, $d = .01$ , $p = .003$ ). PI >> compassion & empathy (physician empathy, $d = .09$ ), and stress & strain (perceived stress, $d = -.03$ ).
(Bonifas & Napoli, 2014)	Trainee social workers	Convenience sample	77	-	Mindfulness curriculum (specific to study)	16 weeks	N/A	PI > wellbeing, satisfaction & flourishing (quality of life, $d = .88$ , $p < .001$ ). PI >> stress & strain (perceived stress, $d = .06$ ).
(Brady et al., 2012)	Psychiatric ward professionals	Convenience sample	16 (23)	-	MBSR adaptation	4 weeks	N/A	PI < stress & strain (stress, $d = -.70$ , $p < .01$ ), burnout (emotional exhaustion, $d = -.50$ ; depersonalisation, $d = -.23$ ; and personal accomplishment, $d = .29$ ). PI > mindfulness & awareness (mindfulness, $d = .64$ , $p < .01$ ; and intrapersonal presence, $d = .54$ , $p = .02$ ).
(Brooker et al., 2013)	Disability professionals	Convenience sample	34 (36)	-	Occupational mindfulness training program	8 weeks	N/A	PI < wellbeing, satisfaction, & flourishing (extrinsic job satisfaction, $p < .05$ ). PI > mindfulness & awareness ( $p < .05$ ), stress & strain ( $p < .05$ ), and wellbeing, satisfaction & flourishing (positive affect, $p < .05$ ; and negative affect, $p < .05$ ). PI >> burnout, compassion & empathy, depression, wellbeing, satisfaction & flourishing.
(Brooker et al., 2014)	Disability professionals	Convenience sample	12	-	Occupational mindfulness training program	8 weeks	N/A	PI > job performance (restraint of patients, and seclusion of patients; $p < .05$ ).
(Burnett & Pettijohn, 2015)	Healthcare employees	Random allocation	20 active & 17 passive	18	MBST	5 weeks	Passive intervention : abstention from work activity.	Passive intervention group: PI >> emotional intelligence & regulation, stress & strain (perceived stress, $d = -.09$ ). Control group: PI >> emotional intelligence & regulation, and stress & strain (perceived stress, $d = -.70$ ).

(Christopher, Christopher, Dunnagan, & Schure, 2006)	Trainee counsellors	Convenience sample	11	-	Mindfulness curriculum (specific to study)	1 term	Control: nothing. N/A	Qualitative interviews: PI < burnout, and stress & strain.
(Cohen & Miller, 2009)	Trainee clinical psychologists	Convenience sample	21 (28)	-	Interpersonal mindfulness training	6 weeks	N/A	PI < anxiety ( $d = -.46, p = .027$ ), and stress & strain (perceived stress, $d = -.53, p < .001$ ). PI > emotional intelligence & regulation (emotional intelligence, $d = .39, p = .020$ ), and relationships (social connectedness, $d = .57, p = .002$ ). PI > depression ( $d = -.11$ ), mindfulness & awareness (mindful attention awareness, $d = .48$ ), and wellbeing, satisfaction & flourishing (life satisfaction, $d = .43, p = .051$ ; searching of meaning in life, $d = -.35$ ; and presence of meaning in life, $d = .12$ ).
(Cohen-Katz, Wiley, Capuano, Baker, Kimmel, et al., 2005)	Nurses	RCT	12 (14)	13	MBSR	8 weeks	Wait-list	PI < burnout ( $p = .050$ ). PI > mindfulness & awareness ( $p = .001$ ). PI > distress & anger.
(Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005)	Nurses	RCT	12 (14)	13	MBSR	8 weeks	Wait-list	Qualitative data analysis ( $n = 12$ ): PI > emotional intelligence & regulation (self-acceptance), mindfulness & awareness (self-care, and self-awareness), relationships, and wellbeing, satisfaction & flourishing (relaxation).
(Dobie, Tucker, Ferrari, & Rogers, 2015)	Mental health professionals	Convenience sample	9	-	MBSR adaptation	8 weeks	N/A	PI < anxiety ( $d = -.86, p = .02$ ), distress ( $p = .002$ ), and stress & strain (stress, $d = -.96, p < .05$ ). PI > mindfulness & awareness (mindfulness, $d = .41$ ). PI > depression ( $d = -.44, p = .06$ ).
(De Vibe et al., 2013)	Trainee doctors	RCT	144	144	MBSR adaptation	6 weeks	Nothing	PI < distress & anger (distress, $d = -.77, p < .001$ ), and stress & strain (stress, $d = -.27, p = .021$ ). PI > wellbeing, satisfaction & flourishing (subjective wellbeing, $d = .43, p < .001$ ). PI > burnout (burnout, $d = -.13$ ), and mindfulness & awareness (act aware, $d = -.04$ ; describe, $d = -.06$ ; observe, $d = .18$ ; non-judging, $d = -.23$ ; and non-reacting, $d = .31$ ).
(de Zoysa, Ruths, Walsh, & Hutton, 2014)	Mental health professionals	Convenience sample	7	-	MBCT (in (Ruths et al., 2013))	8 weeks	N/A	Qualitative interviews: PI > emotional intelligence & regulation (self-regulation).
(Dorian & Killebrew, 2014)	Trainee psychotherapists	Convenience sample	21	-	Mindfulness curriculum (specific to study)	10 weeks	N/A	Qualitative interviews: PI < distress & anger. PI > compassion & empathy (compassion), emotional intelligence & regulation (acceptance), and mindfulness & awareness (awareness, and coping).
(Duchemin, B. A. Steinberg, D. R. Marks, K. Vanover, & M. Klatt, 2015)	Intensive care professionals	RCT	16	16	Mindfulness program (specific to study)	8 weeks	Wait-list	PI < stress & strain ( $p = .040$ ). PI > wellbeing, satisfaction & flourishing (quality of life, $p = .031$ ). PI > anxiety, burnout, depression, and mindfulness & awareness.
(Erogul, Singer, McIntyre, & Stefanov, 2014)	Trainee doctors	RCT	29	30	MBCT	8 weeks	Nothing	PI < stress & strain (perceived stress, $d = -.60, p = .03$ ). PI > compassion & empathy (self-compassion, $d = .88, p < .001$ ). PI > resilience ( $d = .27, p = .05$ ).

(Felton, Coates, & Christopher, 2015)	Trainee counsellors	Convenience sample			Mindfulness curriculum (specific to study)	15 weeks	N/A	Qualitative interviews: PI < stress & strain. PI > compassion & empathy (compassion), emotional intelligence & regulation (acceptance), and mindfulness & awareness (awareness).
(Fisher & Hemanth, 2015)	Clinical psychologists	Convenience sample	8	-	Mindfulness program (specific to study)	10 weeks	N/A	Qualitative interviews: PI > emotional intelligence & regulation (acceptance), and wellbeing, satisfaction & flourishing (relaxation).
(Fortney, Luchterhand, Zakletskaia, Zgierska, & Rakel, 2013)	Primary care clinicians	Convenience sample	28 (32)	-	MBSR adaptation	18 hours (over 5 sessions)	N/A	PI < anxiety ( $d = -.47, p = .006$ ), burnout (emotional exhaustion, $d = -.31, p = .009$ ; depersonalisation, $d = -.22, p = .005$ ; and personal accomplishment, $d = .50, p < .001$ ), depression (depression, $d = -.54, p < .001$ ), and stress & strain (perceived stress, $d = -.54, p = .002$ ; and stress, $d = -.31, p = .002$ ). PI > compassion & empathy (compassion, $d = -.04$ ), resilience (resilience, $d = .17$ ).
(Foureur, Besley, Burton, Yu, & Crisp, 2013)	Nurses & midwives	Convenience sample	28 (40)		MBSR adaptation	1 day (& 8 weeks practice)		PI < distress & anger (distress, $d = -.59, p = .031$ ), and stress & strain (stress, $d = -.65, p = .004$ ). PI > wellbeing, satisfaction & flourishing (sense of coherence, $d = .73, p = .009$ ). PI > anxiety ( $d = -.28, p = .079$ ), and depression ( $d = -.33$ ).
(Galantino, Baime, Maguire, Szapary, & Farrar, 2005)	Healthcare professionals	Convenience sample	84	-	Mindfulness program (specific to study)	8 weeks	N/A	PI < anxiety ( $p = .001$ ), burnout ( $p = .002$ ), depression ( $p = .001$ ), and distress & anger ( $p = .001$ ). PI > compassion & empathy, and stress & strain.
(Gauthier et al., 2015)	Paediatric ICU nurses	Convenience sample	38 (45)	-	Mindfulness program (specific to study)	30 days	N/A	PI < stress & strain (stress, $d = -.40, p = .006$ ). PI > burnout (emotional exhaustion, $d = -.18$ ; depersonalisation, $d = -.13$ ; and personal accomplishment, $d = .12$ ), compassion & empathy (self-compassion, $d = .23$ ), and mindfulness & awareness (mindful attention awareness, $d = .07$ ).
(Gockel et al., 2013)	Trainee social workers	Convenience sample	38	94	MBSR adaptation	10 weeks	N/A	PI > job performance (counselling self-efficacy, $d = .53, p = .005$ ), mindfulness & awareness (mindfulness, $d = .72, p = .034$ ).
(Goodman & Schorling, 2012)	Healthcare professionals	Convenience sample	93	-	Mindfulness for healthcare providers	8 weeks	N/A	PI > burnout (emotional exhaustion, $d = -.29$ ; depersonalisation, $d = -.44$ ; and personal accomplishment, $d = .44$ ), and health (mental health, $d = .78$ ; physical health, $d = -.02$ ).
(Grepmaier, Mitterlehner, Loew, & Nickel, 2007)	Trainee psychotherapists	Convenience sample	58	55	Mindfulness program (specific to study)	9 weeks	Pre-training	PI > job performance (patients' distress, $d = -.93, p < .01$ ).
(Hallman, O'Connor, Hasenau, & Brady, 2014)	Psychiatric service professionals	Convenience sample	12 (13)	-	MBSR	8 weeks	N/A	PI < stress & strain (perceived stress, $d = -.20, p < .05$ ). PI > mindfulness & awareness (mindfulness, $d = .68, p < .05$ ).
(Hemanth & Fisher, 2015)	Clinical psychology trainees	Convenience sample	10	-	Mindfulness program (specific to study)	10 weeks	N/A	Qualitative interviews: PI > compassion & empathy, job performance, relationships, and emotional intelligence & regulation.
(Hopkins & Proeve, 2013)	Trainee psychologists	Convenience sample	11	-	MBCT	8 weeks	N/A	PI > compassion & empathy (emotional concern, $d = -.40$ ; perspective taking, $d = -.37$ ; personal distress, $d = -.23$ ; and fantasy, $d = -.30; p < .01$ ), and mindfulness & awareness (act aware, $d = .11$ ; observe, $d = .43$ ; describe, $d = .18$ ; non-reacting, $d = .77$ ; and non-judging, $d = 1.27; p < .05$ ). PI > stress & strain, (perceived stress, $d = -.67$ ).

(Horner, Piercy, Eure, & Woodard, 2014)	Nurses	Convenience sample	31 (46)	12 (28)	Mindfulness program (specific to study)	10 weeks	Nothing	<p>PI &gt;&lt; burnout, compassion &amp; empathy, mindfulness &amp; awareness, stress &amp; strain, and wellbeing, satisfaction &amp; flourishing (professional quality of life).</p> <p>PI &lt; anxiety (trait, <math>d = -1.41, p = .008</math>), depression (depression with the CESD-10, <math>d = -1.50, p = .002</math>; and depression with the PHQ-9, <math>d = -1.56, p &lt; .001</math>), and stress &amp; strain (perceived stress, <math>d = -1.30, p &lt; .01</math>). PI &gt; health (health responsibility, <math>d = .96</math>; interpersonal relations, <math>d = 1.40</math>; nutrition, <math>d = .34</math>; physical activity, <math>d = .81</math>; spiritual growth, <math>d = .99</math>; stress management, <math>d = 1.17</math>; absenteeism, <math>d = -.50</math>; activity impairment, <math>d = -1.23</math>; presenteeism, <math>d = -1.28</math>; and work productivity loss, <math>d = -1.38; p &lt; .05</math>). PI &gt;&lt; anxiety (state, <math>d = -1.02</math>).</p>
(Johnson et al., 2015)	Healthcare professionals	RCT	20	20	Resilience training	8 weeks	Wait-list	
(Kemper & Khirallah, 2015)	Health professionals	Convenience sample	112	-	Mindfulness in daily life	1 hour	N/A	<p>PI &gt; mindfulness &amp; awareness (cognitive and affective mindfulness, <math>d = .24, p = .004</math>; and mindful attention awareness, <math>d = .20, p &lt; .001</math>), and resilience (resilience, <math>d = .21, p &lt; .001</math>).</p>
(Klatt et al., 2015)	Intensive care IC staff	RCT	34	34	Mindfulness in motion	8 weeks	N/A	<p>PI &gt; resilience (engagement, <math>p = .012</math>; resilience, <math>p = .023</math>; and vigour, <math>p = .033</math>).</p>
(Krasner et al., 2009)	Primary care physicians	Convenience sample	70	-	Mindfulness program (specific to study)	8 weeks	N/A	<p>PI &lt; burnout (emotional exhaustion, <math>d = -.37</math>, depersonalisation, <math>d = -.19</math>; and personal accomplishment, <math>d = .15; p &lt; .001</math>), and distress &amp; anger (distress, <math>d = -.47, p &lt; .001</math>). PI &gt; compassion &amp; empathy (physician empathy, <math>d = .36, p &lt; .001</math>), and mindfulness &amp; awareness (mindfulness, <math>d = .86, p &lt; .001</math>).</p>
(Mackenzie et al., 2006)	Nurses	RCT	16	14	MBSR adaptation	4 weeks	Wait-list	<p>PI &lt; burnout (emotional exhaustion, <math>d = .32, p &lt; .01</math>; depersonalisation, <math>d = -.04, p &lt; .05</math>; and personal accomplishment, <math>d = 1.55, p &lt; .05</math>). PI &gt; wellbeing, satisfaction &amp; flourishing (relaxation dispositions, <math>d = .24, p &lt; .01</math>). PI &gt;&lt; wellbeing, satisfaction &amp; flourishing (intrinsic job satisfaction, <math>d = .17</math>; satisfaction with life, <math>d = -.13</math>; and sense of coherence, <math>d = .16</math>).</p>
(Manotas, Segura, Eraso, Oggins, & McGovern, 2014)	Healthcare professionals	RCT	40 (66)	43 (65)	MBSR adaptation	4 weeks	NR	<p>PI &lt; distress &amp; anger (distress, <math>d = -.61, p = .006</math>), and stress &amp; strain (perceived stress, <math>d = -.68, p &lt; .001</math>). PI &gt; mindfulness &amp; awareness (act aware, <math>d = -.29</math>; observe, <math>d = .23</math>; describe, <math>d = -.28</math>; non judging, <math>d = .32</math>; non reacting, <math>d = .03</math>; and total mindfulness, <math>d = .07; p &lt; .001</math>).</p>
(Martín-Asuero & García-Banda, 2010)	Healthcare professionals	Selected sample	29	-	MBSR adaptation	8 weeks	N/A	<p>PI &lt; depression (rumination, <math>d = -.19, p = .010</math>), and distress &amp; anger (psychological distress, <math>d = -.59, p = .016</math>). PI &gt; wellbeing, satisfaction &amp; flourishing, (negative affect, <math>d = -.26, p = .002</math>). P &gt;&lt; stress &amp; strain (daily stress, <math>d = -.39</math>).</p>
(Martín-Asuero et al., 2014)	Healthcare professionals	RCT	43	25	MBSR adaptation	8 weeks	Wait-list	<p>PI &lt; anxiety (<math>p &lt; .001</math>), burnout (emotional exhaustion, <math>d = -.59</math>; depersonalisation, <math>d = -.32</math>; and personal accomplishment, <math>d = .27; p &lt; .01</math>), depression (<math>p &lt; .05</math>), and distress &amp; anger (distress, <math>d = -.83, p &lt; .001</math>). PI &gt; compassion &amp; empathy (physician empathy, <math>d = .40, p &lt; .05</math>), and mindfulness &amp; awareness (act aware, <math>d = .84</math>; describe, <math>d = .44</math>; observe, <math>d = 1.27</math>; non-reacting, <math>d = 1.21</math>; and non-judging, <math>d = .49; p &lt; .05</math>).</p>

(McConachie, McKenzie, Morris, & Walley, 2014)	Support staff	RCT	66	54	Acceptance and mindfulness workshop	1.5 days	Wait-list	PI < distress & anger (distress, $d = -.35, p < .001$ ). PI >< wellbeing, satisfaction & flourishing (mental wellbeing, $d = .17$ ).
(Mealer et al., 2014)	Intensive care nurses	RCT	13	14	Resilience training program*	12 weeks	Nothing	PI < depression ( $p = .03$ ), and stress & strain (PTSD, $p = .01$ ). PI > resilience ( $p = .01$ ). PI >< anxiety & burnout.
(Moody et al., 2013)	Paediatric oncology staff	RCT	24	23	Mindfulness program (specific to study)	8 weeks	Nothing	PI >< burnout, depression, and stress & strain.
(Moore, 2008)	Trainee clinical psychologists	Convenience sample	16 (23)	-	Mindfulness program (specific to study)	4 weeks	N/A	PI > mindfulness & awareness ( $p = .04$ ) PI >< compassion & empathy, and stress & strain.
(Newsome, Christopher, Dahlen, & Christopher, 2006)	Counsellors	Convenience sample	33	-	Mindfulness curriculum (specific to study)	15 weeks	N/A	Qualitative interviews: PI >< emotional intelligence & regulation (acceptance), mindfulness & awareness (awareness), health, relationships, and wellbeing, satisfaction & flourishing (spirituality).
(Newsome, Waldo, & Gruszka, 2012)	Trainee helping professionals	Convenience sample	31	-	Mindfulness program (specific to study)	6 weeks	N/A	PI < stress & strain (perceived stress, $d = -1.01, p < .0001$ ). PI > compassion & empathy (self-compassion, $d = 1.13, p < .0001$ ), and mindfulness & awareness (mindful attention awareness, $d = .91, p < .001$ ).
(Noone & Hastings, 2010)	Disability support workers	Convenience sample	34	-	Promotion of acceptance in carers and teachers	1.5 days	N/A	PI < distress & anger (distress, $d = -.54, p = .020$ ). PI >< stress & strain (stress, $d = -.13$ ).
(Pflugeisen, Drummond, Ebersole, Mundell, & Chen, 2015)	Physicians	Convenience sample	19 (23)	-	MBSR adaptation	8 weeks	N/A	PI < burnout (emotional exhaustion, $d = -.46$ ; depersonalisation, $d = -.32$ ; and personal accomplishment, $d = .56$ ; $p < .03$ ), and stress & strain (perceived stress, $d = -.87, p = .005$ ). PI > mindfulness & awareness (mindfulness skills, $d = .84, p = .01$ ).
(Pipe et al., 2009)	Nurses	RCT	15	17	MBSR adaptation	4 weeks	Wait-list	PI < distress & anger (psychological distress, $d = -.39, p = .009$ ). PI >< depression ( $d = -.54$ ), job performance (caring efficacy, $d = .48$ ), and relationships (interpersonal sensitivity, $d = .38, p = .29$ ).
(Poulin et al., 2008) [study 1]	Nurses	RCT	16	10	MBSR adaptation	4 weeks	Imagery & progressive muscle relaxation	PI > wellbeing, satisfaction & flourishing (relaxation, $d = -.63, p < .05$ ). PI >< burnout (emotional exhaustion, $d = -.07$ ; depersonalisation, $d = -.16$ ; and personal accomplishment, $d = .73$ ).
(Raab et al., 2015)	Mental health professionals	Convenience sample	22	-	MBSR	8 weeks	N/A	PI > compassion & empathy (self-compassion, $d = .48, p = .003$ ). PI >< burnout (depersonalisation, $d = -.11$ ; emotional exhaustion, $d = -.20$ ; and personal accomplishment, $d = .20$ ), and wellbeing, satisfaction & flourishing (quality of life, $d = .02$ ).
(Rimes & Wingrove, 2011)	Trainee clinical psychologists	Convenience sample	20	-	MBCT	8 weeks	N/A	PI < depression (rumination, $d = -.57, p < .0005$ ). PI > anxiety ( $d = .26, p < .05$ ), compassion & empathy (fantasy, $d = .52$ ; self-compassion, $d = .48$ , empathic concern, $d = .00$ ; personal distress, $d = -.06$ ; and perspective taking, $d = -.03$ ; $p < .05$ ), and mindfulness & awareness (act aware, $d = .10$ ; non non-reacting, $d = .59$ ; non judging, $d = .52$ ; describe, $d = .31$ ; and observe, $d = .38$ ; $p < .001$ ). PI >< stress & strain (perceived stress, $d = -.23$ ).
(Rocco, Dempsey, & Hartman, 2012)	Mental health professionals	Convenience sample	16	-	Calm abiding meditation	8 weeks	N/A	Qualitative interviews: PI > emotional intelligence & regulation (acceptance, and emotion regulation), mindfulness & awareness (awareness), and health (health behaviours).
(Ruths et al., 2013)	Mental health professionals	Convenience sample	27	-	MBCT	8 weeks	N/A	PI < distress & anger ( $p = .003$ ). PI > mindfulness & awareness ( $p = .008$ ). PI >< anxiety, and wellbeing, satisfaction & flourishing (satisfaction with life).

(Shapiro et al., 1998b)	Trainee doctors	RCT	37	36	Stress reduction and relaxation	7 weeks	Wait-list	PI < anxiety (state, $d = -.46$ ; and trait, $d = -.59$ ; $p < .05$ ), depression (depression, $d = -.46$ , $p < .006$ ), and distress & anger (psychological distress, $d = -.69$ , $p < .02$ ). PI > compassion & empathy (empathy, $d = .47$ , $p < .05$ ), and wellbeing, satisfaction & flourishing (spirituality, $d = .32$ , $p < .02$ ).
(Shapiro et al., 2005)	Healthcare professionals	RCT	18	20	MBSR	8 weeks	Wait-list	PI < stress & strain (perceived stress, $d = -.15$ , $p = .04$ ). PI > compassion & empathy (self-compassion, $d = .02$ , $p = .004$ ). PI >> burnout (emotional exhaustion, $d = -.18$ ; depersonalisation, $d = -.74$ ; and personal accomplishment, $d = .64$ ), distress & anger (distress, $d = -.07$ ), and wellbeing, satisfaction & flourishing (satisfaction with life, $d = .15$ ).
(Shapiro et al., 2007)	Trainee psychotherapists	Convenience sample	22	32	MBSR	8 weeks	Psychology course	PI < anxiety (state, $d = -.55$ , $p = .0005$ ; and trait, $d = -.91$ , $p = .0002$ ), depression (rumination, $d = -.41$ , $p = .0006$ ), and stress & strain (perceived stress, $d = -.67$ , $p < .0001$ ). PI > compassion & empathy (self-compassion, $d = .42$ , $p < .0001$ ), mindfulness & awareness (mindful attention awareness, $d = .36$ , $p = .006$ ), and wellbeing, satisfaction & flourishing (positive affect, $d = .57$ , $p = .0002$ ; and negative affect, $d = -.46$ , $p = .04$ ).
(Singh et al., 2015)	Disability professionals	Convenience sample	9	-	Mindfulness-based positive behavioural support	7 days	N/A	PI < stress & strain (perceived stress, $d = -3.89$ , $p < .001$ ), PI > job performance (restraining patients, $p < .001$ ; staff injury, $p < .001$ ; , disciplining patients, $p < .001$ ).
(Singh et al., 2006)	Psychiatric staff	Convenience sample	18 (3 teams)	18 (same as expt group)	Mindfulness-based mentoring	11, 8 or 6 sessions	Control within & between teams	PI > job performance (team functioning, $p < .001$ ).
(Song & Lindquist, 2015)	Trainee nurses	RCT	21 (25)	23 (25)	MBSR	8 weeks	Wait-list	PI < anxiety ( $d = -.50$ , $p = .023$ ), depression ( $d = -.70$ , $p = .002$ ), and stress & strain (stress, $d = -.85$ , $p < .001$ ). PI > mindfulness & awareness (mindful attention awareness, $d = .13$ , $p = .010$ ).
(Stew, 2011)	Trainee occ therapists	Convenience sample	12	-	MBSR adaptation	4 weeks	N/A	Qualitative interviews ( $n = 10$ ): PI > emotional intelligence & regulation (acceptance), and mindfulness & awareness.
(Tarrasch, 2014)	Trainee counsellors and support staff	Convenience sample	19	-	Mindfulness curriculum (specific to study)	2 terms	N/A	Qualitative interviews ( $n = 19$ ) PI > emotional intelligence & regulation (acceptance), mindfulness & awareness (awareness, calmness, coping).
(Van der Riet, Rossiter, Kirby, Dluzewska, & Harmon, 2015)	Trainee nurses	Convenience sample	14	-	Mindfulness program (specific to study)	7 weeks	N/A	Qualitative analysis: PI < stress & strain. PI > mindfulness & awareness (awareness), emotional intelligence & regulation, and relationships.
(West et al., 2014)	Physicians	RCT	35 (37)	37	Small group curriculum*	10 weeks	Nothing	PI >> compassion & empathy (physician empathy, $d = -.05$ ), stress & strain (perceived stress, $d = .13$ ); and wellbeing, satisfaction & flourishing (job satisfaction, $d = -.14$ ).

All statistically significant results are reported. Effect sizes were calculated when means and standard deviations were available, otherwise, just statistically significant differences are offered. Note. < = decreases in; > = increases in; >> = no change in; expt = experimental group; cnt = control group; PI = post-intervention; NR = not-reported; MBCT = mindfulness-based cognitive therapy; MBSR = mindfulness-based stress reduction; MBST = mindfulness-based stress reduction therapy. MM = mindfulness meditation; NCC = neural correlates of consciousness; NR = not recorded; N/A = not applicable; NA = not available; RCT = randomized controlled trial; \* = number in parenthesis is the initial sample size (if different from sample size featured in analysis); \*\* = mindfulness just one component of broader intervention.

**Table 2.** Overview of non-intervention studies.

Authors	Workplace	Meditators	Non-meditators	Analysis	Primary result
(Choi & Koh, 2015)	Nurses	-	330	Correlations	Mindfulness correlation: < stress & strain (job stress, $r = -.279, p < .001$ ). > job satisfaction ( $r = .171, p = .002$ ).
(Christopher et al., 2011)	Counsellors & psychotherapists	13	3	Qualitative	Mindfulness > emotional intelligence & regulation (acceptance, and self-regulation), mindfulness & awareness (awareness), job performance, and relationships.
(Cigolla & Brown, 2011)	Psychotherapists	6	-	Qualitative	Mindfulness > emotional intelligence & regulation (acceptance), mindfulness & awareness (awareness), job performance, relationships, and wellbeing, satisfaction & flourishing (spirituality).
(Di Benedetto & Swadling, 2014)	Psychologists	-	167	Correlation	Mindfulness correlation: < burnout ( $r = -.42, p < .0003$ ).
(Dauenhauer, 2006)	Professional caregivers	-	20	Qualitative	Mindfulness > emotional intelligence & regulation (acceptance, and sensitivity), mindfulness & awareness (awareness), and relationships.
(Gill, Waltz, Suhrbier, & Robert, 2015)	Psychotherapists	7	-	Qualitative	Mindfulness > emotional intelligence & regulation (acceptance), mindfulness & awareness (awareness), job performance, relationships, and wellbeing, satisfaction & flourishing (wellbeing).
(Keane, 2014)	Psychotherapists	-	40	Correlations	Mindfulness (FFMQ, all facets) correlation: > compassion & empathy (IRI Perspective taking; $r$ range .44-.60, $p < .001$ ), mindfulness & awareness (FFMQ Observe, Non-judging, Non-reactivity) correlation: > compassion & empathy (IRI Global empathy; $r$ range .44-.60, $p < .001$ ).
(Kemper et al., 2015)	Health professionals	-	213	Correlations	Mindfulness correlation: stress & strain (perceived stress, $r = -.58, p < 0.001$ ). > health (health, $r = .37, p < .01$ ; sleep quality, $r = -.32, p < .01$ ; and global mental health, $r = .56, p < .001$ ), resilience ( $r = .5, p < .01$ ), and compassion & empathy (self-compassion, $r = .63, p < .001$ ).
(McCollum & Gehart, 2010)	Psychotherapists	13	-	Qualitative	Mindfulness > emotional intelligence & regulation (acceptance), and job performance.
(McCracken & Yang, 2008)	Rehabilitation workers	-	98	Correlations	Mindfulness correlation: < burnout (exhaustion; $r = -.43, p < .05$ ), and stress & strain ( $r = .23, p < .001$ ). > health ( $r = .30, p < .01$ ; vitality, $r = .43, p < .01$ ; social Functioning, $r = .44, p < .001$ ; emotion Functioning, $r = .40, p < .001$ ; and emotion role, $r = .33, p < .001$ ). >> wellbeing, satisfaction & flourishing (job satisfaction).
(Razzaque, Okoro, & Wood, 2015)	Clinical psychologists	-	76	Correlations	Mindfulness correlation: > relationships (therapeutic alliance, $r = .356, p < .01$ ).
(Ryan, Safran, Doran, & Muran, 2012)	Psychotherapists	-	52 (26 dyads)	Correlations	Mindfulness correlation: > relationships (interpersonal functioning, $p < .05$ ; and therapeutic alliance, $p < .05$ ). PI >> job performance (patient distress).
(Simon, Ramsenthaler, Bausewein, Krischke, & Geiss, 2009)	Palliative care professionals	-	10	Qualitative	Mindfulness > emotional intelligence & regulation (acceptance), and job performance.
(Talisman, Harazduk, Rush, Graves, & Haramati, 2015)	Medical training facilitators	62	-	Correlations & qualitative	Mindfulness correlation: < emotional intelligence & regulation (self-affiliation, $r = .413, p < .05$ ). Qualitative interviews: > mindfulness & awareness, compassion & empathy, job performance, and relationships.
(Westphal et al., 2015)	Intensive care nurses	-	50	Correlations	Mindfulness correlation: < anxiety ( $r = -.55, p < .001$ ), burnout (depersonalization; $r = -.37, p < .001$ ; emotional exhaustion; $r = -.52, p < .001$ ), and depression ( $r = .49, p < .001$ ).

Note. < = negative correlation with; > = positive correlation with; >= = no correlation.



An overview of the findings is shown in table 3 below. This shows whether outcomes were either: (a) increased in relation to an MBI; (b) did not change in relation to an MBI (or in exceptional cases, changed in a ‘negative’ direction); or (c) were found in non-intervention studies to be associated with mindfulness (i.e., through regression analyses). A more detailed breakdown of the results is included below in the discussion, featuring tables detailing all the studies that assessed a given outcome.

Table 3. *Summary of Common Outcomes across all Studies*


Outcome	Number of studies assessing	Improvement related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Anxiety	16	9	7, + *1	1
Burnout	25	11	11	3
Compassion & empathy	28	16	9	3
Depression	18	10	7	1
Distress & anger	16	14	2	0
Emotional intelligence & regulation	21	12	2	7
Health	7	3	2	2
Job performance	13	6	1	6
Mindfulness & awareness	39	27	6	6
Relationships	13	5	2	6
Resilience	6	3	2	1
Stress & strain	40	25	12, + *1	3
Wellbeing, satisfaction & flourishing	24	12	11, + *1	2, + *1

*Note:* \* = studies showing *worsening* outcomes in relation to mindfulness. In instances where the total number of studies does not appear to be an accurate product of the other three columns (e.g., in the case of anxiety), this is because some studies used multiple measures with respect to a given outcome, and observed both a significant impact and no significant change.

## Discussion

MBIs generally had a positive impact upon all outcome measures. However, there were some areas in which findings were more equivocal (including burnout, health, resilience, and generic ‘wellbeing’). This discussion will run through the main outcomes in turn, beginning with mindfulness and awareness itself.

Figure 1. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Flow Diagram



## Mindfulness & Awareness

MBIs certainly appear effective at engendering mindfulness, with a small to medium effect size ( $d=.36$ ) as assessed by 33 intervention studies, as shown in table 4 below. The vast majority of these ( $n = 27$ ) showed an increase in mindfulness in relation to an MBI, while six found no significant improvement. However, as positive as these headline figures are, further nuance is provided by digging a little deeper into the results, since a range of scales were used across the studies – scales which construct mindfulness in diverse ways – with some interesting variation. This diversity of scales is both a weakness and a strength. It is a weakness inasmuch as it difficult to draw comparisons across studies. Indeed, inconsistency in the use of scales across studies was a common theme in this review. That said, the diversity of measures does allow us to discern nuances in the development of mindfulness. The most popular tool was Brown and Ryan’s unidimensional (2003) Mindful Attention and Awareness Scale (MAAS), which assesses dispositional mindfulness according to a single core characteristic of mindfulness (open and receptive awareness), which essentially aligns with Kabat-Zinn’s (2003) definition cited above.

By contrast, a number of studies deployed multidimensional scales, most notably Baer, Smith, Hopkins, Krietemeyer, and Toney’s (2006) Five Facets of Mindfulness Questionnaire (FFMQ), which focuses more on mindfulness *skills* (as opposed to the dispositional mindfulness of the MAAS). Here it was difficult to discern a coherent pattern among the studies with respect to the five dimensions/skills. For instance, consider Hopkins and Proeve (2013), Manotas et al. (2014), Martin-Asuero et al. (2014), and Rimes and Wingrove (2011). Their respective effect sizes for the five dimensions varied considerably, as follows: observing (.43, .23, 1.27, .38); describing (.18, -.28, .44, .31); non-judging of inner experience (1.27, .32, .49, .52); non-reactivity to inner experience (.77, .03, 1.21, .59); and acting with awareness (.11, -.29, .84, .10). Thus, there was considerable variation between studies with respect to the

different dimensions; for instance, ‘non-reactivity’ ranged from .03 (Manotas et al., 2014) to 1.21 (Martin-Asuero et al., 2014). Moreover, there was also strong variation within individual studies across the dimensions. For instance, whereas Manotas et al. found small effect sizes for observing (.23) and non-judging (.32), they observed no change with respect to non-reactivity (.03), and actually saw *worsening* skills in describing (-.28) and acting with awareness (-.29). Such variation shows the value of drilling down into the fine-grained details of studies. Furthermore, it highlights the notion that – so far as multidimensional scales are concerned – mindfulness is not a monolithic construct, but rather comprises nuances, upon which there may be differential rates of change and development.

Table 4. *Mindfulness & Awareness Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Five facets of mindfulness questionnaire	(Brooker et al., 2013) (Hopkins & Proeve, 2013) ( <b>Manotas et al., 2014</b> ) ( <b>Martín-Asuero et al., 2014</b> ) (Rimes & Wingrove, 2011)	(De Vibe et al., 2013) ( <b>Duchemin et al., 2015</b> )	(Keane, 2014)
Freiberg mindfulness inventory	(Gockel et al., 2013)		
Kentucky inventory of mindfulness skills	(Dobie et al., 2015) (Moore, 2008) (Pflugeisen et al., 2015)		
Mindful attention awareness scale	( <b>Cohen-Katz, Wiley, Capuano, Baker, Kimmel, et al., 2005</b> ) (Kemper & Khirallah, 2015) (Newsome et al., 2012) (Ruths et al., 2013) (Shapiro et al., 2007) ( <b>Song &amp; Lindquist, 2015</b> )	(J. S. Cohen & Miller, 2009) (Gauthier et al., 2015) (Horner et al., 2014)	
Mindful therapy scale	(Aggs & Bambling, 2010)		
Qualitative interviews	(Beckman et al., 2012) ( <b>Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005</b> ) (Dorian & Killebrew, 2014) (Felton et al., 2015) (Rocco et al., 2012) (Stew, 2011) (Tarrasch, 2014) (Van der Riet et al., 2015)	(Newsome et al., 2006)	(Christopher et al., 2011) (Cigolla & Brown, 2011) (Dauenhauer, 2006) (Gill et al., 2015) (Talisman et al., 2015)
Toronto mindfulness scale	(Brady et al., 2012) (Hallman et al., 2014)		
Two factor mindfulness scale	(Krasner et al., 2009)		

Note. Authors in bold denote RCT studies.

## Anxiety

Turning now to the various wellbeing outcomes, firstly, On balance, mindfulness appears to have a beneficial impact upon anxiety, as shown in table 5 below, with a medium

effect size ( $d=-.51$ ). While nine studies reported an improvement in relation to an MBI, six observed no change (although one further study (Rimes & Wingrove, 2011) actually reported worsening levels of anxiety). In addition, of the non-intervention studies, Westphal et al. (2015) reported an inverse correlation between anxiety and mindfulness. Given the high prevalence and burden of anxiety among healthcare professionals – e.g., a survey of Chinese nurses found the prevalence of clinically-significant anxiety symptoms to be as high as 43.4% – the improvements in anxiety linked to MBIs are noteworthy, modest though they are. As with mindfulness, a range of scales were deployed. The most prominent were Spielberger, Gorsuch, and Lushene's (1970) State-Trait Anxiety Inventory, and Lovibond and Lovibond's (1995) Depression Anxiety Stress Scale (DASS). The multidimensional DASS is particularly useful, since it also covers depression and stress, therefore it enables more ground to be covered with the one scale (thus reducing the empirical demands placed on participants).

Table 5. *Anxiety Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Beck anxiety inventory	(J. S. Cohen & Miller, 2009)		
Burns anxiety inventory	(Barbosa et al., 2013)		
Depression anxiety stress scale [anxiety]	(Dobie et al., 2015) (Fortney et al., 2013) <b>(Song &amp; Lindquist, 2015)</b>	<b>(Duchemin et al., 2015)</b> (Foureur et al., 2013)	
Hospital anxiety & depression scale [anxiety]		<b>(Mealer et al., 2014)</b> (Rimes & Wingrove, 2011)!	(Westphal et al., 2015)
Penn state worry questionnaire		(Ruths et al., 2013)	
Profile of mood states [anxiety]	(Galantino et al., 2005)	<b>(Martín-Asuero et al., 2014)</b>	
State trait anxiety inventory	<b>(Johnson et al., 2015)</b> (Shapiro, G. Schwartz, & G. Bonner, 1998a) (Shapiro et al., 2007)	<b>(Johnson et al., 2015)</b> (Ruths et al., 2013)	

*Note.* Authors in bold denote RCT studies; ! in third column = poorer outcome in relation to mindfulness.

## Burnout

Regarding burnout, the results were more equivocal, as shown in table 6 below. Of the 22 intervention studies examining this, only 11 registered a significant improvement, while equally 11 reported no significant change. Nevertheless, the overall effect size in this outcome was small to medium ( $d=-.33$ ). In addition, three non-intervention studies observed an inverse correlation between burnout and mindfulness.

One possible explanation for the relatively equivocal results with respect to the MBIs may lie in the relatively small sample sizes of many studies. Some intervention studies that did not find a significant improvement in burnout certainly observed trends in the predicted direction (e.g., Mealer et al., 2014; Poulin et al., 2008; Raab et al., 2015; Shapiro et al., 2005), although De Vibe et al. (2013) found trends in the other direction. Larger sample sizes may allow any impact of MBIs on burnout to be clearer. Another possible explanation is the multifaceted nature of the construct. The dominant measure used was the Maslach Burnout Inventory (Maslach, Jackson, & Leiter, 1986), which has three dimensions: emotional exhaustion, cynicism (also known as ‘depersonalisation’), and professional efficacy (or accomplishment). When considering the components separately, a number of studies found that MBIs tended to have a stronger positive effect (albeit still non-significant) on emotional exhaustion compared to the other two components (e.g., Barbosa et al., 2013; Duchemin, Steinberg, Marks, Vanover, & Klatt, 2015; Moody et al., 2013; Poulin et al., 2008).

Table 6. *Burnout outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Copenhagen burnout inventory	(Bazarko et al., 2013)	(Brooker et al., 2013)	(Di Benedetto & Swadling, 2014)
Maslach burnout inventory	(Barbosa et al., 2013) (Brady et al., 2012) <b>(Cohen-Katz, Wiley, Capuano, Baker, Kimmel, et al., 2005)</b> (Fortney et al., 2013) (Galantino et al., 2005) (Krasner et al., 2009) <b>(Mackenzie et al., 2006)</b> <b>(Martín-Asuero et al., 2014)</b> (Pflugeisen et al., 2015)	<b>(De Vibe et al., 2013)</b> <b>(Duchemin et al., 2015)</b> (Gauthier et al., 2015) (Goodman & Schorling, 2012) <b>(Mealer et al., 2014)</b> <b>(Moody et al., 2013)</b> <b>(Poulin et al., 2008)</b> (Raab et al., 2015) <b>(Shapiro et al., 2005)</b>	(Westphal et al., 2015)
Professional quality of life scale [burnout]		(Horner et al., 2014)	
Profile of mood states [fatigue]	<b>(Martín-Asuero et al., 2014)</b>		
Profile of mood states [vigour]	(Galantino et al., 2005) (Krasner et al., 2009)		
Qualitative interviews	(Christopher et al., 2006)		
SF-12-v2 health survey [vitality]			(McCracken & Yang, 2008)

Note: Authors in bold denote RCT studies

## Depression

The results were generally favourable with respect to depression, as shown in table 7 below, with an overall medium effect size ( $d=-.53$ ). Of the 16 intervention studies examining this, while 10 registered a significant improvement, seven reported no significant change. Meanwhile, in terms of non-intervention studies, Westphal et al. (2015) reported an inverse correlation between depression and mindfulness. The relatively favourable results for this outcome are welcome, given the relatively high incidence of depression in HCPs. For instance, a study by Caplan (1994) in the UK found high levels of depression, particularly among GPs, 27% of whom scored as borderline or likely to be depressed. These figures contrast with estimates that around 2.3% of the general UK adult population experience a depressive episode at any one time (i.e., in the past week), with 9% experiencing mixed anxiety and depressive disorder (The Health and Social Care Information Centre, 2009). There are many hypothesised reasons for greater liability to depression among HCPs, including personality traits like perfectionism, burdens of clinical responsibility, and reluctance to seek-treatment (Bright & Krahn, 2011). Whatever the reasons, it is encouraging that, on balance, MBIs appear to help in this regard – although it bears repeating that over one third of intervention studies reported no significant change – reflecting the more established efficacy of MBIs such as MBCT with respect to depression (Segal et al., 2002).

Table 7. *Depression Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Beck depression inventory		(Moody et al., 2013)	
Centre for epidemiological studies – depression	(Johnson et al., 2015)	(J. S. Cohen & Miller, 2009)	
Depression anxiety stress scale [depression]	(Fortney et al., 2013) (Song & Lindquist, 2015)	(Brooker et al., 2013) (Dobie et al., 2015) (Duchemin et al., 2015) (Foureur et al., 2013)	
Emotional Control Questionnaire	(Martín-Asuero & García-Banda, 2010)		
Hospital anxiety & depression scale [depression]	(Mealer et al., 2014)		(Westphal et al., 2015)
Patient health questionnaire	(Johnson et al., 2015)		
Profile of mood states [depression]	(Galantino et al., 2005) (Martín-Asuero et al., 2014)		
Reflection-rumination questionnaire	(Rimes & Wingrove, 2011) (Shapiro et al., 2007)		

---

*Note.* Authors in bold denote RCT studies.

## Stress & Strain

More consistent results were found for stress, by far the outcome receiving the most attention, as shown in table 8 below. Of the 37 intervention studies examining this, 25 registered a significant improvement in relation to an MBI, while 12 reported no significant change (although, in addition, Brooker et al. (2013) observed worsening levels). The global effect size for this outcome was small to medium ( $d=-.42$ ). Three non-intervention studies also observed an inverse correlation between stress and mindfulness. These generally positive results are again welcome: as with the other outcomes, stress is generally higher among HCPs than in the general population: Firth-Cozens (2003) reported that the proportion of HCPs being above threshold levels of stress is around 28% in surveys, compared with about 18% in the general working population. As with depression, a similar range of factors have been implicated in elevated stress levels among HCPs, from long working hours, to the burden of clinical responsibility (Sochos, Bowers, & Kinman, 2012). Unfortunately, as highlighted above, these burdens have only increased over recent years, due to factors such as curbs on healthcare spending meaning that overwork has become even more acute. As noted above, a survey of NHS staff found that 61% reporting feeling stress all or most of the time, and 59% stating that the stress is worse this year than last year (Dudman et al., 2015). Thus, while it is encouraging that MBIs may help alleviate or prevent stress, it is of course vital that these underlying structural causes are also addressed.



Table 8.

*Stress & Strain Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Depression	(Dobie et al., 2015) ( <b>Duchemin et al., 2015</b> )	(Brooker et al., 2013)!	
anxiety stress scale [stress]	(Fortney et al., 2013) (Foureur et al., 2013)		
Derogatis stress profile	(Beddoe & Murphy, 2004) ( <b>Song &amp; Lindquist, 2015</b> )		
Job stress questionnaire			(Choi & Koh, 2015)
Perceived medical school stress	( <b>De Vibe et al., 2013</b> )		
Perceived stress questionnaire		(Martín-Asuero & García-Banda, 2010)	
Posttraumatic diagnostic scale	( <b>Mealer et al., 2014</b> )		
Perceived stress scale	(Bazarko et al., 2013) (J. S. Cohen & Miller, 2009) ( <b>Erogul et al., 2014</b> ) (Fortney et al., 2013) (Hallman et al., 2014) ( <b>Johnson et al., 2015</b> ) ( <b>Manotas et al., 2014</b> ) (Newsome et al., 2012) (Pflugeisen et al., 2015) ( <b>Shapiro et al., 2005</b> ) (Shapiro et al., 2007) (Singh et al., 2015) (Brady et al., 2012)	(Bond et al., 2013) (Bonifas & Napoli, 2014) (Brooker et al., 2013)! (Burnett & Pettijohn, 2015) (Hopkins & Proeve, 2013) ( <b>Moody et al., 2013</b> ) (Moore, 2008) (Rimes & Wingrove, 2011) ( <b>West et al., 2014</b> )	(Kemper et al., 2015)
Mental health professionals stress scale			
Professional quality of life scale [stress]		(Horner et al., 2014)	
Nursing stress scale	(Gauthier et al., 2015)		
Qualitative interviews	(Felton et al., 2015) (Van der Riet et al., 2015) (Christopher et al., 2006)	(Bond et al., 2013) ( <b>Moody et al., 2013</b> )	
Salivary $\alpha$ -Amylase	( <b>Duchemin et al., 2015</b> )		
Salivary cortisol		(Galantino et al., 2005)	
Staff stressor questionnaire		(Noone & Hastings, 2010)	
Stress (survey question)			(McCracken & Yang, 2008)
Stress & tension ratings	(Aggs & Bambling, 2010)		

Note. Authors in bold denote RCT studies; ! in third column = poorer outcome in relation to mindfulness.

**Other Wellbeing Outcomes**

This general pattern of mindfulness being associated with wellbeing was followed across the other outcomes. For example, 15 studies examined the relationship between mindfulness and distress or anger, and generally found it to have a positive impact as shown in table 9 below, with a total effect size between medium and large ( $d=.60$ ): 13 registered an improvement, whereas only two reported no change. Mindfulness was also associated with various more ‘positive’ wellbeing outcomes, although the results overall were equivocal as shown in table 10, with an overall small to medium effect size ( $d=.36$ ). Of the 21 intervention studies examining outcomes in this area, while 12 registered an improvement, 11 reported no change. (The non-additive nature of the numbers in that last sentence reflects the fact that two studies used multiple wellbeing measures, and reported both significant and non-significant outcomes in relation to these.) In addition, two non-intervention studies observed a correlation with mindfulness (while McCracken and Yang (2008, actually observed an inverse correlation). Mindfulness also appeared conducive to health with a medium to large effect size ( $d=.62$ ), although there were fewer studies focusing on such outcomes, as seen in table 11 below. Of the 5 intervention studies examining this, three registered an improvement, while two reported no change; additionally, two non-intervention studies observed a correlation with mindfulness

Table 9. *Distress & Anger Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Brief symptom inventory	( <b>Manotas et al., 2014</b> )	( <b>Cohen-Katz, Wiley, Capuano, Baker, Kimmel, et al., 2005</b> ) (Shapiro et al., 2005)	
Depression anxiety stress scale	(Foureur et al., 2013)		
General health questionnaire	( <b>De Vibe et al., 2013</b> ) (Foureur et al., 2013) ( <b>McConachie et al., 2014</b> ) (Noone & Hastings, 2010) (Ruths et al., 2013)		
Profile of mood states [anger]	(Galantino et al., 2005) (Krasner et al., 2009) ( <b>Martín-Asuero et al., 2014</b> )		
Qualitative interviews	(Dorian & Killebrew, 2014)		
Symptom checklist-90-R	(Martín-Asuero & García-Banda, 2010) ( <b>Pipe et al., 2009</b> ) ( <b>Shapiro et al., 1998a</b> )		

Note. Authors in bold denote RCT studies.

Table 10.

*Wellbeing, Satisfaction & Flourishing Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Brief serenity index	(Bazarko, Cate, Azocar, & Kreitzer, 2013)		
Index of core spiritual experiences	<b>(Shapiro et al., 1998a)</b>		
Job satisfaction scale		<b>(Mackenzie et al., 2006)</b>	
Job satisfaction (survey question)			(McCracken & Yang, 2008)!
Meaning in life questionnaire		(J. S. Cohen & Miller, 2009)	
Minnesota satisfaction questionnaire		(Brooker et al., 2013)!	
Physician job satisfaction scale		<b>(West et al., 2014)</b>	
Positive & negative affect scale	(Brooker et al., 2013) (Martín-Asuero & García-Banda, 2010) (Shapiro et al., 2007)		
Professional quality of life scale	<b>(Duchemin et al., 2015)</b>	(Brooker et al., 2013) (Horner et al., 2014)	
Quality of life index	(Bonifas & Napoli, 2014)		
Quality of life inventory		(Raab et al., 2015)	
Qualitative interviews	(Fisher & Hemanth, 2015)	(Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005)	(Gill et al., 2015)
Qualitative interviews (spirituality)		(Newsome et al., 2006)	(Cigolla & Brown, 2011)
Satisfaction with life scale	<b>(Mackenzie et al., 2006) (Poulin et al., 2008)</b>	(Brooker et al., 2013) (J. S. Cohen & Miller, 2009) (Ruths et al., 2013) <b>(Shapiro et al., 2005)</b> <b>(Mackenzie et al., 2006)</b>	
Sense of coherence	(Foureur et al., 2013)		
Smith relaxation disposition inventory	<b>(Mackenzie et al., 2006) (Poulin et al., 2008)</b>		
Subjective wellbeing scale	<b>(De Vibe et al., 2013)</b>		
Warwick-Edinburgh mental wellbeing scale		<b>(McConachie et al., 2014)</b>	

*Note.* Authors in bold denote RCT studies; ! in third column = poorer outcome in relation to mindfulness; ! in fourth column = inverse correlation with mindfulness.

Table 11. *Health Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Health promoting lifestyle profile	<b>(Johnson et al., 2015)</b>		
Patient reported outcomes measurement information system			(Kemper et al., 2015)
Qualitative interviews	(Rocco et al., 2012)	(Newsome et al., 2006)	
SF-12-v2 health survey [physical health]	(Bazarko et al., 2013)	(Goodman & Schorling, 2012)	(McCracken & Yang, 2008)
Workplace productivity and impairment general health questionnaire	<b>(Johnson et al., 2015)</b>		

*Note.* Authors in bold denote RCT studies.

In addition to these primary wellbeing outcomes, mindfulness was also linked to various factors and qualities *associated* with wellbeing – including relationships, resilience, and emotional intelligence – which may provide an explanation for the generally positive outcomes adumbrated above. Regarding relationships, mindfulness practice seems to have a positive impact as seen in Table 12, with an effect size between small and medium ( $d=.46$ ). Most of the 13 studies analysing this outcome found either improvement or benign association with regard to mindfulness, while only two failed to provide significant results. Similarly, mindfulness was also linked to resilience, although the results were somewhat equivocal: as shown in Table 13, of the five intervention studies examining this, three observed an improvement while two reported no significant change. The overall effect size for this outcome was small ( $d=.21$ ). Meanwhile, Kemper et al. (2015) observed a correlation with mindfulness.

Mindfulness appeared to impact also upon emotional intelligence and regulation, as shown in table 14 below. Of the 14 intervention studies examining this, 12 observed an improvement and only two reported no significant change. Nevertheless, this time no effect size was found ( $d=.18$ ). In addition, seven non-intervention studies observed a correlation with mindfulness. The significance of this particular outcomes is that, as outlined above, a key mechanism through which mindfulness is thought to exert its positive effects is reperceiving (Shapiro et al., 2006), also known as decentring (Fresco et al., 2007). This ability – which means that people are better able to detach themselves from distressing qualia that might otherwise precipitate feelings of stress etc. – could be regarded as an aspect of a more general capacity of emotion regulation (Walsh & Shapiro, 2006). The suggestion is that mindfulness might positively impact on wellbeing in the following way: (a) mindfulness involves introspective practices that facilitate the development of attention and awareness skills; (b) development of these skills leads to enhanced emotional regulation (including abilities such as reperceiving); and (c) emotional regulation is a meta-skill that subserves multiple wellbeing outcomes (while, conversely, poor regulation skills are a transdiagnostic factor underlying diverse psychopathologies; Aldao et al., 2010). Future work may help to elucidate these hypothesised causal chains further, e.g., through longitudinal studies deploying regression analyses.

Table 12. *Resilience Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Brief resilience scale	(Kemper & Khirallah, 2015)		(Kemper et al., 2015)
Connor David resiliency scale	<b>(Klatt et al., 2015) (Mealer et al., 2014)</b>		
Resilience scale		<b>(Erogul et al., 2014)</b> (Fortney et al., 2013)	
Utrecht work engagement scale [vigour]	<b>(Klatt et al., 2015)</b>		

Note: Authors in bold denote RCT studies.

Table 13. *Relationships Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Inventory of interpersonal problems-32			(Ryan et al., 2012)
Symptom checklist-90-R [interpersonal sensitivity]		(Pipe et al., 2009)	
Qualitative interviews	(Beckman et al., 2012) ( <b>Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005</b> ) (Hemanth & Fisher, 2015) (Van der Riet et al., 2015)	(Newsome et al., 2006)	(Christopher et al., 2011) (Cigolla & Brown, 2011) (Dauenhauer, 2006) (Gill et al., 2015) (Talisman et al., 2015)
Social-connectedness scale	(J. S. Cohen & Miller, 2009)		

Note: Authors in bold denote RCT studies.

Table 14. *Emotional Intelligence & Regulation Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Self-report of emotional intelligence	(J. S. Cohen & Miller, 2009)		
Qualitative interviews	( <b>Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005</b> ) (de Zoysa et al., 2014) (Hemanth & Fisher, 2015) (Rocco et al., 2012) (Van der Riet et al., 2015)		(Christopher et al., 2011) (Talisman et al., 2015)
Qualitative interviews (acceptance)	( <b>Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005</b> ) (Dorian & Killebrew, 2014) (Felton et al., 2015) (Fisher & Hemanth, 2015) (Rocco et al., 2012) (Stew, 2011) (Tarrasch, 2014)	(Newsome et al., 2006)	(Christopher et al., 2011) (Cigolla & Brown, 2011) (Dauenhauer, 2006) (Gill et al., 2015) (McCollum & Gehart, 2010) (Simon et al., 2009)
Schutte Self Report Emotional Intelligence Test		(Burnett & Pettijohn, 2015)	
Self-regulation questionnaire	(Bond et al., 2013)		

Note. Authors in bold denote RCT studies

Finally, the impact of mindfulness was not limited to the wellbeing of HCPs, but also was associated with enhanced job performance. The dominant outcome in this respect was compassion and/or empathy, as shown in table 15. Of the 28 intervention studies examining this, 16 observed an improvement while nine reported no significant change, showing an overall effect size between small and medium ( $d=.31$ ); meanwhile, three non-intervention

studies observed a correlation with mindfulness. Mindfulness was also associated with a broad range of other aspects of job performance, as shown in table 16. Of the seven intervention studies examining outcomes in this area, six observed an improvement and only one found no change, with a large global effect size ( $d=.82$ ). Six non-intervention studies also observed a correlation with mindfulness.

Table 15.

*Compassion & Empathy Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Empathy construct rating scale	<b>(Shapiro et al., 1998a)</b>		
Interpersonal reactivity index	(Hopkins & Proeve, 2013)	(Beddoe & Murphy, 2004) (Galantino et al., 2005)	(Keane, 2014)
Jefferson scale of physician empathy	(Barbosa et al., 2013) (Bazarko et al., 2013) (Krasner et al., 2009)	(Bond et al., 2013) <b>(West et al., 2014)</b>	
Jefferson scale of physician empathy [compassion]	<b>(Martín-Asuero et al., 2014)</b>		
Neff compassion scale		(Moore, 2008)	
Professional quality of life scale [compassion]		(Horner et al., 2014)	
Qualitative interviews	(Dorian & Killebrew, 2014) (Felton et al., 2015) (Hemanth & Fisher, 2015)		(Talisman et al., 2015)
Santa Clara brief compassion scale		(Brooker et al., 2013) (Fortney et al., 2013)	
Self-compassion scale	(Bazarko et al., 2013) (Bond et al., 2013) <b>(Erogul et al., 2014)</b> (Newsome et al., 2012) (Raab et al., 2015) (Rimes & Wingrove, 2011) <b>(Shapiro et al., 2005)</b> (Shapiro et al., 2007)	(Brooker et al., 2013) (Gauthier et al., 2015)	(Kemper et al., 2015)

Note. Authors in bold denote RCT studies

Table 16.

*Job Performance Outcomes across all Studies*

Measure	Improvement (positive change) related to mindfulness intervention	No change in relation to mindfulness intervention	Association (benign) with mindfulness
Caring efficiency scale		<b>(Pipe et al., 2009)</b>	
Counsellor activity self-efficacy scale	(Gockel et al., 2013)		
Patient distress [SC-90-R]	(Grepmaier et al., 2007)		
Qualitative interviews	(Hemanth & Fisher, 2015)		(Christopher et al., 2011) (Cigolla & Brown, 2011) (Gill et al., 2015) (McCollum & Gehart, 2010) (Simon et al., 2009) (Talisman et al., 2015)

Restraint of patients	(Brooker et al., 2014) (Singh et al., 2015)
Seclusion of patients	(Brooker et al., 2014)
Treatment team functioning checklist	(Singh et al., 2006)
Verbal redirection [disciplining patients]	(Singh et al., 2015)

---

*Note.* Authors in bold denote RCT studies

## **Summary and Recommendations**

Overall, MBIs had a positive impact upon most outcome measures (although some outcomes were rather equivocal, such as burnout). Moreover, a fairly large evidence-base regarding the use of mindfulness in healthcare settings is gradually accumulating, with 81 papers included in the current review, comprising a total of 3,805 participants. Together, these studies suggest mindfulness can potentially reduce mental health issues (e.g., stress), enhance wellbeing-related outcomes (e.g., job satisfaction), and improve aspects of job performance. These outcomes appear to be fairly evenly distributed across different healthcare professions. For instance, one might speculate that occupations which potentially have greater familiarity with psychological interventions like mindfulness, such as those in the mental health arena, might be more amenable to its effects. However, that appears to not be the case. Of the 81 papers analysed here, 32 (39%) specifically involved people working in mental health. These percentages were roughly reflected in the patterns of findings with respect to the various outcomes. For instance, in terms of anxiety, mental health professionals were involved in three of the nine interventions that reported a significant improvement, and two of the seven that found no such improvement (including one that found a worsening impact). Thus, it appears that mindfulness might be helpful to HCPs generally, regardless of their particular occupational role.

However, there are a number of issues with the research which limits the conclusions that can be drawn. In terms of the QATQS quality assessment, few studies scored highly in all respects (as shown in supplementary table 1). For instance, of the 66 intervention studies, only 26 (39%) involved a control group, while just 20 (30%) conducted an RCT. Without a



control group, it is harder to ascribe any positive changes observed to mindfulness per se. Then, even when controls are included, unless participants are randomised into groups, it is possible that differences in baseline characteristics between the groups generated interaction effects, thereby compromising the results. For example, in Barbosa et al. (2016), the 16 participants who entered the experimental group (reduced to 13 on attrition) did so after an invitation email was sent to the entire student population (of around 1300); by contrast, the control group consisted of individuals who were subsequently selected as matching the composition of the experimental group, and were paid to take part. Thus, it is conceivable – and indeed likely – that the experimental participants already had an interest in mindfulness, although whether they did so was not reported by the study. Furthermore, there were baseline differences in anxiety, with moderate levels among the experimental group (which also perhaps accounts for their interest in participating), compared to mild levels in the control group. Such factors complicate the assessment of the efficacy of MBIs, which is why RCT designs are generally preferable (although one acknowledges that not all researchers may be in a position to implement such a design). Further issues include heterogeneity with respect to both the type of MBI and the outcome measures looked at (which makes it difficult to conduct comparative or meta-analytic assessments, and hence to draw robust conclusions about the research as a whole), and the fact that the research is currently biased towards psychiatric outcome measures (e.g., anxiety, stress, depression), with little attention exploring other outcomes relevant to the work arena, such as work engagement or creativity.

Based on these critiques, the following recommendations can be made vis-à-vis future work in this area. First, where possible, studies should implement an RCT design, ideally with large numbers of participants (determined by a *priori* power calculations drawing on estimated effect size). Second, in addition to a wait-list control protocol, it would be useful if trials included an ‘active’ control group (e.g., an exercise programme). This will better enable

any positive effects to be ascribed to mindfulness per se (i.e., rather than people simply being involved in an absorbing group activity). Third, it would be good to see a diversification of outcome measures, with studies looking beyond ‘negative’ psychiatric issues (e.g., depression and anxiety), and also focusing on more ‘positive’ (i.e., non-clinical) outcomes such as work engagement, social capital, and creativity. Finally, where possible, trials should involve established MBIs (rather than bespoke adaptations), to better enable comparison across studies. However, there is also a need to move beyond MBIs developed for clinical contexts (e.g., MBSR), and to explore MBIs created specifically for the workplace. Nevertheless, despite the issues with the current research base, the evidence of the value of mindfulness for HCPs is strong, and one might speculate that this will only strengthen over the years ahead.

## References

- Aggs, C., & Bambling, M. (2010). Teaching mindfulness to psychotherapists in clinical practice: The mindful therapy programme. *Counselling and Psychotherapy Research*, 10(4), 278-286. doi: 10.1080/14733145.2010.485690
- Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review*, 30(2), 217-237. doi: <http://dx.doi.org/10.1016/j.cpr.2009.11.004>
- Baer, R. A., Smith, G. T., & Allen, K. B. (2004). Assessment of mindfulness by self-report the Kentucky inventory of mindfulness skills. *Assessment*, 11(3), 191-206.
- Baer, R. A., Smith, G. T., Hopkins, J., Krietemeyer, J., & Toney, L. (2006). Using self-report assessment methods to explore facets of mindfulness. *Assessment*, 13(1), 27-45.
- Barbosa, P., Raymond, G., Zlotnick, C., Wilk, J., Toomey, I. R., & Mitchell, I. J. (2013). Mindfulness-based stress reduction training is associated with greater empathy and

- reduced anxiety for graduate healthcare students. *Education for Health: Change in Learning and Practice*, 26(1), 9-14. doi: 10.4103/1357-6283.112794
- Bazarko, D., Cate, R. A., Azocar, F., & Kreitzer, M. J. (2013). The impact of an innovative mindfulness-based stress reduction program on the health and well-being of nurses employed in a corporate setting. *Journal of workplace behavioral health*, 28(2), 107-133.
- Beckman, H. B., Wendland, M., Mooney, C., Krasner, M. S., Quill, T. E., Suchman, A. L., & Epstein, R. M. (2012). The impact of a program in mindful communication on primary care physicians. *Academic Medicine*, 87(6), 815-819. doi: 10.1097/ACM.0b013e318253d3b2
- Beddoe, A. E., & Murphy, S. O. (2004). Does mindfulness decrease stress and foster empathy among nursing students? *Journal of Nursing Education*, 43(7), 305-312.
- Bond, A. R., Mason, H. F., Lemaster, C. M., Shaw, S. W., Mullin, C. S., Holick, E. A., & Saper, R. B. (2013). Embodied health: The effects of a mind-body course for medical students. *Medical Education Online*, 18(1). doi: 10.3402/meo.v18i0.20699
- Bonifas, R. P., & Napoli, M. (2014). Mindfully Increasing Quality of Life: A Promising Curriculum for MSW Students. *Social Work Education*, 33(4), 469-484. doi: 10.1080/02615479.2013.838215
- Botha, E., Gwin, T., & Purpora, C. (2015). The effectiveness of mindfulness based programs in reducing stress experienced by nurses in adult hospital settings: a systematic review of quantitative evidence protocol. *JBIR database of systematic reviews and implementation reports*, 13(10), 21-29. doi: 10.11124/jbisrir-2015-2380
- Brady, S., O'Connor, N., Burgermeister, D., & Hanson, P. (2012). The Impact of Mindfulness Meditation in Promoting a Culture of Safety on an Acute Psychiatric Unit.

*Perspectives in Psychiatric Care*, 48(3), 129-137. doi: 10.1111/j.1744-

6163.2011.00315.x

Bright, R. P., & Krahn, L. (2011). Depression and suicide among physicians. *Current Psychiatry*, 10(4), 16.

Brooker, J. E., Julian, J., Webber, L., Chan, J., Shawyer, F., & Meadows, G. (2013).

Evaluation of an Occupational Mindfulness Program for Staff Employed in the Disability Sector in Australia. *Mindfulness*, 4(2), 122-136. doi: 10.1007/s12671-012-0112-7

Brooker, J. E., Webber, L., Julian, J., Shawyer, F., Graham, A. L., Chan, J., & Meadows, G.

(2014). Mindfulness-based Training Shows Promise in Assisting Staff to Reduce Their Use of Restrictive Interventions in Residential Services. *Mindfulness*, 5(5), 598-603. doi: 10.1007/s12671-014-0306-2

Brooks, S. K., Gerada, C., & Chalder, T. (2011). Review of literature on the mental health of doctors: Are specialist services needed? *Journal of Mental Health*, 20(2), 146-156.

doi: 10.3109/09638237.2010.541300

Brown, K. W., & Ryan, R. M. (2003). The benefits of being present: Mindfulness and its role in psychological well-being. *Journal of Personality and Social Psychology*, 84(4), 822-848.

Burnett, M., & Pettijohn, C. (2015). Investigating the efficacy of mind-body therapies and emotional intelligence on worker stress in an organizational setting: An experimental approach. *Journal of Organizational Culture, Communications and Conflict*, 19(1), 146-158.

Burton, A., Burgess, C., Dean, S., Koutsopoulou, G. Z., & Hugh-Jones, S. (2017). How effective are mindfulness-based interventions for reducing stress among healthcare professionals? A systematic review and meta-analysis. *Stress and Health*, 33, 3-13.

- Caplan, R. P. (1994). Stress, anxiety, and depression in hospital consultants, general practitioners, and senior health service managers. *BMJ*, 309(6964), 1261-1263. doi: 10.1136/bmj.309.6964.1261
- Chambers, R., Gullone, E., & Allen, N. B. (2009). Mindful emotion regulation: An integrative review. *Clinical Psychology Review*, 29(6), 560-572. doi: 10.1016/j.cpr.2009.06.005
- Chiesa, A., Calati, R., & Serretti, A. (2011). Does mindfulness training improve cognitive abilities? A systematic review of neuropsychological findings. *Clin Psychol Rev*, 31(3), 449-464. doi: 10.1016/j.cpr.2010.11.003
- Choi, J. I., & Koh, M. S. (2015). Relations of job stress, burnout, mindfulness and job satisfaction of clinical nurses. *International Journal of Bio-Science and Bio-Technology*, 7(3), 121-128. doi: 10.14257/ijbsbt.2015.7.3.12
- Christopher, J. C., Chrisman, J. A., Trotter-Mathison, M. J., Schure, M. B., Dahlen, P., & Christopher, S. B. (2011). Perceptions of the long-term influence of mindfulness training on counselors and psychotherapists: A qualitative inquiry. *Journal of Humanistic Psychology*, 51(3), 318-349. doi: 10.1177/0022167810381471
- Christopher, J. C., Christopher, S. E., Dunnagan, T., & Schure, M. (2006). Teaching self-care through mindfulness practices: The application of yoga, meditation, and qigong to counselor training. *Journal of Humanistic Psychology*, 46(4), 494-509. doi: 10.1177/0022167806290215
- Cigolla, F., & Brown, D. (2011). A way of being: Bringing mindfulness into individual therapy. *Psychotherapy Research*, 21(6), 709-721. doi: 10.1080/10503307.2011.613076

- Cohen-Katz, J., Wiley, S., Capuano, T., Baker, D. M., Deitrick, L., & Shapiro, S. (2005). The effects of mindfulness-based stress reduction on nurse stress and burnout: a qualitative and quantitative study, part III. *Holistic Nursing Practice, 19*(2), 78-86.
- Cohen-Katz, J., Wiley, S. D., Capuano, T., Baker, D. M., Kimmel, S., & Shapiro, S. (2005). The effects of mindfulness-based stress reduction on nurse stress and burnout, Part II: A quantitative and qualitative study. *Holistic Nursing Practice, 19*(1), 26-35.
- Cohen, J. S., & Miller, L. J. (2009). Interpersonal mindfulness training for well-being: A pilot study with psychology graduate students. *Teachers College Record, 111*(12), 2760-2774.
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior, 38*, 385-396.
- Dauenhauer, J. A. (2006). Mindfulness theory and professional family caregivers in long-term care facilities. *Journal of Aging Studies, 20*(4), 351-365. doi: 10.1016/j.jaging.2005.11.003
- De Vibe, M., Solhaug, I., Tyssen, R., Friberg, O., Rosenvinge, J. H., Sørli, T., & Bjørndal, A. (2013). Mindfulness training for stress management: A randomised controlled study of medical and psychology students. *BMC Medical Education, 13*(1). doi: 10.1186/1472-6920-13-107
- de Zoysa, N., Ruths, F. A., Walsh, J., & Hutton, J. (2014). Mindfulness Based Cognitive Therapy for Mental Health Professionals: A Long-Term Qualitative Follow-up Study. *Mindfulness, 5*(1), 10-17. doi: 10.1007/s12671-012-0141-2
- Di Benedetto, M., & Swadling, M. (2014). Burnout in Australian psychologists: Correlations with work-setting, mindfulness and self-care behaviours. *Psychology, Health and Medicine, 19*(6), 705-715. doi: 10.1080/13548506.2013.861602

- Dobie, A., Tucker, A., Ferrari, M., & Rogers, J. M. (2015). Preliminary evaluation of a brief mindfulness-based stress reduction intervention for mental health professionals. *Australasian Psychiatry*. doi: 10.1177/1039856215618524
- Dorian, M., & Killebrew, J. E. (2014). A Study of Mindfulness and Self-Care: A Path to Self-Compassion for Female Therapists in Training. *Women and Therapy*, 37(1-2), 155-163. doi: 10.1080/02703149.2014.850345
- Draper, B., Kőlves, K., De Leo, D., & Snowden, J. (2014). The impact of patient suicide and sudden death on health care professionals. *General Hospital Psychiatry*, 36(6), 721-725. doi: <http://dx.doi.org/10.1016/j.genhosppsych.2014.09.011>
- Duchemin, A.-M., Steinberg, B. A., Marks, D. R., Vanover, K., & Klatt, M. (2015). A Small Randomized Pilot Study of a Workplace Mindfulness-Based Intervention for Surgical Intensive Care Unit Personnel: Effects on Salivary  $\alpha$ -Amylase Levels. *Journal of Occupational and Environmental Medicine*, 57(4), 393-399. doi: 10.1097/jom.0000000000000371
- Duchemin, A.-M., Steinberg, B. A., Marks, D. R., Vanover, K., & Klatt, M. (2015). A Small Randomized Pilot Study of a Workplace Mindfulness-Based Intervention for Surgical Intensive Care Unit Personnel: Effects on Salivary  $\alpha$ -Amylase Levels. *Journal of Occupational and Environmental Medicine*. doi: 10.1097/JOM.0000000000000371
- Dudman, J., Isaac, A., & Johnson, S. (2015, 10 June 2015). Revealed: How the stress of working in public services is taking its toll on staff, *The Guardian*. Retrieved from <http://www.theguardian.com/society/2015/jun/10/stress-working-public-services-survey>
- Erogul, M., Singer, G., McIntyre, T., & Stefanov, D. G. (2014). Abridged Mindfulness Intervention to Support Wellness in First-Year Medical Students. *Teaching and Learning in Medicine*, 26(4), 350-356. doi: 10.1080/10401334.2014.945025

- Felton, T. M., Coates, L., & Christopher, J. C. (2015). Impact of Mindfulness Training on Counseling Students' Perceptions of Stress. *Mindfulness*, 6(2), 159-169. doi: 10.1007/s12671-013-0240-8
- Firth-Cozens, J. (2003). Doctors, their wellbeing, and their stress : It's time to be proactive about stress—and prevent it. *BMJ : British Medical Journal*, 326(7391), 670-671.
- Fisher, P., & Hemanth, P. (2015). The development, facilitation and initial evaluation of a mindfulness group for a clinical psychology training course. *Clinical Psychology Forum*, 2015(266), 12-16.
- Fortney, L., Luchterhand, C., Zakletskaia, L., Zgierska, A., & Rakel, D. (2013). Abbreviated mindfulness intervention for job satisfaction, quality of life, and compassion in primary care clinicians: A pilot study. *Annals of Family Medicine*, 11(5), 412-420. doi: 10.1370/afm.1511
- Foureur, M., Besley, K., Burton, G., Yu, N., & Crisp, J. (2013). Enhancing the resilience of nurses and midwives: Pilot of a mindfulness-based program for increased health, sense of coherence and decreased depression, anxiety and stress. *Contemporary nurse*, 45(1), 114-125.
- Frank, J. L., Reibel, D., Broderick, P., Cantrell, T., & Metz, S. (2015). The Effectiveness of Mindfulness-Based Stress Reduction on Educator Stress and Well-Being: Results from a Pilot Study. *Mindfulness*, 6(2), 208-216. doi: 10.1007/s12671-013-0246-2
- Fresco, D. M., Moore, M. T., van Dulmen, M. H. M., Segal, Z. V., Ma, S. H., Teasdale, J. D., & Williams, J. M. G. (2007). Initial psychometric properties of the experiences questionnaire: Validation of a self-report measure of decentering. *Behavior Therapy*, 38(3), 234-246. doi: 10.1016/j.beth.2006.08.003
- Galantino, M. L., Baime, M., Maguire, M., Szapary, P. O., & Farrar, J. T. (2005). Association of psychological and physiological measures of stress in health-care professionals



- during an 8-week mindfulness meditation program: Mindfulness in practice. *Stress and Health*, 21(4), 255-261. doi: 10.1002/smi.1062
- Gao, Y.-Q., Pan, B.-C., Sun, W., Wu, H., Wang, J.-N., & Wang, L. (2012). Anxiety symptoms among Chinese nurses and the associated factors: a cross sectional study. *BMC Psychiatry*, 12(1), 141.
- Gauthier, T., Meyer, R. M. L., Greife, D., & Gold, J. I. (2015). An On-the-Job Mindfulness-based Intervention For Pediatric ICU Nurses: A Pilot. *Journal of pediatric nursing*, 30(2), 402-409. doi: 10.1016/j.pedn.2014.10.005
- Gill, M., Waltz, J., Suhrbier, P., & Robert, L. (2015). Non-duality and the Integration of Mindfulness into Psychotherapy: Qualitative Research with Meditating Therapists. *Mindfulness*, 6(4), 708-722. doi: 10.1007/s12671-014-0310-6
- Givens, J. L., & Tjia, J. (2002). Depressed Medical Students' Use of Mental Health Services and Barriers to Use. *Academic Medicine*, 77(9), 918-921.
- Gockel, A., Burton, D., James, S., & Bryer, E. (2013). Introducing Mindfulness as a Self-Care and Clinical Training Strategy for Beginning Social Work Students. *Mindfulness*, 4(4), 343-353. doi: 10.1007/s12671-012-0134-1
- Goodman, M. J., & Schorling, J. B. (2012). A mindfulness course decreases burnout and improves well-being among healthcare providers. *International journal of psychiatry in medicine*, 43(2), 119-128. doi: 10.2190/PM.43.2.b
- Grepmaier, L., Mitterlehner, F., Loew, T., & Nickel, M. (2007). Promotion of mindfulness in psychotherapists in training: Preliminary study. *European Psychiatry*, 22(8), 485-489. doi: 10.1016/j.eurpsy.2007.02.004
- Hallman, I. S., O'Connor, N., Hasenau, S., & Brady, S. (2014). Improving the culture of safety on a high-acuity inpatient child/adolescent psychiatric unit by mindfulness-

- based stress reduction training of staff. *Journal of Child and Adolescent Psychiatric Nursing*, 27(4), 183-189. doi: 10.1111/jcap.12091
- Hemanth, P., & Fisher, P. (2015). Clinical Psychology Trainees' Experiences of Mindfulness: an Interpretive Phenomenological Analysis. *Mindfulness*, 6(5), 1143-1152. doi: 10.1007/s12671-014-0365-4
- Hopkins, A., & Proeve, M. (2013). Teaching mindfulness-based cognitive therapy to trainee psychologists: Qualitative and quantitative effects. *Counselling Psychology Quarterly*, 26(2), 115-130. doi: 10.1080/09515070.2013.792998
- Horner, J. K., Piercy, B. S., Eure, L., & Woodard, E. K. (2014). A pilot study to evaluate mindfulness as a strategy to improve inpatient nurse and patient experiences. *Appl Nurs Res*, 27(3), 198-201. doi: 10.1016/j.apnr.2014.01.003
- Johnson, J. R., Emmons, H. C., Rivard, R. L., Griffin, K. H., & Dusek, J. A. (2015). Resilience Training: A Pilot Study of a Mindfulness-Based Program with Depressed Healthcare Professionals. *EXPLORE: The Journal of Science and Healing*, 11(6), 433-444. doi: 10.1016/j.explore.2015.08.002
- Kabat-Zinn, J. (1982). An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: Theoretical considerations and preliminary results. *General Hospital Psychiatry*, 4(1), 33-47. doi: [http://dx.doi.org/10.1016/0163-8343\(82\)90026-3](http://dx.doi.org/10.1016/0163-8343(82)90026-3)
- Kabat-Zinn, J. (2003). Mindfulness-based interventions in context: Past, present, and future. *Clinical Psychology: Science and Practice*, 10(2), 144-156. doi: 10.1093/clipsy.bpg016
- Keane, A. (2014). The Influence of Therapist Mindfulness Practice on Psychotherapeutic Work: A Mixed-Methods Study. *Mindfulness*, 5(6), 689-703. doi: 10.1007/s12671-013-0223-9

- Kemper, K. J., & Khirallah, M. (2015). Acute Effects of Online Mind–Body Skills Training on Resilience, Mindfulness, and Empathy. *Journal of Evidence-Based Complementary and Alternative Medicine*, 20(4), 247-253. doi: 10.1177/2156587215575816
- Kemper, K. J., Mo, X., & Khayat, R. (2015). Are mindfulness and self-compassion associated with sleep and resilience in health professionals? *Journal of Alternative and Complementary Medicine*, 21(8), 496-503. doi: 10.1089/acm.2014.0281
- Khamisa, N., Oldenburg, B., Peltzer, K., & Ilic, D. (2015). Work related stress, burnout, job satisfaction and general health of nurses. *International journal of environmental research and public health*, 12(1), 652-666.
- Klatt, M. D., Steinberg, B., & Duchemin, A. M. (2015). Mindfulness in motion (Mim): An onsite mindfulness based intervention (mbi) for chronically high stress work environments to increase resiliency and work engagement. *Journal of Visualized Experiments*, 2015(101), 1-11. doi: 10.3791/52359
- Krasner, M. S., Epstein, R. M., Beckman, H., Suchman, A. L., Chapman, B., Mooney, C. J., & Quill, T. E. (2009). Association of an educational program in mindful communication with burnout, empathy, and attitudes among primary care physicians. *JAMA - Journal of the American Medical Association*, 302(12), 1284-1293. doi: 10.1001/jama.2009.1384
- Kuoppala, J., & Kekoni, J. (2013). At the sources of one's well-being: Early rehabilitation for employees with symptoms of distress. *Journal of Occupational and Environmental Medicine*, 55(7), 817-823. doi: 10.1097/JOM.0b013e31828dc930
- Lamothe, M., Rondeau, É., Malboeuf-Hurtubise, C., Duval, M., & Sultan, S. (2016). Outcomes of MBSR or MBSR-based interventions in health care providers: A systematic review with a focus on empathy and emotional competencies. *Complementary Therapies in Medicine*, 24, 19-28.

- Lau, M. A., Bishop, S. R., Segal, Z. V., Buis, T., Anderson, N. D., Carlson, L. . . . , Carmody, J. (2006). The Toronto mindfulness scale: Development and validation. *Journal of Clinical Psychology*, 62(12), 1445–1467.
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour Research and Therapy*, 33(3), 335-343. doi: [http://dx.doi.org/10.1016/0005-7967\(94\)00075-U](http://dx.doi.org/10.1016/0005-7967(94)00075-U)
- Lutz, A., Slagter, H. A., Dunne, J. D., & Davidson, R. J. (2008). Attention regulation and monitoring in meditation. *Trends in Cognitive Sciences*, 12(4), 163-169. doi: 10.1016/j.tics.2008.01.005
- Mackenzie, C. S., Poulin, P. A., & Seidman-Carlson, R. (2006). A brief mindfulness-based stress reduction intervention for nurses and nurse aides. *Applied Nursing Research*, 19(2), 105-109. doi: 10.1016/j.apnr.2005.08.002
- Manotas, M., Segura, C., Eraso, M., Oggins, J., & McGovern, K. (2014). Association of brief mindfulness training with reductions in perceived stress and distress in Colombian health care professionals. *International Journal of Stress Management*, 21(2), 207-225. doi: 10.1037/a0035150
- Martín-Asuero, A., & García-Banda, G. (2010). The mindfulness-based stress reduction program (MBSR) reduces stress-related psychological distress in healthcare professionals. *Spanish Journal of Psychology*, 13(2), 897-905.
- Martín-Asuero, A., Queraltó, J. M., Pujol-Ribera, E., Berenguera, A., Rodríguez-Blanco, T., & Epstein, R. M. (2014). Effectiveness of a mindfulness education program in primary health care professionals: A pragmatic controlled trial. *Journal of Continuing Education in the Health Professions*, 34(1), 4-12. doi: 10.1002/chp.21211

- Maslach, C., Jackson, S., & Leiter, M. (1986). *Maslach Burnout Inventory Manual*. Palo Alto, CA: Consult. Psychol. Press.
- McCollum, E. E., & Gehart, D. R. (2010). Using mindfulness meditation to teach beginning therapists therapeutic presence: A qualitative study. *Journal of Marital and Family Therapy*, 36(3), 347-360. doi: 10.1111/j.1752-0606.2010.00214.x
- McConachie, D. A. J., McKenzie, K., Morris, P. G., & Walley, R. M. (2014). Acceptance and mindfulness-based stress management for support staff caring for individuals with intellectual disabilities. *Research in Developmental Disabilities*, 35(6), 1216-1227. doi: 10.1016/j.ridd.2014.03.005
- McCracken, L. M., & Yang, S. Y. (2008). A Contextual Cognitive-Behavioral Analysis of Rehabilitation Workers' Health and Well-Being: Influences of Acceptance, Mindfulness, and Values-Based Action. *Rehabilitation Psychology*, 53(4), 479-485. doi: 10.1037/a0012854
- Mealer, M., Conrad, D., Evans, J., Jooste, K., Solyntjes, J., Rothbaum, B., & Moss, M. (2014). Feasibility and acceptability of a resilience training program for intensive care unit nurses. *American Journal of Critical Care*, 23(6), e97-e105. doi: 10.4037/ajcc2014747
- Mirsky, A., Anthony, B., Duncan, C., Ahearn, M., & Kellam, S. (1991). Analysis of the elements of attention: A neuropsychological approach. *Neuropsychology Review*, 2(2), 109-145. doi: 10.1007/BF01109051
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine*, 6(7), e1000097. doi: doi:10.1371/journal.pmed.1000097
- Moody, K., Kramer, D., Santizo, R. O., Magro, L., Wyshogrod, D., Ambrosio, J., . . . Stein, J. (2013). Helping the Helpers: Mindfulness Training for Burnout in Pediatric

- Oncology-A Pilot Program. *Journal of pediatric oncology nursing*, 30(5), 275-284.  
doi: 10.1177/1043454213504497
- Moore, P. (2008). Introducing mindfulness to clinical psychologists in training: An experiential course of brief exercises. *Journal of Clinical Psychology in Medical Settings*, 15(4), 331-337. doi: 10.1007/s10880-008-9134-7
- Mossialos, E., Wenzl, M., Osborn, R., & Anderson, C. (2015). International Profiles of Health Care Systems: 2014. Washington: The Commonwealth Fund.
- Murray, M., Murray, L., & Donnelly, M. (2016). Systematic review of interventions to improve the psychological well-being of general practitioners. *BMC Family Practice*, 17(1), 36. doi: 10.1186/s12875-016-0431-1
- National Collaborating Centre for Methods and Tools (2008). *Quality Assessment Tool for Quantitative Studies (QATQS)*. Hamilton, ON: McMaster University.
- Newsome, S., Christopher, J. C., Dahlen, P., & Christopher, S. (2006). Teaching counselors self-care through mindfulness practices. *Teachers College Record*, 108(9), 1881-1900. doi: 10.1111/j.1467-9620.2006.00766.x
- Newsome, S., Waldo, M., & Gruszka, C. (2012). Mindfulness Group Work: Preventing Stress and Increasing Self-Compassion Among Helping Professionals in Training. *Journal for Specialists in Group Work*, 37(4), 297-311. doi: 10.1080/01933922.2012.690832
- Noone, S. J., & Hastings, R. P. (2010). Using Acceptance and Mindfulness-Based Workshops with Support Staff Caring for Adults with Intellectual Disabilities. *Mindfulness*, 1(2), 67-73. doi: 10.1007/s12671-010-0007-4
- Pflugeisen, B. M., Drummond, D., Ebersole, D., Mundell, K., & Chen, D. (2015). Brief Video-Module Administered Mindfulness Program for Physicians: A Pilot Study. *Explore (NY)*. doi: 10.1016/j.explore.2015.10.005

- Pipe, T. B., Bortz, J. J., Dueck, A., Pendergast, D., Buchda, V., & Summers, J. (2009). Nurse leader mindfulness meditation program for stress management: a randomized controlled trial. *Journal of nursing administration*, 39(3), 130-137.
- Posner, M. I., & Petersen, S. E. (1990). The attention system of the human brain. *Annual Review of Neuroscience*, 13(1), 25-42. doi: doi:10.1146/annurev.ne.13.030190.000325
- Poulin, P. A., Mackenzie, C. S., Soloway, G., & Karayolas, E. (2008). Mindfulness training as an evidenced-based approach to reducing stress and promoting well-being among human services professionals. *International Journal of Health Promotion and Education*, 46(2), 72-80.
- Raab, K., Sogge, K., Parker, N., & Flament, M. F. (2015). Mindfulness-based stress reduction and self-compassion among mental healthcare professionals: a pilot study. *Mental Health, Religion and Culture*, 18(6), 503-512. doi: 10.1080/13674676.2015.1081588
- Raffone, A., & Srinivasan, N. (2010). The exploration of meditation in the neuroscience of attention and consciousness. *Cognitive Processing*, 11(1), 1-7. doi: 10.1007/s10339-009-0354-z
- Rasmussen, V., Turnell, A., Butow, P., Juraskova, I., Kirsten, L., Wiener, L., . . . on behalf of the, I. R. C. (2015). Burnout among psychosocial oncologists: an application and extension of the effort–reward imbalance model. *Psycho-Oncology*, n/a-n/a. doi: 10.1002/pon.3902
- Razzaque, R., Okoro, E., & Wood, L. (2015). Mindfulness in Clinician Therapeutic Relationships. *Mindfulness*, 6(2), 170-174. doi: 10.1007/s12671-013-0241-7
- Rimes, K. A., & Wingrove, J. (2011). Pilot study of mindfulness-based cognitive therapy for trainee clinical psychologists. *Behavioural and Cognitive Psychotherapy*, 39(2), 235-241. doi: 10.1017/S1352465810000731

- Rocco, S., Dempsey, S., & Hartman, D. (2012). Teaching calm abiding meditation to mental health workers: A descriptive account of valuing subjectivity. *Contemporary Buddhism, 13*(2), 193-211. doi: 10.1080/14639947.2012.716707
- Ruths, F. A., de Zoysa, N., Frearson, S. J., Hutton, J., Williams, J. M. G., & Walsh, J. (2013). Mindfulness-Based Cognitive Therapy for Mental Health Professionals-a Pilot Study. *Mindfulness, 4*(4), 289-295. doi: 10.1007/s12671-012-0127-0
- Ryan, A., Safran, J. D., Doran, J. M., & Muran, J. C. (2012). Therapist mindfulness, alliance and treatment outcome. *Psychotherapy Research, 22*(3), 289-297. doi: 10.1080/10503307.2011.650653
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-Based Cognitive Therapy for Depression: A New Approach to Preventing Relapse*. New York: Guilford Press.
- Shapiro, S. L., Astin, J. A., Bishop, S. R., & Cordova, M. (2005). Mindfulness-based stress reduction for health care professionals: Results from a randomized trial. *International Journal of Stress Management, 12*(2), 164-176. doi: 10.1037/1072-5245.12.2.164
- Shapiro, S. L., Brown, K. W., & Biegel, G. M. (2007). Teaching self-care to caregivers: Effects of mindfulness-based stress reduction on the mental health of therapists in training. *Training and Education in Professional Psychology, 1*(2), 105-115. doi: 10.1037/1931-3918.1.2.105
- Shapiro, S. L., Carlson, L. E., Astin, J. A., & Freedman, B. (2006). Mechanisms of mindfulness. *Journal of Clinical Psychology, 62*(3), 373-386. doi: 10.1002/jclp.20237
- Shapiro, S. L., Schwartz, G., & Bonner, G. (1998a). Effects of mindfulness-based stress reduction on medical and premedical students. *Journal of Behavioral Medicine, 21*(6), 581-599. doi: 10.1023/A:1018700829825



- Shapiro, S. L., Schwartz, G. E., & Bonner, G. (1998b). Effects of mindfulness-based stress reduction on medical and premedical students. *Journal of Behavioral Medicine*, 21(6), 581-599. doi: 10.1023/A:1018700829825
- Simon, S. T., Ramsenthaler, C., Bausewein, C., Krischke, N., & Geiss, G. (2009). Core attitudes of professionals in palliative care: a qualitative study. *International Journal of Palliative Nursing*, 15(8), 405-411.
- Singh, N. N., Lancioni, G. E., Karazsia, B. T., Myers, R. E., Winton, A. S. W., Latham, L. L., & Nugent, K. (2015). Effects of Training Staff in MBPBS on the Use of Physical Restraints, Staff Stress and Turnover, Staff and Peer Injuries, and Cost Effectiveness in Developmental Disabilities. *Mindfulness*, 6(4), 926-937. doi: 10.1007/s12671-014-0369-0
- Singh, N. N., Singh, S. D., Sabaawi, M., Myers, R. E., & Wahler, R. G. (2006). Enhancing treatment team process through mindfulness-based mentoring in an inpatient psychiatric hospital. *Behavior Modification*, 30(4), 423-441. doi: 10.1177/0145445504272971
- Sochos, A., Bowers, A., & Kinman, G. (2012). Work Stressors, Social Support, and Burnout in Junior Doctors: Exploring Direct and Indirect Pathways. *Journal of Employment Counseling*, 49(2), 62-73. doi: 10.1002/j.2161-1920.2012.00007.x
- Song, Y., & Lindquist, R. (2015). Effects of mindfulness-based stress reduction on depression, anxiety, stress and mindfulness in Korean nursing students. *Nurse Education Today*, 35(1), 86-90. doi: 10.1016/j.nedt.2014.06.010
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *Manual for the State-Trait Anxiety Inventory (Self-Evaluation Questionnaire)*. Palo Alto, CA: Consulting Psychologists Press.

- Stew, G. (2011). Mindfulness training for occupational therapy students. *British Journal of Occupational Therapy*, 74(6), 269-276. doi: 10.4276/030802211X13074383957869
- Talisman, N., Harazduk, N., Rush, C., Graves, K., & Haramati, A. (2015). The impact of mind-body medicine facilitation on affirming and enhancing professional identity in health care professions faculty. *Academic Medicine*, 90(6), 780-784. doi: 10.1097/ACM.0000000000000720
- Tarrasch, R. (2014). Mindfulness Meditation Training for Graduate Students in Educational Counseling and Special Education: A Qualitative Analysis. *Journal of Child and Family Studies*, 1-12. doi: 10.1007/s10826-014-9939-y
- The Health & Social Care Information Centre (2009). *Adult psychiatric morbidity in England, Results of a household survey*. London: The Health & Social Care Information Centre.
- Toppinen-Tanner, S., Ojajarvi, A., Väänäänen, A., Kalimo, R., & Jäppinen, P. (2005). Burnout as a Predictor of Medically Certified Sick-Leave Absences and Their Diagnosed Causes. *Behavioral Medicine*, 31(1), 18-32. doi: 10.3200/BMED.31.1.18-32
- Trowbridge, K., & Mische Lawson, L. (2016). Mindfulness-based interventions with social workers and the potential for enhanced patient-centered care: A systematic review of the literature. *Social Work in Health Care*, 55(2), 101-124. doi: 10.1080/00981389.2015.1094165
- Tyssen, R., Vaglum, P., Grønvold, N. T., & Ekeberg, Ø. (2000). The impact of job stress and working conditions on mental health problems among junior house officers. A nationwide Norwegian prospective cohort study. *Medical Education*, 34(5), 374-384. doi: 10.1046/j.1365-2923.2000.00540.x

- Van der Riet, P., Rossiter, R., Kirby, D., Dluzewska, T., & Harmon, C. (2015). Piloting a stress management and mindfulness program for undergraduate nursing students: Student feedback and lessons learned. *Nurse Education Today*, 35(1), 44-49. doi: 10.1016/j.nedt.2014.05.003
- Walsh, R., & Shapiro, S. L. (2006). The meeting of meditative disciplines and western psychology: A mutually enriching dialogue. *American Psychologist*, 61(3), 227-239. doi: 10.1037/0003-066X.61.3.227
- West, C. P., Dyrbye, L. N., Rabatin, J. T., Call, T. G., Davidson, J. H., Multari, A., . . . Shanafelt, T. D. (2014). Intervention to promote physician well-being, job satisfaction, and professionalism a randomized clinical trial. *JAMA internal medicine*, 174(4), 527-533. doi: 10.1001/jamainternmed.2013.14387
- Westphal, M., Bingisser, M. B., Feng, T., Wall, M., Blakley, E., Bingisser, R., & Kleim, B. (2015). Protective benefits of mindfulness in emergency room personnel. *Journal of Affective Disorders*, 175, 79-85. doi: 10.1016/j.jad.2014.12.038

Supplementary table 1.

*QATQS Scoring Assessment of Intervention Studies*

Authors	Selection bias	Design	Cofounders	blinding	Data collection	Attrition	Global
(Aggs & Bambling, 2010)	3	3	3	3	3	2	3
(Barbosa et al., 2013)	3	3	2	3	2	2	3
(Bazarko et al., 2013)	2	3	3	3	2	1	3
(Beckman et al., 2012)	Q	Q	Q	Q	Q	Q	Q
(Beddoe & Murphy, 2004)	3	3	3	3	2	2	3
(Bond et al., 2013)	3	3	3	3	2	2	3
(Bonifas & Napoli, 2014)	3	3	3	3	1	1	3
(Brady et al., 2012)	3	3	3	3	2	2	3
(Brooker et al., 2013)	3	3	3	3	1	2	3
(Brooker et al., 2014)	3	3	3	3	2	2	3
(Burnett & Pettijohn, 2015)	3	2	2	3	1	2	3
(Christopher et al., 2006)	Q	Q	Q	Q	Q	Q	Q
(J. S. Cohen & Miller, 2009)	3	3	3	3	1	2	3
(Cohen-Katz, Wiley, Capuano, Baker, Kimmel, et al., 2005)	2	2	2	3	1	1	2
(Cohen-Katz, Wiley, Capuano, Baker, Deitrick, et al., 2005)	Q	Q	Q	Q	Q	Q	Q
(Dobie et al., 2015)	3	3	3	3	1	1	3
(De Vibe et al., 2013)	2	1	1	1	1	1	1
(de Zoysa et al., 2014)	Q	Q	Q	Q	Q	Q	Q
(Dorian & Killebrew, 2014)	Q	Q	Q	Q	Q	Q	Q
(Duchemin et al., 2015)	1	1	1	1	1	1	1
(Erogul et al., 2014)	1	1	1	1	1	2	1
(Felton et al., 2015)	Q	Q	Q	Q	Q	Q	Q
(Fisher & Hemanth, 2015)	Q	Q	Q	Q	Q	Q	Q
(Fortney et al., 2013)	3	3	3	3	1	1	3

(Foureur et al., 2013)	3	3	3	3	1	2	3
(Galantino et al., 2005)	3	3	2	3	1	3	3
(Gauthier et al., 2015)	3	3	3	3	1	2	3
(Goodman & Schorling, 2012)	2	3	3	3	1	3	3
Grepmaier et al., 2007)	2	2	1	2	1	1	2
(Hallman et al., 2014)	2	3	3	2	1	1	3
(Hemanth & Fisher, 2015)	Q	Q	Q	Q	Q	Q	Q
(Hopkins & Proeve, 2013)	3	3	3	3	1	2	3
(Horner et al., 2014)	3	3	3	3	1	3	3
(Johnson et al., 2015)	3	3	1	3	1	2	3
(Klatt et al., 2015)	3	3	3	3	1	2	3
(Kemper & Khirallah, 2015)	3	3	3	3	1	2	3
(Krasner et al., 2009)	2	3	3	3	1	2	3
(Kuoppala & Kekoni, 2013)	1	2	2	2	1	1	2
(Mackenzie et al., 2006)	3	2	2	3	1	2	3
(Manotas et al., 2014)	2	2	1	3	1	2	2
(Martín-Asuero & García-Banda, 2010)	2	3	3	3	1	1	3
(Martín-Asuero et al., 2014)	2	2	2	3	1	1	2
(McConachie et al., 2014)	2	1	1	3	1	2	2
(Mealer et al., 2014)	1	1	1	2	1	1	1
(Moody et al., 2013)	1	1	1	2	1	2	1
(Moore, 2008)	1	3	3	2	1	2	3
(Newsome et al., 2006)	1	3	3	3	3	3	3
(Newsome et al., 2012)	1	3	3	2	1	2	3
(Noone & Hastings, 2010)	1	3	3	1	1	3	3
(Pflugeisen et al., 2015)	1	3	3	2	2	1	3
(Pipe et al., 2009)	2	1	2	1	1	1	1
(Poulin et al., 2008)	1	2	1	1	1	3	2
(Raab et al., 2015)	1	3	3	2	1	2	3
(Rimes & Wingrove, 2011)	1	3	3	1	1	3	3
(Rocco et al., 2012)	Q	Q	Q	Q	Q	Q	Q

(Ruths et al., 2013)	2	3	3	1	1	1	3
(Shapiro et al., 1998b)	1	1	2	1	1	1	1
(Shapiro et al., 2005)	1	1	2	2	1	3	2
(Shapiro et al., 2007)	1	2	1	2	1	1	1
(Singh et al., 2015)	1	3	3	1	1	3	2
(Singh et al., 2006)	1	3	3	1	2	2	3
(Song & Lindquist, 2015)	1	1	2	1	1	1	1
(Stew, 2011)	Q	Q	Q	Q	Q	Q	Q
(Tarrasch, 2014)	Q	Q	Q	Q	Q	Q	Q
(Van der Riet et al., 2015)	Q	Q	Q	Q	Q	Q	Q
(West et al., 2014)	1	1	1	1	1	2	1

*Note.* Q = qualitative study.