



OPEN

A broad spectrum of genomic changes in latinamerican patients with EXT1 / EXT2-CDG

SUBJECT AREAS:
DISEASE GENETICS
CANCER GENOMICSReceived
7 January 2014Accepted
22 July 2014Published
18 September 2014Correspondence and
requests for materials
should be addressed to
C.G.A.
(asteggianocarla@
hotmail.com)M. A. Delgado¹, G. Martinez-Domenech¹, P. Sarrion², R. Urreiziti², L. Zecchini³, H. H. Robledo⁴, F. Segura⁵, R. Dodelson de Kremer¹, S. Balcells², D. Grinberg² & C. G. Asteggiano^{1,6,7}

¹Centro de Estudio de las Metabolopatías Congénitas (CEMECO), Facultad de Ciencias Médicas, Universidad Nacional de Córdoba, Hospital de Niños de la Santísima Trinidad, Córdoba, Argentina, ²Universitat de Barcelona, IBUB, Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Departament de Genètica, Facultat de Biologia, Barcelona, España, ³Servicio de Traumatología, Hospital de Niños de la Santísima Trinidad, Córdoba, Argentina, ⁴Servicio de Bioimágenes, Hospital de Niños de la Santísima Trinidad, Córdoba, Argentina, ⁵II^{da} Cátedra de Ortopedia y Traumatología, Facultad de Ciencias Médicas, Universidad Nacional de Córdoba, Argentina, ⁶Cátedra de Farmacología, Facultad de Medicina, Universidad Católica de Córdoba, Argentina, ⁷Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

Multiple osteochondromatosis (MO), or EXT1/EXT2-CDG, is an autosomal dominant O-linked glycosylation disorder characterized by the formation of multiple cartilage-capped tumors (osteochondromas). In contrast, solitary osteochondroma (SO) is a non-hereditary condition. EXT1 and EXT2, are tumor suppressor genes that encode glycosyltransferases involved in heparan sulfate elongation. We present the clinical and molecular analysis of 33 unrelated Latin American patients (27 MO and 6 SO). Sixty-three percent of all MO cases presented severe phenotype and two malignant transformations to chondrosarcoma (7%). We found the mutant allele in 78% of MO patients. Ten mutations were novel. The disease-causing mutations remained unknown in 22% of the MO patients and in all SO patients. No second mutational hit was detected in the DNA of the secondary chondrosarcoma from a patient who carried a nonsense EXT1 mutation. Neither EXT1 nor EXT2 protein could be detected in this sample. This is the first Latin American research program on EXT1/EXT2-CDG.

Multiple osteochondromatosis (MO; MIM# 133700, 133701), also known as EXT1/EXT2-CDG in the Congenital Disorder of Glycosylation (CDG) nomenclature^{1,2} is an autosomal dominant disease. MO is genetically heterogeneous and 70–90% of patients present mutations in one of two genes: EXT1 (MIM 608177) (8q24.11-q24.13)³ or EXT2 (MIM 608210) (11p12-p11)^{4,5}. Both are ubiquitously expressed tumor-suppressor genes of the EXT gene family. All members of this gene family have been cloned encode glycosyltransferases involved in the adhesion and/or polymerization of heparan sulfate chains (HS)^{6–10}.

Heparan sulfate proteoglycans (HSPG) are ubiquitously expressed at cell surfaces and in extracellular matrices. They are composed of a core protein and one or more heparan sulfate glycosaminoglycan chains (linear polysaccharides formed by alternating N-acetylated or N-sulfated glucosamine units and uronic acid) that interact with numerous proteins, including growth factors, morphogens and extracellular matrix proteins¹¹. Each HS binds to a serine unit of a proteoglycan core protein via O-linked-xylosylation binding^{11,12}. The truncated HSPG disturb specific growth-factor-binding in chondrocytes, resulting in abnormal signaling and altered endochondral ossification, thus leading to MO¹³.

MO is characterized by the formation of multiple cartilaginous tumors (osteochondromas), that mainly affect the metaphyses of long bones or the surface of flat bones^{14–18}. Complications may involve bone and surrounding tissue deformities, fractures or mechanical joint problems, vascular compression, arterial thrombosis, aneurysm, pseudoaneurysm formation, and venous thrombosis. Pain, acute ischemia, signs of phlebitis or nerve compression occur alongside the most severe complication, the malignant transformation of osteochondroma to secondary peripheral chondrosarcoma (0.5–5% of patients)^{16–23}. EXT1 and EXT2 have been analysed using different techniques to search for point mutations and structural alterations. Intragenic deletions involving single or multiple exons of EXT1 or EXT2 genes have been found in about 10% of cases^{24–29}. Additionally, the promoter of EXT1 was analysed in some cases. The EXT1 core promoter region was reported to map to approximately –917 bp upstream of the EXT1 start codon, within a 123-bp region³⁰. One SNP within this region, rs34016643,



was shown to have a significant effect on *EXT1* promoter activity (the C-allele resulting in a 56% rise in promoter activity) compared to the G-wild-type allele³⁰. The presence of an additional MO-causing gene has been proposed to explain the absence of an *EXT1* or *EXT2* mutation in a small percentage of MO patients (15–30%)^{17,31,32}. To date, more than 600 different *EXT1* and 345 *EXT2* mutations have been found worldwide and an update on all reported mutations is deposited at <http://medgen.ua.ac.be/LOVD>²⁰.

This study represents the first Latin American research program in MO, with a broad spectrum of genomic changes detected, including 10 novel pathogenic mutations identified in *EXT1/EXT2*-CDG patients. Twenty-one different mutant alleles in the *EXT1* or *EXT2* genes were found in a cohort of 27 MO patients, most of them with a severe phenotype, including two patients with malignant transformation to chondrosarcoma. No mutation was found in six MO patients after performing sequencing and MLPA analyses.

Results

Phenotypic characterization. We observed multiple osteochondromas in 27 out of 33 patients, who ranged from 3 to 55 years at diagnosis. Orthopedic deformities of the forearm, shortening of limbs, ankle, varus or valgus of the knee, short metacarpal bones, scoliosis, synostosis, arthritis, and vessel or nerve compression were some common manifestations. The lesions were located in the femur (71%), tibia (67%), humerus (67%), fibula (62%), radius (52%) and pelvis (29%), a frequent site of malignant transformation to chondrosarcoma. Phenotypic data were available for 78% of the MO patients (n = 21), of whom four presented with a moderate phenotype (15% of all MO patients) and 17 with a severe presentation of the disease (63% of all MO patients) (Table 1, Figure 1A). A severe phenotype ranging from grade IS to IVS was observed in most of the MO patients (Figure 1B). Seventy six percent of them presented an age of onset below 5 years and 59% manifested familial inheritance (Table 1). Two patients developed malignant transformation as a large chondrosarcoma on the pelvis that led to severe vascular and organ compression: P06, a 32 year-old female with a type IV severe phenotype, reported by Delgado et al. and P38, a 42 year-old male with a type IIS severe phenotype (Table 1)²².

Gene sequence and dose analyses of *EXT1* and *EXT2* exons. Exons and flanking regions of the *EXT1* and *EXT2* genes were sequenced from the genomic DNA of the 33 patients and MLPA analysis was performed in DNA samples of those with negative results for sequencing analysis. The mutant allele was found in 78% of the MO patients including one patient with solitary presentation (P36) (Table 1). We identified 21 pathogenic mutations, 15 in *EXT1* and 6 in *EXT2* (five nonsense, six frame-shift, four missense, three splice-site mutations, and three large deletions identified by MLPA) listed in Table 2. Six of the *EXT1* mutations were novel (p.Val78Glyfs*111, p.Leu264Pro, p.Lys306*, p.Arg346Thr, c.1164 + 1C > A, and p.Gln583Arg) as were four of the *EXT2* mutant alleles (p.Asp307-Valfs*45, p.Trp394*, p.Asp539Glnfs*5 and a deletion of exon 4 to 14).

Bioinformatic predictions for the *EXT1* missense mutations suggested a pathogenic role for these genomic changes. In particular, the p.Leu264Pro change was considered “*disease causing*” by Mutation Taster (score: 0.999, amino acid sequence changed, protein features might be affected with potential luminal loss and splice site changes) and “*probably damaging*” by PolyPhen2 (score: 0.997 sensitivity: 0.27; specificity: 0.98), while ESE Finder predicted an increased level of the enhancer splicing proteins SF2/ASF (score changed from −0.21685 to 1.24048), and SF2/ASF (score changed from 0.4979 to 1.75265). The novel missense mutation, p.Arg346Thr, change from a basic amino acid (Arg) to a non-polar one (Thr) and it was predicted to be “*disease causing*” by Mutation Taster (score 0.999, amino acid sequence changed, protein features might be affected with potential

luminal loss and splice site changes) and “*probably damaging*” by PolyPhen2 (score: 0.993, sensitivity: 0.47; specificity: 0.96). ESE Finder predicted diminished levels of enhancer splicing protein SRSF2 (SC35). The other novel *EXT1* missense mutation, p.Gln583Arg, is a change from an uncharged polar amionacid (Gln) to a basic one (Arg) and it was predicted to be “*disease causing*” by Mutation Taster (score: 0.999, amino acid sequence changed, protein features might be affected with potential luminal loss and splice site changes) although PolyPhen2 predicted it to be “*benign*” (score: 0.002 sensitivity: 0.99; specificity: 0.30). The Protein Homology Fold Recognition Engine Phyre2, (<http://www.sbg.bio.ic.ac.uk/phyre2/html>, last accessed March 2014) was used to predict the effect of missense mutations on 3D structure and the missense mutation p.Arg346Thr removes two fragments of alpha helix between aa 345 and 347, and from aa 635 to 639, and a beta sheet from aa 361 to 368 in *EXT1* protein. The p.Leu264Pro mutation adds an alpha helix structure from aa 161 to 166 and removes an alpha helix from aa 344 to 346, while removing a beta sheet structure from aa 360 to 365 and introducing a segment of beta sheet from aa 724 to 726. The other novel missense mutation, p.Gln583Arg removes two fragments of alpha helix between aa 39 and 41, and from aa 635 to 639 in the *EXT1* protein.

In silico analyses for one novel intronic mutation (c.1164 + 1 G > A) predicted the use of cryptic donor splice sites: Human Splice Finder (<http://www.umd.be/HSF/>, last accessed March 2014), considered the use of a cryptic donor splice site (score: 91.85%) located 74 nucleotides downstream from the wild-type sequence, while NetGene2 (<http://www.cbs.dtu.dk/cgi-bin/webface?jobid=netgene2/> last accessed March 2014) predicted the use of a cryptic donor splicing site (score: 0.76) 201 nucleotides downstream from the wild-type (score:0.83).

In three patients large deletions were detected by MLPA (Tables 1 and 2). In *EXT1* (exon 1, P36) and a deletion of 11 exons (6–16) in (P12), and the third one was a deletion in *EXT2* (exon 6, P04). Normal MLPA profiles were obtained for 19 patients.

No mutation was found in 12 cases (6 MO and 6 SO) after performing sequencing and MLPA analyses. Most of these patients did not have a positive family history of osteochondromatosis (Table 1).

Analysis of the *EXT1* promoter. We sequenced 435 bp upstream of the *EXT1* gene including the 123-bp region described to contain the basic promoter elements³⁰ in samples from patients and 9 controls, but no mutation was detected. We found that four patients (P18, P21, P34 and P41), and one control individual, were heterozygous carriers of the C allele of SNP rs34016643, which has been previously shown to have a significant effect on *EXT1* promoter activity, with the C-allele resulting in a 56% rise in promoter activity compared to the G (wild-type) allele³⁰. No pathogenic mutation was identified in *EXT1* or *EXT2* in three of these four patients, while patient P41 bore a nonsense mutation (c.1219C > T, p.Gln407*) in exon 4 of the *EXT1* gene (Table 1).

Loss of heterozygosity analysis in a chondrosarcoma. We had access to a chondrosarcoma sample from P06. We have detected the heterozygous p.L283* mutation in *EXT1* in the tumor sample. We further analysed both genes by MLPA and we did not detect any dose alteration in the chondrosarcoma from this patient. The patient was heterozygous for the single nucleotide polymorphism rs11546829 in exon 3 of the *EXT1* gene. Loss of heterozygosity for this marker was not observed in the analysis of DNA in the tumor tissue.

Discussion

This work represents the first clinical, biochemical and molecular research on multiple hereditary osteochondromatosis (*EXT1/EXT2*-CDG) in Latin American patients. Thirty-three unrelated patients



Table 1 | Overview of EXT1 and EXT2 mutations and the phenotype found in this cohort

Patient	Sex	Gene	DNA	Deduced protein change	EXT1 promoter SNP rs34016643		MLPA	Family History	Phenotype	Age of onset	Other clinical features
					WT	rs34016643					
P01	male	EXT2	c.1182G > A	p.Trp394*	WT	WT	NA	No	MO/IIIS	1,5 yo	Vertebral location
P02	fem	EXT1	c.1469delT	p.Leu490Argfs*9	WT	WT	NA	No	MO/IIIS	5 yo	Surgery/Sinostoses
P03	fem	ND	WT	WT	Normal	No	SO	5 yo	Exostoses in humerus
P04	male	EXT2	ex 6 del	Unknown	WT	WT	Abnormal	No	MO/IIIS	5 m	Shortening of limbs
P05	male	EXT1	c.752delT	p.Leu251*	WT	WT	NA	Yes	MO/IVS	4 yo	Surgery/Axis deviations (cubito and radius)
P06 ¹	fem	EXT1	c.848T > A	p.Leu283*	WT	WT	Normal	Yes	MO/IVS	12 yo	Chondrosarcoma /Surgery
P07	fem	ND	WT	WT	Normal	No	SO	6 yo	Surgery
P08	fem	EXT1	c.1037G > C	p.Arg346Thr	WT	WT	NA	No	MO/IIIS	3 yo	Scoliosis
P12	male	EXT2	ex 4-14del	Unknown	WT	WT	Abnormal	Yes	MO/IVS	2 m	Surgery/Scapular and ribs location.
P13	male	EXT2	c.920_929del10insTG	p.Asp307Valfs*45	WT	WT	NA	Yes	MO/IIIS	2 m	Abnormal karyotype (18 q deletion)
P14	fem	EXT1	c.369_370delAG	p.Lys126Asnfs*62	WT	WT	Normal	Yes	MO/IS	1 yo	Scapular osteochondromas
P15	male	EXT1	c.232insG	p.Val78Glyfs*111	WT	WT	NA	No	MO/M	8 yo	Deformity of the heel
P16	male	EXT1	c.916A > T	p.Lys306*	WT	WT	NA	No	MO/M	4 yo	Decreased bone density
P17	male	EXT1	c.791T > C	p.Leu264Pro	WT	WT	NA	Yes	MO/IIIS	1 m	Restricted joint motion
P18	male	ND	G/C	G/C	Normal	No	SO	14 yo	Ribs location
P19	male	EXT2	c.626 + 1G > A	-	WT	WT	Normal	Yes	MO/NA	8 yo	NA
P21	male	ND	G/C	G/C	Normal	No	MO/NA	9 yo	Ribs location
P24	male	ND	WT	WT	Normal	No	MO/IIIS	2 yo	Surgery/Shortening and deformities of limbs
P25	fem	EXT1	c.1748A > G	p.Gln583Arg	WT	WT	Normal	Yes	MO/IIIS	5 yo	Vertebral location
P26	fem	ND	WT	WT	Normal	No	MO/M	11 yo	Surgery/Scoliosis
P27	fem	EXT2	c.1616_1623del8ins10	p.Asp539Glnfs*5	WT	WT	Normal	No	MO/NA	2 m	Deformity of the hip
P28	male	EXT1	c.1164 + 1G > A	-	WT	WT	Normal	Yes	MO/NA	10 m	Surgery/Shortening and deformities of limbs
P29	fem	EXT1	c.1722 + 1G > A	-	WT	WT	Normal	No	MO/IVS	1 m	Scoliosis
P30	fem	ND	WT	WT	Normal	No	SO	4 yo	Restricted joint motion
P31	male	ND	WT	WT	Normal	No	SO	8 yo	Surgery/Bilateral valgus
P32	fem	ND	WT	WT	Normal	No	SO	9 yo	NA
P34	fem	ND	G/C	G/C	Normal	No	MO/IS	3 yo	Bilateral valgus, vertebral
P36	fem	EXT1	ex 1 del	Unknown	WT	WT	Abnormal	No	MO/M	10 yo	NA
P37	male	EXT1	c.248_249insA	p.Gln84Alafs*105	WT	WT	NA	No	MO/NA	5 m	Distrophy in ribs
P38 ¹	male	ND	WT	WT	Normal	Yes	MO/IIIS	10 yo	Chondrosarcoma /Severe vascular compression, phlebitis
P39	fem	ND	WT	WT	Normal	NA	MO/NA	12 yo	Deformity of ankles
P40	male	EXT1	c.1018C > T	p.Arg340Cys	WT	WT	NA	Yes	MO/IIIS	2 m	Scapular osteochondromas
P41	male	EXT1	c.1219C > T	p.Gln407*	G/C	G/C	NA	Yes	MO/IIIS	1 yo	Pelvic and Scapular Osteochondromas

Novel mutations are indicated in bold.

¹Patients with malignant transformations to chondrosarcoma.

(WT) wild type = G/G. (ND) No mutation detected by sequencing and MLPA analysis; (NA) Not Available; (S) Severe phenotype; (M) Mild phenotype; (MO) Multiple osteochondroma; (SO) Solitary osteochondroma.




Figure 1 | Genotype-phenotype association in MO patients (n = 27). (A) Graph showing the proportion of severe phenotype (blue), mild phenotype (red) and patients with phenotype not available (green) and the distribution of *EXT1* and *EXT2* mutations or no mutations identified (NM) within each category. (B) Grade of phenotype severity among severely affected patients and distribution of *EXT1* and *EXT2* mutations or no mutations identified (NM) within each category.

were studied, 27 of which presented with MO. The mutant allele was identified in 21 of these patients (78%). *EXT1* mutations (71%) were more common than *EXT2* mutations (29%) and most of the *EXT1* mutations were located in the first six exons. These results are consistent with recent studies reporting that *EXT1* is responsible for ~65–75% of MO cases^{20,27}.

The 67% of *EXT1* mutations (10/15), were located in exon 1 or 2, which encode the exostosin domain of the *EXT1* protein (from amino acid 111 to 396). Most of these patients presented a severe

phenotype (67%). Twenty per cent of *EXT1* mutations (n = 3) were located in the glycosyltransferase domain (from amino acid 480 to 729) (Table 2). In contrast, *EXT2* mutations (n = 6) were more frequent in the last exons. Thirty three per cent of the *EXT2* mutations (n = 2) were found in the exons that encode the glycosyltransferase domain (from exon 10 to 14) (Table 2). The structure of the different *EXT1* and *EXT2* protein domains was analyzed using Phyre2 to predict a decrease or loss of protein function according to the detected mutations and the altered structure of protein

Table 2 | List of mutations in *EXT1* or *EXT2* gene in MO patients

Gene	Patient	Exon-Intron	DNA	Deduced protein change	Mutation Type	Publication
<i>EXT1</i>	P37	Ex 1	c.248_249insA,	p.Gln84Alafs*105	Frameshift	Francannet, et al 2001
<i>EXT1</i>	P15	Ex 1	c.232dupG	p.Val78Glyfs*111	Frameshift	This study
<i>EXT1</i>	P14	Ex 1	c.369_370delAG	p.Lys126Asnfs*62	Frameshift	Ciavarella, et al, 2013
<i>EXT1</i>	P05	Ex 1	c.752delT	p.Leu251*	Nonsense	Ciavarella, et al, 2013
<i>EXT1</i>	P17	Ex 1	c.791T > C	p.Leu264Pro	Missense	This study
<i>EXT1</i>	P06 ¹	Ex 1	c.848T > A	p.Leu283*	Nonsense	Delgado, et al, 2012
<i>EXT1</i>	P16	Ex 1	c.916A > T	p.Lys306*	Nonsense	This study
<i>EXT1</i>	P36	Ex1	ex1del	Unknown	Deletion	LOVD
<i>EXT1</i>	P40	Ex 2	c.1018C > T	p.Arg340Cys	Missense	Philippe, et al, 1997
<i>EXT1</i>	P08	Ex 2	c.1037G > C	p.Arg346Thr	Missense	This study
<i>EXT1</i>	P28	In 3-4	c.1164 + 1G > A	--	Splice site	This study
<i>EXT1</i>	P41	Ex 4	c.1219C > T	p.Gln407*	Nonsense	LOVD
<i>EXT1</i>	P02	Ex 6	c.1469delT	p.Leu490Argfs*9	Frameshift	Anh, 1995
<i>EXT1</i>	P29	In 8-9	c.1722 + 1G > A	--	Splice site	Jennes I et al, 2008
<i>EXT1</i>	P25	Ex 9	c.1748A > G	p.Gln583Arg	Missense	This study
<i>EXT2</i>	P19	In 3-4	c.626 + 1G > A	--	Splice site	Vink et al. 2004
<i>EXT2</i>	P13	Ex 5	c.920_929del10insTG	p.Asp307Valfs*45	Frameshift	This study
<i>EXT2</i>	P04	Ex 6	ex6del	Unknown	Deletion	Leube, et al, 2008
<i>EXT2</i>	P01	Ex 8	c.1182G > A	p.Trp394*	Nonsense	This study
<i>EXT2</i>	P27	Ex 10	c.1616_1623del8ins10	p.Asp539Glnfs*5	Frameshift	This study
<i>EXT2</i>	P12	Ex 4-14	ex4-14del	Unknown	Deletion	This study

Novel mutations are indicated in bold. We considered as new mutations those not published and/or not mentioned in the LOVD databases. LOVD:<http://medgen.ua.ac.be/LOVDv.2.0/>

¹Patient with malignant transformation to chondrosarcoma.



domains³² (Protein Homology Fold Recognition Engine, <http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index>). Regarding the 3-D prediction for two novel *EXT1* missense mutations, p.Leu264Pro and p.Arg346Thr, the Phyre2 bioinformatic tool showed that both mutations produce impairment in protein folding or alterations in the exostosin or glycosyltransferase domains. For the third one, p.Gln583Arg, Phyre2 showed the removal of alpha helix structures with possible alterations in *EXT1* glycosyltransferase domain. A type II severe phenotype (IIS) was found in patients carrying the two first missense mutations (P17 and P08), and a severe phenotype (IIIS) was observed in P25 carrying the p.Gln583Arg mutation.

Six out of the 14 mutations in the *EXT1* gene (p.Val78Glyfs*111, p.Leu264Pro, p.Lys306*, p.Arg346Thr, c.1164 + 1C > A and p.Gln583Arg) and four of the six *EXT2* mutant alleles (p.Asp307Valfs*45, p.Trp394*, p.Asp539Glnfs*5, and exon4-14del) were novel. Although some mutation hotspots have been reported²⁰ (<http://medgen.ua.ac.be/LOVDv.2.0/>), we did not observe recurrent *EXT1* or *EXT2* mutations in patients in this study. The missense mutation c.1018C > T (p.Arg340Cys) observed in P40 and the exon 6 deletion c.1469delT (p.Leu490Argfs*9) found in P02 were previously described to cause the impairment of heparan sulfate synthesis^{20,33}.

The MLPA analysis in gDNA of patient P36, showed the complete deletion of *EXT1* exon 1. This patient had previously been reported as an SO case because he only had one lesion, but this detection in germline DNA allow us to change the diagnosis into MO patient with very mild symptoms (only a single osteochondroma lesion)³³ (Table 1 and 2). Two molecular defects in *EXT2* were detected by MLPA: the deletion of exon 6 (P04) and of 10 exons (from exon 4 to exon 14) in patient P12. This patient also carried an abnormal karyotype (an 18 q deletion).

Splice-site mutations were detected in *EXT1* in two patients. One of them, c.1164 + 1G > A in intron 3, was a novel mutation observed in patient P28. This novel mutation was analyzed *in silico* and the use of alternative cryptic donor sites was predicted. The phenotype in these patients was severe, presenting deformity of the limbs, valgus, restricted joint movement and scoliosis. Furthermore, one splice site mutation previously described was detected in P19 in *EXT2*. Clinical data were not available for this patient²⁵. There are several possible explanations for the lack of identification of mutations in some of the MO patients (22%). The mutation may have been in the *EXT1* or *EXT2* genes but in regions that were not analyzed. We did not look for mutations in deep intronic regions or in the 5' and 3' UTR sequences. Instead, the promoter region was genotyped and no mutation was detected. A recent study described a regulatory role for a G/C SNP (rs3401643) located at position -1158 bp, within a USF1 transcription factor binding site³⁰. These authors observed that the presence of the C-allele resulted in a ~56% increase in *EXT1* promoter activity. The effect of this allele in the four patients of the present study who are heterozygous for it will require further studies. It is well established that methylation of cytosine residues in the promoter region leads to transcription repression in tumor suppressor genes; nevertheless this does not seem to be the case for *EXT1* and *EXT2* promoters in osteochondromas or in chondrosarcomas^{34,35}. Finally, the possible existence of other genes responsible for MO should also be considered. A putative *EXT3* gene, located on the short arm of chromosome 19, has been proposed to explain the absence of an *EXT1* or *EXT2* mutation in a small percentage of MO patients (15–30%). Nevertheless, the existence of this third locus is generally accepted to be a false linkage result.

Inactivating mutations in the *EXT1* and *EXT2* genes were previously reported as the most common event in MO patients resulting in the formation of non-functional *EXT1* or *EXT2* proteins with a variable degree of expression in tissues^{27,36}. We observed 11 truncating mutations that create premature stop codons presenting a high grade of severity in patient's phenotype (Table 1 and 2). One of these

patients (P06) presented malignant transformation to chondrosarcoma and we detected the p.Leu283* mutation in the *EXT1* gene. Very low or null levels of *EXT1* and *EXT2* proteins were detected by Western blot in this patient. However, in these experiments, the bands corresponding to GAPDH (control protein) were very weak and the lack of additional sample precluded repetition. We think that in spite of the technical problems, this observation should be reported to allow comparisons with other studies. Obviously, further cases should be analyzed to confirm these findings. The loss of *EXT2* protein suggests that *EXT1* mutations probably interfere with the function of exostosin's complexes in the Golgi, inactivating the holoenzyme, degrading the whole protein, or interfering in some other function in the Golgi³⁷.

Several studies have suggested that MO patients present a more severe phenotype due to *EXT1* mutations than *EXT2* mutations^{16,18,21} while other studies could not confirm this observation^{23,36}. Pedrini et al 2011 recently performed a genotype–phenotype association study in a large cohort of MO patients and identified some specific correlation according to a new clinical classification system³¹. Our patients presented some of the most common manifestations, including orthopedic deformities of the forearm, ankle, varus or valgus of the knee, arthritis, vessels and nerve compression and very short stature (below the third percentile). The bones most often affected were tibia, femur, radius, humerus and fibula. Nevertheless, we observed a severe phenotype (12% type IS, 53% type IIS, 12% type IIIS and 23% type IVS) in 63% of MO patients (Figure 1). The remaining 15% presented with a moderate phenotype without a family history of the disease. We observed that the grade of severity differed between the proband and other affected members in the family, according to previously reported intra-familial variability^{18,38}. Nevertheless, no family history for MO was reported in 56% of MO patients. Patients with a mutation in the *EXT2* gene showed a smaller number of affected bones (data not shown) consistent with a recent study³⁹. The most frequently observed skeletal deformations in our patients were shortening of limbs, varus or valgus knee, short metacarpal bones, scoliosis, shortened stature and synostosis, with no evidence of differences between the grade of severity in the phenotype observed in patients with *EXT1* or *EXT2* (Figure 1). Genotype–phenotype correlations are difficult to establish in MO patients because most of the *EXT1* and *EXT2* variants are private mutations²⁰.

Malignant transformation to a chondrosarcoma is the most important complication in MO, and has been estimated to occur in 0.5–5% of patients¹⁷. Patients P06 and P38 developed malignant tumors, which gives a frequency of malignant transformation of 7% in our cohort of patients. Patient P06 bore the pathogenic mutation c.848T > A (p.Leu283*) in the first exon of the *EXT1* gene²², while no mutation was detected in P38, neither in *EXT1* nor *EXT2* (Table 1). It has been shown that hereditary osteochondromas and secondary chondrosarcomas are associated with a second mutational hit in the *EXT* genes^{40,41}. We thus investigated this possibility in DNA extracted from the osteochondroma tissue resected from P06 by Sanger sequence and MLPA but we found no evidence of a somatic mutation as a second hit in any of these genes. The presence of genetic rearrangements at the *EXT1* and *EXT2* loci (as the second mutational hit) in P06 osteochondromas and secondary chondrosarcomas was ruled out.

In conclusion, we have identified the disease-causing mutation in 21 out of 27 MO patients, including 10 mutations described for the first time. No mutation was identified in SO cases. Structural analyses predicted a disruption of important domains of *EXT1* proteins bearing missense mutations. A potentially functional promoter polymorphism was found in three patients with no other mutation, in one patient with a disease-causing mutation and in one control. No second hit was identified in a sample from a chondrosarcoma. Further studies are needed to identify the molecular bases of the



disease in 22% of the patients of this cohort and to understand the mechanisms underlying the malignant transformation process.

Methods

Patients and control individuals. We investigated 33 patients (18 males and 15 females), from unrelated pedigrees with osteochondromatosis from Chile and Argentina (27 MO and 6 SO). Nine control samples from healthy subjects were included in the promoter studies. Diagnosis was made on the basis of clinical manifestations and confirmed by physical and/or radiographic examinations at the Orthopedic and Imaging Departments, Children's Hospital of Córdoba, National University of Córdoba, Argentina. DNA and tissues samples from patients and their relatives were obtained together with their informed consent in accordance with the Helsinki Declaration as revised in 2000. The study was approved by the Ethics Committee (CIEIS) Act N° 95/2007. Genomic DNA was obtained from peripheral blood leukocytes and tissue samples from discarded tissues obtained by surgery, using the Wizard Genomic DNA purification Kit (Promega, Madison, WI). DNA was extracted according to the manufacturer's instructions.

Clinical studies and phenotypic data. Clinical variables were analyzed according to a scale established by the Musculoskeletal Tumoral Society with some modifications¹⁸. This scale includes the evaluation of all palpable lesions, patient's height, deformities, and functional limitations. Lesion quality and the severity of the disease were assessed according to age of onset (before/after 3 years), number of exostoses (more/less than 10 osteochondromas), vertebral location of the exostoses (absence or presence), stature (above/below 10th percentile), and functional rating (good or fair). The degree of severity was classified as mild (M) or severe (S). Four subcategories were defined in patients with a severe phenotype (from types IS to IVS)^{16,18}.

Genotyping and mutation analysis. The 11 *EXT1* and 13 *EXT2* coding exons and their intronic flanking regions were amplified by PCR from genomic DNA. Primer sequences and PCR conditions were as described by Sarrion et al 2013⁴². All fragments, except those corresponding to exon 1 of *EXT1*, could be amplified by PCR simultaneously. Exon 1 of *EXT1* and exon 2 of *EXT2* were split into several overlapping fragments, to obtain amplification products that did not exceed 650 bp. PCR was performed in a 50- μ l reaction volume, containing ~100 ng of genomic DNA, 1–2 mM MgCl₂, 0.2 mM of each dNTP, 0.4 μ M of each forward and reverse primer and 0.7 U of GoTaqR Flexi polymerase (Promega, Madison, WI). All PCR programs included an initial denaturation of 4 min at 95°C, followed by 35 cycles of 30 sec at 95°C, 30 sec at annealing temperature (Ta) and 1 min at 72°C. An extension at 72°C was then performed for 5 min. The annealing temperature was 60°C for all primer combinations, except during the amplification of overlapping regions of exon 1 of *EXT1*. For these primer combinations, Ta was set at 55°C for ex1.1 and 57°C for ex1.2 and ex1.3. The *EXT1* promoter region, between positions –1285 and –851, was also analyzed by sequencing. The PCR reaction was performed as described above with a Ta of 55°C. All PCR products were purified using a PCR purification kit (GE Healthcare) and sequenced with BigDye 3.1 (Applied Biosystems Life Technologies). The sequences were analyzed in an ABI PRISM 3730 DNA Analyzer (Applied Biosystems Life Technologies).

The presence of all detected mutations was confirmed by digestion with the appropriate restriction enzyme. Novel mutations were confirmed by analyzing 100 control alleles. The mutations were given the official HGVS nomenclature (www.hgvs.org). The reference sequences were NM_000127.2 for *EXT1* and NM_000401.3 for *EXT2*.

MLPA. The number of copies of the *EXT1* and *EXT2* exons present in the patient's genomic DNA was analyzed using the multiplex ligation-dependent probe amplification (MLPA) technique designed by MRC-Holland (code #P215-B1 EXT, MRC-Holland, Amsterdam, The Netherlands) following the manufacturer's instructions. PCR products were run on an ABI 3730 DNA Analyzer capillary sequencer (Applied Biosystems, Forster City, CA, USA). Peaks were analyzed using Coffalyser v9.4 software (MRC-Holland Vs 05; 30-08-2007). The proportion of each peak relative to the height of all peaks was calculated for each sample and then compared to proportions for the corresponding peak averaged for a set of at least ten normal DNA samples. Samples with ratios between 0.7 and 1.3 were considered as bearing a normal copy number. Ratios below 0.7 were considered to correspond to deletions, and above 1.3 to duplications. Each positive result was confirmed in a second independent MLPA reaction.

Assessment of functionality of missense mutations. In order to assess the possible pathogenic effect of the new missense mutations, the changes were analyzed using three in-silico online tools: PolyPhen-2 (Polymorphism Phenotyping v2; <http://genetics.bwh.harvard.edu/pph2/>, last accessed March 2014), Mutation Taster (<http://www.mutationtaster.org/>, last accessed March 2014), and ESE Finder 3.0 (ESE: Exonic Splicing Enhancer; http://rulai.cshl.edu/cgi-bin/tools/ESE3/ese_finder.cgi?process=home, last accessed March 2014). Protein Homology Fold Recognition Engine (<http://www.sbg.bio.ic.ac.uk/phyre2/html>, last accessed March 2014) was used to predict the implications of missense mutations for EXT1 3D structure Human Splice Finder (<http://www.umd.be/HSF/>, last accessed March 2014) and NetGene2 (<http://www.cbs.dtu.dk/cgi-bin/webface?jobid=netgene2>, last accessed March 2014) online tools were used to assess the possible effect of novel intronic mutations on splicing.

Ethics statement. The methods were carried out in accordance with the approved guidelines and in accordance with the Helsinki Declaration as revised in 2000. The study was approved by the Ethics Committee (CIEIS) Act N° 95/2007.

- Martinez-Duncker, I., Asteggiano, C. & Freeze, H. H. [Congenital Disorders of Glycosylation] [59–83] [Biochemistry Research Trends, Nova Science Publishers Inc Nueva York, 2012].
- Jaeken, J. Congenital disorders of glycosylation. *Ann. N. Y. Acad. Sci.* **1214**, 190–198 (2010).
- Ahn, J. et al. Cloning of the putative tumour suppressor gene for hereditary multiple exostoses (EXT1). *Nat. Genet.* **11**, 137–143 (1995).
- Stickens, D. et al. The EXT2 multiple exostoses gene defines a family of putative tumour suppressor genes. *Nat. Genet.* **14**, 25–32 (1996).
- Wuyts, W. et al. Positional cloning of a gene involved in hereditary multiple exostoses. *Hum. Mol. Genet.* **5**, 1547–1557 (1996).
- Busse, M. et al. Contribution of EXT1, EXT2, and EXTL3 to heparan sulfate chain elongation. *J. Biol. Chem.* **282**, 32802–32810 (2007).
- Kim, B. T. et al. Human tumor suppressor EXT gene family members EXTL1 and EXTL3 encode alpha 1,4-N-acetylglucosaminyltransferases that likely are involved in heparan sulfate/heparin biosynthesis. *Proc. Natl. Acad. Sci.* **98**, 7176–7181 (2001).
- Van Hul, W. et al. Identification of a third EXT-like gene (EXTL3) belonging to the EXT gene family. *Genomics.* **15**; 47, 230–7 (1998).
- Wuyts, W. et al. Identification and characterization of a novel member of the EXT gene family, EXTL2. *Eur J Hum Genet.* **5**, 382–9 (1997).
- Wise, C. A., Clines, G. A., Massa, H., Trask, B. J. & Lovett, M. Identification and localization of the gene for EXTL, a third member of the multiple exostoses gene family. *Genome Res.* **7**, 10–16 (1997).
- Gallagher, J. & Lyon, M. Structure and function of heparan sulphate proteoglycans. *Biochem. J.* **236**, 313–325 (1986).
- Carlsson, P., Presto, J., Spillmann, D., Lindahl, U. & Kjell n, L. Heparin/heparan sulfate biosynthesis: processive formation of N-sulfated domains. *J. Biol. Chem.* **283**, 20008–20014 (2008).
- Shi, X. & Zaia, J. Organ-specific heparan sulfate structural phenotypes. *J. Biol. Chem.* **284**, 11806–11814 (2009).
- Zak, B. M., Crawford, B. E., & Esko, J. D. Hereditary multiple exostoses and heparan sulfate polymerization. *Biochim. Biophys. Acta* **1573**, 346–355 (2002).
- De Andrea, C. E. & Hogendoorn, P. C. Epiphyseal growth plate and secondary peripheral chondrosarcoma: the neighbours matter. *J. Pathol.* **226**, 219–228 (2012).
- Alvarez, C., Tredwell, S., De Vera, M. & Hayden, M. The genotype-phenotype correlation of hereditary multiple exostoses. *Clin. Genet.* **70**, 122–130 (2006).
- Bov e, J. V. Multiple osteochondromas. *Orphanet J. Rare Dis.* **13**, 3 (2008).
- Francannet, C. et al. Genotype-phenotype correlation in hereditary multiple exostoses. *J. Med. Genet.* **38**, 430–434 (2001).
- Hameetman, L., Bov e, J. V., Taminiau, A. H., Kroon, H. M. & Hogendoorn, P. C. Multiple osteochondromas: clinicopathological and genetic spectrum and suggestions for clinical management. *Hered. Cancer Clin. Pract.* **2**, 161–173 (2004).
- Jennes, I. et al. Multiple osteochondromas: mutation update and description of the multiple osteochondromas mutation database (MOdb). *Hum. Mutat.* **30**, 1620–1627 (2009).
- Alvarez, C. M., De Vera, M. A., Heslip, T. R. & Casey, B. Evaluation of the anatomic burden of patients with hereditary multiple exostoses. *Clin. Orthop. Relat. Res.* **462**, 73–79 (2007).
- Delgado, M. A. et al. A novel nonsense mutation of the EXT1 gene in an Argentinean patient with multiple hereditary exostoses: a case report. *J. Bone Joint Surg. Am.* **94**, e76 (2012).
- Kitsoulis, P. et al. Osteochondromas: review of the clinical, radiological and pathological features. *In Vivo* **22**, 633–646 (2008).
- Jennes, I. et al. Mutation screening of EXT1 and EXT2 by denaturing high-performance liquid chromatography, direct sequencing analysis, fluorescence in situ hybridization, and a new multiplex ligation-dependent probe amplification probe set in patients with multiple osteochondromas. *J. Mol. Diagn.* **10**, 85–92 (2008).
- Vink, G. R. et al. Mutation screening of EXT1 and EXT2 by direct sequence analysis and MLPA in patients with multiple osteochondromas: splice site mutations and exonic deletions account for more than half of the mutations. *Eur. J. Hum. Genet.* **13**, 470–474 (2005).
- Lonie, L. et al. Determination of the mutation spectrum of the EXT1/EXT2 genes in British Caucasian patients with multiple osteochondromas, and exclusion of six candidate genes in EXT negative cases. *Hum. Mutat.* **27**, 1160 (2006).
- Pedrini, E. et al. Novel EXT1 and EXT2 mutations identified by DHPLC in Italian patients with multiple osteochondromas. *Hum. Mutat.* **26**, 280 (2005).
- White, S. J., Sterrenburg, E., van Ommen, G. J., den Dunnen, J. T. & Breuning, M. H. An alternative to FISH: detecting deletion and duplication carriers within 24 hours. *J. Med. Genet.* **40**, e113 (2003).
- Wuyts, W. et al. Mutations in the EXT1 and EXT2 genes in hereditary multiple exostoses. *Am. J. Hum. Genet.* **62**, 346–354 (1998).
- Jennes, I. et al. Identification and functional characterization of the human EXT1 promoter region. *Gene* **492**, 148–159 (2012).



31. Pedrini, E. *et al.* Genotype phenotype correlation study in 529 patients with multiple hereditary exostoses: identification of “protective” and “risk” factors. *J. Bone Joint Surg. Am.* **93**, 2294–302 (2011).
32. Ciavarella, M. *et al.* 20 novel point mutations and one large deletion in EXT1 and EXT2 genes: report of diagnostic screening in a large Italian cohort of patients affected by hereditary multiple exostosis. *Gene* **515**, 339–348 (2013).
33. McCormick, C., Duncan, G., Goutsos, K. T. & Tufaro, F. The putative tumor suppressors EXT1 and EXT2 form a stable complex that accumulates in the Golgi apparatus and catalyzes the synthesis of heparan sulfate. *Proc. Natl. Acad. Sci. USA* **97**, 668–673(2000).
34. Ropero, S. *et al.* Epigenetic loss of the familial tumor-suppressor gene exostosin-1 (EXT1) disrupts heparan sulfate synthesis in cancer cells. *Hum. Mol. Genet.* **13**, 2753–2765 (2004).
35. Hameetman, L. *et al.* Decreased EXT expression and intracellular accumulation of heparan sulphate proteoglycan in osteochondromas and peripheral chondrosarcomas. *J. Pathol.* **211**, 399–409 (2007).
36. Signori, E. *et al.* A combined analytical approach reveals novel EXT1/2 gene mutations in a large cohort of Italian multiple osteochondromas patients. *Genes Chrom. Cancer* **46**, 470–477 (2007).
37. Bernard, M. A. *et al.* Diminished levels of the putative tumor suppressor proteins EXT1 and EXT2 in exostosis chondrocytes. *Cell Motil. Cytoskeleton* **48**, 149–162 (2001).
38. Hennekam, R. C. Hereditary multiple exostoses. *J. Med. Genet.* **28**, 262–266 (1991).
39. Stancheva-Ivanova, M. K. *et al.* Clinical and molecular studies of EXT1/EXT2 in Bulgaria. *J. Inherit. Metab. Dis.* **34**, 917–921 (2011).
40. Bovée, J. V. EXT1 hit for mouse osteochondroma. *Proc. Natl. Acad. Sci. USA.* **107**, 1813–1814 (2010).
41. Hecht, J. T. *et al.* Hereditary multiple exostoses (EXT): mutational studies of familial EXT1 cases and EXT-associated malignancies. *Am. J. Hum. Genet.* **60**, 80–86 (1997).
42. Sarrión, P. *et al.* Mutations in the EXT1 and EXT2 genes in Spanish patients with multiple osteochondromas. *Sci. Rep.* **3**, 1346 (2013).

Acknowledgments

The authors thank the patients and their families for their cooperation. They wish to acknowledge Joanne Ferrier for revising the English. The authors are also grateful for the support from the Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), which is an initiative of the ISCIII. Grant support: Scientific and Technical National Research Council (CONICET); FONCyT PICT 2006–2350 and BID N° 2437/OC-AR PICT2010-2824, Catholic University of Cordoba Grant 2010/2011/2012/2013, Spanish Ministry of Education and Science (SAF2011-25431, PIB2010AR00473) and the Catalan Government (2009SGR971).

Author contributions

Study design: S.B., D.G., C.G.A. Collection and data samples: M.A.D., G.M.-D., P.S., L.Z., H.H.R., F.S. Performance of experiments: M.A.D., G.M.-D., C.G.A. Data interpretation and analysis: M.A.D., G.M.-D., P.S., R.U., R.D.-K., S.B., D.G., C.G.A. Draft composition: M.A.D., G.M.-D., S.B., D.G. C.G.A. conceived of the study, and participated in its design, coordination and helped to draft the manuscript. All authors reviewed the manuscript.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Delgado, M.A. *et al.* A broad spectrum of genomic changes in latinamerican patients with EXT1/EXT2-CDG. *Sci. Rep.* **4**, 6407; DOI:10.1038/srep06407 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>