

Search for CP Violation in the Decays $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$

B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ X. Prudent,¹ V. Tisserand,¹ A. Zghiche,¹ J. Garra Tico,² E. Grauges,² L. Lopez,³ A. Palano,³ M. Pappagallo,³ G. Eigen,⁴ B. Stugu,⁴ L. Sun,⁴ G. S. Abrams,⁵ M. Battaglia,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ J. A. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ D. Lopes Pegna,⁵ G. Lynch,⁵ L. M. Mir,⁵ T. J. Orimoto,⁵ I. L. Osipenkov,⁵ M. T. Ronan,^{5,*} K. Tackmann,⁵ T. Tanabe,⁵ W. A. Wenzel,⁵ P. del Amo Sanchez,⁶ C. M. Hawkes,⁶ A. T. Watson,⁶ H. Koch,⁷ T. Schroeder,⁷ D. Walker,⁸ D. J. Asgeirsson,⁹ T. Cuhadar-Donszelmann,⁹ B. G. Fulsom,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ M. Barrett,¹⁰ A. Khan,¹⁰ M. Saleem,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ K. Yu. Todyshev,¹¹ M. Bondioli,¹² S. Curry,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² P. Lund,¹² M. Mandelkern,¹² E. C. Martin,¹² D. P. Stoker,¹² S. Abachi,¹³ C. Buchanan,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ F. Liu,¹⁴ O. Long,¹⁴ B. C. Shen,¹⁴ G. M. Vitug,¹⁴ L. Zhang,¹⁴ H. P. Paar,¹⁵ S. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ A. Cunha,¹⁶ B. Dahmes,¹⁶ T. M. Hong,¹⁶ D. Kovalskyi,¹⁶ J. D. Richman,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. J. Flacco,¹⁷ C. A. Heusch,¹⁷ J. Kroseberg,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ M. G. Wilson,¹⁷ L. O. Winstrom,¹⁷ E. Chen,¹⁸ C. H. Cheng,¹⁸ F. Fang,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ R. Andreassen,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ K. Mishra,¹⁹ M. D. Sokoloff,¹⁹ F. Blanc,²⁰ P. C. Bloom,²⁰ S. Chen,²⁰ W. T. Ford,²⁰ J. F. Hirschauer,²⁰ A. Kreisel,²⁰ M. Nagel,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ J. G. Smith,²⁰ K. A. Ulmer,²⁰ S. R. Wagner,²⁰ J. Zhang,²⁰ A. M. Gabareen,²¹ A. Soffer,^{21,†} W. H. Toki,²¹ R. J. Wilson,²¹ F. Winklmeier,²¹ D. D. Altenburg,²² E. Feltresi,²² A. Hauke,²² H. Jasper,²² J. Merkel,²² A. Petzold,²² B. Spaan,²² K. Wacker,²² V. Klose,²³ M. J. Kobel,²³ H. M. Lacker,²³ W. F. Mader,²³ R. Nogowski,²³ J. Schubert,²³ K. R. Schubert,²³ R. Schwierz,²³ J. E. Sundermann,²³ A. Volk,²³ D. Bernard,²⁴ G. R. Bonneaud,²⁴ E. Latour,²⁴ V. Lombardo,²⁴ Ch. Thiebaut,²⁴ M. Verderi,²⁴ P. J. Clark,²⁵ W. Gradl,²⁵ F. Muheim,²⁵ S. Playfer,²⁵ A. I. Robertson,²⁵ J. E. Watson,²⁵ Y. Xie,²⁵ M. Andreotti,²⁶ D. Bettoni,²⁶ C. Bozzi,²⁶ R. Calabrese,²⁶ A. Cecchi,²⁶ G. Cibinetto,²⁶ P. Franchini,²⁶ E. Luppi,²⁶ M. Negrini,²⁶ A. Petrella,²⁶ L. Piemontese,²⁶ E. Prencipe,²⁶ V. Santoro,²⁶ F. Anulli,²⁷ R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ S. Pacetti,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,‡} M. Piccolo,²⁷ M. Rama,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Contri,²⁸ M. Lo Vetere,²⁸ M. M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ K. S. Chaisanguanthum,²⁹ M. Morii,²⁹ J. Wu,²⁹ R. S. Dubitzky,³⁰ J. Marks,³⁰ S. Schenk,³⁰ U. Uwer,³⁰ D. J. Bard,³¹ P. D. Dauncey,³¹ R. L. Flack,³¹ J. A. Nash,³¹ W. Panduro Vazquez,³¹ M. Tibbetts,³¹ P. K. Behera,³² X. Chai,³² M. J. Charles,³² U. Mallik,³² J. Cochran,³³ H. B. Crawley,³³ L. Dong,³³ V. Eyges,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ A. E. Rubin,³³ Y. Y. Gao,³⁴ A. V. Gritsan,³⁴ Z. J. Guo,³⁴ C. K. Lae,³⁴ A. G. Denig,³⁵ M. Fritsch,³⁵ G. Schott,³⁵ N. Arnaud,³⁶ J. Béquilleux,³⁶ A. D'Orazio,³⁶ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ V. Lepeltier,³⁶ F. Le Diberder,³⁶ A. M. Lutz,³⁶ S. Pruvot,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ J. Serrano,³⁶ V. Sordini,³⁶ A. Stocchi,³⁶ W. F. Wang,³⁶ G. Wormser,³⁶ D. J. Lange,³⁷ D. M. Wright,³⁷ I. Bingham,³⁸ J. P. Burke,³⁸ C. A. Chavez,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ D. E. Hutchcroft,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸ A. J. Bevan,³⁹ K. A. George,³⁹ F. Di Lodovico,³⁹ R. Sacco,³⁹ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ D. A. Hopkins,⁴⁰ S. Paramesvaran,⁴⁰ F. Salvatore,⁴⁰ A. C. Wren,⁴⁰ D. N. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² D. Bailey,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² Y. M. Chia,⁴² C. L. Edgar,⁴² G. D. Lafferty,⁴² T. J. West,⁴² J. I. Yi,⁴² J. Anderson,⁴³ C. Chen,⁴³ A. Jawahery,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ J. M. Tuggle,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ E. Salvati,⁴⁴ S. Saremi,⁴⁴ R. Cowan,⁴⁵ D. Dujmic,⁴⁵ P. H. Fisher,⁴⁵ K. Koeneke,⁴⁵ G. Sciolla,⁴⁵ M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ M. Zhao,⁴⁵ Y. Zheng,⁴⁵ S. E. Mclachlin,^{46,*} P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ M. Simard,⁴⁹ P. Taras,⁴⁹ F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ G. De Nardo,⁵¹ F. Fabozzi,^{51,§} L. Lista,⁵¹ D. Monorchio,⁵¹ C. Sciacca,⁵¹ M. A. Baak,⁵² G. Raven,⁵² H. L. Snoek,⁵² C. P. Jessop,⁵³ K. J. Knoepfel,⁵³ J. M. LoSecco,⁵³ G. Benelli,⁵⁴ L. A. Corwin,⁵⁴ K. Honscheid,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ J. P. Morris,⁵⁴ A. M. Rahimi,⁵⁴ J. J. Regensburger,⁵⁴ S. J. Sekula,⁵⁴ Q. K. Wong,⁵⁴ N. L. Blount,⁵⁵ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ J. A. Kolb,⁵⁵ M. Lu,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ N. Gagliardi,⁵⁶ A. Gaz,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ A. Pompili,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ E. Ben-Haim,⁵⁷ H. Briand,⁵⁷ G. Calderini,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ Ph. Leruste,⁵⁷ J. Malcès,⁵⁷ J. Ocariz,⁵⁷ A. Perez,⁵⁷ J. Prendki,⁵⁷ L. Gladney,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ E. Manoni,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰

S. Bettarini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ A. Cervelli,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. A. Mazur,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ G. Rizzo,⁶⁰ J. J. Walsh,⁶⁰ J. Biesiada,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ E. Baracchini,⁶² F. Bellini,⁶² G. Cavoto,⁶² D. del Re,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² P. D. Jackson,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Renga,⁶² C. Voena,⁶² M. Ebert,⁶³ T. Hartmann,⁶³ H. Schröder,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ G. Castelli,⁶⁴ B. Franek,⁶⁴ E. O. Olaiya,⁶⁴ W. Roethel,⁶⁴ F. F. Wilson,⁶⁴ S. Emery,⁶⁵ M. Escalier,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ X. R. Chen,⁶⁶ H. Liu,⁶⁶ W. Park,⁶⁶ M. V. Purohit,⁶⁶ R. M. White,⁶⁶ J. R. Wilson,⁶⁶ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ P. Bechtel,⁶⁷ R. Claus,⁶⁷ J. P. Coleman,⁶⁷ M. R. Convery,⁶⁷ J. C. Dingfelder,⁶⁷ J. Dorfan,⁶⁷ G. P. Dubois-Felsmann,⁶⁷ W. Dunwoodie,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ M. T. Graham,⁶⁷ P. Grenier,⁶⁷ C. Hast,⁶⁷ W. R. Innes,⁶⁷ J. Kaminski,⁶⁷ M. H. Kelsey,⁶⁷ H. Kim,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ S. Li,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ D. B. MacFarlane,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ I. Ofte,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ T. Pulliam,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ D. Su,⁶⁷ M. K. Sullivan,⁶⁷ K. Suzuki,⁶⁷ S. K. Swain,⁶⁷ J. M. Thompson,⁶⁷ J. Va'vra,⁶⁷ A. P. Wagner,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ K. Yi,⁶⁷ C. C. Young,⁶⁷ V. Ziegler,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ T. S. Miyashita,⁶⁸ B. A. Petersen,⁶⁸ L. Wilden,⁶⁸ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ R. Bula,⁶⁹ J. A. Ernst,⁶⁹ V. Jain,⁶⁹ B. Pan,⁶⁹ M. A. Saeed,⁶⁹ F. R. Wappler,⁶⁹ S. B. Zain,⁶⁹ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ J. L. Ritchie,⁷¹ A. M. Ruland,⁷¹ C. J. Schilling,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ M. Pelliccioni,⁷³ M. Bomben,⁷⁴ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ L. Lanceri,⁷⁴ L. Vitale,⁷⁴ V. Azzolini,⁷⁵ N. Lopez-March,⁷⁵ F. Martinez-Vidal,⁷⁵ D. A. Milanes,⁷⁵ A. Oyanguren,⁷⁵ J. Albert,⁷⁶ Sw. Banerjee,⁷⁶ B. Bhuyan,⁷⁶ K. Hamano,⁷⁶ R. Kowalewski,⁷⁶ I. M. Nugent,⁷⁶ J. M. Roney,⁷⁶ R. J. Sobie,⁷⁶ P. F. Harrison,⁷⁷ J. Ilic,⁷⁷ T. E. Latham,⁷⁷ G. B. Mohanty,⁷⁷ H. R. Band,⁷⁸ X. Chen,⁷⁸ S. Dasu,⁷⁸ K. T. Flood,⁷⁸ J. J. Hollar,⁷⁸ P. E. Kutter,⁷⁸ Y. Pan,⁷⁸ M. Pierini,⁷⁸ R. Prepost,⁷⁸ S. L. Wu,⁷⁸ and H. Neal⁷⁹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

²³Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²⁴Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

²⁵University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁶Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

²⁹Harvard University, Cambridge, Massachusetts 02138, USA

³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

- ³¹Imperial College London, London, SW7 2AZ, United Kingdom
³²University of Iowa, Iowa City, Iowa 52242, USA
³³Iowa State University, Ames, Iowa 50011-3160, USA
³⁴Johns Hopkins University, Baltimore, Maryland 21218, USA
³⁵Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
³⁶Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France
³⁷Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁸University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁹Queen Mary, University of London, E1 4NS, United Kingdom
⁴⁰University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴¹University of Louisville, Louisville, Kentucky 40292, USA
⁴²University of Manchester, Manchester M13 9PL, United Kingdom
⁴³University of Maryland, College Park, Maryland 20742, USA
⁴⁴University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁵Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁶McGill University, Montréal, Québec, Canada H3A 2T8
⁴⁷Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁸University of Mississippi, University, Mississippi 38677, USA
⁴⁹Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
⁵⁰Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵¹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵²NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁴Ohio State University, Columbus, Ohio 43210, USA
⁵⁵University of Oregon, Eugene, Oregon 97403, USA
⁵⁶Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁷Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie–Paris6, Université Denis Diderot–Paris7, F-75252 Paris, France
⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁹Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁶⁰Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶¹Princeton University, Princeton, New Jersey 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁸Stanford University, Stanford, California 94305-4060, USA
⁶⁹State University of New York, Albany, New York 12222, USA
⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
⁷¹University of Texas at Austin, Austin, Texas 78712, USA
⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁵IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁶University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁷Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
⁷⁸University of Wisconsin, Madison, Wisconsin 53706, USA
⁷⁹Yale University, New Haven, Connecticut 06511, USA

(Received 31 August 2007; published 15 February 2008)

We measure time-integrated CP -violating asymmetries of neutral charmed mesons in the modes $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ with the highest precision to date by using $D^0 \rightarrow K^- \pi^+$ decays to correct detector asymmetries. An analysis of 385.8 fb^{-1} of data collected with the *BABAR* detector yields values of $a_{CP}^{KK} = (0.00 \pm 0.34(\text{stat}) \pm 0.13(\text{syst}))\%$ and $a_{CP}^{\pi\pi} = (-0.24 \pm 0.52(\text{stat}) \pm 0.22(\text{syst}))\%$, which agree with standard model predictions.

Evidence for quantum-mechanical oscillations in neutral charmed mesons has recently been reported [1,2], increasing the importance of understanding the relative behaviors in this particle-antiparticle system. Unknown processes could contribute significantly to these oscillations, and there are many theoretical scenarios in which particle-antiparticle asymmetries are also expected. Charge-parity (CP) violation in time-integrated decay rates of charmed mesons at levels as large as 1% has not yet been experimentally ruled out [3], and at this level would be evidence of unknown physical phenomena [4,5]. The CP -even decays $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ [6] are Cabibbo suppressed, with the two neutral charmed mesons, D^0 and \bar{D}^0 , sharing the final states. CP -violating asymmetries in these modes are predicted to be $\mathcal{O}(0.001\% - 0.01\%)$ in the standard model of particle physics [7], yet have not been measured precisely due to limited sample sizes and relatively large systematic effects [8].

We search for CP violation in decays of charmed mesons produced from charm-quark pairs in the reaction $e^+ e^- \rightarrow c\bar{c}$ by measuring the asymmetries in the partial decay widths Γ ,

$$a_{CP}^{KK} = \frac{\Gamma(D^0 \rightarrow K^- K^+) - \Gamma(\bar{D}^0 \rightarrow K^+ K^-)}{\Gamma(D^0 \rightarrow K^- K^+) + \Gamma(\bar{D}^0 \rightarrow K^+ K^-)}, \quad (1)$$

$$a^{\pi\pi} = \frac{\Gamma(D^0 \rightarrow \pi^- \pi^+) - \Gamma(\bar{D}^0 \rightarrow \pi^+ \pi^-)}{\Gamma(D^0 \rightarrow \pi^- \pi^+) + \Gamma(\bar{D}^0 \rightarrow \pi^+ \pi^-)}. \quad (2)$$

In this construction, a_{CP}^{hh} , $h = K, \pi$, includes all CP violating contributions, direct and indirect [4]. The presence of direct CP violation in one or both modes would be signaled by a nonvanishing difference between the modes, $a_{CP}^{KK} - a_{CP}^{\pi\pi} \neq 0$. Indirect CP -violating asymmetries in these modes arising from D^0 - \bar{D}^0 oscillations have been measured in analyses of decay-time distributions [9], most recently with a precision of 0.30% [2].

Precise quantification of asymmetry in D^0 -flavor assignment, called tagging, has long been considered the primary experimental challenge in these measurements. We develop a new technique for measuring and correcting this asymmetry using only the recorded data. However, forward-backward (FB) asymmetry in $c\bar{c}$ production may be more significant at the center-of-mass energy of $e^+ e^-$ collisions in *BABAR*, $\sqrt{s} \approx 10.6$ GeV. This production asymmetry will create a difference in the numbers of reconstructed D^0 and \bar{D}^0 events due to the FB detection asymmetries coming from the boost of the center-of-mass system (c.m.s.) relative to the laboratory.

The production asymmetry has two physical components. Interference in $e^+ e^- \rightarrow c\bar{c}$ as mediated by either a virtual γ or a virtual Z^0 contributes at the percent level at this energy, and is well understood. In addition, asymmetries induced by higher-order QED effects are expected to have polar-angle dependence and to peak sharply in the

forward and backward directions [10]. Although well considered for μ -pair production [11], the precise shape of this contribution for D production is not known.

We use a data sample corresponding to an integrated luminosity of 385.8 fb^{-1} collected with the *BABAR* detector [12] at the PEP-II $e^+ e^-$ collider at SLAC. The production vertices of charged particles are measured with a silicon-strip detector (SVT), and their momenta are measured by the SVT and a drift chamber (DCH) in a 1.5 T magnetic field. Information from a Cherenkov-radiation detector, along with energy-deposition measurements from the SVT and DCH, provide K - π discrimination.

We analyze neutral D mesons produced from $D^{*+} \rightarrow D^0 \pi_s^+$; the charge of the π_s , a low-momentum (soft) pion, indicates the flavor of the D^0 at production. To correct for asymmetry in this flavor tag, we measure the relative detection efficiency for soft pions in recorded data using the decay $D^0 \rightarrow K^- \pi^+$ with (tagged) and without (nontagged) soft-pion flavor tagging. The only detector asymmetry present in reconstruction of the signal modes is due to the tagging π_s , since the CP final states are reconstructed identically for D^0 and \bar{D}^0 .

We reconstruct the four decay chains $D^0 \rightarrow K^- \pi^+$; $D^{*+} \rightarrow D^0 \pi_s^+$, $D^0 \rightarrow K^- \pi^+$; $D^{*+} \rightarrow D^0 \pi_s^+$, $D^0 \rightarrow K^- K^+$; and $D^{*+} \rightarrow D^0 \pi_s^+$, $D^0 \rightarrow \pi^- \pi^+$. We require D^0 candidates to have center-of-mass momenta greater than 2.4 GeV/ c , which removes almost all B decays. Each D^0 daughter must satisfy a likelihood-based particle-identification selection and must have at least two position measurements in each of the z and ϕ coordinates of the SVT. We require π_s^\pm candidates to have a lab momentum greater than 100 MeV/ c and at least six position measurements in the SVT.

For $h = K, \pi$, we accept candidates with an invariant mass $1.79 < m_{hh} < 1.93$ GeV/ c^2 and, for final states with a π_s , an invariant mass difference $0.140 < \Delta m < 0.152$ GeV/ c^2 , where $\Delta m \equiv m_{hh\pi_s} - m_{hh}$. For each D^0 candidate, we constrain the $h^+ h^-$ tracks to originate from a common vertex; for applicable final states, we also require the D^0 and π_s to originate from a common vertex within the $e^+ e^-$ interaction region. We select candidates for which the χ^2 probability of the vertex fit of the two D^0 daughters is greater than 0.005. For the KK and $\pi\pi$ modes, final asymmetries are calculated using events for which the polar angle of the D^0 momentum in the c.m.s. with respect to the beam axis satisfies $|\cos\theta_{D^0}^{c.m.s.}| < 0.8$.

We statistically separate signal from background in the selected events by calculating signal weights based on an optimized likelihood function [13]. The likelihood function is composed of probability density functions (PDFs) that are fitted to the mass distributions using the maximum likelihood technique. For the nontagged sample, a one-dimensional PDF is fitted to the $m_{K\pi}$ distribution; for the tagged samples, two-dimensional PDFs are fitted to the m_{hh} and Δm distributions. Two-dimensional PDFs are used

for the tagged samples to account for possible asymmetries in the background from correctly reconstructed D^0 decays with a misassociated π_s candidate; this background category peaks in m_{hh} but does not peak in Δm . The PDFs in this analysis are nearly identical to those used in an analysis of the decay $D^0 \rightarrow K^+ \pi^-$ [1], since the signal shapes and background sources are very similar. Although the PDFs are motivated by studies of simulated events, all of the shape parameters are varied in the fits to recorded data. Our selection of PDFs is treated as a source of systematic uncertainty. Because the signal shape is indistinguishable for D^0 and \bar{D}^0 distributions, we use the same signal PDF to describe both flavors of a mode and fit it to them simultaneously to reduce statistical uncertainties. The KK and $\pi\pi$ invariant mass distributions for D^0 and \bar{D}^0 , with fitted PDFs overlaid, are shown in Fig. 1. This analysis is sensitive only to ratios of D^0 -signal yields to \bar{D}^0 -signal yields, and not to absolute yields, so the final results are relatively insensitive to the exact forms of the PDFs.

The decay $D^0 \rightarrow K^- \pi^+$ is chosen as a calibration mode because it provides an easily reconstructed independent sample with high statistics. However, detector asymmetries in reconstruction of the D^0 final state cannot be ignored [see Figs. 2(a) and 2(b)]. These must be corrected to isolate the soft-pion asymmetry.




FIG. 1. Invariant mass distributions of the KK final state tagged as (a) D^0 and (b) \bar{D}^0 , and the $\pi\pi$ final state tagged as (c) D^0 and (d) \bar{D}^0 . Distributions of data (points with error bars) in the signal region $0.1434 < \Delta m < 0.1474$ GeV/c^2 are overlaid with fitted PDFs (dashed line, shaded areas). The white regions under the central peaks represent signal events, the light gray misassociated π_s^\pm events, and the dark gray remaining nonpeaking background. The data are shown over ranges extended beyond the fitted regions to illustrate the physical background shapes.

Using the nontagged $K\pi$ sample, we produce a map of the relative reconstruction efficiency between D^0 and \bar{D}^0 in this final state in terms of the momenta of both D^0 daughters, shown by components in Figs. 2(a) and 2(b). For each D^0 daughter, we consider the momentum magnitude and polar angle in the lab with respect to the beam axis; these components are correlated. The daughters are, however, factorizable from one another. By considering the normalized product of the K and π efficiency-map components, we obtain a four-dimensional relative-efficiency map for correcting $D^0 \rightarrow K^- \pi^+$ relative to $\bar{D}^0 \rightarrow K^+ \pi^-$. The presence of prompt D^0 decays not originating from a D^{*+} in the nontagged sample extends the kinematic boundaries of the map but does not otherwise affect it.

This $K\pi$ map is used to weight the D^0 candidates in the slow-pion tagged $K\pi$ sample, eliminating asymmetries due to the D^0/\bar{D}^0 daughters. Because all charm production is subject to the same production asymmetries, these are simultaneously removed from the tagged $K\pi$ sample by this correction. After the weights have been applied, the remaining asymmetry in the sample is due to the relative soft-pion efficiency.

We produce a map of the relative soft-pion efficiency in terms of the pion-momentum magnitude and polar angle in the lab [Fig. 2(c)]. Charm production is azimuthally uniform, and ϕ is found to be uncorrelated with other momentum variables. Therefore, the ϕ dependence is accounted for by an integrated scale factor. The uncertainties shown [Fig. 2(d)] are due to the statistical uncertainties in the sample yields. Signal-mode D^0 yields are weighted with this π_s map to correct for the soft-pion tagging




FIG. 2. $K\pi$ efficiency-map components obtained from the nontagged D^0 daughters (a) K and (b) π , and (c) π_s efficiency map with (d) statistical errors from the tagged $K\pi$ sample. Maps are produced from the ratios of candidate numbers of D^0 to \bar{D}^0 .

TABLE I. Signal yields in reconstructed modes. Listed uncertainties are statistical only. Corrections are applied only to D^0 samples, but all postcorrection samples are restricted to the phase space of the correction map.

Final state	Raw yields		Corr. used	Postcorrection yields	
	D^0	\bar{D}^0		D^0	\bar{D}^0
$K\pi$	$3\,363\,000 \pm 6000$	$3\,368\,000 \pm 6000$	None
$K\pi\pi_s$	$705\,100 \pm 1000$	$703\,500 \pm 1000$	$K\pi$ Map	633 300	630 100
$KK\pi_s$	$65\,730 \pm 340$	$63\,740 \pm 330$	π_s Map	65 210	63 490
$\pi\pi\pi_s$	$32\,210 \pm 310$	$31\,930 \pm 310$	π_s Map	31 900	31 760

asymmetry. The signal modes (with remaining production asymmetries) can thus be analyzed for evidence of CP violation. In Table I, we list the raw and postcorrection yields for the calibration and signal samples in this analysis. In calculating these corrections, histogram bins near kinematic boundaries with fewer than 5000 events are removed.

CP violation would appear as an asymmetry in D^0/\bar{D}^0 yields, independent of any kinematic variables. Because of the FB asymmetry in production, we calculate yield asymmetries as a function of $\cos\theta = \cos\theta_{D^0}^{\text{c.m.s.}}$ and decompose these into even and odd parts. We define

$$a^\pm(\cos\theta) = \frac{n_{D^0}(\pm|\cos\theta|) - n_{\bar{D}^0}(\pm|\cos\theta|)}{n_{D^0}(\pm|\cos\theta|) + n_{\bar{D}^0}(\pm|\cos\theta|)}, \quad (3)$$

$$a_{CP} = a_{CP}(\cos\theta) \approx (a^+(\cos\theta) + a^-(\cos\theta))/2, \quad (4)$$

$$a_{FB}(\cos\theta) \approx (a^+(\cos\theta) - a^-(\cos\theta))/2, \quad (5)$$

where n_{D^0} and $n_{\bar{D}^0}$ are the numbers of signal events for D^0 and \bar{D}^0 after applying the weights discussed above, a_{CP} is the even component, and $a_{FB}(\cos\theta)$ the odd component. Equations (4) and (5) are approximate as second-order terms in a^\pm have been omitted. The even part, representing CP -violating effects, would provide evidence of a uniform yield asymmetry. The odd part represents the production asymmetry, including higher-order QED contributions. From the several values of a_{CP} obtained as a function of $|\cos\theta|$, we obtain a central value from a χ^2 minimization.

We consider three sources of systematic error to be significant. One source is the choice of PDFs used to describe the signal and background distributions, which affects the statistical background subtraction. We estimate this systematic uncertainty by substituting different background shapes in m and Δm and an alternative two-dimensional signal shape in the fits to the tagged samples.

TABLE II. Summary of systematic uncertainties.

Category	Δa_{CP}^{KK}	$\Delta a_{CP}^{\pi\pi}$
2-Dim. PDF shapes	$\pm 0.04\%$	$\pm 0.05\%$
π_s correction	$\pm 0.08\%$	$\pm 0.08\%$
a_{CP} extraction	$\pm 0.09\%$	$\pm 0.20\%$
Quadrature sum	$\pm 0.13\%$	$\pm 0.22\%$

Another source is the binning choices made and dependences in the π_s -efficiency correction. We estimate the size of this uncertainty by varying the number of bins and the required number of events per bin in histograms used to calculate efficiencies, and by adding a ϕ dependence to the efficiency correction. We find the largest uncertainty here arises from the particular choice of binning in the π_s -efficiency map. Because the systematic uncertainty in applying the π_s -efficiency correction is the same for both modes, we evaluate its size using the larger signal sample. Finally, we consider the procedure for extracting a_{CP} . We vary the binning and the accepted range of $|\cos\theta|$; the largest uncertainty comes from the latter. All other sources of systematic uncertainty are highly suppressed because the final states are reconstructed identically for D^0 and \bar{D}^0 . We summarize the contributions to the total systematic uncertainty in Table II. The smaller $\pi\pi$ sample size influences the calculation of its systematic uncertainty.

For KK , we measure $a_{CP}^{KK} = (0.00 \pm 0.34(\text{stat}) \pm 0.13(\text{syst}))\%$. For $\pi\pi$, we measure $a^{\pi\pi} = (-0.24 \pm 0.52(\text{stat}) \pm 0.22(\text{syst}))\%$. Statistical uncertainties of




FIG. 3. CP -violating asymmetries in (a) KK and (b) $\pi\pi$, and forward-backward asymmetries in (c) KK and (d) $\pi\pi$. In (a) and (b), the dashed lines represent the central values and the hatched regions the 1σ intervals, obtained from χ^2 minimizations.

0.1% in the π_s correction have been included in the final statistical uncertainty values. The even and odd asymmetries for each mode as a function of $|\cos\theta|$ are shown in Fig. 3. We conclude from the χ^2 minimizations in Figs. 3(a) and 3(b) that there is no evidence of CP violation in either of the Cabibbo-suppressed two-body modes of D^0 decay. This result is in agreement with standard model predictions. It also provides a new constraint on theories beyond the standard model [5], some of which predict significant levels of CP violation in these modes. The asymmetries observed in Figs. 3(c) and 3(d) represent the two standard model asymmetries discussed. Although an exact prediction of these forward-backward asymmetries does not exist, the observed values do not contradict expectations. Thus, although we report both the most precise measurements of time-integrated CP asymmetries in charm decays and the first measurements of the FB asymmetry in charm-pair production at $\sqrt{s} \approx 10.6$ GeV, we do not report evidence of unknown phenomena at work in the neutral charmed meson system.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MEC (Spain), and STFC (U.K.). Individuals have received support from the Marie Curie EIF (European Union) and the A.P. Sloan Foundation.

*Deceased.

†Now at Tel Aviv University, Tel Aviv, 69978, Israel.

‡Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

§Also with Università della Basilicata, Potenza, Italy.

||Also with Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain.

- [1] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **98**, 211802 (2007).
- [2] M. Staric *et al.* (Belle Collaboration), Phys. Rev. Lett. **98**, 211803 (2007).
- [3] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **94**, 122001 (2005); B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **71**, 091101 (2005); D. Cronin-Hennessy *et al.* (CLEO Collaboration), Phys. Rev. D **72**, 031102 (2005); **75**, 119904 (2007).
- [4] Y. Grossman, A.L. Kagan, and Y. Nir, Phys. Rev. D **75**, 036008 (2007).
- [5] G. Burdman and I. Shipsey, Annu. Rev. Nucl. Part. Sci. **53**, 431 (2003).
- [6] Unless otherwise indicated, particle types and decay processes imply also their charge conjugates.
- [7] F. Buccella, M. Lusignoli, G. Miele, A. Pugliese, and P. Santorelli, Phys. Rev. D **51**, 3478 (1995); S. Bianco, F.L. Fabbri, D. Benson, and I. Bigi, Riv. Nuovo Cimento Soc. Ital. Fis. **26**, No. 7, 1 (2003).
- [8] W.M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [9] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **91**, 121801 (2003); S.E. Csorna *et al.* (CLEO Collaboration), Phys. Rev. D **65**, 092001 (2002).
- [10] Photon-box amplitudes interfering with the single-photon amplitude and Bremsstrahlung amplitudes interfering among themselves create a forward-backward asymmetry in $e^+e^- \rightarrow c\bar{c}$.
- [11] F.A. Berends, K.J.F. Gaemers, and R. Gastmans, Nucl. Phys. **B63**, 381 (1973); R. W. Brown, K. O. Mikaelian, V. K. Cung, and E. A. Paschos, Phys. Lett. B **43**, 403 (1973).
- [12] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [13] M. Pivk and F.R. Le Diberder, Nucl. Instrum. Methods Phys. Res., Sect. A **555**, 356 (2005).