

Measurements of Branching Fractions and Time-Dependent CP -Violating Asymmetries in $B \rightarrow \eta'/K$ Decays

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges-Pous,² A. Palano,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ A. W. Borgland,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomoisky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. E. Latham,⁹ F. F. Wilson,⁹ T. Cuhadar-Donszelmann,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ D. Thiessen,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² V. N. Ivanchenko,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ Sh. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ A. Lu,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzki,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ S. Yang,¹⁹ S. Jayatilleke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ J. Zhang,²¹ L. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² J. L. Harton,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² B. Spaan,²³ D. Altenburg,²⁴ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ E. Feltresi,²⁴ A. Hauke,²⁴ H. M. Lacker,²⁴ E. Maly,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ A. Sarti,²⁷ F. Anulli,²⁸ R. Baldini-Feroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ G. Crosetti,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² M. J. Charles,³³ G. J. Grenier,³³ U. Mallik,³³ A. K. Mohapatra,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ J. Lamsa,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ T. C. Petersen,³⁵ M. Pierini,³⁵ S. Plaszczynski,³⁵ M. H. Schune,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹ R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ M. A. Winter,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ M. T. Naisbit,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³ V. B. Koptchev,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ G. Cerizza,⁴⁶ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,*} F. Fabozzi,^{50,*} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³

T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malclès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ M. Bondioli,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ G. Simi,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ V. Miftakov,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,^{61,62} A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ G. De Nardo,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ J. Strube,^{54,67} D. Su,⁶⁷ M. K. Sullivan,⁶⁷ J. Va'vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ H. Kim,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,^{74,†} L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ F. Martinez-Vidal,^{2,75} R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ J. Wu,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ M. G. Greene,⁸⁰ and H. Neal⁸⁰

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Dipartimento di Fisica and INFN, Università di Bari, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵Institute of Physics, University of Bergen, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸Institute for Particle Physics, University of California at Santa Cruz, Santa Cruz, California 95064, USA

¹⁹California Institute of Technology, Pasadena, California 91125, USA

²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA

²¹University of Colorado, Boulder, Colorado 80309, USA

²²Colorado State University, Fort Collins, Colorado 80523, USA

- ²³*Institut für Physik, Universität Dortmund, D-44221 Dortmund, Germany*
- ²⁴*Institut für Kern- und Teilchenphysik, Technische Universität Dresden, D-01062 Dresden, Germany*
- ²⁵*Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁶*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁷*Dipartimento di Fisica and INFN, Università di Ferrara, I-44100 Ferrara, Italy*
- ²⁸*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁹*Dipartimento di Fisica and INFN, Università di Genova, I-16146 Genova, Italy*
- ³⁰*Harvard University, Cambridge, Massachusetts 02138, USA*
- ³¹*Physikalisches Institut, Universität Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³²*Imperial College London, London, SW7 2AZ, United Kingdom*
- ³³*University of Iowa, Iowa City, Iowa 52242, USA*
- ³⁴*Iowa State University, Ames, Iowa 50011-3160, USA*
- ³⁵*Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France*
- ³⁶*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
- ³⁷*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁸*Queen Mary, University of London, E1 4NS, United Kingdom*
- ³⁹*Royal Holloway and Bedford New College, University of London, Egham, Surrey TW20 0EX, United Kingdom*
- ⁴⁰*University of Louisville, Louisville, Kentucky 40292, USA*
- ⁴¹*University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴²*University of Maryland, College Park, Maryland 20742, USA*
- ⁴³*University of Massachusetts, Amherst, Massachusetts 01003, USA*
- ⁴⁴*Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
- ⁴⁵*McGill University, Montréal, Québec, Canada H3A 2T8*
- ⁴⁶*Dipartimento di Fisica and INFN, Università di Milano, I-20133 Milano, Italy*
- ⁴⁷*University of Mississippi, University, Mississippi 38677, USA*
- ⁴⁸*Laboratoire René J. A. Lévesque, Université de Montréal, Montréal, Québec, Canada H3C 3J7*
- ⁴⁹*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*
- ⁵⁰*Dipartimento di Scienze Fisiche and INFN, Università di Napoli Federico II, I-80126 Napoli, Italy*
- ⁵¹*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵²*University of Notre Dame, Notre Dame, Indiana 46556, USA*
- ⁵³*The Ohio State University, Columbus, Ohio 43210, USA*
- ⁵⁴*University of Oregon, Eugene, Oregon 97403, USA*
- ⁵⁵*Dipartimento di Fisica and INFN, Università di Padova, I-35131 Padova, Italy*
- ⁵⁶*Laboratoire de Physique Nucléaire et de Hautes Energies, Universités Paris VI et VII, F-75252 Paris, France*
- ⁵⁷*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁵⁸*Dipartimento di Fisica and INFN, Università di Perugia, I-06100 Perugia, Italy*
- ⁵⁹*Dipartimento di Fisica, Scuola Normale Superiore and INFN, Università di Pisa, I-56127 Pisa, Italy*
- ⁶⁰*Prairie View A&M University, Prairie View, Texas 77446, USA*
- ⁶¹*Princeton University, Princeton, New Jersey 08544, USA*
- ⁶²*Dipartimento di Fisica and INFN, Università di Roma La Sapienza, I-00185 Roma, Italy*
- ⁶³*Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁴*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom*
- ⁶⁵*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁶*University of South Carolina, Columbia, South Carolina 29208, USA*
- ⁶⁷*Stanford Linear Accelerator Center, Stanford, California 94309, USA*
- ⁶⁸*Stanford University, Stanford, California 94305-4060, USA*
- ⁶⁹*State University of New York, Albany, New York 12222, USA*
- ⁷⁰*University of Tennessee, Knoxville, Tennessee 37996, USA*
- ⁷¹*University of Texas at Austin, Austin, Texas 78712, USA*
- ⁷²*University of Texas at Dallas, Richardson, Texas 75083, USA*
- ⁷³*Dipartimento di Fisica Sperimentale and INFN, Università di Torino, I-10125 Torino, Italy*
- ⁷⁴*Dipartimento di Fisica and INFN, Università di Trieste, I-34127 Trieste, Italy*
- ⁷⁵*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*
- ⁷⁶*Vanderbilt University, Nashville, Tennessee 37235, USA*
- ⁷⁷*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*
- ⁷⁸*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*
- ⁷⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁸⁰*Yale University, New Haven, Connecticut 06511, USA*

(Received 8 February 2005; published 17 May 2005)

We present measurements of the $B \rightarrow \eta' K$ branching fractions; for $B^+ \rightarrow \eta' K^+$ we measure also the time-integrated charge asymmetry \mathcal{A}_{ch} , and for $B^0 \rightarrow \eta' K_S^0$ the time-dependent CP -violation parameters

S and C . The data sample corresponds to $232 \times 10^6 B\bar{B}$ pairs produced by e^+e^- annihilation at the $Y(4S)$. The results are $\mathcal{B}(B \rightarrow \eta' K^+) = (68.9 \pm 2.0 \pm 3.2) \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \eta' K^0) = (67.4 \pm 3.3 \pm 3.2) \times 10^{-6}$, $\mathcal{A}_{\text{ch}} = 0.033 \pm 0.028 \pm 0.005$, $S = 0.30 \pm 0.14 \pm 0.02$, and $C = -0.21 \pm 0.10 \pm 0.02$, where the first error quoted is statistical and the second is systematic.

DOI: 10.1103/PhysRevLett.94.191802

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Measurements of time-dependent CP asymmetries in B^0 meson decays through a Cabibbo-Kobayashi-Maskawa (CKM) favored $b \rightarrow c\bar{c}s$ amplitude [1] have provided a crucial test of the mechanism of CP violation in the standard model (SM) [2]. Such decays to a charmonium state plus a K^0 meson are dominated by a single weak phase. Decays of B^0 mesons to charmless hadronic final states, such as ϕK^0 , $K^+K^-K^0$, $\eta'K^0$, π^0K^0 , and $f_0(980)K^0$, proceed mostly via a single penguin (loop) amplitude with the same weak phase [3], but CKM-suppressed amplitudes and multiple particles in the loop introduce other weak phases whose contribution is not negligible [4–8].

For the decay $B^0 \rightarrow \eta' K^0$, these additional contributions are expected to be small, so the time-dependent asymmetry measurement for this decay provides an approximate measurement of $\sin 2\beta$. Theoretical bounds for the small deviation ΔS between the time-dependent CP -violating parameter S measured in this decay and in the charmonium K^0 decays have been calculated with an SU(3) analysis [4,5]. Such bounds have been improved by measurements of B^0 decays to a pair of neutral light pseudoscalar mesons [9,10]. From these and other measurements, improved SU(3) based bounds have been derived [6], with the conclusion that ΔS is expected to be less than 0.10 (with a theoretical uncertainty less than ~ 0.03). QCD factorization calculations conclude that ΔS is even smaller [7]. A significantly larger ΔS could arise from non-SM amplitudes [8].

The time-dependent CP -violating asymmetry in the decay $B^0 \rightarrow \eta' K^0$ has been measured previously by the BABAR [11] and Belle [12] experiments. In this Letter we update our previous measurements with an improved analysis and a data sample 4 times larger. We also measure the $B^0 \rightarrow \eta' K^0$ and $B^+ \rightarrow \eta' K^+$ branching fractions [13], and for $B^+ \rightarrow \eta' K^+$ the time-integrated charge asymmetry $\mathcal{A}_{\text{ch}} = (\Gamma^- - \Gamma^+)/(\Gamma^- + \Gamma^+)$ where $\Gamma^\pm = \Gamma(B^\pm \rightarrow \eta' K^\pm)$. In the SM \mathcal{A}_{ch} is expected to be small; a nonzero value would signal direct CP violation in this channel.

The data were collected with the BABAR detector [14] at the PEP-II asymmetric e^+e^- collider [15]. An integrated luminosity of 211 fb^{-1} , corresponding to $232 \times 10^6 B\bar{B}$ pairs, was recorded at the $Y(4S)$ resonance (center-of-mass energy $\sqrt{s} = 10.58 \text{ GeV}$). Charged particles are detected and their momenta measured by the combination of a silicon vertex tracker (SVT), consisting of five layers of double-sided detectors, and a 40-layer central drift chamber, both operating in the 1.5 T magnetic field of a solenoid. Charged-particle identification (PID) is provided by the average energy loss in the tracking devices and by an

internally reflecting ring-imaging Cherenkov detector (DIRC) covering the central region. Photons and electrons are detected by a CsI(Tl) electromagnetic calorimeter.

From a candidate $B\bar{B}$ pair we reconstruct a B^0 decaying into the flavor eigenstate $f = \eta' K_S^0 (B_{CP})$. We also reconstruct the vertex of the other B meson (B_{tag}) and identify its flavor. The difference $\Delta t \equiv t_{CP} - t_{\text{tag}}$ of the proper decay times t_{CP} and t_{tag} of the CP and tag B mesons, respectively, is obtained from the measured distance between the B_{CP} and B_{tag} decay vertices and from the boost ($\beta\gamma = 0.56$) of the e^+e^- system. The Δt distribution is given by

$$F(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} \{1 \mp \Delta w \pm (1 - 2w)[S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t)]\}. \quad (1)$$

The upper (lower) sign denotes a decay accompanied by a B^0 (\bar{B}^0) tag, τ is the mean B^0 lifetime, Δm_d is the mixing frequency, and the mistag parameters w and Δw are the average and difference, respectively, of the probabilities that a true B^0 is incorrectly tagged as a \bar{B}^0 or vice versa. The tagging algorithm [16] has seven mutually exclusive tagging categories of differing response purities (including one for untagged events that we retain for yield determinations). The measured analyzing power, defined as efficiency times $(1 - 2w)^2$ summed over all categories, is $(30.5 \pm 0.6)\%$, as determined from a large sample of B decays to fully reconstructed flavor eigenstates (B_{flav}). The parameter C measures direct CP violation. If $C = 0$, then $S = \sin 2\beta + \Delta S$.

Monte Carlo simulations [17] of the signal decay modes, $B\bar{B}$ backgrounds, and detector response are used to tailor the event selection criteria. We reconstruct B meson candidates by combining a K^+ or a K_S^0 with an η' meson. We reconstruct η' mesons through the decays $\eta' \rightarrow \rho^0 \gamma$ ($\eta'_{\rho\gamma}$) and $\eta' \rightarrow \eta \pi^+ \pi^-$ with $\eta \rightarrow \gamma \gamma$ ($\eta'_{(\gamma\gamma)\pi\pi}$) or $\eta \rightarrow \pi^+ \pi^- \pi^0$ ($\eta'_{(3\pi)\pi\pi}$). For the K^+ track we require an associated DIRC Cherenkov angle between -5 and $+2$ standard deviations (σ) from the expected value for a kaon. We select $K_S^0 \rightarrow \pi^+ \pi^-$ decays by requiring the $\pi^+ \pi^-$ invariant mass to be within 12 MeV of the nominal K^0 mass and by requiring a flight length with significance $> 3\sigma$. We select $K_S^0 \rightarrow \pi^0 \pi^0$ decays by requiring that the $\pi^0 \pi^0$ invariant mass be within 30 MeV of the nominal K^0 mass. Daughter pions from η' decays are required to have PID information inconsistent with proton, electron, and kaon hypotheses. The photon energy E_γ must be greater than 30 MeV for π^0 candidates, 50 (100) MeV for η candidates for the $\eta'_{(\gamma\gamma)\pi\pi} K^0$ ($\eta'_{(\gamma\gamma)\pi\pi} K^+$) samples,

and greater than 100 MeV for $\eta'_{\rho\gamma}$ candidates. We make the following requirements on the invariant mass (in MeV): $490 < m_{\gamma\gamma} < 600$ for $\eta \rightarrow \gamma\gamma$, $120 < m_{\gamma\gamma} < 150$ for π^0 ($100 < m_{\gamma\gamma} < 155$ in $K_S^0 \rightarrow \pi^0\pi^0$), $510 < m_{\pi\pi} < 1000$ for ρ^0 , $520 < m_{\pi\pi} < 570$ for $\eta \rightarrow \pi^+\pi^-\pi^0$, $945 < m_{\eta'} < 970$ for $\eta'_{\eta(\gamma\gamma)\pi\pi}$, and $930 < m_{\eta'} < 980$ for $\eta'_{\rho\gamma}$.

A B meson candidate is characterized kinematically by the energy-substituted mass $m_{\text{ES}} \equiv \sqrt{(\frac{1}{2}s + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - \mathbf{p}_B^2}$ and the energy difference $\Delta E \equiv E_B^* - \frac{1}{2}\sqrt{s}$, where (E_0, \mathbf{p}_0) and (E_B, \mathbf{p}_B) are four-momenta of the $Y(4S)$ and the B candidate, respectively, and the asterisk denotes the $Y(4S)$ rest frame. We require $|\Delta E| \leq 0.2$ GeV and $5.25 \leq m_{\text{ES}} \leq 5.29$ GeV.

To reject the dominant background from continuum $e^+e^- \rightarrow q\bar{q}$ events ($q = u, d, s, c$), we use the angle θ_T between the thrust axis of the B candidate and that of the rest of the tracks and neutral clusters in the event, calculated in the $Y(4S)$ rest frame. The distribution of $\cos\theta_T$ is sharply peaked near ± 1 for combinations drawn from jetlike $q\bar{q}$ pairs and is nearly uniform for the isotropic B decays; we require $|\cos\theta_T| < 0.9$. From Monte Carlo simulations of $B^0\bar{B}^0$ and B^+B^- events, we find evidence for a small (1%–2%) $B\bar{B}$ background contribution for the channels with $\eta' \rightarrow \rho^0\gamma$, so we have added a $B\bar{B}$ component to the fit described below for those channels.

We use an unbinned, multivariate maximum-likelihood fit to extract signal yields and CP -violation parameters. We indicate with j the species of event: signal, $q\bar{q}$ combinatorial background, or $B\bar{B}$ background. We use four discriminating variables: m_{ES} , ΔE , Δt , and a Fisher discriminant \mathcal{F} [18]. The Fisher discriminant combines four variables: the angles with respect to the beam axis of the B momentum and B thrust axis in the $Y(4S)$ rest frame, and the zeroth and second angular moments of the energy flow, excluding the B candidate, about the B thrust axis [19].

For each species j and tagging category c , we define a total probability density function (PDF) for event i as

$$\mathcal{P}_{j,c}^i \equiv \mathcal{P}_j(m_{\text{ES}}^i) \mathcal{P}_j(\Delta E^i) \mathcal{P}_j(\mathcal{F}^i) \mathcal{P}_j(\Delta t^i, \sigma_{\Delta t}^i; c), \quad (2)$$

where $\sigma_{\Delta t}^i$ is the error on Δt for event i . With n_j defined to be the number of events of the species j and $f_{j,c}$ the fraction of events of species j for each category c , we write the extended likelihood function for all events belonging to category c as

$$\mathcal{L}_c = \exp\left(-\sum_j n_{j,c}\right) \times \prod_i^{N_c} (n_{\text{sig}} f_{\text{sig},c} \mathcal{P}_{\text{sig},c}^i + n_{q\bar{q}} f_{q\bar{q},c} \mathcal{P}_{q\bar{q}}^i + n_{B\bar{B}} f_{B\bar{B},c} \mathcal{P}_{B\bar{B}}^i), \quad (3)$$

where $n_{j,c}$ is the yield of events of species j found by the fitter in category c and N_c the number of events of category c in the sample. We fix both $f_{\text{sig},c}$ and $f_{B\bar{B},c}$ to $f_{B_{\text{flav}},c}$ the

values measured with the large B_{flav} sample [20]. The total likelihood function \mathcal{L}_d for decay mode d is given as the product over the seven tagging categories. Finally, when combining decay modes we form the grand likelihood $\mathcal{L} = \prod_d \mathcal{L}_d$.

The PDF $\mathcal{P}_{\text{sig}}(\Delta t, \sigma_{\Delta t}; c)$, for each category c , is the convolution of $F(\Delta t; c)$ [Eq. (1)] with the signal resolution function (sum of three Gaussians) determined from the B_{flav} sample. The other PDF forms are the sum of two Gaussians for $\mathcal{P}_{\text{sig}}(m_{\text{ES}})$ and $\mathcal{P}_{\text{sig}}(\Delta E)$, the sum of three Gaussians for $\mathcal{P}_{q\bar{q}}(\Delta t; c)$, a conjunction of two Gaussians with different widths below and above the peak for $\mathcal{P}_j(\mathcal{F})$ [a small “tail” Gaussian is added for $\mathcal{P}_{q\bar{q}}(\mathcal{F})$], a linear dependence for $\mathcal{P}_{q\bar{q}}(\Delta E)$, and the function $x\sqrt{1-x^2} \times \exp[-\xi(1-x^2)]$ for $\mathcal{P}_{q\bar{q}}(m_{\text{ES}})$, with $x \equiv 2m_{\text{ES}}/\sqrt{s}$.

For the signal and $B\bar{B}$ background components we determine the PDF parameters from simulation. We study large control samples of B decays to charmed final states of similar topology to verify the simulated resolutions in ΔE and m_{ES} , adjusting the PDFs to account for any differences found. For the $q\bar{q}$ background we use $(m_{\text{ES}}, \Delta E)$ sideband data to obtain initial values and ultimately leave them free to vary in the final fit.

We compute the branching fractions and charge asymmetry from fits made without Δt or flavor tagging, applied to candidates with $\eta'_{\eta(\gamma\gamma)\pi\pi}$ and $\eta'_{\rho\gamma}$ combined with K^+ or $K_S^0 \rightarrow \pi^+\pi^-$. The free parameters in the fit are the signal and $q\bar{q}$ background yields, the peak position and lower and upper width parameters of $\mathcal{P}_j(\mathcal{F})$ for signal and $q\bar{q}$ background, the tail fraction for $\mathcal{P}_{q\bar{q}}(\mathcal{F})$, the slope of $\mathcal{P}_{q\bar{q}}(\Delta E)$ and ξ , the width of the core Gaussian of $\mathcal{P}_{\text{sig}}(\Delta E)$, the mean of the core Gaussian of $\mathcal{P}_{\text{sig}}(m_{\text{ES}})$, $n_{B\bar{B}}$ for $B \rightarrow \eta'_{\rho\gamma}K$, and the signal and background \mathcal{A}_{ch} for charged modes.

Table I lists the quantities used to determine the branching fraction. Equal production rates of B^+B^- and $B^0\bar{B}^0$ pairs have been assumed. To study biases from the likelihood fit, we apply the method to simulated samples constructed to contain the signal and background populations expected for data. The resulting yield biases, from unmodeled correlations in the signal PDF, are about 4% for the measurements with $\eta' \rightarrow \rho^0\gamma$ and negligible for those with $\eta'_{\eta(\gamma\gamma)\pi\pi}$. The purity estimate in Table I is given by the ratio of the signal yield to the effective background plus signal, the latter being defined as the square of the error on the yield.

In Fig. 1 we show projections onto m_{ES} and ΔE for a subset of the data for which the signal likelihood (computed without the variable plotted) exceeds a mode-dependent threshold that optimizes the sensitivity.

For the time-dependent analysis, we require $|\Delta t| < 20$ ps and $\sigma_{\Delta t} < 2.5$ ps. We improve the sample size by combining the five decay chains listed in Table II in a single fit with 98 free parameters: S , C , signal yields (5), $\eta'_{\rho\gamma}K^0 B\bar{B}$ background yields (2), continuum background

TABLE I. Signal yield, purity P (%), reconstruction efficiency ϵ (%), daughter branching fraction product, measured branching fraction (\mathcal{B}) in units of 10^{-6} , and \mathcal{A}_{ch} .

| Mode | Yield | P | ϵ | $\prod B_i$ | \mathcal{B} | $\mathcal{A}_{\text{ch}}(10^{-2})$ |
|--|---------------|-----|------------|-------------|---------------|------------------------------------|
| $\eta'_{\eta(\gamma\gamma)\pi\pi}K^+$ | 609 ± 28 | 78 | 23 | 0.175 | 66 ± 3 | -0.1 ± 4.4 |
| $\eta'_{\rho\gamma}K^+$ | 1347 ± 57 | 41 | 26 | 0.295 | 72 ± 3 | 5.5 ± 3.6 |
| $\eta'K^+$ | Combined | | | | 69 ± 2 | 3.3 ± 2.8 |
| $\eta'_{\eta(\gamma\gamma)\pi\pi}K^0_{\pi^+\pi^-}$ | 198 ± 16 | 77 | 23 | 0.060 | 61 ± 5 | |
| $\eta'_{\rho\gamma}K^0_{\pi^+\pi^-}$ | 457 ± 30 | 51 | 26 | 0.102 | 73 ± 5 | |
| $\eta'K^0$ | Combined | | | | 67 ± 3 | |

yields (5) and fractions (30), background Δt , m_{ES} , ΔE , \mathcal{F} PDF parameters (54). The parameters τ and Δm_d are fixed to world-average values [21].

Table II gives the yields, S and C , and Fig. 2 the Δt projections and asymmetry of the combined neutral modes for events selected as for Fig. 1.

The major systematic uncertainties affecting the branching fraction measurements reflect the imperfect knowledge of the η' branching fractions (3.4%) [21], and of the reconstruction efficiency (0.8% per charged track, 1.5% per photon, and 2.1% per K_S^0) estimated from auxiliary studies. We take one-half of the measured yield bias (0%–2%) as a systematic error. Bias and systematic uncertainties for \mathcal{A}_{ch} have been estimated from the values obtained for the background component in the fit to the data. We apply the resulting correction of +0.016 and include its systematic error of 0.005.

For the time-dependent measurements, we find approximately equal (0.01) systematic uncertainties from several sources: variation of the signal PDF shape parameters within their errors, SVT alignment, position and size of the beam spot, $B\bar{B}$ background, modeling of the signal Δt distribution, and interference between the CKM-suppressed $\bar{b} \rightarrow \bar{u}c\bar{d}$ amplitude and the favored $b \rightarrow c\bar{u}d$ amplitude for some tag-side B decays [22]. The B_{flav} sample is used to determine the errors associated with




FIG. 1 (color online). The B candidate m_{ES} and ΔE projections for $\eta'K^+$ (a),(b) and $\eta'K_S^0$ (c),(d). Points with error bars represent the data, the solid line the fit function, and the dashed line its background component.

TABLE II. Results with statistical errors for the $B^0 \rightarrow \eta'K_S^0$ time-dependent fits.

| Mode | Signal yield | S | C |
|--|--------------|------------------|------------------|
| $\eta'_{\eta(\gamma\gamma)\pi\pi}K^0_{\pi^+\pi^-}$ | 188 ± 15 | 0.01 ± 0.28 | -0.18 ± 0.18 |
| $\eta'_{\rho\gamma}K^0_{\pi^+\pi^-}$ | 430 ± 26 | 0.44 ± 0.19 | -0.30 ± 0.13 |
| $\eta'_{\eta(3\pi)\pi\pi}K^0_{\pi^+\pi^-}$ | 54 ± 8 | 0.79 ± 0.47 | 0.11 ± 0.35 |
| $\eta'_{\eta(\gamma\gamma)\pi\pi}K^0_{\pi^0\pi^0}$ | 44 ± 9 | -0.04 ± 0.57 | -0.65 ± 0.42 |
| $\eta'_{\rho\gamma}K^0_{\pi^0\pi^0}$ | 94 ± 23 | -0.45 ± 0.68 | 0.41 ± 0.40 |
| Combined fit | 804 ± 40 | 0.30 ± 0.14 | -0.21 ± 0.10 |

the signal Δt resolutions, tagging efficiencies, and mistag rates, and published measurements [21] for τ_B and Δm_d . Summing all systematic errors in quadrature, we obtain 0.02 for S and C .

In conclusion, we have used samples of about 2000 $\eta'K^+$ and 800 $\eta'K_S^0$ events to measure the branching fractions, the time-integrated charge asymmetry and the time-dependent asymmetry parameters S and C . The measured branching fractions are $\mathcal{B}(B^+ \rightarrow \eta'K^+) = (68.9 \pm 2.0 \pm 3.2) \times 10^{-6}$ and $\mathcal{B}(B^0 \rightarrow \eta'K^0) = (67.4 \pm 3.3 \pm 3.2) \times 10^{-6}$, and the charge asymmetry is $\mathcal{A}_{\text{ch}} = 0.033 \pm 0.028 \pm 0.005$. These precise branching fraction measurements help the theoretical understanding of these decays [23]. The measured charge asymmetry is consistent with zero, with 90% C.L. interval $[-0.012, 0.078]$, and constrains the amount of possible direct CP violation in $B^+ \rightarrow \eta'K^+$ decays.

The measured time-dependent CP violation parameters in $B^0 \rightarrow \eta'K_S^0$ are $S = 0.30 \pm 0.14 \pm 0.02$ and $C = -0.21 \pm 0.10 \pm 0.02$. Our result for S differs from that measured by *BABAR* in $B^0 \rightarrow J/\psi K_S^0$ [16] by 3.0 standard deviations; it also represents an improvement by a factor of 2.4 (1.9) in precision over the published results of *BABAR*




FIG. 2 (color online). Projections onto Δt for $\eta'K_S^0$ of the data (points with error bars), fit function (solid line), and background function (dashed line), for (a) B^0 and (b) \bar{B}^0 tagged events, and (c) the asymmetry between B^0 and \bar{B}^0 tags.

[11] (Belle [12]). All these measurements supersede our previous published results [11].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A.P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

*Also with Università della Basilicata, Potenza, Italy.

†Deceased.

- [1] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **89**, 201802 (2002); K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **66**, 071102(R) (2002).
- [2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] Y. Grossman and M.P. Worah, Phys. Lett. B **395**, 241 (1997); D. Atwood and A. Soni, Phys. Lett. B **405**, 150 (1997).
- [4] Y. Grossman *et al.*, Phys. Rev. D **68**, 015004 (2003).
- [5] C.-W. Chiang, M. Gronau, and J.L. Rosner, Phys. Rev. D **68**, 074012 (2003).
- [6] M. Gronau, J.L. Rosner, and J. Zupan, Phys. Lett. B **596**, 107 (2004).
- [7] M. Beneke and M. Neubert, Nucl. Phys. **B675**, 333 (2003).
- [8] D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
- [9] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **93**, 181806 (2004).
- [10] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **70**, 032006 (2004).
- [11] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **91**, 161801 (2003).
- [12] K. Abe *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 261602 (2003).
- [13] Charge conjugate decay modes are implied unless explicitly stated.
- [14] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [15] PEP-II Conceptual Design Report No. SLAC-R-418, 1993.
- [16] B. Aubert *et al.* (*BABAR* Collaboration), hep-ex/0408127.
- [17] The *BABAR* detector Monte Carlo simulation is based on GEANT4: S. Agostinelli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [18] R. A. Fisher, Ann. Eugenics **7**, 179 (1936).
- [19] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **92**, 061801 (2004).
- [20] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **66**, 032003 (2002).
- [21] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [22] O. Long, M. Baak, R.N. Cahn, and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).
- [23] See, for example, M. Beneke and M. Neubert, Nucl. Phys. **B651**, 225 (2003); E. Kou and A. I. Sanda, Phys. Lett. B **525**, 240 (2002); C-W. Chiang and J.L. Rosner, Phys. Rev. D **65**, 074035 (2002), and references therein; M-Z. Yang and Y-D. Yang, Nucl. Phys. **B609**, 469 (2001); E. V. Shuryak and A. R. Zhitnitsky, Phys. Rev. D **57**, 2001 (1998).