



WORKING PAPERS

Col·lecció d'Economia E19/395

Assortative multisided assignment games. The extreme core points

Javier Martinez de Albéniz
Carlos Rafels
Neus Ybern



UNIVERSITAT DE
BARCELONA

Assortative multisided assignment games. The extreme core points

Abstract: We analyze assortative multisided assignment games, following Sherstyuk (1999) and Martínez-de-Albéniz et al. (2019). In them players' abilities are complementary across types (i.e. supermodular), and also the output of the essential coalitions is increasing depending on types.

We study the extreme core points and show a simple mechanism to compute all of them. In this way we describe the whole core. This mechanism works from the original data array and the maximum number of extreme core points is obtained.

JEL Codes: C71.

Keywords: Assortative market, assignment game, multisided assignment game, core, extreme core allocations.

F. Javier Martinez de Albéniz
Universitat de Barcelona

Carlos Rafels
Universitat de Barcelona

Neus Ybern
Universitat Politècnica de Catalunya

Acknowledgements: Financial support from research grant ECO2017-86481-P (Ministerio de Economía y Competitividad, AEI/FEDER, UE) is gratefully acknowledged, and from the Generalitat de Catalunya, through grant 2017SGR778.

1. Introduction

Economic markets with indivisible goods have been considered using worthy matching models. In this setting there are different but related models. In a two-sided matching game or assignment game there are essential coalitions formed from two different types of agents and these essential coalitions are the singletons and doubletons containing one agent of each type. The bilateral assignment game comes initially from Shapley (1955), but Shapley and Shubik (1971) is the paper most cited. In it the authors introduce and analyze a housing market as a bilateral assignment market. We refer now to another seminal paper, Becker (1973). In it, pursuing a general theory of marriage, Becker introduces a special class of assignment games, the two-sided assortative ones. In some assignment problems Becker displays the well-known effect of *mating of the likes*. Finally, Crawford and Knoer (1981) develops a model of labor market by using matching and assignment tools. This last model easily allows to motivate the relevance to study m -sided matching games, with $m \geq 3$. It is easy to think of situations where m types of different skills' workers are needed to achieve valuable essential coalitions. Precisely the main purpose of this paper is to analyze m -sided assortative games.

In all these previous models the most relevant set solution is the core. Roughly speaking the core is formed by all those allocations in which no coalition of agents can improve its reward on its own. Although the core of two-sided assignment games has been extensively studied, and important properties are known: non-emptiness, its lattice structure, the side-optimal core allocations, etc., the core of m -sided assignment games, $m \geq 3$, has not got the same attention. There are only a few papers on it. Many

difficulties arise when moving from two-sided to m -sided problems. Most of the results fall in the negative which in our opinion does not mean the subject is unimportant.

In this paper we develop positive results. Firstly we analyze a simple mechanism to describe the whole core of any assortative m -sided assignment game. Our method characterizes for the first time all the extreme core allocations of any assortative m -sided matching game. The procedure can be applied for the two-sided case as well as the generic m -sided case. The mechanism depends only on the assignment array data, with no need to compute the characteristic function of the game. We give also the maximum number of extreme core allocations, $m \cdot (m!)^{n-1}$, where m is the number of sectors and n is the number of agents in each sector. As a by-product we obtain the number of extreme core allocations when we deal only with two sectors, 2^n . Finally our mechanism is an extension of the one recently published for the two-sided assortative assignment games (Martínez-de-Albéniz et al., 2019). It is simplified in some features and proofs are completely different. The two-sided assortative case was also analyzed in Eriksson et al. (2000) where they show that the core is ordered in payoffs inside each sector. We prove that this property remains true for the general m -sided case.

Multisided assignment games were analyzed for the first time in Quint (1991). After showing a three-sided example with an empty core, Quint presents a class of games with the property that the core is non-empty, i.e. balanced. Stuart (1997) proposes another balanced class of multisided assignment games, not related to Quint's class (none of them includes the other). A proof of the non-emptiness is provided, but no description or characterization of the core is given in any of the two models.

Sherstyuk (1999) introduces another important class of m -sided match-

ing games. She analyzes for the first time the assortative multisided assignment games. The definition of this class relies on two conditions imposed on the assignment array: supermodularity and monotonicity. Both conditions assume that agents in each sector can be ranked by some trait or ability. Supermodularity is a complementary property of agents' ability across types. Monotonicity means that ability is aligned with the worth generated by the essential coalitions.

Assortative multisided assignment games form a large class of m -sided assignment games: a full-dimensional cone. In Sherstyuk's paper it is proved the non-emptiness of the core and she describes some extreme core allocations, $m!$ of them, by using the associated characteristic function.

2. Preliminaries on the multisided assignment markets

A multisided assignment market $(N^1, N^2, \dots, N^m; A)$ is formed by m non-empty pairwise disjoint finite sets of agents, $N^k = \{1^k, 2^k, \dots, n_k^k\}$ ¹ for $k \in M = \{1, \dots, m\}$ and a non-negative m -dimensional array $A = (a_E)_{E \in \prod_{k=1}^m N^k}$. Each entry a_E represents some measure of the joint productivity of agents in $E = (i_1, i_2, \dots, i_m) \in \prod_{k=1}^m N^k$, one of each set when they are matched together. We assume that we need exactly one agent of each type to realize the value of a transaction. Each set N^k is called a sector and corresponds to a different type of agents, having different skills. Any m -tuple of agents $E = (i_1, \dots, i_m) \in \prod_{k=1}^m N^k$ is called an *essential coalition* and we use E either as the m -tuple or as the set of elements formed by its components. In the case of two sectors, $m = 2$, matrix A is known as the

¹To simplify notation, when no confusion arises, we will drop the superscript to describe the agents in N^k , i.e. $N^k = \{1, 2, \dots, n_k\}$. Its cardinality is $|N^k| = n_k$.

assignment matrix (Shapley and Shubik, 1972). When the number of agents is the same in each sector $|N^1| = |N^2| = \dots = |N^m|$ the assignment market is said to be square.

A *matching* μ among N^1, \dots, N^m is a set of essential coalitions such that any agent belongs at most to one coalition in μ , and $|\mu| = \min_{k \in M} |N^k|$. An agent who does not belong to any of the essential coalitions of μ is unmatched by μ . The set of all matchings is denoted by $\mathcal{M}(N^1, \dots, N^m)$. A matching μ is *optimal* if it maximizes $\sum_{E \in \mu} a_E$ over the set $\mathcal{M}(N^1, \dots, N^m)$. The set of all optimal matchings is denoted by $\mathcal{M}_A^*(N^1, \dots, N^m)$.

Shapley and Shubik (1971) associates any bilateral assignment market with a cooperative game², the *assignment game*. In the multisided assignment game (Quint, 1991), the set of players is $N = \bigcup_{k=1}^m N^k$ and the characteristic function w_A is defined for any $S \subseteq N$ such that $S \cap N^k \neq \emptyset$ for all $k \in M$, by

$$w_A(S) = \max_{\mu \in \mathcal{M}(S \cap N^1, \dots, S \cap N^m)} \sum_{E \in \mu} a_E, \quad \text{and } 0 \text{ otherwise.}$$

Notice that any essential coalition evaluates its worth by exactly the corresponding entry, and any other coalition determines its worth by essential coalition combinations its members can form.

The agents of a multisided assignment market may divide among themselves their worth, $w_A(N)$, in any way they like. Thus an *allocation* is a non-negative vector $x = (x^1, x^2, \dots, x^m) \in \prod_{k=1}^m \mathbb{R}_+^{n_k}$. Vector $x^k \in \mathbb{R}_+^{n_k}$ is interpreted as the payoffs to agents in N^k , i.e. x_i^k is the payoff associated to

²In a cooperative game (N, v) , the set of players is given by $N = \{1, \dots, n\}$ and v is a function that assigns a real number $v(S)$ for any coalition $S \subseteq N$ with $v(\emptyset) = 0$. Its core is defined as $C(v) := \{x \in \mathbb{R}^n \mid \sum_{i \in N} x_i = v(N) \text{ and for all } S \subseteq N, \sum_{i \in S} x_i \geq v(S)\}$. A game is named balanced if its core is non-empty.

player i of sector k . For any essential coalition $E = (i_1, \dots, i_m) \in \prod_{k=1}^m N^k$ we write $x(E) = \sum_{k=1}^m x_{i_k}^k$.

The core of the multisided assignment game $C(w_A)$ is described for any fixed optimal matching $\mu \in \mathcal{M}_A^*(N^1, \dots, N^m)$ as those allocations $x \in \prod_{k=1}^m \mathbb{R}_+^{n_k}$ satisfying

$$\begin{aligned} x(E) &= a_E \quad \text{for all } E \in \mu, \\ x(E) &\geq a_E \quad \text{for all } E \notin \mu, \end{aligned}$$

and unassigned agents by μ receive a zero payoff in any core allocation.

In the two-sided case, Shapley and Shubik (1971) proves that the core of any assignment game is always non-empty, but in the multisided case, $m \geq 3$, it is known (Kaneko and Wooders, 1982, or Quint, 1991) that the core may be empty.

Becker (1973) introduces two-sided assortative assignment markets. For multisided assignment markets, we assume that the elements of each sector are ordered by some trait and then N^k for $k \in M$ is an ordered set with the natural order. Therefore $\prod_{k=1}^m N^k$ is a lattice and for any pair of essential coalitions $E, E' \in \prod_{k=1}^m N^k$ we can define $E \vee E'$ as the maximum component-wise and $E \wedge E'$ as the minimum component-wise.

A multisided assignment market $(N^1, N^2, \dots, N^m; A)$ is an *assortative market* if it satisfies:

a) supermodularity:³

$$a_E + a_{E'} \leq a_{E \vee E'} + a_{E \wedge E'} \quad \text{for all } E, E' \in \prod_{k=1}^m N^k. \quad (1)$$

³Notice that this condition implies that array entries form a supermodular function in the lattice $N^1 \times N^2 \times \dots \times N^m$ with the usual order (see Topkis, 1998).

b) monotonicity (non-decreasing rows, columns, etc.):

$$a_E \leq a_{E'} \quad \text{for all } E \leq E', \quad E, E' \in \Pi_{k=1}^m N^k. \quad (2)$$

Whenever these two conditions are met, array A is called assortative.

From the supermodularity condition, in a multisided assortative assignment market at least one optimal matching $\mu \in \mathcal{M}_A^*(N^1, \dots, N^m)$ is *monotone*,⁴ i.e.

$$\text{for any } E, E' \in \mu, \quad \text{either } E \leq E' \text{ or } E' \leq E.$$

When the assortative assignment market is square, $|N^1| = |N^2| = \dots = |N^m| = n$ there is only one monotone matching which is placed in the main diagonal. If we denote the following essential coalitions: $E_i = (i, i, \dots, i)$, for $i = 1, 2, \dots, n$, this monotone matching is $\mu = \{E_1, E_2, \dots, E_n\}$. This is, by the previous observation, optimal in the square supermodular case, maybe not unique.

From now on, we concentrate in the square case, since any non-square assortative array could be analyzed by adding null rows of entries at the beginning of the array, to make it square. In this way we preserve supermodularity and the monotonicity conditions.

We give some new features of any square multisided assortative assignment market. To this end, the *central strip* in a square multisided assignment market are those essential coalitions

$$E = (i_1, i_2, \dots, i_m) \quad \text{such that} \quad \max_{k \in M} i_k - \min_{k \in M} i_k \leq 1.$$

⁴Notice that if there are two essential coalitions E, E' of μ that are not comparable, we can use supermodularity to obtain a new optimal matching with $E \vee E'$ and $E \wedge E'$. Sherstyuk (1999) calls such a matching consecutive.

or equivalently those essential coalitions such that

$$E_{i-1} \leq E \leq E_i \quad \text{for } i = 2, \dots, n. \quad (3)$$

Theorem 2.1. *For any square multisided assortative assignment market $(N^1, N^2, \dots, N^m; A)$ we have:*

(a) *The main diagonal of the assignment array A is an optimal matching (maybe not unique).*

(b) *An allocation $x \in \prod_{k=1}^m \mathbb{R}_+^{n_k}$ belongs to the core $C(w_A)$ if and only if⁵*

$$(b1) \quad x(E) = a_E \quad \text{for all } E = E_1, E_2, \dots, E_n, \quad (4)$$

$$(b2) \quad x(E) \geq a_E \quad \text{for all } E \in \prod_{k=1}^m N^k \text{ such that} \\ E_{i-1} < E < E_i \quad \text{for } i = 2, \dots, n. \quad (5)$$

(c) *At any core allocation $x \in C(w_A)$ we have for all $k \in M$*

$$0 \leq x_1^k \leq x_2^k \leq \dots \leq x_n^k.$$

Proof. Item (a) follows by our previous comments. To prove (b) assume that $x \in \prod_{k=1}^m \mathbb{R}_+^{n_k}$ satisfies (4) and (5). We prove that $x(E) \geq a_E$ for all essential coalitions $E = (i_1, i_2, \dots, i_m)$ by induction on $r = \max_{k \in M} i_k - \min_{k \in M} i_k$. Assume the induction hypothesis: If E is such that $\max_{k \in M} i_k - \min_{k \in M} i_k \leq r$ then $x(E) \geq a_E$. Notice that for $r = 1$ the inequalities are just (4) and (5). Let $E = (i_1, i_2, \dots, i_m)$ such that $\max_{k \in M} i_k - \min_{k \in M} i_k = r \geq 2$. Denote $j = 1 + \min_{k \in M} i_k$. Then, by supermodularity, $a_E + a_{E_j} \leq a_{E \wedge E_j} + a_{E \vee E_j}$. Clearly $E \wedge E_j$ belongs to the central strip, $E \vee E_j$ satisfies the induction hypothesis, and $x(E_j) = a_{E_j}$. Therefore, $a_E \leq x(E \wedge E_j) + x(E \vee E_j) -$

⁵We denote $E < E'$ for $E \leq E'$ and $E \neq E'$.

$x(E_j) = x(E)$. To see (c), assume for instance $x \in C(w_A)$. Then for $i = 1, \dots, n-1$ we have $x(E_i) = a_{E_i}$, and take the essential coalition E' given by $(i+1, i, \dots, i)$. Then we have $\sum_{k=1}^m x_i^k = a_{E_i}$, and $x_{i+1}^1 + \sum_{k=2}^m x_i^k \geq a_{E'}$. Thus, $0 \leq a_{E'} - a_{E_i} \leq x_{i+1}^1 - x_i^1$. \square

Notice that item (b) means that only the central strip of array A is necessary to determine the core conditions. Item (c) means that in any square assortative market, payoffs in the core are such that for any sector, agents are ranked in the same way.

Remark 2.1. *Looking at the proof of Theorem 2.1, notice that the proof of items (a) and (b) only uses the supermodularity condition (1) of the assignment array.*

Item (c) is implied by the monotonicity condition (2) and the fact that we have an optimal matching in the main diagonal. It could be interesting to know which conditions on the array A characterize the results of the above theorem.

A different proof of item (b) in the supermodular two-sided case can be found in Martínez-de-Albéniz and Rafels (2014). The fact that payoffs to agents in the core are ordered is known for two-sided assortative matrices (see Eriksson et al., 2000).

3. Extreme core allocations

Now we give a simple procedure to obtain all the extreme core points. To this end, for notational convenience we introduce, for any square assortative multisided assignment market, an auxiliary agent 0 for any sector. We denote $E_0 = (0, 0, \dots, 0)$ with $a_{E_0} = 0$ and also for any E such that $E_0 < E < E_1$ we denote $a_E = 0$.

A *path* p is a sequence of essential coalitions connecting the initial one E_0 with the last one E_n passing through all essential coalitions E_0, E_1, \dots, E_n where $E_i = (i, i, \dots, i)$ for $i = 0, 1, \dots, n$. Moreover, between E_{i-1} and E_i , $i = 1, \dots, n$, the essential coalitions are such that from one essential coalition to the next one we change the agent of only one sector, moving from agent $i - 1$ to agent i . Then path p is

$$p = (E_0, \dots, E_1, \dots, E_{i-1}, E_i^1, E_i^2, \dots, E_i^{m-1}, E_i, \dots, E_n),$$

where $E_{i-1} < E_i^1 < E_i^2 < \dots < E_i^{m-1} < E_i$, for $i = 1, 2, \dots, n$. As a consequence, these paths are included in the central strip, see (3). Given a path p , notice that each block $E_{i-1} < E_i^1 < E_i^2 < \dots < E_i^{m-1} < E_i$, for $i = 1, 2, \dots, n$ can also be described by a particular permutation $\theta^i \in \Theta(M)$ indicating the order of the sectors that are sequentially increased. The set of all paths is denoted by \mathcal{P}_n^m .

For each path $p \in \mathcal{P}_n^m$ we associate an allocation vector, which we name the p -vector, $x^p \in \prod_{k=1}^m \mathbb{R}_+^{n_k}$ by solving the linear equations given by all the places of the selected path

$$x^p(E) = a_E \quad \text{for } E \text{ belonging to } p, \quad (6)$$

where we use $(x^p)_0^k = 0$, for $k = 1, \dots, m$, that is any auxiliary agent 0 gets a null payoff.

For each path p the above linear system has a unique non-negative solution. We prove uniqueness and non-negativeness by induction over n . Firstly notice that if $n = 1$ there are $m!$ different paths between E_0 and E_1 , but vector x^p is $a_{E_1} e_k$ for some $k \in M$ where e_k is the canonical vector. Assume that the solution is unique and non-negative up to E_{i-1} , and without loss of generality assume that the next essential coalition E_i^1 of path p is

$(i, i-1, \dots, i-1)$. Then by (6) we have

$$\sum_{k=1}^m x_{i-1}^k = a_{E_{i-1}}, \quad \text{and}$$

$$x_i^1 + \sum_{k=2}^m x_{i-1}^k = a_{E_i^1},$$

where we have dropped the superscript p for the path. Then, using the monotonicity (2) and the induction hypothesis we obtain

$$x_i^1 = x_{i-1}^1 + (a_{E_i^1} - a_{E_{i-1}}) \geq x_{i-1}^1 \geq 0.$$

Therefore for each path $p \in \mathcal{P}_n^m$ we have a unique and non-negative p -vector.

Now let us write $\mathcal{E}xt(C(w_A))$ the set of all extreme core points.⁶ We prove next that any extreme core point is linked to a path, that is, there is a correspondence between paths and extreme core points. This is our following theorem, but we need some lemmas and notation.

Lemma 3.1. *Let $(N^1, N^2, \dots, N^m; A)$ be a square multisided assortative assignment market. For any extreme core point $x \in C(w_A)$ we have $x_1^{k^*} = a_{E_1}$ for some $k^* \in M$ and $x_1^k = 0$ for all $k \in M \setminus \{k^*\}$.*

Proof. Suppose, on the contrary, that there are two sectors, $k', k'' \in M$ such that $x_1^{k'} > 0$ and $x_1^{k''} > 0$ and define $\varepsilon = \min\{x_1^{k'}, x_1^{k''}\} > 0$. Now define $y, z \in \prod_{k=1}^m \mathbb{R}_+^{n_k}$ as follows, for $t = 1, \dots, n$,

$$y_t^k = \begin{cases} x_t^k, & \text{for } k \in M \setminus \{k', k''\}, \\ x_t^{k'} + \varepsilon, & \text{for } k = k', \\ x_t^{k''} - \varepsilon, & \text{for } k = k'', \end{cases} \quad z_t^k = \begin{cases} x_t^k, & \text{for } k \in M \setminus \{k', k''\}, \\ x_t^{k'} - \varepsilon, & \text{for } k = k', \\ x_t^{k''} + \varepsilon, & \text{for } k = k''. \end{cases}$$

⁶If $X \subseteq \mathbb{R}^n$ is a convex set, an element of this convex set $x \in X$ is an extreme point if $x = \frac{1}{2}y + \frac{1}{2}z$ for some $y, z \in X$, then $x = y = z$.

Clearly by Theorem 2.1(c) and the definition of ε these are non-negative vectors, and since $y(E) = x(E)$ and $z(E) = x(E)$ for all essential coalitions E , we have $y, z \in C(w_A)$. As a consequence $x = \frac{1}{2}y + \frac{1}{2}z$ with $y \neq x$ and $z \neq x$, getting a contradiction with the fact that x is an extreme core point. \square

Now we introduce for any $i \in \{1, 2, \dots, n\}$ the submarket given by all the first i agents from any sector, and the corresponding restricted array. Formally, that is $(N_i^1, N_i^2, \dots, N_i^m; A^i)$ where $N_i^k = \{1, \dots, i\}$ for all $k \in M$ and A^i is given by $A^i = (a_E)_{E \in \prod_{k=1}^m N_i^k}$. Each of these markets is assortative and an optimal matching is given by the main diagonal when the original market is assortative and square.

Next we relate the extreme core points of these markets with our original square multisided assortative assignment market. To this end, for each $x \in C(w_A)$ we denote by \bar{x}^i the restriction of vector x to the coordinates of $\prod_{k=1}^m N_i^k$, i.e.

$$\bar{x}^i = (x_1^1, \dots, x_i^1, x_1^2, \dots, x_i^2, \dots, x_1^m, \dots, x_i^m) \in \prod_{k=1}^m \mathbb{R}_+^{N_i^k}. \quad (7)$$

Clearly $\bar{x}^i \in C(w_{A^i})$ for all $i \in \{1, 2, \dots, n\}$ if $x \in C(w_A)$.

In our next lemma we prove that whenever we take an extreme core point we obtain an extreme core point, given by the restriction, for all submarkets previously defined.

Lemma 3.2. *Let $(N^1, N^2, \dots, N^m; A)$ be a square multisided assortative assignment market, and $x \in C(w_A)$ be an extreme core point. Then, $\bar{x}^i \in C(w_{A^i})$ is an extreme core point for $i = 1, \dots, n - 1$.*

Proof. Suppose on the contrary that $i^* \in \{1, \dots, n - 1\}$ is the first index such that \bar{x}^{i^*} is not an extreme point of $C(w_{A^{i^*}})$. By Lemma 3.1, $i^* > 1$.

Since we are assuming $\bar{x}^{i^*} \in C(w_{A^{i^*}})$ but not an extreme core point, there are two points $y^*, z^* \in C(w_{A^{i^*}})$ such that

$$\bar{x}^{i^*} = \frac{1}{2}y^* + \frac{1}{2}z^* \quad \text{with} \quad y^* \neq \bar{x}^{i^*} \quad \text{and} \quad z^* \neq \bar{x}^{i^*}. \quad (8)$$

Notice that for all $i < i^*$ we have $y_i^k = z_i^k = x_i^k$ for all $k \in M$, because the corresponding restriction \bar{x}^{i^*-1} gives an extreme core point.

Now define the following vectors $y, z \in \prod_{k=1}^m \mathbb{R}^{n_k}$ as follows: for all $k \in M$,

$$y_i^k = \begin{cases} x_i^k, & \text{for } i = 1, \dots, i^* - 1, \\ x_i^k + \varepsilon^k, & \text{for } i = i^*, \dots, n, \end{cases} \quad z_i^k = \begin{cases} x_i^k, & \text{for } i = 1, \dots, i^* - 1, \\ x_i^k - \varepsilon^k, & \text{for } i = i^*, \dots, n. \end{cases}$$

where $\varepsilon^k = (y^*)_{i^*}^k - x_{i^*}^k$ for all $k \in M$. Notice that because of (8), at least one ε^k must be different from zero, and we have $(z^*)_{i^*}^k - x_{i^*}^k = -\varepsilon^k$ for all $k \in M$. Moreover

$$\sum_{k \in M} \varepsilon^k = \sum_{k \in M} (y^*)_{i^*}^k - x_{i^*}^k = y^*(E_{i^*}) - x(E_{i^*}) = a_{E_{i^*}} - a_{E_{i^*}} = 0. \quad (9)$$

We claim $y, z \in C(w_A)$ and $x = \frac{1}{2}y + \frac{1}{2}z$ with $y \neq x$ and $z \neq x$.

Firstly we show $y \geq 0$ and $z \geq 0$. Clearly $y_i^k \geq 0$ and $z_i^k \geq 0$ for $i = 1, \dots, i^* - 1$, and all $k \in M$. Moreover, for all $k \in M$ we have $y_n^k \geq y_{n-1}^k \geq \dots \geq y_{i^*}^k$ and $z_n^k \geq z_{n-1}^k \geq \dots \geq z_{i^*}^k$, and to conclude notice that $y_{i^*}^k = x_{i^*}^k + \varepsilon^k = (y^*)_{i^*}^k \geq 0$ and also $z_{i^*}^k \geq 0$.

Secondly, $y(E_i) = a_{E_i}$, $z(E_i) = a_{E_i}$ for $i = 1, \dots, n$, by their definitions.

Finally, we show that $y(E) \geq a_E$ and $z(E) \geq a_E$ for all essential coalitions E in the central strip. For all essential coalitions in the central strip such that $E_{i^*} \leq E$, by (9) $y(E) = x(E) + \sum_{k \in M} \varepsilon^k = x(E) \geq a_E$ and analogously $z(E) \geq a_E$. By its definition $y(E) = z(E) = x(E) \geq a_E$ for all essential coalitions E , in the central strip such that $E \leq E_{i^*-1}$. For the case $E_{i^*-1} < E < E_{i^*}$, we claim that $y(E) = y^*(E)$ and $z(E) = z^*(E)$, since we have that $y_{i^*}^k = (y^*)_{i^*}^k$ and $z_{i^*}^k = (z^*)_{i^*}^k$ for all $k \in M$.

By Theorem 2.1(b) we have $y, z \in C(w_A)$ and $x = \frac{1}{2}y + \frac{1}{2}z$ with $y \neq x$ and $z \neq x$, contradicting x is an extreme core point. \square

These two lemmas allow to establish our main theorem.

Theorem 3.1. *Let $(N^1, N^2, \dots, N^m; A)$ be a square multisided assortative assignment market. In it, p -vectors coincide with extreme core points, i.e.*

$$\mathcal{E}xt(C(w_A)) = \{x^p\}_{p \in \mathcal{P}_n^m}.$$

Proof. We prove first that for all path $p \in \mathcal{P}_n^m$ we have $x^p \in C(w_A)$. To this end we prove $x^p(E) \geq a_E$ for all $E_{i-1} < E < E_i$ for all $i = 1, \dots, n$. By Theorem 2.1(b) this is enough to justify $x^p \in C(w_A)$.

Without loss of generality we assume that the essential coalitions of path p between E_{i-1} and $E_i, i = 1, \dots, n$, are given by

$$E_{i-1}, (i, i-1, \dots, i-1), (i, i, i-1, \dots, i-1), \dots, E_i, \quad (10)$$

that is, they follow the natural order of sectors, first moves the first sector, second the second sector and so forth. We denote by $E_i^t = (i, \dots, i, \overset{t}{i}, i-1, i-1, \dots, i-1), 1 \leq t \leq m-1$, the essential coalition in the previous path such that t is the position of the last i agent, $i = 1, \dots, n$. As a matter of notation, $E_i^0 = E_{i-1}$ and $E_i^m = E_i$.

Given any essential coalition $E = (i_1, i_2, \dots, i_m)$ with $E_{i-1} < E < E_i, i = 1, \dots, n$, we define $r(E) = \#\{k \mid i_k = i\}$, the number of i agents in the essential coalition E . Now, we prove $x^p(E) \geq a_E$ with $E_{i-1} < E < E_i$ by induction on the number $r(E)$. Clearly $1 \leq r(E) \leq m-1$. If $r(E) = 1$ let l be the position of the only i . If $l = 1$ there is nothing to prove, and if $l > 1$ notice that $E \wedge E_i^{l-1} = E_{i-1}$ and $E \vee E_i^{l-1} = E_i^l$. Therefore, by supermodularity and the way essential coalitions of path p have been chosen,

$a_E + a_{E_i^{l-1}} \leq a_{E_{i-1}} + a_{E_i^l}$, and then $a_E + x^p(E_i^{l-1}) \leq x^p(E_{i-1}) + x^p(E_i^l)$. Now clearly $x^p(E) \geq a_E$. Assume our induction hypothesis is true up to $r-1$ and let E be such that $r(E) = r$. There are then r positions with agent i and let l be the last of these positions. Then $a_E + a_{E_i^{l-1}} \leq a_{E \wedge E_i^{l-1}} + a_{E_i^l}$, by supermodularity. We can apply the induction hypothesis to $E \wedge E_i^{l-1}$ since it has $r(E \wedge E_i^{l-1}) = r-1$ positions with an i . Now $a_E + x(E_i^{l-1}) \leq a_{E \wedge E_i^{l-1}} + a_{E_i^l} \leq x(E \wedge E_i^{l-1}) + x(E_i^l)$, and therefore $a_E \leq x(E)$ to finish with this part of the proof.

Moreover, vector x^p for $p \in \mathcal{P}_n^m$ is an extreme core point. To see it, just notice that if it were the midpoint of two other core points, these core points must satisfy with equality all the entries of path p . By uniqueness of the solution, they coincide with x^p . We have established that each path gives an extreme core point.

Now we prove that any extreme core point is associated to some path. Let $x \in C(w_A)$ be an extreme core point. Then by Lemma 3.2, \bar{x}^i is also an extreme core point of $C(w_{A_i})$ for all $i \in \{1, \dots, n\}$, see (7) for notations.

Suppose on the contrary that x is not a p -vector for any path $p \in \mathcal{P}_n^m$, and let $i^* \in \{1, \dots, n\}$ be the first index such that \bar{x}^{i^*} is not a p -vector for any $p \in \mathcal{P}_{i^*}^m$. Notice that $|N_{i^*}^1| = |N_{i^*}^2| = \dots = |N_{i^*}^m| = i^*$.

Clearly, by Lemma 3.1, $i^* > 1$ since any path between E_0 and E_1 gives a_{E_1} to some agent and zero to the others. Vector \bar{x}^{i^*-1} is a p -vector for some path $p_{i^*-1} \in \mathcal{P}_{i^*-1}^m$ and consider the set of paths in $\mathcal{P}_{i^*}^m$ that coincide with p_{i^*-1} for all essential coalitions in the central strip $E \leq E_{i^*-1}$. Denote this set by \mathcal{B}_{i^*} .

Consider now the set given by convex hull of the p -vectors corresponding to paths in \mathcal{B}_{i^*} , that is $Conv\{x^p\}_{p \in \mathcal{B}_{i^*}}$. This is a non-empty, compact and convex set and clearly vector \bar{x}^{i^*} cannot be a convex combination of these

core points $\{x^p\}_{p \in \mathcal{B}_{i^*}}$. Then we can apply the separating hyperplane theorem (see Boyd and Vandenberghe, 2004) to this point and set. Therefore there exists vector

$$r = (r_1^1, r_2^1, \dots, r_{i^*}^1, r_1^2, r_2^2, \dots, r_{i^*}^2, \dots, r_1^m, r_2^m, \dots, r_{i^*}^m) \in \prod_{k=1}^m \mathbb{R}^{n_{i^*}^k}$$

such that

$$r \cdot \bar{x}^{i^*} < r \cdot x^p \quad \text{for all } p \in \mathcal{B}_{i^*}. \quad (11)$$

Let $\theta \in \Theta(M)$ be an ordering of sectors M such that $r_{i^*}^{\theta(1)} \geq r_{i^*}^{\theta(2)} \geq \dots \geq r_{i^*}^{\theta(m)}$, and define the following sequence of sets: $S_0 = \emptyset, S_1 = \{\theta(1)\}, S_2 = \{\theta(1), \theta(2)\}, \dots, S_m = M$.

For each $S \subseteq M$ we associate the corresponding essential coalition

$$E^S = (i_1, i_2, \dots, i_m) \quad \text{with } i_k = i^* \text{ if } k \in S \text{ and } i_k = i^* - 1 \text{ if } k \notin S.$$

Notice that $E^{S_0} = E_{i^*-1} = (i^* - 1, i^* - 1, \dots, i^* - 1)$ and $E^{S_m} = E_{i^*} = (i^*, i^*, \dots, i^*)$ and take a path $\bar{p} \in \mathcal{B}_{i^*}$ such that $E^{S_1}, E^{S_2}, \dots, E^{S_{m-1}}$ are the essential coalitions of the path \bar{p} between E_{i^*-1} and E_{i^*} . Then the p -vector associated to the above path $\bar{p} \in \mathcal{B}_{i^*}$ satisfies

$$x^{\bar{p}}(E^{S_k}) = a_{E^{S_k}} \quad \text{for } k = 0, 1, \dots, m. \quad (12)$$

The previous system (12) gives

$$(x^{\bar{p}})_{i^*}^{\theta(k)} = (x^{\bar{p}})_{i^*-1}^{\theta(k)} + a_{E^{S_k}} - a_{E^{S_{k-1}}} \quad \text{for } k = 1, 2, \dots, m.$$

By construction of path \bar{p} we have that

$$(x^{\bar{p}})_i^k = (\bar{x}^{i^*})_i^k = x_i^k \quad \text{for } 1 \leq i \leq i^* - 1 \quad \text{and all } k \in M. \quad (13)$$

Now,

$$\begin{aligned}
r \cdot x^{\bar{p}} &= \sum_{k=1}^m \sum_{i=1}^{i^*-1} r_i^k \cdot (x^{\bar{p}})_i^k + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot (x^{\bar{p}})_{i^*}^{\theta(k)} \\
&= \sum_{k=1}^m \sum_{i=1}^{i^*-1} r_i^k \cdot x_i^k + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot \left((x^{\bar{p}})_{i^*-1}^{\theta(k)} + a_{ES_k} - a_{ES_{k-1}} \right) \\
&= \sum_{k=1}^m \sum_{i=1}^{i^*-1} r_i^k \cdot x_i^k + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot (x^{\bar{p}})_{i^*-1}^{\theta(k)} \\
&\quad + \sum_{k=1}^{m-1} \left(r_{i^*}^{\theta(k)} - r_{i^*}^{\theta(k+1)} \right) \cdot a_{ES_k} - r_{i^*}^{\theta(1)} \cdot a_{ES_0} + r_{i^*}^{\theta(m)} \cdot a_{ES_m} \\
&\leq \sum_{k=1}^m \sum_{i=1}^{i^*-1} r_i^k \cdot x_i^k + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot (x^{\bar{p}})_{i^*-1}^{\theta(k)} \\
&\quad + \sum_{k=1}^{m-1} \left(r_{i^*}^{\theta(k)} - r_{i^*}^{\theta(k+1)} \right) \cdot x(ES_k) - r_{i^*}^{\theta(1)} \cdot x(ES_0) + r_{i^*}^{\theta(m)} \cdot x(ES_m) \\
&= \sum_{k=1}^m \sum_{i=1}^{i^*-1} r_i^k \cdot x_i^k + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot (x^{\bar{p}})_{i^*-1}^{\theta(k)} + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot (x(ES_k) - x(ES_{k-1})) \\
&= \sum_{k=1}^m \sum_{i=1}^{i^*-1} r_i^k \cdot x_i^k + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot (x^{\bar{p}})_{i^*-1}^{\theta(k)} + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot x_{i^*}^{\theta(k)} - \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot x_{i^*-1}^{\theta(k)} \\
&= r \cdot \bar{x}^{i^*} + \sum_{k=1}^m r_{i^*}^{\theta(k)} \cdot \left((x^{\bar{p}})_{i^*-1}^{\theta(k)} - x_{i^*-1}^{\theta(k)} \right) = r \cdot \bar{x}^{i^*},
\end{aligned}$$

where the inequality comes from $x \in C(w_A)$ and the fact that $r_{i^*}^{\theta(k)} - r_{i^*}^{\theta(k+1)} \geq 0$ for $k = 1, \dots, m-1$, and the last equality by (13).

We have reached a contradiction with (11). Consequently any extreme core point is a p -vector. \square

Once we have established the main result of the paper, we move to some related questions. We have just proved that paths from E_0 to E_n characterize the extreme core allocations of any square assortative multisided

assignment market. We discuss now which is the maximum number of extreme core allocations.

Take an arbitrary square assortative multisided game with m sectors and n agents in each sector. We claim the maximum number of extreme core allocations is

$$m \cdot (m!)^{n-1}. \quad (14)$$

Indeed, as any path is composed of n subpaths, one for each subpart from E_{i-1} to E_i , for $i = 1, \dots, n$, we easily obtain that the total number of paths from E_0 to E_n is given by $(m!)^n$. Since we are interested in counting how many extreme core allocations, we have to take into account that at the beginning of any path, that is, from E_0 to E_1 , only m different allocations are possible. At this part $m!$ paths collapse at most into m different vectors, precisely those vectors where the worth a_{E_1} is allocated to a particular agent and give a zero payoff to the rest of agents, see Lemma 3.1. By all these arguments, formula (14) is justified.

For the special case where⁷ array A satisfies

$$a_E + a_{E'} = a_{E \vee E'} + a_{E \wedge E'} \quad \text{for any essential coalitions } E, E', \quad (15)$$

the formula (14) reduces to m if $a_{E_1} > 0$ or to 1 if $a_{E_1} = 0$.

As a numerical illustration, take the following $2 \times 2 \times 2$ array A , with $N^k = \{1^k, 2^k\}$ for $k = 1, 2, 3$, which is a valuation array,

$$A = \begin{pmatrix} 10 & 11 \\ 12 & 13 \end{pmatrix} \quad \begin{pmatrix} 14 & 15 \\ 16 & 17 \end{pmatrix}.$$

In it the rows correspond to agents in the first sector, columns to agents in the second sector and matrices to agents in the third sector. Then, for

⁷These are supermodular and submodular arrays, and they are called *valuation* arrays.

example, $a_{(1,2,2)} = 15$. Its extreme core allocations are

$$x_1 = (10, 12; 0, 1; 0, 4),$$

$$x_2 = (0, 2; 10, 11; 0, 4),$$

$$x_3 = (0, 2; 0, 1; 10, 14).$$

They can be computed by applying the p -vectors mechanism. Notice that to apply this mechanism we have to check the monotonicity condition (2), not implied by the fact that the array is a valuation.

Moreover any square valuation array A , monotonic or not, is *fully-optimal* in the sense that all its matchings are optimal, i.e. $\mathcal{M}_A^*(N^1, \dots, N^m) = \mathcal{M}(N^1, \dots, N^m)$. Any pair of non-comparable essential coalitions E, E' in any matching can be changed by $E \vee E'$ and $E \wedge E'$ without losing efficiency. The converse is not true,⁸ as the next example shows. The $2 \times 2 \times 2$ array A ,

$$A = \begin{pmatrix} 3 & 6 \\ 6 & 6 \end{pmatrix} \quad \begin{pmatrix} 6 & 6 \\ 6 & 9 \end{pmatrix}$$

is a fully-optimal multisided assignment matrix, but not a valuation, since $a_{(1,1,1)} + a_{(2,1,2)} = 3 + 6 = 9 < a_{(1,1,2)} + a_{(2,1,1)} = 12$.

Moreover, it has an empty core, since being a fully-optimal matrix, any core allocation must satisfy with equality all the array' entries, but, as the reader can check, they form a non-compatible linear system of equations.

Another important feature of a valuation array is that its entries can always be arranged monotonically by a suitable permutation of the agents. Therefore they can be seen as assortative markets. The way to see which permutation is suitable is the following. Take any core element and from it

⁸For two-sided square assignment matrices, valuation and fully-optimal are equivalent.

derive a permutation of agents in each sector such that arranges the components in a non-decreasing way. Notice that this core element satisfies with equality all entries in the array. In this way we obtain an assortative array, that is, where the monotonicity property also holds. As a consequence we can apply our results to any square valuation array. This fact simplifies the assertions made in Sherstyuk (1999) since there is no need to distinguish valuation markets from assortative ones.

It is easy to generate examples in which the maximum number of extreme core points given in (14) is attained.

Consider the following $2 \times 2 \times 2$ array A ,

$$A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \quad \begin{pmatrix} 3 & 7 \\ 5 & 10 \end{pmatrix}.$$

Notice that all inequalities of the supermodular property concerning non-comparable essential coalitions are strict. There are $3 \cdot (3!)^1 = 18$ different extreme core allocations, that correspond to different paths.

$$\begin{aligned} x_1 &= (1, 2; 0, 2; 0, 6), & x_2 &= (0, 1; 1, 3; 0, 6), \\ \mathbf{x}_3 &= (\mathbf{0}, \mathbf{1}; \mathbf{0}, \mathbf{2}; \mathbf{1}, \mathbf{7}), & x_4 &= (1, 2; 0, 5; 0, 3), \\ \mathbf{x}_5 &= (\mathbf{0}, \mathbf{1}; \mathbf{1}, \mathbf{6}; \mathbf{0}, \mathbf{3}), & x_6 &= (0, 1; 0, 5; 1, 4), \\ x_7 &= (1, 3; 0, 1; 0, 6), & x_8 &= (0, 2; 1, 2; 0, 6), \\ \mathbf{x}_9 &= (\mathbf{0}, \mathbf{2}; \mathbf{0}, \mathbf{1}; \mathbf{1}, \mathbf{7}), & \mathbf{x}_{10} &= (\mathbf{1}, \mathbf{4}; \mathbf{0}, \mathbf{1}; \mathbf{0}, \mathbf{5}), \\ x_{11} &= (0, 3; 1, 2; 0, 5), & x_{12} &= (0, 3; 0, 1; 1, 6), \\ x_{13} &= (1, 3; 0, 5; 0, 2), & \mathbf{x}_{14} &= (\mathbf{0}, \mathbf{2}; \mathbf{1}, \mathbf{6}; \mathbf{0}, \mathbf{2}), \\ x_{15} &= (0, 2; 0, 5; 1, 3), & \mathbf{x}_{16} &= (\mathbf{1}, \mathbf{4}; \mathbf{0}, \mathbf{4}; \mathbf{0}, \mathbf{2}), \\ x_{17} &= (0, 3; 1, 5; 0, 2), & x_{18} &= (0, 3; 0, 4; 1, 3). \end{aligned}$$

The six vectors in boldface correspond to the $m! = 3! = 6$ vectors given in Sherstyuk (1999).

References

- [1] Becker, G.S., 1973. A Theory of Marriage: Part I. *The Journal of Political Economy* 81(4), 813–846.
- [2] Boyd, S., Vandenberghe, L., 2004. *Convex optimization*. Cambridge University Press, Cambridge, U.K.
- [3] Crawford, V.P., Knoer, E.M., 1981. Job matching with heterogeneous firms and workers. *Econometrica*, 49(2), 437–450.
- [4] Eriksson, K., Karlander, J., Öller, L.-E., 2000. Becker’s assortative assignments: stability and fairness. *Mathematical Social Sciences* 39, 109–118.
- [5] Kaneko, M., Wooders, M.H., 1982. Cores of partitioning games. *Mathematical Social Sciences* 3(4), 313–327.
- [6] Martínez-de-Albéniz, F.J., Rafels, C., 2014. Cooperative assignment games with the inverse Monge property. *Discrete Applied Mathematics* 162, 42–50.
- [7] Martínez-de-Albéniz, F.J., Rafels, C., Ybern, N., 2019. Solving Becker’s assortative assignments and extensions, *Games and Economic Behavior* 113, 248–261.
- [8] Quint, T., 1991. The core of an m -sided assignment game. *Games and Economic Behavior* 3, 487–503.
- [9] Shapley, L.S., 1955. Markets as cooperative games. *Rand Research Memorandum P-629*. Rand Corp., Santa Monica, pp. 1–7.

- [10] Shapley, L.S., Shubik, M., 1971. The assignment game I: the core. International Journal of Game Theory 1, 111–130.
- [11] Sherstyuk, K., 1999. Multisided matching games with complementarities. International Journal of Game Theory 28, 489–509.
- [12] Stuart, H.W., 1997. The supplier-firm-buyer game and its m -sided generalization. Mathematical Social Sciences 34(1), 21–27.
- [13] Topkis, D. M. (1998). Supermodularity and complementarity. Princeton University Press.