

# Accepted Manuscript

Validity and Reliability of the Decision Regret Scale in Cancer Patients Receiving Adjuvant Chemotherapy

Caterina Calderon, Ph.D, Pere Joan Ferrando, Urbano Lorenzo-Seva, Oliver Higuera, Teresa Ramon y Cajal, Jacobo Rogado, Margarida Mut-Lloret, Alejandra Rodriguez-Capote, Carlos Jara, Paula Jimenez-Fonseca

PII: S0885-3924(18)31086-8

DOI: <https://doi.org/10.1016/j.jpainsymman.2018.11.017>

Reference: JPS 9974

To appear in: *Journal of Pain and Symptom Management*

Received Date: 30 August 2018

Revised Date: 16 November 2018

Accepted Date: 18 November 2018

Please cite this article as: Calderon C, Ferrando PJ, Lorenzo-Seva U, Higuera O, Ramon y Cajal T, Rogado J, Mut-Lloret M, Rodriguez-Capote A, Jara C, Jimenez-Fonseca P, Validity and Reliability of the Decision Regret Scale in Cancer Patients Receiving Adjuvant Chemotherapy, *Journal of Pain and Symptom Management* (2019), doi: <https://doi.org/10.1016/j.jpainsymman.2018.11.017>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**TITLE: Validity and Reliability of the Decision Regret Scale in Cancer Patients Receiving Adjuvant Chemotherapy****Abstract**

**Objectives** Decisional regret is an indicator of satisfaction with the treatment decision and can help to identify those patients who need more support and evaluate the efficacy of decision support interventions. The objectives of this study are, 1) to evaluate the psychometric properties of the Decision Regret Scale and 2) to analyze the moderating effect of psychological distress on functional status and regret in patients with cancer following adjuvancy.

**Methods** A prospective, multicenter cohort of 403 patients who completed the Decision Regret Scale (DRS), Health-related Quality of Life (EORTC QLQ-C30), and Brief Symptom Inventory (BSI). The evaluation was conducted six months after receiving adjuvant treatment in patients with resected cancer.

**Results** After treatment, most participants (51.9%) experienced no decision regret; 33.7% felt mild regret, and 14.4% exhibited high levels of regret. The Spanish version of the DRS demonstrated satisfactory properties: it had a strong, clear unidimensional factorial structure with substantial loadings. Decisional regret was related with lower scores on functional, symptom, and quality of life scales, and higher levels of psychological distress (all  $p=0.001$ ). Psychological distress was found to have a moderating effect on the relationship between functional state and decision regret.

**Conclusions** The Spanish version of the DRS is a reliable, valid tool to evaluate regret and post-decisional quality in clinical practice and further highlights the potential clinical implications of psychological distress for the relation between physical status and regret.

**Keywords:** cancer; decision regret; functional status; psychological distress; validity; reliability

## **Introduction**

Decision-making as to the advisability of receiving or rejecting adjuvant-chemotherapy that seeks to lessen the risk of recurrence after tumor resection is complex for cancer patients<sup>1</sup>. Medical oncologists provide their patients with information about the risks and benefits of chemotherapy to facilitate an informed decision<sup>2</sup>.

Regret is a negative emotional reaction resulting from the untoward effects of a given choice. Evaluating regret in medical decision-making is uncommon, given the lack of reliable, valid measures. Although a systematic review of regret measures yielded ten instruments<sup>3</sup>, only three were applicable to decision-making in the context of medical care: *Regret Scale*<sup>4</sup> that appraises regret in men with prostate cancer; *Anticipatory Regret Questionnaire*<sup>5</sup> that captures a sense of regret in the future about donating blood, and *Decision Regret Scale (DRS)*<sup>6</sup> that gauges regret in patients who have already made a medical decision<sup>6</sup>.

Regret has been correlated with low degrees of satisfaction with preoperative information, depression, anxiety, and stress<sup>7</sup>; with negative body image and psychological distress<sup>8</sup> in patients with breast cancer, and with impaired quality of life (QoL) in patients with prostate cancer<sup>9</sup>. The psychological factors involved in regret include perception of health and psychological distress, both of which are common in individuals with cancer beginning adjuvant treatment<sup>10</sup>. With this background, this study seeks to analyze the psychometric properties of DRS in patients with resected cancer treated with adjuvant chemotherapy and to examine the moderating effect of psychological distress between functional status and regret.

## **Methods**

### *Research Design and Study Population*

A multi-institutional, prospective, observational research design was used to examine the incidence of decision regret with respect to adjuvant therapy and evaluate the psychometrics of DRS. Within this design, DRS data were collected at a single time point (see below); consequently, all analyses performed in this study are cross-sectional. The study pooled consecutive patients recruited at 14 teaching hospitals in Spain from June 2015 to December 2017 and was approved by the Ethics Review Board at each institution and by the Spanish Agency of Medicines and Medical Devices (AEMPS). STROBE guidelines were used to ensure the reporting of this study (Additional file 1)<sup>11</sup>.

Inclusion criteria were: aged >18 years, have a histologically confirmed, non-advanced, resected solid tumor, be eligible for adjuvant treatment.

#### *Sample characteristics*

A total of 403 cancer patients (55.3% women and 44.7% men) completed the assessment 6 months after receiving adjuvant-chemotherapy. Mean age was 58.4 years (range 25-84); most were married or partnered (79.4%), unemployed (58.8%), and had attained a primary level of education (61%). The most common tumor sites were colon (43.9%) and breast (31%), stages I-II (53.6%). All received chemotherapy and 30.8% received associated radiotherapy.

*DRS adaptation.* The DRS was first developed in English<sup>6</sup>. It was adapted to Spanish by forward-backward translation<sup>12</sup> of the English version by two independent bilingual translators; both of whom are native Spanish speakers and fluent in English. The two Spanish versions were translated back into English by a professional translator with experience in medical translation and by two medical doctors who had not been involved in the forward translation. Another two bilingual translators blind to the original English version back-translated the revised Spanish version; finally, the study directors compared and synthesized the back-translation with the original questionnaire, culminating in a final Spanish version of the DRS.

#### *Data collection*

Data collection was similar for all hospitals. Following a full explanation of study objectives and procedures, candidates were invited to participate and complete the questionnaire at home. Study participation was voluntary and anonymous. Participants completed the questionnaires individually, with no time limit. This visit was structured according to standard practice at each center, although it was agreed that as a minimum, risk of relapse, options for adjuvant treatment, risk of adverse effects, and possible treatment efficacy should be discussed with the patients. Participant's flow chart is given in Fig.1.

Insert Figure 1 here

*Decision Regret Scale.* The DRS is a five-item, self-report scale to evaluate decisional regret. Items are scored on five-point Likert scales, ranging from 1 to 5. Scores were reversed for items 2 and 4; mean scores were obtained and then converted by subtracting 1 and multiplying by 25. Scores range from 0-100, with higher scores

indicating greater regret. Internal consistency reliability estimates ( $\alpha=0.81-0.92$ ) for oncology patients<sup>6</sup>.

Health-related Quality of Life. The EORTC QLQ-C30 (V.3.0) is a cancer-specific measure of HRQoL<sup>13</sup>. It consists of 30 items that assess quality of life (QoL) as regards functional scales, and a global QoL scale<sup>14</sup>. All scale scores are linearly transformed to a 0–100 scale. Higher scores represent a higher level of functioning or QoL, and more symptom burden. Cronbach's alpha estimates range from 0.74 to 0.88<sup>13</sup>.

Brief Symptom Inventory-18. The BSI-18<sup>15</sup> is a self-report inventory designed to evaluate psychological distress, using a 5-point scale from 0-4. Raw scores are converted to T-scores based on gender-specific normative data. Higher scores indicate greater psychological distress<sup>15</sup>. Cronbach's alpha estimates range from 0.72 to 0.84<sup>16</sup>.

#### *Data analysis*

The internal psychometric properties of the DRS scale were examined using a three-stage series of analyses. First, basic item descriptive statistics were explored. Second, a confirmatory factor analysis (CFA) was performed to study the scale's expected unidimensionality, as well as the pattern of item-trait relations. Third, provided that the DRS behaved as essentially unidimensional, the reliability of its scores was scrutinized. The unidimensional CFA solution in step 2 above was fitted using robust weighted least squares estimation with mean and variance corrected fit statistics as implemented in the Mplus program<sup>17</sup>. Model fit and appropriateness were assessed with three groups of measures. First, model residuals and relative fit were evaluated with SRMS (Standardized Root Mean Square Residual) and Root Mean Squared Error of Approximation (RMSEA) statistics. Second, relative comparative fit was examined with the comparative fit index (CFI) (as a relative measure of fit with respect to the null independence model). Finally, additional indices of appropriateness were also obtained to verify the strength and replicability of the solution (h index), as well as closeness to unidimensionality (Explained Common Variance ECV index)<sup>18</sup> with the FACTOR software program<sup>19</sup>. As for reference values, CFI values  $\geq 0.95$  are indicative of good model fit<sup>20</sup>, whereas SRMR values  $\leq 0.08$  and RMSEA values  $\leq 0.06$  are considered satisfactory fitting models<sup>21,22</sup>. Finally, once the proposed structure had been fitted and found suitable, DRS score reliability was assessed using the omega coefficient<sup>23</sup>. Validity analysis proceeded in two stages. First, the product-moment correlations between DRS scores and EORTC scores, BSI scores, and toxicity degree were obtained.

Second, multiple linear regression was used to examine the potential moderating effects between QoL and psychological distress on decision regret. Validity analyses were conducted using IBM-SPSS 23.0 statistical software package (SPSS, INC., Chicago, III) for Windows.

## **Results**

### *Descriptive analyses*

The mean DRS score was 10.6 (SD=15.4). Most participants 51.9% ( $n=209$ ) experienced no decision regret. Degrees of decisional regret were not significantly influenced by gender ( $F_{(1,401)}=0.503$ ,  $p=0.478$ ), age ( $\leq 60$  years vs  $>60$  years;  $F_{(1,401)}=1.310$ ,  $p=0.253$ ), or education (primary level vs  $\geq$ high school;  $F_{(1,401)}=0.133$ ,  $p=0.716$ ). Likewise, decisional regret was not significantly affected by treatment (chemotherapy vs chemo- and radiotherapy) ( $F_{(1,401)}=1.203$ ,  $p=0.273$ ).

DRS item score distributions were unimodal and asymmetrical (positively skewed). Item-total correlations were all  $>0.40$ . Provided that the scale behaves unidimensionally, this result indicates that the items have adequate discriminating power and are strongly related to the construct they measure (see table 1).

Insert Table 1 here

### *CFA analysis*

Given (a) the ordered-categorical nature of the response variables, (b) the skewed item distributions, and (c) the relatively high item discriminating power, we opted to use the underlying-variables approach and fit the FA model to the inter-item polychoric correlation matrix<sup>24</sup>. The initial fit of the unidimensional model was unacceptable by all standards. Inspection of the results, however, clearly revealed that the sole source of misfit was the correlated residual between items 2 and 4. This result, confirmed by cross-validation, is to be expected since (a) these are the two reverse-keyed items (i.e., a method effect) and (b) they share specific content. After freeing the residual covariance, an almost perfect fit was obtained: SRMS=0.01; RMSEA=0.0; CFI=1. The remaining indices of appropriateness were also relatively acceptable. The ECV value was 0.81, indicating that the solution could indeed be considered essentially unidimensional, whereas the h index was 0.91, suggesting that the solution was both robust and replicable<sup>26</sup>. Note that all loadings are well above their corresponding standard error, and those of items 1, 3, and 5 are particularly high. This result is consistent with the substantial item-total correlations found above and suggest that the DRS items are

strongly related to the construct they measure. The omega reliability estimate for the raw scores was 0.87, which is remarkably high, given the limited number of items.

### *Validity analyses*

After treatment, higher levels of decisional regret correlated significantly with lower scores on the functional ( $r=-0.316$ ,  $p=0.001$ ) and global QoL ( $r= -0.257$ ,  $p=0.001$ ) subscales. Higher levels of decisional regret, however, correlated significantly with higher symptom ( $r=0.278$ ,  $p=0.001$ ) and psychological distress ( $r=0.297$ ,  $p=0.001$ ) scores. Of the four interactions tested, one was significant — the moderational effect of psychological distress on the relationship between functional status and regret ( $B= -.015$ , 95% CI [0.02, 0.05],  $\beta= -.96$ ,  $t=2.87$ ,  $p=0.004$ ). The analysis indicated that 12.2% of the variation in regret was explained by functional status and the interaction effects between functional status and psychological distress ( $F_{(3,397)}=19.51$ ,  $p<0.001$ ) (see table 2 and Figure 2) Thus, when functional status is high, there are minimal differences in regret experienced by both patients suffering from high and those with low levels of psychological distress.

Insert Table 2 and Figure 2 here

Given the positively skewed distribution (i.e., floor effects) of DRS scores, as well as measurement error in the reported measures, the validity results discussed thus far are likely to be somewhat attenuated. Therefore, these results can be better considered as lower limits, and the latent relations between the constructs of interest (decisional regret, psychological distress...) are likely to be stronger than the product-moment correlations and proportions of shared variance based on what the 'proxy' measures suggest.

### *Discussion*

This study presents important findings. First, few patients with cancer (14.4%) regret having undergone adjuvant treatment after surgery for a non-metastatic tumor, despite its side effects. Our results are consistent with previous studies that have evaluated decision regret in oncological patients and suggest that most patients are satisfied with their treatment and present low levels of decisional regret<sup>8,9,25</sup>. Studies using the DRS to assess decision regret found that 19.5% of women with breast cancer regretted reconstruction following mastectomy<sup>8</sup> and only 4% of men with prostate cancer regretted their decision one year after surgery.

As for internal psychometrics, the Spanish version of the DRS displayed good properties. It had a clear, strong unidimensional factorial structure with substantial loadings and the scores derived from it demonstrated acceptable reliability even for individual assessment. These results are remarkable, taking into account that they were obtained with only 5 items. Despite these good properties, however, there is still room for further improvement. The two negatively keyed items had weaker discriminating power than the remaining ones and shared residual variance that had to be accounted for in the model to achieve an acceptable level of fit. One hypothesis is that participants (Spanish patients) confuse the wording of these items and follow the pattern of response for positively worded statements<sup>26</sup>. Other researchers have suggested that the inclusion of the reverse worded items increases the risk of inattention and confusion<sup>27,28</sup>. Replacing negatively worded items with alternative items to measure decision regret would help eliminate error and improve the scale's variance.

As regards external validity, convergent validity was acceptable and a pattern of significant correlations with health outcomes was observed in the expected direction. This is particularly relevant, since adjuvant chemotherapy is the treatment of choice after surgery for certain locally advanced cancers, such as breast and colon cancer, and treatment tends to have immediate, negative repercussions for patients' physical status, symptomatology, and global QoL<sup>29</sup>, although their adverse effects are usually temporary<sup>29</sup>. Our results indicate that the patients most likely to regret treatment are those with worse QoL, including physical status, presence of symptoms, and greater psychological distress. Other authors found that regret was greater in patients with heart disease with worse perceived physical health<sup>30</sup>, and regret was associated with depression, anxiety, and stress in women with breast cancer<sup>31</sup>. We did not find differences in regret by gender, age, educational level, or type of treatment. It must be remembered that in medical contexts, patients tend to provide socially acceptable responses on questionnaires that rate their attitude toward the quality of care and treatment received<sup>7</sup>. We have attempted to avoid this problem by guaranteeing participants' anonymity.

The moderating role of psychological distress provides a more complete view of the relation between physical status and regret. Thus, when physical status is adequate, no significant differences are seen among cancer patients with respect to regret, whatever their level of psychological distress. In contrast, when health is perceived as poor, those patients with high levels of psychological distress report significantly higher degrees of



decisional regret. These findings correlate in part with the Monitoring Process Model<sup>32</sup>, according to which a person's perception of their own health impacts how they confront and respond to it. To the best of our knowledge, there are no other studies that analyze the moderating role of psychological distress between physical status and decision regret in oncology patients. In future studies, it would be interesting to ascertain if this regret is reversible; i.e., whether decision regret decreases if/when patients' physical and/or psychological status improve(s).

This study has a series of **limitations**. First, the sample is heterogeneous, to enable subgroup analyses to be performed based on tumor site. Second, because the study was designed to include a single evaluation of regret, information about causality cannot be examined, nor can we offer a possible explanation as to how regret evolves over time. Third, it may be that the results of our study cannot be extrapolated to patients with advanced tumors, whose clinical situation and prognosis differ markedly. Finally, we must be cautious when interpreting these results, bearing in mind that all the patients eligible to participate did so voluntarily, which may have introduced a self-selection bias. Likewise, a limitation of clinical significance is the absence of appropriately matched comparison samples of patients to assess regret regarding their decision not to proceed with chemotherapy. Future research is required to explore the contribution of other psychological factors that may influence decision regret, e.g., outcome expectations and fear of recurrence.

### *Conclusions*

The results of this study indicate that the Spanish version of the DRS exhibits satisfactory psychometric properties, with a clear and strong unidimensional factorial structure, and acceptable reliability and validity. The moderating effect of psychological distress in the relation between physical status and regret tells us that the effect of physical status on regret is more accentuated if patients display greater psychological distress. This result points to the need to perform individualized interventions that promote improvement of physical status following chemotherapy. In general, after completing adjuvant treatment, there are fewer visits to the oncologist. However, some individuals, due to their deteriorated physical and psychological condition as a result of residual toxicity of chemotherapy, will require more frequent visits to help them improve their status, boost recovery, and lessen the likelihood of post-chemotherapy

rejection. Maintaining a certain level of psychological wellbeing will likely benefit patients in their recovery.

ACCEPTED MANUSCRIPT

## REFERENCES

1. Meyers BM, Cosby R, Quereshy F, et al. Adjuvant Chemotherapy for Stage II and III Colon Cancer Following Complete Resection: A Cancer Care Ontario Systematic Review. *Clin Oncol* 2017; 29: 459–465.
2. Verhoeff SR, van Erning FN, Lemmens VEPP, et al. Adjuvant chemotherapy is not associated with improved survival for all high-risk factors in stage II colon cancer. *Int J Cancer* 2016; 139: 187–193.
3. Joseph-Williams N, Edwards A, Elwyn G. The importance and complexity of regret in the measurement of ‘good’ decisions: a systematic review and a content analysis of existing assessment instruments. *Heal Expect* 2011; 14: 59–83.
4. Clark JA, Wray NP, Ashton CM. Living With Treatment Decisions: Regrets and Quality of Life Among Men Treated for Metastatic Prostate Cancer. *J Clin Oncol* 2001; 19: 72–80.
5. Godin G, Sheeran P, Conner M, et al. Factors explaining the intention to give blood among the general population. *Vox Sang* 2005; 89: 140–149.
6. Brehaut JC, O’Connor AM, Wood TJ, et al. Validation of a Decision Regret Scale. *Med Decis Mak* 2003; 23: 281–292.
7. Beach MC, Roter DL, Wang N-Y, et al. Are physicians’ attitudes of respect accurately perceived by patients and associated with more positive communication behaviors? *Patient Educ Couns* 2006; 62: 347–54.
8. Sheehan J, Sherman KA, Lam T, et al. Regret associated with the decision for breast reconstruction: The association of negative body image, distress and surgery characteristics with decision regret. *Psychol Health* 2008; 23: 207–219.
9. Davison BJ, So AI, Goldenberg SL. Quality of life, sexual function and decisional regret at 1 year after surgical treatment for localized prostate cancer. *BJU Int* 2007; 100: 780–785.
10. Calderon C, Jiménez-Fonseca P, Ferrando PJ, et al. Psychometric properties of the Shared Decision-Making Questionnaire (SDM-Q-9) in oncology practice. *Int J Clin Heal Psychol*. Epub ahead of print 2018. DOI: 10.1016/j.ijchp.2017.12.001.
11. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008; 61: 344–349.
12. Hambleton RK, Merenda PF, Spielberger CD. *Adapting educational and psychological tests for cross-cultural assessment*. Mahwah, NJ: Lawrence Erlbaum, 2005.
13. Aaronson NK, Ahmedzai S, Bergman B, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; 85: 365–76.
14. Fayers PM, Aaronson NK, Bjordal K, Groenvold M, Curran D BA. *The EORTC QLQ-C30 Scoring Manual (3rd edition)*. Brussels: 2001.
15. Derogatis LR. *BSI 18, Brief Symptom Inventory 18: Administration, scoring and procedures manual*. Minneapolis: NCS Pearson, Inc., 2001.
16. Merport A, Recklitis CJ. Does the Brief Symptom Inventory-18 Case Rule Apply in Adult Survivors of Childhood Cancer?: Comparison With the Symptom Checklist-90. 2012; 37: 650–659.
17. Muthén, L. K. Muthén B. *Mplus user’s guide. Fifth Edition*. Los Angeles, 2007.

18. Ferrando PJ, Lorenzo-Seva U. Assessing the Quality and Appropriateness of Factor Solutions and Factor Score Estimates in Exploratory Item Factor Analysis. *Educ Psychol Meas* 2017; 001316441771930.
19. Lorenzo-Seva U, Ferrando PJ. FACTOR 9.2: A Comprehensive Program for Fitting Exploratory and Semiconfirmatory Factor Analysis and IRT Models. *Appl Psychol Meas* 2013; 37: 497–498.
20. Schermelleh-Engel K, Moosbrugger H, Müller H. Evaluating the Fit of Structural Equation Models: Tests of Significance and Descriptive Goodness-of-Fit Measures. *Methods Psychol Res Online* 2003; 23–74.
21. Hair JF, Black WC, Babin BJ. RE Anderson Multivariate data analysis: A global perspective.
22. Schermelleh-Engel K, Kerwer M, Klein AG. Evaluation of model fit in nonlinear multilevel structural equation modeling. *Front Psychol* 2014; 5: 181.
23. McDonald RP. *Test theory: A unified treatment*. L. Erlbaum Associates, Mahwah, N.J. 1999.
24. Ferrando PJ, Lorenzo-Seva U. *Unrestricted item factor analysis and some relations with Item Response Theory*. Technical Report. Department of Psychology, Universitat Rovira i Virgili, Tarragona. <http://psico.fcep.urv.es/utilitats/factor/>, 2013.
25. Martinez KA, Li Y, Resnicow K, et al. Decision Regret following Treatment for Localized Breast Cancer. *Med Decis Mak* 2015; 35: 446–457.
26. Woods CM. Careless responding to reverse-worded items: Implications for confirmatory factor analysis. *J Psychopathol Behav Assess* 2006; 28: 186.
27. Sonderen E van, Sanderman R, Coyne JC. Ineffectiveness of Reverse Wording of Questionnaire Items: Let's Learn from Cows in the Rain. *PLoS One* 2013; 8: e68967.
28. Weijters B, Baumgartner H, Schillewaert N. Reversed item bias: An integrative model. *Psychol Methods* 2013; 18: 320–334.
29. Biswas T, Efird JT, Prasad S, et al. The survival benefit of neoadjuvant chemotherapy and pCR among patients with advanced stage triple negative breast cancer. *Oncotarget* 2017; 8: 112712–112719.
30. Hickman RL, Pinto MD, Lee E, et al. Exploratory and confirmatory factor analysis of the decision regret scale in recipients of internal cardioverter defibrillators. *J Nurs Meas* 2012; 20: 21–34.
31. Sheehan J, Sherman KA, Lam T, et al. Association of information satisfaction, psychological distress and monitoring coping style with post-decision regret following breast reconstruction. *Psychooncology* 2007; 16: 342–351.
32. Miller S. *Monitoring versus blunting styles of coping with cancer influence the information patients want and need about their disease. Implications for cancer screening and management*. 1995. Epub ahead of print 1 August 1995. DOI: 10.1002/1097-0142(19950715)76:23.0.CO;2-K.

**Table 1.** Item Descriptive Statistics of the Spanish DRS (n=403)

Questions	Mean	SD	Skews	R <sub>item-total</sub>
It was the right decision	1.22	0.57	2.7 (.1)	0.62
I regret the choice	1.65	1.33	1.8 (.1)	0.59
I would make the same choice if I had to do it over again	1.28	0.76	3.4 (.1)	0.61
The choice did me a lot of harm	1.73	1.23	1.6 (.1)	0.60
The decision was a wise one	1.26	0.55	2.8 (.1)	0.62

*Note.* SD: standard deviation. Score ranges from 1 (*strongly agree*) to 5 (*strongly disagree*).

In brackets the standard error for the skew statistics. R<sub>item-total</sub> = item-total corrected correlation.

**Table 2.** Moderational effects between quality-of-life scales and psychological distress on decision regret; multiple linear regression analyses summary

Model	<i>Unstandardized coefficients</i>		<i>Standardized coefficient</i>	<i>t</i>	<i>p</i>
	<i>B</i>	<i>Std. Error</i>	$\beta$		
1 (Constant)	2.187	12.987		0.168	0.866
Functional status	-0.166	0.055	-0.205	-3.019	0.003
PD	0.332	0.148	0.153	2.243	0.025
2 (Constant)	-79.959	31.343		-2.551	0.011
Functional status	0.852	0.358	1.056	2.379	0.018
PD	1.485	0.427	0.682	3.477	0.001
Functional*PD	-0.015	0.005	-0.964	-2.874	0.004

*Note.* PD: Psychological distress



