

Journal Pre-proof

Is temperature reducing the transmission of COVID-19 ?

Aurelio Tobías, Tomás Molina



PII: S0013-9351(20)30446-1

DOI: <https://doi.org/10.1016/j.envres.2020.109553>

Reference: YENRS 109553

To appear in: *Environmental Research*

Received Date: 14 April 2020

Revised Date: 16 April 2020

Accepted Date: 16 April 2020

Please cite this article as: Tobías, A., Molina, Tomá., Is temperature reducing the transmission of COVID-19 ?, *Environmental Research* (2020), doi: <https://doi.org/10.1016/j.envres.2020.109553>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc.

IS TEMPERATURE REDUCING THE TRANSMISSION OF COVID-19 ?

Aurelio Tobías (1), Tomás Molina (2)

1. Institute of Environmental Assessment and Water Research (IDAEA), Spanish Council for Scientific Research (CSIC), Barcelona, Spain.
2. Department of Applied Physics, University of Barcelona, Barcelona, Spain.

Correspondence to:

Aurelio Tobías

Institute of Environmental Assessment and Water Research (IDAEA),
Spanish Council for Scientific Research (CSIC)

C/ Jordi Girona 18-26.

08034 Barcelona, Spain.

e-mail: aurelio.tobias@idaea.csic.es

Keywords:

Temperature, SARS-CoV-2, COVID-19, incidence rate, time-series regression,
Barcelona, Spain.

Conflict of interest:

None.

Acknowledgments:

To Joe Brew for providing Online Appendix Figure 1.

Dear Editor,

The first confirmed case of SARS-CoV-2 in the city of Barcelona, Spain, was identified on February 25th, 2020. Since then, the epidemic spread countrywide, and Spain has become the second most affected country worldwide (130,759 cases, on April 5th) after the United States, and the second in the number of deaths (12,418 deaths, on April 5th) after Italy (Our World in Data, 2020). The regions most affected are Madrid and Catalonia, where Barcelona is located (Supplementary Figure 1).

The transmission of coronaviruses can be affected by several factors, including the climate (Hemmes et al., 1962). Therefore, it is essential to understand the role that weather conditions could have on the transmission of COVID-19. However, there is still a lack of evidence. A previous study in the four major Chinese cities suggested that the SARS outbreak was significantly associated with the temperature, increasing the daily incidence rate with lower temperatures (Tan et al., 2005). Recent preprint studies using national data from countries affected by COVID-19 reported that low temperature is beneficial to the viral transmission (Triplett 2020; Wang et al., 2020a; Wang et al., 2020b). However, these should be taken with caution since have not yet been peer reviewed.

We collected daily incident diagnosed cases confirmed from PCR tests in the Barcelona's Health Region (*Regió Sanitària de Barcelona*), between March 2nd and April 5th, 2020. Data was collected from the COVID-19 registry of PCR tests by the Health Evaluation and Quality Agency of Catalonia (AQuAs, 2020). Daily maximum temperature was provided by the Meteorological Service of Catalunya (MeteoCat). We fitted a quasi-Poisson regression model allowing for features specific to the infectious disease context (Imai et al., 2015). We adjusted for linear and quadratic trends, weekends, and the lockdown period (after March 16th), and accounting for residual autocorrelation by adjusting for the logarithm of lagged outcome (Brumback et al., 2000). Since the incubation the median incubation period for COVID-19 is just over 5-6 days (Lauer et al., 2020), we used a one week distributed lag model (Gasparrini et al., 2010).

The incidence rate of a positively diagnosed case of COVID-19 ranged from 0 to 60% and daily maximum temperature between 12.2°C and 22.8°C (Supplementary Figure


2). An average increase of 1°C of maximum temperature decreased the incidence rate by -7.5% (95%CI=[-12.3; -2.6]) on the same day (lag 0) (Figure 1). Afterward, the temperature effect almost disappears, being substantially reduced from lag 1 to lag 6. We evaluated the linearity of the associations without observing evidence of departure from linearity (Supplementary Figure 3).

Temperature is an essential factor in people's living environment, which can play a significant role in public health in terms of epidemic development and control (McMichael et al., 2008). There is a specific temperature that best fits the benefit of the virus and that lower temperatures contribute to the transmission because it is when the vulnerability increases. Thus, the arrival of summer could reduce the transmission of the COVID-19. Our result agreed with those already reported in other geographical locations, showing how the number of diagnosed cases increase below a maximum temperature of 10 °C and linearly decreasing afterward (Triplett 2000; Wang et al., 2020b). However, these are cross-sectional studies at the country scale, which might have some measurement error due to high spatial variability when estimating a fixed temperature exposure for large geographic areas (Soares et al., 2012). Our study considers a smaller area, mainly defined by Barcelona's metropolitan area, where the daily variability of maximum temperature can be estimated more accurately. However, we have not been able to account for socioeconomic status or other meteorological factors, such as ultraviolet radiation or absolute humidity. While daily mortality could be a more robust indicator of the virus behavior, because it is not affected by the number of people tested, mortality may have different incident lag, from days to weeks from the diagnostic. We used instead of the incidence rate of positive diagnosed COVID-19 confirmed with a PCR test.

Although we adjusted for time-trends and accounted for residual autocorrelation, typical for in infectious diseases (Imai et al., 2015), we are concerned that our finding could still be due to a spurious relationship due to the small number of days considered in our analysis. Our result has to be further explored; however, it could provide a first clue to understanding the temperature-transmission relation of COVID-19 since it shares genetic similarities with SARS (Benvenuto et al., 2020).

Journal Pre-proof

Figure 1. Risk increase of daily incidence rate of positive diagnosed cases of COVID-19 for a rise of 1 °C of maximum temperature from lag 0 to lag 6.



References

- AQuAs, 2020. Registre de test de COVID-19 realitzats a Catalunya. Segregació per sexe i àrea bàsica de salut (ABS). Dades Obertes Catalunya. Generalitat de Catalunya. [Accesed April 5th, 2020]. Available from: <https://analisi.transparenciacatalunya.cat/Salut/Registre-de-test-de-COVID-19-realitzats-a-Cataluny/xuwf-dxjd>
- Benvenuto, D., et al., 2020. The 2019-new coronavirus epidemic: Evidence for virus evolution. *J Med Virol.* 92, 455-459.
- Brumback, B. A., et al., 2000. Transitional Regression Models, with Application to Environmental Time Series. *Journal of the American Statistical Association.* 95.
- Our World in Data, 2020. Coronavirus Disease (COVID-19) – Statistics and Research. Oxford Martin School, The University of Oxford, Global Change Data Lab. [Accessed April 5th, 2020]. Available from: <https://ourworldindata.org/coronavirus/>
- Gasparrini, A., et al., 2010. Distributed lag non-linear models. *Stat Med.* 29, 2224-34.
- Hemmes, J. H., et al., 1962. Virus survival as a seasonal factor in influenza and poliomyelitis. *Antonie Van Leeuwenhoek.* 28, 221-33.
- Imai, C., et al., 2015. Time series regression model for infectious disease and weather. *Environ Res.* 142, 319-27.
- Lauer, S. A., et al. The Incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Annals of Internal Medicine*, 2020; DOI: 10.7326/M20-0504
- McMichael, A. J., et al., 2008. International study of temperature, heat and urban mortality: the 'ISOTHURM' project. *Int J Epidemiol.* 37, 1121-31.
- Soares, P. M. M., et al., 2012. WRF high resolution dynamical downscaling of ERA-Interim for Portugal. *Climate Dynamics.* 39, 2497-2522.
- Tan, J., et al., 2005. An initial investigation of the association between the SARS outbreak and weather: with the view of the environmental temperature and its variation. *J Epidemiol Community Health.* 59, 186-92.
- Triplett, M., 2020. Evidence that higher temperatures are associated with lower incidence of COVID-19 in pandemic state, cumulative cases reported up to March 27. medRxiv preprint. doi: <https://doi.org/10.1101/2020.04.02.20051524>
- Wang, J., et al., 2020a. High Temperature and High Humidity Reduce the Transmission of COVID-19. *SSRN Electronic Journal.* doi: <http://dx.doi.org/10.2139/ssrn.3551767>
- Wang, M., et al., 2020b. Temperature significantly change COVID-19 transmission in 429 cities. medRxiv preprint. doi: <https://doi.org/10.1101/2020.02.22.20025791>

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: