

Correction to "Oxidative Cleavage of Cellobiose by Lytic Polysaccharide Monooxygenase (LPMO)-Inspired Copper Complexes"

Andrea. C. Neira, Paulina R. Martínez-Alanis, Gabriel Aullón, Marcos Flores-Alamo, Paulino Zerón, Anna Company, Juan Chen, Johann B. Kasper, Wesley R. Browne, Ebbe Nordlander, and Ivan Castillo*

ACS Omega 2019, 4, 6, 10729–10740 DOI: [10.1021/acsomega.9b00785](https://doi.org/10.1021/acsomega.9b00785)



Cite This: ACS Omega 2020, 5, 28403–28403



Read Online

ACCESS |



Metrics & More



Article Recommendations

It should be specified that in the preparative scale oxidations of cellobiose, the total volume of the reaction mixture was 2 mL at a phosphate buffer concentration of 200 mM. The column employed for HPLC-MS analysis was Poroshell 120 EC-C18 (2.7 μm), and samples were filtered through a short plug of Celite prior to analysis to eliminate the copper complexes.

We identified the peak in Figure S19 in the Supporting Information as a leaching product from the HPLC column when $\text{H}_2\text{O}_2/\text{Et}_3\text{N}$ is employed as oxidant for cellobiose degradation. Upon further analysis, the peak giving rise to $m/z = 118$ and 235 is consistent with protonated triethylamine *N*-oxide $[\text{Et}_3\text{NOH}]^+$ and $[(\text{Et}_3\text{NO})_2\text{H}]^+$, respectively.

Received: October 7, 2020

Published: October 23, 2020

