

Impact of COVID-19 outbreak by income: hitting hardest the most deprived

Jose Miguel Baena-Díez^{a,b,c}, Maria Barroso^{c,d}, Sara Isabel Cordeiro-Coelho^a, Jorge L. Díaz^{c,e}, María Grau^{c,f,g*}.

^aPrimary Care Centre La Marina, Catalan Institute of Health, Barcelona, Spain

^bIDIAP Jordi Gol, Catalan Institute of Health, Barcelona, Spain

^cCardiovascular Epidemiology and Genetics, IMIM – Hospital del Mar Medical Research Institute, Barcelona, Spain

^dEastfield Health, Ashburton, New Zealand

^eDocent Unit Preventive Medicine and Public Health Parc de Salut Mar – Barcelona Public Health Agency – Pompeu-Fabra University, Barcelona, Spain

^fConsortium for Biomedical Research in Epidemiology and Public Health (CIBERESP), Barcelona, Spain

^gDepartment of Medicine, University of Barcelona, Spain

*Corresponding author: María Grau, Cardiovascular Epidemiology and Genetics, IMIM – Hospital del Mar Medical Research Institute
Telf. +34 933160800; email. mgrau@imim.es

ABSTRACT

Background. The impact of the 2019 coronavirus disease (COVID-19) has many facets. This ecological study analyzed age-standardized incidence rates by economic level in Barcelona.

Methods. We evaluated confirmed cases of COVID-19 in Barcelona (Spain) between February 26 and April 19, 2020. Districts were classified according to most recent (2017) mean income data. The reference for estimating age-standardized cumulative incidence rates was the 2018 European population. The association between incidence rate and mean income by district was estimated with the Spearman rho.

Results. The lower the mean income, the higher the COVID-19 incidence (Spearman rho =0.83; p-value =0.003). Districts with the lowest mean income had the highest incidence of COVID-19 per 10 000 inhabitants; in contrast, those with the highest income had the lowest incidence. Specifically, the district with the lowest income had 2.5 times greater incidence of the disease, compared with the highest-income district [70 (95% confidence interval 66-73) vs 28 (25-31), respectively].

Conclusions. The incidence of COVID-19 showed an inverse socioeconomic gradient by mean income in the 10 districts of the City of Barcelona. Beyond healthcare for people with the disease, attention must focus on a health strategy for the whole population, particularly in the most deprived areas.

Key words: Healthcare disparities, Socioeconomic factors, Epidemiology, COVID-19, Coronavirus infections

Introduction

The outbreak of the 2019 coronavirus disease (COVID-19) in China's Hubei province quickly spread nearly all over the world, with staggering medical, social and economic consequences.¹ Worldwide, the pandemic accounts for more than 10 million confirmed cases and 500 000 deaths. While the US, Brazil, Russia or UK presented the highest absolute numbers, Belgium, UK, and Spain had the highest death rates.^{2,3} Thus, the figures made available in Spain by the Ministry of Health on June 30, 2020, showed 249 271 confirmed cases (5986 cases x 100 000 inhabitants), of which 125 183 were hospitalized and 11 664 required intensive care. Finally, this disease accounted for 28 355 deaths, with a case-fatality rate of 11.4%.⁴

Several authors have posed a possible socioeconomic gradient in the COVID-19 outbreak^{5, 6, 7} and differences in knowledge, attitudes, and practices towards COVID-19 by socioeconomic status.^{8, 9} Health inequalities have been pointed out in the context of the pandemic, not only in low- and middle-income countries, but also in high-income countries among deprived populations (e.g., having low income or lacking health insurance).^{10, 11, 12, 13} In addition, the COVID-19 crisis has revealed the fragility of ageing societies to cope with infectious diseases in modern times. Indeed, individuals older than 70 years accounted for more than 85% of registered deaths in Italy.¹⁴ However, few studies in high income countries have analysed the age-adjusted impact of the COVID-19 pandemic in deprived and non-deprived areas, as measured by average income.

The objective of this study was to analyse the differences in COVID-19 age-standardized incidence rate by mean income of the 10 districts of the City of Barcelona.

Methods

For this ecological study, all cases of COVID-19 confirmed by polymerase chain reaction (PCR) for which data were available to ascertain the patient's primary care centre in the City of Barcelona were extracted from the official COVID-19 registry of the Catalan Government's Department of Health. The primary care centres were then classified according to their corresponding district in order to obtain the total count of cases by district from February 26, when the first case reported in the city,¹⁵ until April 19, 2020. The most recent (2017) information on the average income by district was extracted from official data provided by the Barcelona City Council. The estimate used was the Family Available Income per capita, which considers the income available for consumption and accumulated savings per person.¹⁶ This analysis followed the regulations in European Union law on data protection and privacy for all individuals within the European Union (GDPR/2018), the Declaration of Helsinki on ethical principles for medical research involving human subjects.

The crude incidence rates were estimated per 10,000 individuals, assuming the age distribution (<25 years, ≥25 and <40 years, ≥40 and <65 years, ≥65 years) of COVID-19 cases in Barcelona was similar to that reported for Spain.¹⁷ To standardize the incidence rates by age, we used the direct method with the 2018 European population as the reference.¹⁸ Age-adjusted cumulative incidence was calculated with the corresponding 95% confidence interval (95%CI). To ascertain the statistical association between the age-adjusted incidence rate and the mean income by district, Spearman rho was adjusted. The analysis was performed with R statistical software (version 3.6.3) and Epidat software (version 3.1).

Results

Our estimates showed an overall cumulative incidence rate in Barcelona of 49.24 cases per 10,000 inhabitants between February 26 (date of first reported case in the city) until April 19, 2020. The ecological analysis to assess case distribution by district found that the lower the mean income, the higher the COVID-19 incidence (Spearman $\rho = 0.83$; p value = 0.003) (Figure 1). The districts with the lowest per capita income had the highest incidence of COVID-19 per 10 000 inhabitants, whereas the districts with the highest mean income had the lowest incidence (Figure 2). Specifically, the incidence rate in the most deprived district (Nou Barris) was 2.5 times higher than the district with the highest socio-economic level (Sarrià-Sant Gervasi). There are two exceptions on the gradient, Sant Andreu and Ciutat Vella. The divergent results in the Sant Andreu district could be explained by the high percentage (25%) of residents younger than 25 years, who are less vulnerable to COVID-19. In the district of Ciutat Vella (“Old City”), the results reflect the reality of this high-tourism area: low mean income in a relatively young population (>30% aged 25 to 39 years) and a high presence of non-resident foreign population (>25%). This also makes COVID-19 surveillance more difficult, as the public health system’s primary care settings are oriented to residents, while visitors would be served by emergency rooms and private clinics.

Discussion

The COVID-19 incidence rate from February 26 to April 19, 2020, followed a socio-economic gradient in Barcelona. Thus, the highest age-standardized incidence rate was observed in the most socioeconomically deprived district. This rate was 2.5 times greater than that observed in the district with the highest mean income. In addition, we

observed a sustained and significant upward trend in the incidence of COVID-19 according to declining mean income by district.

Regarding our results, a report has shown substantial variation in the rates for COVID-19 hospitalizations and deaths across the New York City boroughs. The Bronx –the borough with the highest proportion of racial/ethnic minorities, the most persons living in poverty, and the lowest levels of educational attainment– had higher rates of hospitalization and death related to COVID-19.⁶ In a recent report of the UK Biobank, striking gradients in risk of hospitalization for COVID-19 were noted according to race and a metric of socioeconomic deprivation.¹²

Improvements in the health literacy of citizens, defined as people's knowledge and capacity to obtain, process, and understand health information and services to make appropriate health decisions, could help to reduce the risk of infection spreading and increase understanding of the need for social responsibility and adherence to disease prevention measures.¹⁹ Along this line, Zhong et al. showed that Chinese residents of relatively high socioeconomic status, and particularly women, had good knowledge, optimistic attitudes, and appropriate practices towards COVID-19 during the initial rapid rise of the COVID-19 outbreak.⁸ In addition, racial disparities in knowledge, attitudes and practices regarding COVID-19 were described in the USA.⁹ Thus, understanding community risk and making decisions about community mitigation, including social distancing and strategic health care resource allocation, requires monitoring the numbers of COVID-19 cases, deaths, and changes in incidence in small, well characterized areas of the general population.²⁰

Our results pointed out that efforts to contain an epidemic cannot ignore health equity issues. As the most deprived areas in Barcelona had the highest COVID-19

incidence, disease-control efforts should be more intensive in districts with the most marginalized and vulnerable population. In addition, ensuring equal treatment opportunities for all is key, but financial protection during outbreak also matters greatly.²¹ The link between poverty and disease has already been explored in depth. If this vicious cycle is not broken, local problems of health inequity will remain or could even be exacerbated in areas experiencing an epidemic.¹¹ Previous studies have observed that the prevalence and severity of the COVID-19 pandemic is magnified because of the pre-existing epidemics of chronic disease,^{22,23} which also are socially patterned and associated with the social determinants of health.^{24,25} For instance, in a large cohort in Louisiana (USA), 76.9% of the patients hospitalized with COVID-19 and 70.6% of those who died were black, whereas only 31% of this ethnic group receives routine health care. Black patients in that cohort had higher prevalence of obesity, diabetes, hypertension, and chronic kidney disease at baseline than white patients.²⁶

Bambra et al. point out the rise of a syndemic, defined as a set of closely intertwined and mutual enhancing health problems that significantly affect the overall health status of a population within the context of a perpetuating configuration of noxious social conditions.⁶ This concept is rooted in Dahlgren and Whitehead's classic model of the determinants of health, which shows that individual lifestyles are embedded in social norms and networks and in living and working conditions, which in turn are related to the wider socioeconomic and cultural environment.²⁷ Thus, inequalities in chronic conditions arise as a result of inequalities in exposure to the social determinants of health: the conditions in which people 'live, work, grow and age'. These include working conditions, unemployment, and access to essential goods

and services (egg, water, sanitation and food), housing, and healthcare.⁶ Several plausible explanations can be suggested for the low rates observed in the districts with the highest socio-economic level. First, the work environment mediates a large part of the social gradient in the incidence of several common cardiovascular risk factors as shown in the Gazele cohort.²⁸ The authors discuss the need to include working conditions in policies aimed to reduce social inequalities in health. According to the Economic Co-operation and Development (OECD) report, workers possessing higher levels of skills were significantly more likely to telework when the COVID-19 crisis generated lockdowns that completely changed the working arrangements for millions of workers. Although employers were encouraged to develop home-based teleworking, the likelihood of this option was lower for those without tertiary education, with limited numeracy and literacy skills, and lacking adequate access to technology. The pandemic also exacerbates existing labour market inequalities, and the extent to which these inequalities could further worsen amidst intensified technology adoption in the pandemic's aftermath.²⁹ As Hamidi et al. pointed out, commuter transportation systems are the most significant risk factor for the spread of the COVID-19 pandemic, even greater than population density.³⁰ A case-study performed in Italy shows a direct relationship between daily cases and mobility choices in the preceding 21 days.³¹ Thus, workers with no possibility for home-based teleworking, many of them doing unskilled labour, were likely to have high-risk exposure during the pandemic. Finally, the strong links between housing and health could also be contributing to inequalities in COVID-19 incidence and outcome.^{6,32} Poor-quality housing is associated with overcrowding, which results in higher infection rates,³³ and lower socioeconomic groups have increased exposure to poor-quality or insecure housing and therefore have a higher rate

of negative health consequences.³⁴ In our study, the use of second homes for the confinement period also could have contributed to the low rates observed in the more privileged districts of Barcelona.

Our study has several strengths and limitations. The results were standardized by age because individuals aged 65 and older are more vulnerable to COVID-19 infection; however, we assumed the age distribution of COVID-19 cases of the whole Spanish territory for our population area. Nonetheless, the distribution in Spain is similar to that of several other western European countries (e.g. Italy,³⁵ Netherlands,³⁶ Portugal³⁷ or Sweden³⁸); thus, the distribution in Barcelona is likely to be similar despite potential internal differences across Spain. In addition, the age distribution could differ between the city districts, particularly in Ciutat Vella, with its large population of foreign non-residents and tourists. Despite age-standardization, some residual confounding may exist in the incidence rate calculation, particularly in districts with a high percentage of younger residents (e.g., Sant Andreu and Ciutat Vella). The ecological nature of our analysis, using aggregated data, has intrinsic limitations because variation within districts (i.e. small areas) is not considered. More in-depth analysis should be considered to improve our knowledge on this topic and inform public health strategies for disease prevention.

Conclusion

The incidence of COVID-19 presented an inverse socioeconomic gradient in the city of Barcelona according to average income by district. Attention should be focused not only on care for people with the disease but also on devising a health strategy for the whole population that promotes and supports good public hygiene practices and health

literacy, particularly adapted to the most deprived areas. Further studies are required to analyse whether the mortality from this infectious disease shows a similar pattern.

Acknowledgments The authors are grateful to Oriol Baena for the assistance with the graphic plots and to Elaine Lilly PhD for the English review.

Funding Dr. Maria Grau's was sponsored by Carlos III Health Institute FEDER (CPH17/00012)

Competing interest None

References

- 1 Peckham R. COVID-19 and the anti-lessons of history. *Lancet*. 2020;395:850–852.
- 2 Johns Hopkins Coronavirus Resource Center. Available at: <https://coronavirus.jhu.edu/map.html>
- 3 Vanden Driessche S. Excess mortality during the Covid-19 pandemic: Early evidence from England and Wales. *Soc Sci Med*. 2020;258:113101.
- 4 Ministry of Health - Spanish Government. Update #152. Coronavirus disease 2019 (COVID-19) [Actualización nº 152. Enfermedad por el coronavirus (COVID-19)]. 28.06.2020. Available at: https://www.mscbs.gob.es/profesionales/saludPublica/ccayes/alertasActual/nCov-China/documentos/Actualizacion_150_COVID-19.pdf

- 5 Chung RY, Dong D, Li MM. Socioeconomic gradient in health and the covid-19 outbreak. *BMJ*. 2020;**369**:m1329.
- 6 Bambra C, Riordan R, Ford J, Matthews F. The COVID-19 pandemic and health inequalities. *J Epidemiol Community Health*. 2020;jech-2020-214401.
- 7 Khalatbari-Soltani S, Cumming RG, Delpierre C, Kelly-Irving M. Importance of collecting data on socioeconomic determinants from the early stage of the COVID-19 outbreak onwards. *J Epidemiol Community Health*. 2020;jech-2020-214297. doi:10.1136/jech-2020-214297
- 8 Zhong BL, Luo W, Li HM, *et al*. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci*. 2020;**16**:1745–1752.
- 9 Alobuia WM, Dalva-Baird NP, Forrester JD, Bendavid E, Bhattacharya J, Kebebew E. Racial disparities in knowledge, attitudes and practices related to COVID-19 in the USA. *J Public Health (Oxf)*. 2020;fdaa069. doi:10.1093/pubmed/fdaa069
- 10 Wadhwa RK, Wadhwa P, Gaba P, *et al*. Variation in COVID-19 Hospitalizations and Deaths Across New York City Boroughs. *JAMA*. 2020;10.1001/jama.2020.7197.
- 11 The Lancet. Redefining vulnerability in the era of COVID-19. *Lancet*. 2020;395:1089.
- 12 Patel AP, Paranjpe MD, Kathiresan NP, Rivas MA, Khera AV. Race, Socioeconomic Deprivation, and Hospitalization for COVID-19 in English

- participants of a National Biobank. Preprint. medRxiv.
2020;2020.04.27.20082107.
- 13 Mahajan UV, Larkins-Pettigrew M. Racial demographics and COVID-19 confirmed cases and deaths: a correlational analysis of 2886 US counties. *J Public Health (Oxf)*. 2020;fdaa070. doi:10.1093/pubmed/fdaa070
- 14 Richardson SJ, Carroll CB, Close J, *et al*. Research with older people in a world with COVID-19: identification of current and future priorities, challenges and opportunities. *Age Ageing*. 2020;afaa149. doi:10.1093/ageing/afaa149
- 15 Interactive Map of COVID-19 Cases [Mapa interactiu de casos per ABS]. Available at: <http://aquas.gencat.cat/ca/actualitat/ultimes-dades-coronavirus/mapa-per-abs/>
- 16 Family Available Income per capita in Barcelona (2017) [Distribució territorial de la renda familiar disponible per càpita a Barcelona (2017)]. Ajuntament de Barcelona. Available at: https://ajuntament.barcelona.cat/barcelonaeconomia/sites/default/files/RFD_2017_BCN.pdf
- 17 COVID-19 in Spain. Report #26, April 27, 2020 [Informe sobre la situación de COVID-19 en España nº 26. 27 de abril de 2020]. Available at: <https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAVE/EnfermedadesTransmisibles/Documents/INFORMES/Informes%20COVID-19/Informe%20n%C2%BA%2026.%20Situaci%C3%B3n%20de%20COVID-19%20en%20Espa%C3%B1a%20a%2027%20de%20abril%20de%202020.pdf>
- 18 Eurostat, your key to European statistics. Available at: <https://ec.europa.eu/eurostat/>

- 19 Xu C, Zhang X, Wang Y. Public Health Emergency, Health Literacy and Social Panic: Mapping of COVID-19 Based on the Web Search Data. *J Med Internet Res.* 2020;10.2196/18831.
- 20 CDC COVID-19 Response Team. Geographic Differences in COVID-19 Cases, Deaths, and Incidence - United States, February 12-April 7, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69:465–471.
- 21 Wang Z, Tang K. Combating COVID-19: health equity matters. *Nat Med.* 2020;26:458. 6
- 22 Uribarri A, Núñez-Gil IJ, Aparisi A, *et al.* Impact of renal function on admission in COVID-19 patients: an analysis of the international HOPE COVID-19 (Health Outcome Predictive Evaluation for COVID 19) Registry. *J Nephrol.* 2020;10.1007/s40620-020-00790-5.
- 23 Koh J, Shah SU, Chua PEY, Gui H, Pang J. Epidemiological and Clinical Characteristics of Cases During the Early Phase of COVID-19 Pandemic: A Systematic Review and Meta-Analysis. *Front Med (Lausanne).* 2020;7:295.
- 24 Aguilar-Palacio I, Martinez-Beneito MA, Rabanaque MJ, *et al.* Diabetes mellitus mortality in Spanish cities: Trends and geographical inequalities. *Prim Care Diabetes.* 2017;11(5):453-460.
- 25 Sivén SS, Niiranen TJ, Aromaa A, Koskinen S, Jula AM. Social, lifestyle and demographic inequalities in hypertension care. *Scand J Public Health.* 2015;43(3):246-253.
- 26 Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and Mortality among Black Patients and White Patients with Covid-19. *N Engl J Med.* 2020;382(26):2534-2543.

- 27 Dahlgren G, Whitehead M, European strategies for tackling social inequities in health: Levelling up Part 2 [online]. Available at:
http://www.euro.who.int/__data/assets/pdf_file/0018/103824/E89384.pdf
- 28 Meneton P, Hoertel N, Wiernik E, et al. Work environment mediates a large part of social inequalities in the incidence of several common cardiovascular risk factors: Findings from the Gazel cohort. *Soc Sci Med.* 2018;**216**:59-66.
- 29 Espinoza R, Reznikova L. Who can log in? The importance of skills for the feasibility of teleworking arrangements across OECD countries. OECD Social, Employment and Migration Working Papers No. 242. OECD 2020. Available at:
<https://www.oecd-ilibrary.org/docserver/3fl15a10-en.pdf?expires=1593681356&id=id&accname=guest&checksum=90922E71CD060212E61E5ACFB2B30DCC>
- 30 Hamidi S, Sabouri S, Ewing R. Does Density Aggravate the COVID-19 Pandemic? *J Am Plann Assoc.* 2020 (in press)
- 31 Carteni A, Di Francesco L, Martino M. How mobility habits influenced the spread of the COVID-19 pandemic: Results from the Italian case study. *Sci Total Environ.* 2020;**741**:140489.
- 32 Hernández D, Swope CB. Housing as a Platform for Health and Equity: Evidence and Future Directions. *Am J Public Health.* 2019;**109**:1363–1366.
- 33 Gibson M, Petticrew M, Bambra C, Sowden AJ, Wright KE, Whitehead M. Housing and health inequalities: a synthesis of systematic reviews of interventions aimed at different pathways linking housing and health. *Health Place.* 2011;**17**(1):175-184.

- 34 McNamara CL, Balaj M, Thomson KH, Eikemo TA, Bambra C. The contribution of housing and neighbourhood conditions to educational inequalities in non-communicable diseases in Europe: findings from the European Social Survey (2014) special module on the social determinants of health. *Eur J Public Health*. 2017;**27**(suppl_1):102-106.
- 35 National Institute of Health [Istituto Superiore di Sanita]. Available at: <https://www.epicentro.iss.it/coronavirus/>
- 36 National Institute for Public Health and the Environment Ministry of Health, Welfare and Sport [Rijksinstituut voor Volksgezondheid en Milieu Ministerie van Volksgezondheid, Welzijn en Spor]. Available at: <https://www.rivm.nl/>
- 37 National Health Authority [Direção-Geral da Saúde]. Available at: <https://www.dgs.pt/>
- 38 Folkhälsomyndigheten. The Public Health Agency of Sweden. Available at: <https://www.folkhalsomyndigheten.se/the-public-health-agency-of-sweden/>

FIGURE LEGENDS

Figure 1. Average income and age-standardized incidence rates by district.

Figure 2. Map showing the 10 districts of Barcelona, classified by quartiles (Q) of mean income in euros (Q1 <17 418; Q2 \geq 17 418 and <18 910; Q3 \geq 18 910 and <25 874; Q4 \geq 25 874) and showing the age-standardized incidence rate (95% confidence interval). Base layer extracted from icgc.cat.

Figure 1.




Figure 2.

