



Scientific Letter

A New and More Sensitive Method to Integrate the Desaturation Distance Ratio During a 6-Minute Walking Test in Chronic Respiratory Diseases: Physiological Correlates

Un método nuevo y más sensible para integrar la relación entre distancia y desaturación durante la prueba de la marcha de seis minutos en las enfermedades respiratorias crónicas: correlaciones fisiológicas

Dear Editor,

Exercise limitation is common in many respiratory diseases like chronic obstructive pulmonary disease (COPD), interstitial lung diseases (ILD), and pulmonary arterial hypertension (PAH).^{1–3} The six-minute walking test (6MWT) is widely used to assess exercise response in clinical practice.⁴ Also, previous studies have shown that 6MWT reflects equally well to the conventional constant-load test on the cycle ergometer the degree of impairment in hemodynamic response to exercise.⁵ The desaturation distance ratio (DDR) has recently been proposed as a potential integrative marker of exercise response that considers the distance walked during the test (6MWD) and the oxygen desaturation area estimated from the difference between the transcutaneous oxygen saturation value (SpO₂) measured every minute vs. a theoretical maximum of 100%.^{6,7} We hypothesized that the difference between SpO₂ and the SpO₂ value determined at rest would better reflect the response to exercise in different diseases. This study tests this hypothesis by comparing the DDR values obtained using the original and the new method in patients ($n = 308$) with three prevalent chronic respiratory diseases (COPD, ILD, and PAH). Besides, we sought to investigate the correlation of DDR with resting lung function measurements in these three diseases.

In this retrospective analysis, we included 104 patients with COPD, 104 with ILD, and 100 with PAH, all diagnosed according to international guidelines,^{8–10} who performed a 6MWT between October 2014 and May 2015. All patients were clinically stable, and none was participating in any rehabilitation program or clinical trial. We excluded patients on long-term oxygen therapy, mechanical ventilation, and/or tracheostomized. The study was approved by the Ethics Committee of our hospital (HCB/2017/0469).

Demographic and anthropometric parameters were recorded and resting lung function was measured following international standards.^{11,12} Reference values were those of Roca et al.^{13,14} In patients with PAH, pulmonary hemodynamics were determined by right heart catheterization, also following standard procedures.⁸ All participants performed a 6MWT according to the international guidelines.⁴ Reference values were those of Enright et al.¹⁵ During the test, SpO₂ and heart rate (HR) were monitored continuously by Pulsox[®]-300 (Minolta Co., Tokyo, Japan). Absolute changes in

SpO₂ (Δ SpO₂) were calculated by subtracting the values at baseline from those determined immediately after 6MWT was finished.¹⁶ DDR was calculated as the desaturation area (DA) over the 6MWD.⁷ DA was calculated using two different methods, the one proposed originally by Ijiri et al.⁷ (vs. a theoretical maximal SpO₂ value of 100%), and the one tested in this study (vs. the actual SpO₂ value determined in each patient at rest, before the 6MWT starts). Results are presented as mean \pm standard deviation for those distributed normally and as median and (25–75% percentile) for those non-normally distributed. Categorical variables are presented as proportions. ANOVA was used to compare groups. Correlations were assessed using Spearman's correlation test, and multivariable linear regression was used to explore the relationship of DDR with resting lung function variables, adjusted by age and BMI, in each of the three diseases studied here. A p value < 0.05 was considered statistically significant.

As shown in **Table 1**, most COPD patients were male, with a mean age of 66 ± 8 years, and had moderate-severe airflow limitation and moderately decreased carbon monoxide diffusing capacity (DL_{CO}). ILD patients were of similar age (68 ± 11), but they included a higher proportion of females (45%) and showed a mild restrictive pattern with a moderate DL_{CO} reduction. PAH patients were younger and mainly females, with normal spirometry and lung volumes but a moderately reduced DL_{CO}; their mean pulmonary arterial pressure was 47 (39–54) mmHg (**Table 1**). The 6MWD was similar in all patients and generally well preserved. SpO₂ at rest was lower in COPD and decreased during exercise in all diseases (**Fig. 1**). Using the calculation method initially proposed by Ijiri^{6,7} neither the DA nor DDR was significantly different across diseases (**Table 1**). By contrast, considering the resting SpO₂ value in each individual as its own reference to calculate DA (and, therefore, DDR). Multivariable linear regression analysis showed that DL_{CO} %pred. and resting SpO₂ were independent determinants of DDR in all three diseases. Besides, we observed that, in COPD, the severity of airflow limitation (FEV₁%ref) was also an independent DDR predictor.

The main findings of this study are that: (1) the calculation of DDR considering the resting SpO₂ is more sensitive than the initially proposed method (that considered a theoretical reference value of 100% SpO₂^{6,7}); and (2) DL_{CO} and resting SpO₂ were independent determinants of DDR shared by the three diseases studied here, plus the severity of airflow limitation in patients with COPD.

A few previous studies have investigated DDR (using the original method^{6,7}) in patients with ILD⁶ or COPD.¹⁷ Like in these previous studies, we found that the new DDR calculation in patients with ILD and COPD was significantly related to DL_{CO} and, to a lesser extent, to FEV₁. To our knowledge, this is the first study to investigate DDR in patients with PAH, to contrast it with patients with ILD and COPD, and to investigate potential DDR correlations with lung function at rest in these three diseases.

Table 1

Characteristics of participants and parameters determined during the 6MWT. Results are expressed as the mean ± SD if data is normally distributed or as the median (25–75% percentile) if not.

	COPD (n = 104)	ILD (n = 104)	PAH (n = 100)
Age, years	66 ± 8	68 ± 11	57 ± 16*
Gender, n (% males)	85 (82)	57 (55)	26 (26)*
BMI, kg/m ²	27 ± 5	29 ± 5	27 ± 5*
Smoking exposure, pack-years	46 (35–71)	1 (0–34)	0 (0–14)*
FVC, % predicted	75 ± 16	69 ± 18	85 ± 18*
FEV ₁ , % predicted	46 ± 17	71 ± 17	78 ± 19*
FEV ₁ /FVC, ratio	45 ± 12	76 ± 8	70 ± 10*
TLC, %predicted	106 (94–121)	70 (55–82)	90 (82–100)*
DL _{CO} , %predicted	51 (38–62)	46 (33–59)	54 (42–68)*
mPAP, mmHg	–	–	47 (39–54)
6MWD, meters	456 ± 98	473 ± 113	452 ± 119
Borg dyspnea score at rest	0 (0–2)	0 (0–1)	0 (0–1)
ΔBorg dyspnea score with exercise	2 (1; 4)	3 (1; 4)	3 (1; 4)
SpO ₂ at rest, %	95 (93–96)	96 (94–97)	96 (95–97)*
SpO ₂ at end, %	91 (86–93)	91 (82–94)	89 (83–93)
ΔSpO ₂ (at rest - at end), %	–4 (–8; –2)	–6 (–13; –2)	–7 (–11; –4)*
Original DA	55 (40–77)	54 (35–98)	61 (40–83)
Original DDR	0.121 (0.09–0.17)	0.118 (0.07–0.21)	0.133 (0.10–0.23)
New DA	25 (11–38)	28 (14–62)	34 (21–55)*
New DDR	0.052 (0.021–0.090)	0.063 (0.029–0.144)	0.075 (0.044–0.133)*

Abbreviations: BMI, body mass index; FVC, forced vital capacity; FEV₁, forced expiratory volume in the first second; TLC, total lung capacity; DL_{CO}, diffusion capacity for carbon monoxide; mPAP, mean pulmonary arterial pressure; 6MWD, six-minute walking distance; SpO₂, peripheral capillary oxygen saturation; DA, desaturation area; DDR, desaturation distance ratio.

* p < 0.05.




Fig. 1. SpO₂ value at rest and during the 6MWT in the three chronic pulmonary diseases explored here. For further explanations, see text.

We could not explore the potential prognostic value of DDR in these diseases because this study was retrospective. However, DL_{CO} is a well-established prognostic factor in different chronic respiratory diseases,^{10,18} so its significant and independent relationship with DDR suggests that DDR may also be a good prognostic factor. This will have to be tested prospectively, but it is of note that Hsieh et al. have recently reported that the distance-desaturation product, an index similar to DDR, predicted 6-year mortality in a prospective study of 69 patients with stable non-CF bronchiectasis.¹⁹ Also, In 81 patients with IPF, Lettieri et al. found that each component of the 6MWT independently predicted mortality in IPF with greater accuracy than spirometry. However, a composite of both parameters, the distance-saturation product, provides slightly greater accuracy and represents a novel measure for assessing survival in these patients.²⁰ On the other hand, DDR can be useful to assess the effects of a therapeutic intervention, such as the impact of supplementary oxygen during exercise or other.

Our study has strengths and limitations. Among the former, the description of a new method to calculate DDR in a large number of well-characterized patients with COPD, ILD, and PAH. Among the latter, its retrospective design. Additionally, we did not have information about the patient's exacerbations, which would have allowed a better characterization of the sample.

In conclusion, this study describes a new and more sensitive method to calculate DDR in patients with COPD, ILD, and PAH and shows that DL_{CO} and resting SpO₂ (plus airflow limitation in patients with COPD) were independent determinants of the new DDR. In terms of clinical implication, being new DDR a more

sensitive method, it could offer information about response to drug's treatment and to rehabilitation programs.

Authors' contributions

X. Alsina-Restoy: Conceptualization, Formal analysis, Methodology, Writing – original draft, Writing – review & editing. F. Burgos: Conceptualization, Formal analysis, Methodology, Writing – original draft, Writing – review & editing. R. Torres-Castro: Formal analysis, Writing – original draft, Writing – review & editing; Y. Torralba-García: Writing – original draft, Writing – review & editing; E. Arismendi: Writing – original draft, Writing – review & editing; J.A. Barberà: Writing – original draft, Writing – review & editing; A. Agustí: Formal analysis, Writing – original draft, Writing – review & editing; I. Blanco: Conceptualization, Formal analysis, Methodology, Writing – original draft, Writing – review & editing.

Funding

None.

Conflict of interests

The authors declare no conflict of interest directly or indirectly related to the manuscript contents except FB (which declares them in his COI).

References

1. Dodson MW, Brown LM, Elliott CG. Pulmonary arterial hypertension. *Heart Fail Clin.* 2018;14:255–69, <http://dx.doi.org/10.1016/j.hfc.2018.02.003>.
2. Rabe KF, Watz H. Chronic obstructive pulmonary disease. *Lancet.* 2017;389:1931–40, [http://dx.doi.org/10.1016/S0140-6736\(17\)31222-9](http://dx.doi.org/10.1016/S0140-6736(17)31222-9).
3. Antoniou KM, Margaritopoulos GA, Tomassetti S, Bonella F, Costabel U, Poletti V. Interstitial lung disease. *Eur Respir Rev.* 2014;23:40–54, <http://dx.doi.org/10.1183/09059180.00009113>.
4. Holland AE, Spruit MA, Troosters T, Puhan MA, Pepin V, Saey D, et al. An official European respiratory society/American thoracic society technical standard: field walking tests in chronic respiratory disease. *Eur Respir J.* 2014;44:1428–46, <http://dx.doi.org/10.1183/09031936.00150314>.
5. Vasilopoulou MK, Vogiatzis I, Nasis I, Spetsioti S, Cherouveim E, Koskolou M, et al. On- and off-exercise kinetics of cardiac output in response to cycling and walking in COPD patients with GOLD Stages I–IV. *Respir Physiol Neurobiol.* 2012;181:351–8, <http://dx.doi.org/10.1016/j.resp.2012.03.014>.
6. Pimenta SP, da Rocha RB, Baldi BG, Kawassaki AdM, Kairalla RA, Carvalho CRR. Desaturation – distance ratio: a new concept for a functional assessment of interstitial lung diseases. *Clinics (Sao Paulo).* 2010;65:841–6, <http://dx.doi.org/10.1590/S1807-59322010000900005>.
7. Ijiri N, Kanazawa H, Yoshikawa T, Hirata K. Application of a new parameter in the 6-minute walk test for manifold analysis of exercise capacity in patients with COPD. *Int J Chron Obstruct Pulmon Dis.* 2014;9:1235–40, <http://dx.doi.org/10.2147/COPD.S71383>.
8. Galiè N, Humbert M, Vachiery JL, Gibbs S, Lang I, Torbicki A, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Respir J.* 2015;46:903–75, <http://dx.doi.org/10.1183/13993003.01032-2015>.
9. Singh D, Agusti A, Anzueto A, Barnes PJ, Bourbeau J, Celli BR, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease: the GOLD science committee report 2019. *Eur Respir J.* 2019;53:1900164, <http://dx.doi.org/10.1183/13993003.00164-2019>.
10. Wells AU, Hirani N. Interstitial lung disease guideline: the British Thoracic Society in collaboration with the Thoracic Society of Australia and New Zealand and the Irish Thoracic Society. *Thorax.* 2008;63 Suppl. V:v1–58, <http://dx.doi.org/10.1136/thx.2008.101691>.
11. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J.* 2005;26:319–38, <http://dx.doi.org/10.1183/09031936.05.00034805>.
12. MacIntyre N, Crapo RO, Viegi G, Johnson DC, van der Grinten CP, Brusasco V, et al. Standardisation of the single-breath determination of carbon monoxide uptake in the lung. *Eur Respir J.* 2005;26:720–35, <http://dx.doi.org/10.1183/09031936.05.00034905>.
13. Roca J, Burgos F, Sunyer J, Saez M, Chinn S, Antó JM, et al. Reference values for forced spirometry. *Eur Respir J.* 1998;11:1354–62, <http://dx.doi.org/10.1183/09031936.98.11061354>.
14. Roca J, Rodriguez-Roisin R, Cobo E, Burgos F, Perez J, Clausen JL. Single breath carbon monoxide diffusing capacity prediction equations from a mediterranean population. *Am Rev Respir Dis.* 1989;141:1026–32.
15. Enright PL, Sherrill D. Reference equations for the six-minute walk in healthy adults. *Am J Respir Crit Care Med.* 1998;158:1384–7.
16. Johnson MJ, Close L, Gillon SC, Molassiotis A, Lee PH, Farquhar MC. Use of the modified Borg scale and Numerical Rating Scale to measure chronic breathlessness: a pooled data analysis. *Eur Respir J.* 2016;47:1861–4.
17. Fujimoto Y, Oki Y, Kaneko M, Sakai H, Misu S, Yamaguchi T, et al. Usefulness of the desaturation–distance ratio from the six-minute walk test for patients with COPD. *Int J COPD.* 2017;12:2669–75, <http://dx.doi.org/10.2147/COPD.S143477>.
18. Hsu VM, Chung L, Hummers LK, Shah A, Simms R, Bolster M, et al. Risk factors for mortality and cardiopulmonary hospitalization in systemic sclerosis patients at risk for pulmonary hypertension, in the pharos registry. *J Rheumatol.* 2019;46:176–83, <http://dx.doi.org/10.3899/jrheum.180018>.
19. Hsieh MH, Fang YF, Chung FT, Lee CS, Chang YC, Liu YZ, et al. Distance-saturation product of the 6-minute walk test predicts mortality of patients with non-cystic fibrosis bronchiectasis. *J Thorac Dis.* 2017;9:3168–76, <http://dx.doi.org/10.21037/jtd.2017.08.53>.
20. Lettieri CJ, Nathan SD, Browning RF, Barnett SD, Ahmad S, Shorr AF. The distance-saturation product predicts mortality in idiopathic pulmonary fibrosis. *Respir Med.* 2006;100:1734–41, <http://dx.doi.org/10.1016/j.rmed.2006.02.004>.

Xavier Alsina-Restoy^{a,b}, Felip Burgos^{a,b,c},
Rodrigo Torres-Castro^{a,d}, Yolanda Torralba-García^{a,b,c},
Ebymar Arismendi^{a,c}, Joan Albert Barberà^{a,b,c}, Àlvar Agustí^{a,b,c,1},
Isabel Blanco^{a,b,c,*,1}

^a Servei de Pneumologia, Institut Clínic Respiratori, Hospital Clínic de Barcelona, Universitat de Barcelona, Spain

^b Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

^c Biomedical Research Network in Respiratory Diseases (CIBERES), Madrid, Spain

^d Department of Physical Therapy, Faculty of Medicine, University of Chile, Santiago, Chile

* Corresponding author.

E-mail address: iblanco2@clinic.cat (I. Blanco).

¹ Co-senior authors.