



Enfermería Clínica

www.elsevier.es/enfermeriaclinica



ORIGINAL ARTICLE

Pre-pregnancy overweight and obesity prevalence and relation to maternal and perinatal outcomes

Elena González-Plaza^{a,b,*}, Jordi Bellart^{a,c}, Maria Àngels Martínez-Verdú^{a,b},
Àngela Arranz^{a,b}, Leila Luján-Barroso^{b,d}, Gloria Seguranyes^{b,e}

^a Maternal-Fetal Medicine Department at BCNatal, Barcelona Clinic Hospital, 08028 Barcelona, Spain

^b Department of Nursing: Public, Mental and Maternity and Child Health, School of Nursing, Faculty of Medicine and Health Science, University of Barcelona, 08907 L'Hospitalet del Llobregat, Barcelona, Spain

^c Department of Medicine, Faculty of Medicine and Health Science, University of Barcelona, 08036 Barcelona, Spain

^d Unit of Nutrition and Cancer, Cancer Epidemiology Research Programme, Bellvitge Biomedical Research Institute, Catalan Institute of Oncology, 08907 L'Hospitalet de Llobregat, Spain

^e Research Group on Sexual and Reproductive Health Care "GRASSIR", 08007 Barcelona, Spain

Received 27 August 2020; accepted 25 April 2021

KEYWORDS

Pregnancy;
Obesity;
Overweight;
Body Mass Index;
Pregnancy complications

Abstract

Objective: To identify the prevalence of pre-pregnancy overweight/obesity in pregnant women and its relationship with socio-demographic factors and to describe the maternal and perinatal outcomes in a Barcelona hospital (Spain).

Method: A descriptive cross-association study, with retrospective data collection, was performed Barcelona Hospital. The data of 5447 pregnant women who delivered at ≥ 23 weeks of gestation were included. Body Mass Index (BMI) data were categorised into World Health Organization classifications. p values $< .05$ (two-tailed) were considered significant. Logistic regression models were performed.

Results: The prevalence of pre-pregnancy obesity was 8.4% and 18.9% for overweight. Gestational diabetes was more frequent in pre-pregnancy overweight/obesity (OR 1.92: 95% CI 1.54–2.40 and OR 3.34: 95% CI 2.57–4.33), as were preeclampsia (OR 2.08: 95% CI 1.55–2.79 and OR 3.35: 95% CI 2.38–4.71), induction of labour (OR 1.19: 95% CI 1.02–1.38 and OR 1.94: 95% CI 1.57–2.10), caesarean section (OR 1.41: 95% CI 1.21–1.65 and OR 2.68: 95% CI 2.18–3.29), prematurity (OR 1.28: 95% CI 1–1.65 and OR 1.79: 95% CI 1.32–2.44) and macrosomia (OR 1.87: 95% CI 1.43–2.46 and OR 2.03: 95% CI 1.40–2.93).

Conclusions: One in four pregnant women had pre-pregnancy overweight or obesity. This study shows the relationship between pre-pregnancy overweight or obesity with adverse maternal and perinatal outcomes.

© 2021 Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: eplaza@clinic.cat (E. González-Plaza).

PALABRAS CLAVE

Embarazo;
Obesidad;
Sobrepeso;
Índice de masa corporal;
Complicaciones en el embarazo

Prevalencia de sobrepeso y obesidad preconcepcional en mujeres gestantes y relación con los resultados maternos y perinatales

Resumen

Objetivo: Identificar la prevalencia de obesidad o sobrepeso preconcepcional y su relación con los factores sociodemográficos, y describir los resultados maternos y perinatales en un hospital de Barcelona (España).

Método: Estudio descriptivo de asociación cruzada, con recogida de datos retrospectiva en un hospital de la ciudad de Barcelona (España). Se analizó la información de 5.447 embarazadas con parto ≥ 23 semanas de gestación. El índice de masa corporal fue categorizado según la clasificación de la Organización Mundial de la Salud. Se realizó un análisis estadístico bilateral asumiendo un valor α igual a 0,05. Se realizaron modelos de regresión logística.

Resultados: La prevalencia de obesidad preconcepcional fue del 8,4% y la del sobrepeso, del 18,9%. Las gestantes con obesidad o sobrepeso preconcepcional presentaron con mayor frecuencia diabetes gestacional (OR 1,92; IC 95% 1,54 a 2,40 y OR 3,34; IC 95% 2,57 a 4,33), preeclampsia (OR 2,08; IC 95% 1,55 a 2,79 y OR 3,35; IC 95% 2,38 a 4,71), inducción del parto (OR 1,19; IC 95% 1,02 a 1,38 y OR 1,94; IC 95% 1,57 a 2,10), cesárea (OR 1,41; IC 95% 1,21 a 1,65 y OR 2,68; IC 95% 2,18 a 3,29), prematuridad (OR 1,28; IC 95% 1 a 1,65 y OR 1,79; IC 95% 1,32 a 2,44) y macrosomía (OR 1,87; IC 95% 1,43 a 2,46 y OR 2,03; IC 95% 1,40 a 2,93).

Conclusiones: Una de cada 4 gestantes presentó sobrepeso u obesidad preconcepcional. Se observó relación entre el sobrepeso u obesidad preconcepcional con la presentación de resultados maternos y perinatales adversos.

© 2021 Elsevier España, S.L.U. Todos los derechos reservados.

What is known?

Obesity is an increasingly prevalent public health problem in our society. Preconception obesity is a risk factor for complications during pregnancy and delivery, as well as for the appearance of perinatal complications.

What this contribute?

The prevalence of obesity and preconception overweight observed in a Barcelona hospital was lower than in other Spanish cities. This study points out the relationship between some sociodemographic factors and pregnancy in women with an inadequate Body Mass Index. Preconception overweight and obesity were associated with adverse maternal and perinatal outcomes.

Introduction

Obesity is an increasingly more prevalent health problem in our society.¹ Women who begin pregnancy with a Body Mass Index (BMI) $> 25 \text{ kg/m}^2$ are at higher risk of maternal adverse outcomes.² In Europe, the prevalence of pre-pregnancy overweight and obesity is between 26.8% and 54%.³ However, there are scarce data concerning pre-pregnancy obesity women who lives in the Mediterranean region.⁴

At the same time, socio-demographic and obstetric factors seem to be related to women beginning pregnancy with overweight and obesity.⁵

The socio-demographic characteristics, as well as the diet and lifestyles of pregnant women of southern Europe, could differ from the women of central-northern Europe, and therefore, it could influence perinatal outcomes.

The aims of this study were to identify the prevalence of pre-pregnancy overweight/obesity in pregnant women and its relation with socio-demographic factors and to describe the maternal and perinatal outcomes in a hospital of Barcelona city (Spain).

Method

Study design and setting.

We conducted a descriptive cross association study, with retrospective data collection, of all pregnant women who gave birth in a high maternity complexity hospital in "Clinic Hospital of Barcelona" from January 1, 2015 to December 31, 2016 in Barcelona (Spain).

Study population

The inclusion criteria were women with delivery after 23 weeks of gestational age (GA). Multiple pregnancies and women in whom the pre-pregnancy BMI was not available in the electronic medical record were excluded from the study.

Study variables

The following variables were collected: woman's age in complete years, pre-pregnancy BMI in kg/m² based on the weight reported by the woman herself⁶ and classified into the following categories: underweight (BMI < 18.5 kg/m²); normal weight (BMI between 18.5 and 24.9 kg/m²); overweight (BMI between 25 and 29.9 kg/m²) and obesity (BMI ≥ 30 kg/m²),⁷ country of origin (Spanish, foreign), educational level (primary, secondary, higher), employed (yes, no), previous births (yes, no), gestational diabetes was defined as women with diabetes onset during pregnancy,⁸ preeclampsia was defined as women with hypertension onset during pregnancy,⁹ type of onset of labour (induction, spontaneous, elective caesarean section), type of delivery (spontaneous vaginal birth, instrumental, caesarean section). The variable of GA of the newborn was categorised as premature (<37 GA), term (37–41.6 GA) and post-term (≥42 GA).¹⁰ The weight of the neonate was classified as underweight (<2500 g), normal weight (2500–3999 g) and macrosome (≥4000 g).

Data collection

The data was obtained, retrospectively, from the hospital's computerised medical record system. In February 2017, the hospital information area carried out an automated and anonymised extraction of all the records of women with labour assisted during the two years of study, and the data was downloaded in an "Excel" spreadsheet to which only the research team had access.

Statistical analysis

Descriptive data are presented as number and percentage, and the mean and standard deviation (SD).

Bivariate analysis was performed between the socio-demographic variables and the pre-pregnancy BMI. For the comparison of the categorical variables, Chi-Square test was used. To compare quantitative variables, variance analysis (ANOVA) was performed.

Adjusted multinomial logistic regression was performed with the objective of identifying the socio-demographic and obstetric factors related to the pre-pregnancy BMI, where BMI was assumed a nominal variable taking the normal weight category as the reference group of women. Logistic regression models were performed to evaluate the perinatal and maternal results related to pre-pregnancy BMI of pregnant women. Adjusted Odds Ratio (OR) and 95% confidence intervals (95% CI) were calculated for each model. All statistical tests were bilateral and *p* values <0.05 were considered significant. Descriptive analyses were carried out with the statistical package IBM SPSS version 25®, and multivariate analyses were performed with the statistical package SAS version 9.4® (SAS Institute, Cary, NC, USA).

Ethical considerations

Authorisation was obtained from the Ethics and Clinical Research Committee of the Clinic Hospital of Barcelona,




Figure 1 Study flow chart. GA: gestational age; BMI: body mass index.

Code: HCB/2017/0309. At all times the anonymity and confidentiality of the data were preserved in accordance with the Spanish Organic Law 3/2018, of December 5, about Protection of Personal Data and guarantee of digital rights. The informed consent was exempt because the data were obtained from medical records.

Results

A sample of 6236 women who were assisted at delivery was obtained. Of these, 348 women with multiple births, 25 with delivery before 23 GA, and 416 women in whom the pre-pregnancy BMI did not appear in the computerised medical record were excluded. Finally, the data of 5447 pregnant women (87.3%) were analysed (Fig. 1).

The mean pre-pregnancy BMI was 23.4 kg/m² (95% CI 23.3–23.5). The prevalence of women with pre-pregnancy overweight was 18.9% (*n* = 1032, 95% CI 17.3–20.2), and the prevalence of women with pre-pregnancy obesity was 8.4% (*n* = 458, 95% CI 7.6–9.7). Thus, the percentage of the studied women who presented pre-pregnancy overweight or obesity was 27.3% (*n* = 1490, 95% CI 26.2–28.5).

Table 1 shows the socio-demographic and obstetric characteristics and their relationship with the pre-pregnancy weight status of the pregnant women included in the study. Table 2 shows the results of multinomial logistic regression between the socio-demographic and obstetric characteristics and pre-pregnancy BMI.

It was found that there was relationship between maternal age, country of origin, educational level, employment status, and having previous births, with women's pre-pregnancy BMI.

Pregnant women with a lower mean age were more frequently underweight. Foreign women had more frequently overweight compared to women of Spanish origin. Women with a secondary or higher education presented less frequently pre-pregnancy overweight and obese than pregnant

Table 1 Socio-demographic and obstetric characteristics and relationship with the pre-pregnancy weight status of pregnant women.

| | Total n = 5447 | Underweight n (%) n = 278 (5.1) | Normal weight n (%) n = 3679 (67.5) | Overweight n (%) n = 1032 (18.9) | Obesity n (%) n = 458 (8.4) | p value |
|--------------------------|-------------------|---------------------------------------|-------------------------------------------|----------------------------------------|-----------------------------------|----------|
| <i>Age: years</i> | | | | | | |
| Means (SD) | 32.9 (5.4) | 31.5 (5.9) | 33.1 (5.2) | 32.7 (5.6) | 32.3 (5.8) | <0.001* |
| <i>Country of origin</i> | n = 5447 | n = 278 | n = 3679 | n = 1032 | n = 458 | |
| Spanish | 3527 (64.8) | 186 (5.3) | 2467 (69.9) | 589 (16.7) | 285 (8.1) | |
| Foreign | 1920 (35.2) | 92 (4.8) | 1212 (63.1) | 443 (23.1) | 173 (9) | <0.001** |
| <i>Educational level</i> | n = 5218 | n = 265 | n = 3554 | n = 983 | n = 416 | |
| Primary | 655 (12.6) | 29 (4.4) | 336 (51.3) | 192 (29.3) | 98 (15) | |
| Secondary | 1567 (30) | 66 (4.2) | 950 (60.6) | 374 (23.9) | 177 (11.3) | |
| Higher | 2996 (57.4) | 170 (5.7) | 2268 (75.7) | 417 (13.9) | 141 (4.7) | <0.001** |
| <i>Employed</i> | n = 5282 | n = 269 | n = 3589 | n = 1000 | n = 424 | |
| Yes | 3862 (73.1) | 189 (4.9) | 2758 (71.4) | 665 (17.2) | 250 (6.5) | <0.001** |
| No | 1420 (26.9) | 80 (5.6) | 831 (58.5) | 335 (23.6) | 174 (12.3) | |
| <i>Previous births</i> | n = 5447 | n = 278 | n = 3679 | n = 1032 | n = 458 | |
| Yes | 1766 (32.4) | 78 (4.4) | 1079 (61.1) | 398 (22.5) | 211 (11.9) | <0.001** |
| No | 3681 (67.6) | 200 (5.4) | 2600 (70.6) | 634 (17.2) | 247 (6.7) | |

n = number; SD = standard deviation.

* = ANOVA's test.

** = Chi-Square's test.

Table 2 Results of multinomial logistic regression between the socio-demographic and obstetric characteristics and pre-pregnancy BMI.

| | Underweight OR (95% CI) | Overweight OR (95% CI) | Obesity OR (95% CI) |
|--------------------------|----------------------------|---------------------------|------------------------|
| <i>Age</i> | 0.95 (0.92–0.97) | 1.02 (1.00–1.03) | 1.01 (0.99–1.03) |
| <i>Country of origin</i> | | | |
| Spanish | Reference | Reference | Reference |
| Foreign | 0.91 (0.69–1.21) | 1.23 (1.05–1.44) | 0.84 (0.67–1.05) |
| <i>Educational level</i> | | | |
| Primary | Reference | Reference | Reference |
| Secondary | 0.90 (0.57–1.44) | 0.75 (0.61–0.94) | 0.73 (0.55–0.97) |
| Higher | 1.25 (0.79–1.96) | 0.37 (0.29–0.45) | 0.26 (0.19–0.36) |
| <i>Employed</i> | | | |
| Yes | 0.78 (0.57–1.07) | 0.81 (0.68–0.96) | 0.63 (0.49–0.79) |
| No | Reference | Reference | Reference |
| <i>Previous births</i> | | | |
| Yes | 1.06 (0.80–1.41) | 1.36 (1.16–1.58) | 1.86 (1.50–2.31) |
| No | Reference | Reference | Reference |

OR = Odds Ratio adjusted; 95% CI: 95% confidence interval.

women with a primary education. Overweight or obese were also less frequent among employed women. Finally, women who had had previous births more frequently started pregnancy with pre-pregnancy overweight or obesity than nulliparous women (Table 2).

Regarding the relationship between BMI with maternal complications, gestational diabetes and preeclampsia during pregnancy were more frequent among women with pre-pregnancy overweight and obesity (Table 3).

Regarding labour outcomes, it was observed that women with pre-pregnancy overweight and obesity started labour, more frequently, by induction or by elective caesarean section (Table 4).

Concerning the perinatal outcomes, the prevalence of premature new-borns was higher in women with pre-pregnancy overweight and obesity. Concerning neonate weight, the new-borns of women with pre-pregnancy overweight more frequently presented macrosomia, while in

Table 3 Relationship between the pre-pregnancy weight status of pregnant women and the prevalence of gestational diabetes and preeclampsia.

| | | Gestational diabetes | | p value | OR (95% CI) |
|---------------|-------------|----------------------|-------------|---------|------------------|
| | | Yes n (%) | No n (%) | | |
| Total | 5447 | 501 (9.2) | 4946 (90.8) | | |
| Underweight | 278 (5.1) | 15 (5.4) | 263 (94.6) | | 0.75 (0.44–1.28) |
| Normal weight | 3679 (67.5) | 261 (7.1) | 3418 (92.9) | | Reference |
| Overweight | 1032 (18.9) | 132 (12.8) | 900 (87.2) | | 1.92 (1.54–2.40) |
| Obesity | 458 (8.4) | 93 (20.3) | 365 (79.7) | <0.001* | 3.34 (2.7–4.33) |
| | | Preeclampsia | | p value | OR (95% CI) |
| | | Yes | No | | |
| Total | 5447 | 264 (4.8) | 5183 (95.2) | | |
| Underweight | 278 (5.1) | 11 (4) | 267 (96) | | 1.13 (0.60–2.11) |
| Normal weight | 3679 (67.5) | 130 (3.5) | 3549 (96.5) | | Reference |
| Overweight | 1032 (18.9) | 73 (7.1) | 959 (92.9) | | 2.08 (1.55–2.79) |
| Obesity | 458 (8.4) | 50 (10.9) | 408 (89.1) | <0.001* | 3.35 (2.38–4.71) |

n = number.

* = Chi-Square's test; OR = Odds Ratio; 95% CI: 95% confidence interval.

Table 4 Relationship between pre-pregnancy weight status of pregnant women and labour outcomes.

| | Total n (%) | Type of onset of labour | | | | |
|---------------|----------------|------------------------------------|--------------|------------------|--------------------|------------------|
| | | Spontaneous n (%) | Induction | | Elective caesarean | |
| | | | n (%) | OR (95% CI) | n (%) | OR (95% CI) |
| Total | 5447 | 3379 (62) | 1612 (29.6) | | 456 (8.4) | |
| Underweight | 278 (5.1) | 191 (68.7) | 67 (24.1) | 0.79 (0.60–1.06) | 20 (7.2) | 0.96 (0.60–1.55) |
| Normal weight | 3679 (67.5) | 2372 (64.5) | 1049 (28.5) | Reference | 258 (7) | Reference |
| Overweight | 1032 (18.9) | 612 (59.3) | 321 (31.1) | 1.19 (1.02–1.38) | 99 (9.6) | 1.49 (1.16–1.91) |
| Obesity | 458 (8.4) | 204 (44.5) | 175 (38.2) | 1.94 (1.57–2.10) | 79 (17.2) | 3.56 (2.67–4.76) |
| | n (%) | Type of delivery | | | | |
| | | Spontaneous vaginal birth n (%) | Instrumental | | Caesarean section | |
| | | | n (%) | OR (95% CI) | n (%) | OR (95% CI) |
| Total | 5447 | 3610 (66.3) | 451 (8.3) | | 1386 (25.4) | |
| Underweight | 278 (5.1) | 203 (73) | 25 (9) | 0.99 (0.64–1.52) | 50 (18) | 0.75 (0.54–1.03) |
| Normal weight | 3679 (67.5) | 2530 (68.8) | 316 (8.6) | Reference | 833 (22.6) | Reference |
| Overweight | 1032 (18.9) | 649 (62.9) | 81 (7.8) | 1 (0.77–1.29) | 302 (29.3) | 1.41 (1.21–1.65) |
| Obesity | 458 (8.4) | 228 (49.8) | 29 (6.3) | 1.02 (0.68–1.52) | 201 (43.9) | 2.68 (2.18–3.29) |

n = number; OR = Odds Ratio; 95% CI: confidence interval.

women with pre-pregnancy obesity neonates were more frequently underweight or presented macrosomia (Table 5).

Discussion

This exploratory study shows that the prevalence of women with pre-pregnancy overweight was 18.9% and with pre-pregnancy obesity was 8.4%, that is, that 27.3% of the women in this study had an inadequate BMI to start the pregnancy.

In concordance with previous studies, was observed that foreign women, unemployed women, or with lower educational level had more odds to present pre-pregnancy overweight and obesity. Moreover, perinatal complications were more frequent in women with a BMI > 25 kg/m².¹¹

Our prevalence of pre-pregnancy overweight of 18.9% and 8.4% of pre-pregnancy obesity were lower than the prevalence reported in other Spanish studies.^{12–14} Likewise, our prevalence of pre-pregnancy obesity was 8.4%, which was lower than the 9.6% of 301 women studied in Aragón,¹² the

Table 5 Relationship between pre-pregnancy weight status of pregnant women and perinatal outcomes.

| | n (%) | Term n (%) | Premature | | Post-term | |
|---------------|-------------|---------------|-----------|------------------|-----------|------------------|
| | | | n (%) | OR (95% CI) | n (%) | OR (95% CI) |
| Total | 5447 | 4890 (89.8) | 419 (7.7) | | 138 (2.5) | |
| Underweight | 278 (5.1) | 250 (89.9) | 17 (6.1) | 0.88 (0.53–1.47) | 11 (4) | 1.45 (0.77–2.73) |
| Normal weight | 3679 (67.5) | 3322 (90.3) | 256 (7) | Reference | 101 (2.7) | Reference |
| Overweight | 1032 (18.9) | 920 (89.1) | 91 (8.8) | 1.28 (1–1.65) | 21 (2) | 0.75 (0.47–1.21) |
| Obesity | 458 (8.4) | 398 (86.9) | 55 (12) | 1.79 (1.32–2.44) | 5 (1.1) | 0.41 (0.17–1.02) |

| | n (%) | Neonatal weight | | | | | |
|---------------|-------------|-----------------|-----------|------------------|-----------|------------------|--|
| | | Normal weight | | Underweight | | Macrosoma | |
| | | n (%) | n (%) | OR (95% CI) | n (%) | OR (95% CI) | |
| Total | 5447 | 4675 (85.8) | 477 (8.8) | | 295 (5.4) | | |
| Underweight | 278 (5.1) | 251 (90.3) | 21 (7.6) | 0.87 (0.55–1.38) | 6 (2.2) | 0.46 (0.20–1.05) | |
| Normal weight | 3679 (67.5) | 3204 (87.1) | 308 (8.4) | Reference | 167 (4.5) | Reference | |
| Overweight | 1032 (18.9) | 860 (83.3) | 88 (8.5) | 1.06 (0.83–1.36) | 84 (8.1) | 1.87 (1.43–2.46) | |
| Obesity | 458 (8.4) | 360 (78.6) | 60 (13.1) | 1.73 (1.29–2.33) | 38 (8.3) | 2.03 (1.40–2.93) | |

n = number; OR = Odds Ratio; 95% CI: 95% confidence interval.

17.1% of 6558 women in Canary Islands¹³ and the 13.3% of 16 609 women in Basque Country.¹⁴

Regarding the prevalence of overweight in other European countries, the value in the present study was like that reported in Croatia (19%).³ However, these values are much lower than those reported in Northern Ireland (29.8%).³ Prevalence of 8.4% of pre-pregnancy obesity in our study was higher than in Croatia (7.8%) but lower than the United Kingdom (20.4–25.6%).³

Like other studies, socio-demographic and obstetric factors were related with the pre-pregnancy BMI of women.^{5,15} In Spain, the BMI of adult Spanish women generally increases with age.^{12,13} In this study, foreign women presented more frequently pre-pregnancy overweight compared to Spanish women.¹⁶ In addition, women with a lower education level, or who were unemployed or who had had previous births were more frequently overweight and obese pre-pregnancy.^{5,15}

Regarding data on pregnancy complications, gestational diabetes and preeclampsia were more frequent among women with pre-pregnancy overweight or obesity as in the exploratory study of Yang¹⁷ and in the cohort study of Schummers.¹⁸

Regarding labour outcomes, it was observed that the probability of induction increased with an increase in pre-pregnancy BMI, reaching an OR of 1.94 in women with pre-pregnancy obesity, like what has been described in previous studies.^{19,20}

In this study, the percentage of women with pre-pregnancy obesity who underwent caesarean section (43.9%) was higher than that in a retrospective cohort study conducted in Canary Islands (18.7%),¹³ in a historical cohort study in Basque Country (25.7%)¹⁴ and with a descriptive study conducted in Castilla-La Mancha (34.9%).²¹

In line with the results of a meta-analysis by Dai et al., in 2018, maternal pre-pregnancy overweight and obesity in our study increased the probability of macrosomia in the

newborn²²; and the macrosomia increases the intrapartum complications and neonatal sequelae.²³ As in the study by Kim et al., in 2017, prematurity being more frequent in women with pre-pregnancy overweight and obesity.²⁴ The high rate of prematurity observed in our women could have increased the rates of underweight in our newborns, particularly in obese women who increased the probability of having an underweight neonate with an OR of 2.

Therefore, the identification of women with excess weight prior to conception should be implemented, and care should be adapted to the individual characteristics of women to improve maternal and perinatal outcomes. In addition, sensitive and proactive care must be provided, recognising and reporting possible risks to women, with counselling about diet and physical activity to facilitate an adequate weight before and during pregnancy seeking well-being in women.²⁵ Thus, midwives and obstetricians should implement actions promoting health,²⁶ these include diet and physical activity to achieve optimal health conditions during pregnancy.^{2,27}

At present, there is not much data available about pre-pregnancy overweight and obesity and its relation to maternal outcomes in Spain. The study has been carried out in a hospital that attends high obstetric complexity in Barcelona, a city that has a diverse cultural population.

To our knowledge, this is the largest study to describe prevalence in a Mediterranean city in Spain providing pre-pregnancy weight status of 5447 women and allowing comparison with other studies at an international level.

However, this study did not get data of 416 women (12.7% of the sample) in whom the pre-pregnancy BMI was not available in the electronic clinical records. In addition, although gestational weight gain is an independent factor that can influence maternal and perinatal outcomes, we were not able to provide this gain, because this variable was not always available, in the electronic clinical records, so caused heterogeneity of the hospital registry at the end

of pregnancy.²⁸ Therefore, we suggest improving the quality of our hospital records, and obstetricians and midwives should report data on pregnant women's weight gain.²⁹

Since this was a descriptive study, a relationship was observed between socio-demographic and obstetric factors and the presentation of a high BMI, so the causal inference is limited. Indeed, studies on the multifactorial origin of obesity involve a prospective cohort design that allows controlling for other variables that may influence the results such as lifestyle, culture, or religion.

This results provide knowledge about the prevalence of women with pre-pregnancy overweight and obesity in a high maternity complexity hospital of Barcelona (Spain), and that this prevalence was lower than majority of countries on Europe.

This study shows the relationship between some socio-demographic factors with begin the pregnancy with an inadequate BMI, and the relationship between pre-pregnancy overweight or obesity with adverse maternal and perinatal outcomes.

Funding

This study had been funded in partially by the "Nurse and Society Foundation" as part of the Nurse Research Projects Grants (PR-389/2019) Barcelona, Spain.

Elena González-Plaza received a research grant from "La Pedrera Foundation" (Nurse intensification grant) Barcelona, Spain.

Acknowledgments

We are grateful to Mrs. Mariuxi Burgos for reading this manuscript.

References

1. Ma RCW, Schmidt MI, Tam WH, McIntyre HD, Catalano PM. Clinical management of pregnancy in the obese mother: before conception, during pregnancy, and post partum. *Lancet Diabetes Endocrinol.* 2016;4:1037–49, [http://dx.doi.org/10.1016/S2213-8587\(16\)30278-9](http://dx.doi.org/10.1016/S2213-8587(16)30278-9).
2. Vitner D, Harris K, Maxwell C, Farine D. Obesity in pregnancy: a comparison of four national guidelines. *J Matern Neonatal Med.* 2018;7058:1–11, <http://dx.doi.org/10.1080/14767058.2018.1440546>.
3. Euro-Peristat Project. Core indicators of the health and care of pregnant women and babies in Europe in 2015. www.europeristat.com. Updated November 2018 [accessed October 2019].
4. Devlieger R, Benhalima K, Damm P, Van Assche, Mathieu C, Mahmood T, et al. Maternal obesity in Europe: where do we stand and how to move forward? A scientific paper commissioned by European Board and College of Obstetrics and Gynaecology (EBCOG). *Eur J Obstet Gynecol Reprod Biol.* 2016;201:203–8, <http://dx.doi.org/10.1016/j.ejogrb.2016.04.005>.
5. Bogaerts A, Van den Bergh B, Nuyts E, Martens E, Witters I, Devlieger R. Socio-demographic and obstetrical correlates of pre-pregnancy body mass index and gestational weight gain. *Clin Obes.* 2012;2:150–9, <http://dx.doi.org/10.1111/cob.12004>.

6. Headen I, Cohen AK, Mujahid M, Abrams B. The accuracy of self-reported pregnancy-related weight: a systematic review. *Obes Rev.* 2017;18:350–69, <http://dx.doi.org/10.1111/obr.12486>.
7. World Health Organization. Obesity and overweight. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>. Updated February 16, 2018 [accessed October 2019].
8. International Association of Diabetes and Pregnancy Study Groups Consensus Panel, Metzger BE, Gabbe SG, et al. International association of diabetes and pregnancy study groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. *Diabetes Care.* 2010;33:676–82, <http://dx.doi.org/10.2337/dc09-1848>.
9. Sibai B, Dekker G, Kupfermanc M. Pre-eclampsia. *Lancet.* 2005;365:785–99, [http://dx.doi.org/10.1016/S0140-6736\(05\)17987-2](http://dx.doi.org/10.1016/S0140-6736(05)17987-2).
10. Jiménez R, Figueras J. Prematuridad. In: Cruz M, editor. *Tratado de Pediatría.* 10a ed. Madrid: Ergón; 2011., ISBN 978-84-8473-904-3 p. 69–80.
11. Zhao YN, Li Q, Li YCh. Effects of body mass index and fat percentage on gestational complications and outcomes. *J Obstet Gynaecol Res.* 2014;40:705–10, <http://dx.doi.org/10.1111/jog.12240>.
12. Ramón-Arhués E, Martínez B, Martín S. Gestational weight gain and postpartum weight retention in a cohort of women in Aragon Spain. *Nutr Hosp.* 2017;34:4–8, <http://dx.doi.org/10.20960/nh.749>.
13. Bautista-Castaño I, Henriquez-Sanchez P, Alemán-Perez N, García-Salvador JJ, Gonzalez-Quesada A, García-Hernández JA, et al. Maternal obesity in early pregnancy and risk of adverse outcomes. *PLoS One.* 2013;8:1–6, <http://dx.doi.org/10.1371/journal.pone.0080410>.
14. Melchor I, Burgos J, Del Campo A, Airtzaguen A, Gutiérrez J, Melchor JC. Effect of maternal obesity on pregnancy outcomes in women delivering singleton babies: a historical cohort study. *J Perinat Med.* 2019;47:625–30, <http://dx.doi.org/10.1515/jpm-2019-0103>.
15. Sutherland G, Brown S, Yelland J. Applying a social disparities lens to obesity in pregnancy to inform efforts to intervene. *Midwifery.* 2013;29:338–43, <http://dx.doi.org/10.1016/j.midw.2012.01.016>.
16. Gaillard R, Durmuş B, Hofman A, MacKenbach JP, Steegers EAP, Jaddoe VWV. Risk factors and outcomes of maternal obesity and excessive weight gain during pregnancy. *Obesity.* 2013;21:1046–55, <http://dx.doi.org/10.1002/oby.20088>.
17. Yang W, Han F, Gao X, Chen Y, Ji L, Cai X. Relationship between gestational weight gain and pregnancy complications or delivery outcome. *Sci Rep.* 2017;7:12531, <http://dx.doi.org/10.1038/s41598-017-12921-3>.
18. Schummers L, Hutcheon JA, Bodnar LM, Lieberman E, Himes KP. Risk of adverse pregnancy outcomes by prepregnancy body mass index: a population-based study to inform prepregnancy weight loss counseling. *Obstet Gynecol.* 2015;125:133–43, <http://dx.doi.org/10.1097/AOG.0000000000000591>.
19. Carlson NS, Lowe NK. Intrapartum management associated with obesity in nulliparous women. *J Midwifery Women's Health.* 2014;59:43–53, <http://dx.doi.org/10.1111/jmwh.12073>.
20. Dammer U, Bogner R, Weiss C, Faschingbauer F, Pretschner J, Beckmann MW, et al. Influence of body mass index on induction of labor: a historical cohort study. *J Obstet Gynaecol Res.* 2018;44:697–707, <http://dx.doi.org/10.1111/jog.13561>.
21. Ballesta-Castillejos A, Gómez-Salgado J, Rodríguez-Almagro J, Ortiz-Esquinas, Hernández-Martínez A. Relationship between maternal body mass index and obstetric and perinatal complications. *J Clin Med.* 2020;9:707, <http://dx.doi.org/10.3390/jcm9030707>.
22. Dai RX, He XJ, Hu CL. Maternal pre-pregnancy obesity and the risk of macrosomia: a meta-analysis. *Arch*

- Gynecol Obstet. 2018;297:139–45, <http://dx.doi.org/10.1007/s00404-017-4573-8>.
23. Araujo Júnior E, Peixoto AB, Zamarian AC, Elito Júnior J, Tonni G. Macrosomia. *Best Pract Res Clin Obstet Gynaecol.* 2017;38:83–96, <http://dx.doi.org/10.1016/j.bpobgyn.2016.08.003>.
 24. Kim T, Burn SC, Bangdiwala A, Pace S, Rauk P. Neonatal morbidity and maternal complication rates in women with a delivery body mass index of 60 or higher. *Obstet Gynecol.* 2017;130:988–93, <http://dx.doi.org/10.1097/AOG.0000000000002316>.
 25. Goldstein RF, Abell SK, Ranasinha S, Misso M, Boyle JA, Black MH, et al. Association of gestational weight gain with maternal and infant outcomes a systematic review and meta-analysis. *JAMA.* 2017;317:2207–25, <http://dx.doi.org/10.1001/jama.2017.3635>.
 26. Institute of Medicine and National Research Council. In: Rasmussen KM, Yaktine AL, editors. *Weight gain during pregnancy: reexamining the guidelines.* The National Academies Press: Washington (DC); 2009.
 27. Jones C, Jomeen J. Women with a BMI \geq 30 kg/m² and their experience of maternity care: a meta ethnographic synthesis. *Midwifery.* 2017;53:87–95, <http://dx.doi.org/10.1016/j.midw.2017.07.011>.
 28. International Weight Management in Pregnancy (I-WIP) Collaborative group. Effect of diet and physical activity based interventions in pregnancy on gestational weight gain and pregnancy outcomes: meta-analysis of individual participant data from randomised trials. *BMJ.* 2017;358:j3119, <http://dx.doi.org/10.1136/bmj.j3119>.
 29. Harrison CL, Skouteris H, Boyle J, Teede HJ. Preventing obesity across the preconception, pregnancy and postpartum cycle: Implementing research into practice. *Midwifery.* 2017;52:64–70, <http://dx.doi.org/10.1016/j.midw.2017.06.003>.