

1 **Emotional eating and cognitive restraint mediate the association between**
2 **sleep quality and BMI in young adults**

3
4 María Fernanda Zerón-Rugiero^{a,b}, Álvaro Hernández^{c,d,e}, Trinitat Cambras^f,
5 Maria Izquierdo-Pulido^{a,b#}.
6

7 ^aDepartment of Nutrition, Food Science and Gastronomy, Campus de
8 l'Alimentació Torribera, University of Barcelona, Av. Prat de la Riba, 171, 08921
9 Santa Coloma de Gramenet, Barcelona, Spain. ^bINSA-UB, Nutrition and Food
10 Safety Research Institute, University of Barcelona, Av. Prat de la Riba, 171,
11 08921 Santa Coloma de Gramenet, Barcelona, Spain. ^cCenter for Fertility and
12 Health, Norwegian Institute of Public Health, Marcus Thranes Gate 2, 0473, Oslo,
13 Norway. ^dCIBER Physiopathology of Obesity and Nutrition, Instituto de Salud
14 Carlos III, Madrid, Spain. ^eBlanquerna School of Health Sciences, Universitat
15 Ramon Llull, Carrer de Padilla, 326, 08025, Barcelona, Spain. ^fDepartment of
16 Biochemistry and Physiology, School of Pharmacy and Food Science, University
17 of Barcelona, Av. de Joan XXIII, 27-31, 08028 Barcelona, Spain. [#] Corresponding
18 author.
19

20 Contact details:

21 Dr. Maria Izquierdo-Pulido. Torribera Campus, Av. Prat de la Riba, 171 – Edifici
22 Verdaguer, 08921, Santa Coloma de Gramenet, Barcelona, Spain. E-mail:
23 maria_izquierdo@ub.edu. Phone number: (+34) 934037293.
24

25 **Abstract**

26 This cross-sectional study was designed to investigate whether diet quality and
27 eating behaviors could mediate the association between sleep quality and body
28 mass index (BMI) in young adults. For all participants (n=925; aged 21.4±2.5
29 77.8% women) we evaluated: BMI, sleep quality, diet quality, and eating behavior
30 dimensions (emotional eating, cognitive restraint, and uncontrolled eating).
31 Linear regression models were used to test associations between exposure and
32 outcome variables. Path analysis was conducted with all potential mediators and
33 covariates entered at the same time. Results showed that emotional eating (β =
34 0.04 [95% CI: 0.03;0.06]), cognitive restraint (β = 0.03 [95% CI: 0.01;0.04]),
35 uncontrolled eating (β = 0.02 [95% CI: 0.01;0.04]) and diet quality (β = -0.14 [95%
36 CI: -0.19;-0.08]) were significantly associated with sleep quality. Additionally, BMI
37 was significantly associated with PSQI score (β = 0.09 [95% CI: 0.01;0.17]),
38 emotional eating (β = 0.89 [95% CI: 0.60;1.18]), and cognitive restraint (β = 1.37
39 [95% CI: 1.02;1.71]). After testing for mediation, results revealed that emotional
40 eating and cognitive restraint evidenced a significant mediating effect on the
41 association between sleep quality and BMI. Additionally, diet quality was
42 significantly associated with emotional eating (β = -0.35 [95% CI: -0.56;-0.13]),
43 cognitive restraint (β = 0.53 [95% CI: 0.27;0.79]), and uncontrolled eating (β = -
44 0.49 [95% CI: -0.74;-0.25]). In conclusion, young adults with poor sleep quality
45 are more likely to deal with negative emotions with food, which, in turn, could be
46 associated with higher cognitive restraint, becoming a vicious cycle that has a
47 negative impact on body weight. Our results also emphasize the role of eating
48 behaviors as determinants of diet quality, highlighting the importance of
49 considering sleep quality and eating behaviors when designing obesity
50 prevention strategies in this population.

51

52 **Keywords:** sleep quality, body mass index, emotional eating, cognitive restraint,
53 eating behavior, diet quality.

54 **1. Introduction**

55 College is a period of transition in which many young adults learn to make
56 independent decisions that, among other things, include choosing *what* and *when*
57 to eat, whether or not to exercise, and how much to sleep (Richards & Specker,
58 2021). Unfortunately, this period usually coincides with academic and social
59 demands, as well as life situations that can disrupt their daily habits (Dinis &
60 Bragança, 2018; Wickham, Amarasekara, Bartonicek, & Conner, 2020). In fact,
61 research has shown that young adults who attend college have late sleep
62 schedules and irregular sleep/wake patterns (Izidorczyk, Sitnik-Warchulska,
63 Lizińczyk, & Lipiarz, 2019; Richards & Specker, 2021; Silva et al., 2016; Zerón-
64 Rugerio, Cambras, & Izquierdo-Pulido, 2019), which in turn lead to poor sleep
65 quality (Pilz, Keller, Lenssen, & Roenneberg, 2018). Noteworthy, poor sleep
66 quality affects around 50 – 60% of college students (Chen & Chen, 2019; Dinis &
67 Bragança, 2018; Memon et al., 2021) and has been highlighted as a significant
68 predictor of poor mental health and low well-being (Wickham et al., 2020), obesity
69 (Krističević, Štefan, & Sporiš, 2018), and unhealthy dietary habits (Yamamoto et
70 al., 2018).

71

72 Poor sleep quality can influence 'what' and 'how much' people eat through
73 several mechanisms including alterations in hormones related to appetite (mainly
74 ghrelin and leptin) and increased neuronal activity in response to food intake
75 (Burrows, Fenton, & Duncan, 2020; M.-P. St-Onge, 2017). Not surprisingly,
76 evidence has shown that when people have not slept well, they tend to eat more
77 calories the next day (~350 kcal more) (Al Khatib, Harding, Darzi, & Pot, 2017;
78 M.-P. St-Onge, 2017). In addition, these hormonal and neuronal alterations are

79 accompanied by a greater preference for high-calorie foods, which is mainly
80 driven by hedonic mechanisms (Al Khatib et al., 2017; Goldstein & Walker, 2014).
81 Consequently, poor sleep quality has been highlighted as a modifiable risk factor
82 for obesity (Chaput & Dutil, 2016; Krističević et al., 2018; M.-P. St-Onge, 2017).
83 In fact, Kristicevic et al (Krističević et al., 2018) reported that young adults with
84 poor sleep quality have ~1.40 more odds of being overweight or obese.

85

86 Along these lines, emerging evidence has revealed that the link between poor
87 sleep quality and obesity could also be driven by eating behaviors (Blumfield, Bei,
88 Zimberg, & Cain, 2018; Lundahl & Nelson, 2015; Saleh-ghadimi, Dehghan,
89 Farhangi, & Asghari-jafarabadi, 2019). Blumfield et al (Blumfield et al., 2018)
90 demonstrated that disinhibited eating behavior, understood as the need to
91 overeat in the presence of palatable foods or other disinhibiting stimuli like
92 emotional stress, mediated the association between poor sleep quality and
93 obesity among adults. The authors postulated that in the context of sleep
94 impairment, an increased drive for exciting rewards in combination with
95 disinhibited eating behavior may exacerbate the food seeking behavior,
96 especially for palatable foods (Blumfield et al., 2018), which are those that are
97 known to increase the feelings of calmness and satisfaction (Gibson, 2006). What
98 is equally interesting is that lack of sleep could also act as a stressor and increase
99 food consumption in people prone to emotional eating (Dweck, Jenkins, & Nolan,
100 2014).

101

102 Furthermore, a recent study noted that greater dietary restraint is also associated
103 with poor sleep quality (given by higher sleep fragmentation) (Barragán et al.,

104 2021). However, the potential role of dietary restraint as a mediator of the
105 association between sleep quality and obesity remains scarcely studied.
106 Noteworthy, greater dietary restraint has been associated with overweight and
107 obesity among adolescents and young adults (Racine, 2018; Ramírez-Contreras,
108 Farrán-Codina, Izquierdo-Pulido, & Zerón-Rugerio, 2021). Although this fact
109 might seem paradoxical, it has been suggested that greater dietary restraint is
110 associated with increased bottom-up reward reactivity to food stimuli, which may
111 explain why people who attempt to diet are vulnerable to binge eating and weight
112 gain (Bryant, Rehman, Pepper, & Walters, 2019; Martin-Garcia et al., 2016;
113 Racine, 2018). It should be noted that young adulthood is a period of increased
114 risk for excessive weight gain (Deliens, Deforche, Chapelle, & Clarys, 2019;
115 Fedewa, Das, Evans, & Dishman, 2014; Nelson, Story, Larson, Neumark-
116 Sztainer, & Lytle, 2008), at the same time that it is a crucial stage of life for
117 establishing long-term health behaviors (Fedewa et al., 2014; Nelson et al., 2008;
118 Wickham et al., 2020).

119
120 Taking into account the aforementioned, it is relevant to study the role that dietary
121 intake and eating behaviors can have in the association between sleep quality
122 and obesity among young adults. Our objective was to investigate whether diet
123 quality and eating behaviors could mediate the association between sleep quality
124 and BMI. We hypothesized that poor diet quality and eating behaviors that
125 promote an overconsumption of energy will mediate the association between
126 poor sleep quality and higher BMI. In addition, we aimed to study which were the
127 dietary habits and eating behavior traits that were significantly associated with
128 sleep quality.

129 **2. Methodology**

130 **2.1 Study design, settings, participants and protocol**

131 Young adults (aged 18–30 years) were recruited among undergraduate and
132 postgraduate students at the University of Barcelona (Spain). Participants were
133 invited to take part in this cross-sectional study during the school year, between
134 2017 and 2019. Exclusion criteria consisted of inability to provide the information
135 required for the development of the study or having previously been diagnosed
136 with chronic diseases such as type 2 diabetes, hypertension and/or
137 cardiovascular disease. Based on these criteria, a total of 938 young adults were
138 eligible and provided written informed consent before joining this study. We
139 further excluded participants who were out of the age range, resulting in a final
140 analytical cohort of 925 participants. All study procedures were conducted
141 according to the Declaration of Helsinki and were approved by the Ethics
142 Committee the University of Barcelona (IRB00003099).

143

144 **2.2 Outcome variables**

145 *2.2.1 Anthropometric parameters*

146 Body mass index was estimated through self-reported height and weight.
147 Participants were asked in a questionnaire ‘What is your current weight? (kg)’
148 and ‘What is your current height? (cm)’, and subsequently BMI was calculated as
149 weight (kg) divided by squared height (m). Note that self-reported BMI had a very
150 high agreement with BMI values among a similar population (Zerón-Rugério et
151 al., 2019).

152

153

154 *2.2.2 Diet quality*

155 Diet quality was evaluated through the Mediterranean Diet Quality Index
156 (KIDMED) (Serra-Majem et al., 2004). This test is based on the principles that
157 sustain Mediterranean dietary patterns and those that undermine them. This
158 questionnaire consists of 16 items which are answered as “Yes” or “No”
159 questions. Subsequently, items denoting a lower adherence are assigned a value
160 of –1, while those related to higher adherence are scored +1. The total score
161 ranges from 0 to 12, where higher scores indicate greater adherence to the
162 Mediterranean Diet. In addition, according to the score, adherence to the
163 Mediterranean diet was classified as “poor” (≤ 3 points), “average” (4-7 points), or
164 “good” (≥ 8 points).

165

166 *2.2.3 Eating behavior*

167 The validated Spanish version of the Three Factor Eating Questionnaire (TFEQ-
168 R21) (Martin-Garcia et al., 2016) was used to assess the following dimensions of
169 eating behavior:

170

- 171 i. Emotional eating: understood as the need to overeat when people are unable
172 to cope with emotionally negative situations. This dimension was evaluated
173 through six items.
- 174 ii. Cognitive restraint: understood as the conscious effort of the individual to
175 control what he/she eats to maintain or lose weight. This dimension was
176 evaluated through six items.

177 iii. Uncontrolled eating: understood as the tendency to overeat in response to the
178 loss of control over food intake. This dimension was evaluated through nine
179 items.

180

181 The TFEQ-R21 consists of 21 items which are rated on a four-point Likert scale
182 ranging from 1 (“Definitely true”) to 4 (“Definitely false”). Scores are calculated
183 separately for each dimension as a mean of all items, where the higher the score
184 the greater the emotional eating, the cognitive restraint, and/or the uncontrolled
185 eating (Cappelleri et al., 2009; Martin-Garcia et al., 2016). Note that the main
186 strength of the TFEQ is the existence of vast evidence suggesting an important
187 and robust role for the TFEQ traits in obesity, eating styles, eating disorders and
188 associated factors (Bryant et al., 2019).

189

190 **2.3 Exposure variable**

191 Sleep quality was assessed with the Spanish validated version of the Pittsburgh
192 Sleep Quality Index (PSQI) (Royuela & Macías, 1997). It consists of 19 items,
193 which are grouped to into seven components of sleep quality: subjective sleep
194 quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbance,
195 use of sleeping medications, and daytime dysfunction. Total PSQI scores were
196 calculated as the sum of the seven components. Scores range from 0 to 21,
197 where higher scores indicate poor sleep quality. Additionally, according to the
198 score, sleep quality was classified as “good” (PSQI<5 points) and “poor” (PSQI
199 ≥ 5 points).

200

201

202 **2.4 Covariates**

203 The level of physical activity was measured in Metabolic Equivalents of Task
204 (METs) using the short version of the International Physical Activity
205 Questionnaire (IPAQ), which is validated for Spanish population (Román-Viñas,
206 Ribas-Barba, Ngoa, & Serra-Majem, 2013). In this case, the higher the score, the
207 more intense the level of physical activity. Additionally, the chronotype was
208 evaluated through the midpoint of sleep on free-days (MSF) (Roenneberg, Wirz-
209 Justice, & Mellow, 2003).

210

211 **2.5 Statistical analyses**

212 Normality was confirmed in all variables by histograms and Q-Q plots. Variables
213 are described by means and standard deviations for continuous variables and
214 proportions for categorical data. First, we tested the association between sleep
215 quality and BMI with the outcome variables (diet quality, emotional eating,
216 cognitive restraint and uncontrolled eating) using linear regression analyses.
217 Then, we tested whether those variables (emotional eating and cognitive
218 restraint) that were significantly associated with the BMI were significant
219 mediators of the association between sleep quality and BMI (**Figure 1**). All
220 potential mediators were entered at the same time, using the PROCESS
221 macro (Hayes, 2018) version 3.3 for SPSS.

222

223 In addition, we tested the association of the 16 dietary habits evaluated in the
224 KIDMED questionnaire with sleep quality (PSQI score) and the scores of the
225 eating behavior dimensions (emotional eating, cognitive restraint, and
226 uncontrolled eating). To do this, we used general linear models (GLMs) to

227 calculate adjusted differences in PSQI, emotional eating, cognitive restraint, and
228 uncontrolled eating scores based on KIDMED items (reference category “yes”).
229 Finally, we corrected P-values using the Benjamini–Hochberg method, assuming
230 a False Discovery Rate (FDR) of 5%.

231

232 All analyses were adjusted for age, gender, BMI, diet quality, physical activity,
233 and chronotype (except when the covariate was the dependent variable of the
234 model). All analyses were carried out in R Software, version 3.4.1. (R Core Team,
235 2019), except for the mediation models which were performed with SPSS version
236 25 (IBM SPSS Statistics). A statistical significant test was considered when
237 $p < 0.05$.

238

239 **3. Results**

240 Nine hundred and twenty-five participants (aged 21.4 ± 2.5 years) were included
241 in this cross-sectional study, of whom 77.8% were women (**Table 1**). Overall,
242 59.2% of the participants showed a poor sleep quality. Regarding BMI, most of
243 the participants were classified as normal-weight (78.7%), while 12.0% were
244 overweight/obese and 9.1% were underweight. Furthermore, half of the
245 participants (53.5%) showed poor or average adherence to the Mediterranean
246 diet, while the other half (46.5%) had high adherence to this dietary pattern. **Table**
247 **1** also summarizes the average standardized scores for emotional eating,
248 cognitive restraint, and uncontrolled eating behaviors. Our results also showed
249 that the majority of the participants (57.3%) performed a moderate level of
250 physical activity, while 16.6% reported a low level of physical activity and 26.1%
251 a high level of physical activity.

252 As shown in **Table 2**, PSQI score was significantly associated with diet quality
253 and eating behaviors. Accordingly, a 1-point increment in the PSQI score was
254 significantly associated with less points in the Mediterranean Diet Quality index
255 (-0.14 [95% CI: -0.19; -0.08]). Conversely, a 1-point increment in the PSQI score
256 was associated with higher emotional eating (0.04 [95% CI: 0.03; 0.06]), cognitive
257 restraint (0.03 [95% CI: 0.01; 0.04]), and uncontrolled eating (0.02 [95% CI: 0.01;
258 0.04]) behaviors.

259

260 Regarding BMI, we observed that it was associated with cognitive restraint and
261 emotional eating (**Table 2**). In which case, the results revealed that a 1-point
262 increment in the emotional eating score was associated with 0.89 kg/m² [95% CI:
263 0.06; 1.18] more BMI, while a 1-point increment in the cognitive restraint score
264 was associated with 1.37 kg/m² [95% CI: 1.02; 1.17] more BMI. Furthermore, we
265 observed that BMI was significantly associated with PSQI score ($\beta = 0.09$ [95%
266 CI: 0.01; 0.17], $p = 0.021$) (**Figure 2**).

267

268 *3.1 Emotional eating and cognitive restraint mediate the association between*
269 *sleep quality and BMI*

270 The mediation model that highlights the effect of sleep quality on BMI via
271 emotional eating and cognitive restraint is shown in **Figure 2**. As observed, paths
272 a_1 and a_2 of the structural model were statistically significant ($\beta = 0.05$ [95% CI:
273 0.03; 0.06], $p < 0.001$ and $\beta = 0.03$ [95% CI: 0.02; 0.04], $p < 0.001$, respectively).
274 Consequently, poor sleep quality (higher PSQI score) was significantly
275 associated with both mediators (emotional eating and cognitive restraint).
276 Likewise, our results revealed that paths b_1 and b_2 of the structural model were

277 statistically significant. As such, greater emotional eating and greater cognitive
278 restraint were significantly associated with higher BMI ($\beta= 0.71$ [95% CI: 0.42;
279 1.00], $p<0.001$ and $\beta= 1.22$ [95% CI: 0.87; 1.57], $p<0.001$, respectively).

280

281 Regarding c'-path, we observed that it was not statistically significant (**Figure 2**).
282 Accordingly, our results revealed that there was a significant indirect association
283 between poor sleep quality and higher BMI via emotional eating and cognitive
284 restraint. The summary of indirect effects are shown in **Table 3**. Note that
285 approximately 14.8% of the variance in BMI was accounted by emotional eating
286 and cognitive restraint ($R= 0.385$; $R^2= 0.148$; $p<0.0001$). Additionally, when
287 examining unstandardized beta weights, we observed that cognitive restraint was
288 the strongest predictor of BMI ($\beta= 1.22$, $p<0.001$) (**Figure 2**).

289

290 *3.2 Dietary habits are significantly associated with sleep quality.*

291 Regarding the association between sleep and dietary habits, our results revealed
292 that higher PSQI score was associated with three of the sixteen habits that
293 characterize the Mediterranean diet (**Figure 3**). Specifically, we observed that
294 attending fast-food restaurants more than once a week was associated with
295 higher the PSQI score ($p=0.034$). In contrast, regular consumption of fish and
296 nuts was associated with lower PSQI score ($p=0.041$ and $p=0.045$, respectively).

297

298 *3.3 Dietary habits are associated with emotional eating, cognitive restraint, and*
299 *uncontrolled eating behaviors*

300 Our results also revealed a significant association between eating behavior
301 dimensions and diet quality. In particular, we showed that emotional eating ($\beta= -$

302 0.35 [95% CI: -0.56; -0.13], $p=0.002$) and uncontrolled eating ($\beta= -0.49$ [95% CI:
303 -0.74; -0.25], $p<0.001$) were negatively associated with KIDMED score. It should
304 be noted that consuming fast-food more than once a week ($p<0.001$) and/or
305 eating sweets or candy every day ($p<0.001$ and $p=0.043$) were associated with
306 higher emotional and uncontrolled eating scores (**Table 4**). In addition, we
307 showed that eating commercially baked goods or pastries for breakfast was
308 significantly associated with emotional eating ($p=0.034$).

309

310 On the other hand, we observed that cognitive restraint was positively associated
311 with KIDMED score ($\beta= 0.53$ [95% CI: 0.27; 0.79], $p<0.001$). In this regard, our
312 results showed that the consumption of a second serving of fruit ($p=0.016$), a
313 daily serving of fresh/cooked vegetables ($p<0.001$), and/or a second serving of
314 vegetables ($p<0.001$) was significantly associated with higher cognitive restraint.
315 Conversely, lower cognitive restraint was found among those who consumed
316 commercially baked goods or pastries for breakfast ($p<0.001$).

317

318 **4. Discussion**

319 The main contribution of our research work is that emotional eating and cognitive
320 restraint are significant mediators of the association between sleep quality and
321 BMI, among young adults. Blumfield et al (Blumfield et al., 2018) demonstrated
322 that disinhibited eating behavior mediated the association between poor sleep
323 quality and obesity. This is consistent with our finding on the role of emotional
324 eating as a mediator of the association between sleep quality and BMI. Our
325 results suggest that when sleep is neglected and participants are unable to cope
326 with emotionally negative situations (sadness, anger, anxiety, nervousness, etc.),

327 they turn to food for comfort. This is also in line with another study that showed
328 that greater emotional eating scores were associated with poor sleep quality
329 among young women (Dweck et al., 2014). Importantly, the authors also
330 demonstrated that short sleep duration *per se* could act as a stressor, increasing
331 food consumption in those participants who were prone to emotional eating
332 (Dweck et al., 2014). Note that eating sweet and fatty foods is known to alleviate
333 behavioral signs of distress (Gibson, 2006; Penaforte, Minelli, Rezende, & Japur,
334 2019). Interestingly, our results showed that the consumption of fast-food,
335 commercially baked goods or pastries, and sweets or candy is associated with a
336 higher emotional eating score. Therefore, it is plausible that participants with poor
337 sleep quality are using food as a coping mechanism (Dinis & Bragança, 2018).

338

339 Furthermore, our results revealed that cognitive restraint was the strongest
340 predictor of BMI. According to the literature, this is possible since chronic food
341 restriction could alternate with episodes of overeating, which could lead to weight
342 gain (Herman & Mack, 1975; Martin-Garcia et al., 2016; Polivy & Herman, 1985).
343 This weight gain could also lead to greater restriction, becoming a vicious circle
344 (Herman & Mack, 1975; Martin-Garcia et al., 2016; Polivy & Herman, 1985).
345 Therefore, it is plausible that poor sleep quality increases the risk for overeating
346 in response to emotional stress, leading young adults to make a conscious effort
347 to control food intake to manage their body weight. Interestingly, we observed
348 that young adults with poor sleep quality reported taking small portions of food to
349 control their weight, limited food intake to avoid gaining weight, and/or avoided
350 some foods because they made them fat.

351

352 In line with the aforementioned, we demonstrated that participants with the
353 highest cognitive restraint score were those who did not consume commercially
354 baked goods or pastries for breakfast. In parallel, healthy eating habits, such as
355 consuming two servings of fruits and vegetables per day were associated with
356 greater cognitive restraint. These findings are consistent with other studies which
357 demonstrated that perceptions about the healthiness of foods may be highly
358 relevant to food intake among participants with greater cognitive restraint
359 (Provencher, Polivy, & Herman, 2009; Racine, 2018). Note that perceptions
360 about the healthiness or “fatteningness” of foods may bias the estimates of the
361 caloric content of foods and, unsurprisingly, this could be counterproductive as a
362 weight management strategy (Provencher et al., 2009). In fact, experimental
363 studies have shown that young adults who usually choose low-calorie or “healthy”
364 foods for weight control are more likely to underestimate the caloric content of
365 food and end up consuming more calories (Provencher et al., 2009; Racine,
366 2018). Consequently, evidence has shown that young adults with greater dietary
367 restraint are more likely to be overweight (Racine, 2018; Ramírez-Contreras et
368 al., 2021). Likewise, in our study, a 1-point increment in the cognitive restraint
369 score was associated with ~ 1.22 kg/m² more BMI. Although, the role of cognitive
370 restraint in the relationship between sleep quality and BMI may differ based on
371 age. In this regard, Barragan et al (Barragán et al., 2021) showed that among
372 subjects aged 20–73 years the association between poor sleep quality and BMI
373 was moderated by cognitive restraint.

374

375 Among other relevant findings, we observed that greater uncontrolled eating was
376 significantly associated with poor sleep quality. It is worth noticing that the impact

377 of sleep loss extends to primary reward-motivated behaviors, including a desire
378 for appetizing food, a greater preference for higher caloric foods, and a higher
379 tendency to overeat (Goldstein & Walker, 2014; Greer, Goldstein, & Walker,
380 2013; M. P. St-Onge et al., 2012). This, according to a recent meta-analysis (Al
381 Khatib et al., 2017), implies that lack of sleep increases motivation to search for
382 food to obtain a reward. This could explain why, in our study, participants with
383 poor sleep quality felt that they could not stop eating once they had started and
384 also that it was difficult to stop eating before finishing all the food on their plate. It
385 is also worth noting that higher uncontrolled eating was associated with unhealthy
386 eating habits, such as eating sweets and candy daily, and consuming fast-food
387 more than once a week.

388
389 Our results also revealed that poor sleep quality was significantly associated with
390 lower adherence to the Mediterranean diet, which is consistent with other studies
391 (Ferranti et al., 2016; Godos et al., 2019). However, a novel finding of our study
392 is that, a regular consumption of nuts, important food staple in the Mediterranean
393 diet (at least 2-3 times/week) was associated with higher sleep quality. It should
394 be noted that nuts are high in folate, magnesium, fiber, and tryptophan (Burrows
395 et al., 2020; Mossavar-Rahmani et al., 2017). The latter is converted by the body
396 into serotonin and melatonin. Note that melatonin is essential for sleep regulation
397 (Burrows et al., 2020). Thus, it is plausible that the combination of several
398 nutrients present in nuts is related to a higher sleep quality (Burrows et al., 2020;
399 Marie-Pierre St-Onge, Mikic, & Pietrolungo, 2016; Yamamoto et al., 2018;
400 Zuraikat, Wood, Barragán, & St-Onge, 2021). Therefore, future intervention

401 studies should focus on studying the potential role of nuts as a sleep-promoting
402 foods.

403

404 Regarding other characteristics of the Mediterranean diet and its association with
405 sleep quality, we observed that regular fish consumption (at least 2-3 times/week)
406 was also associated with higher sleep quality, which is in line with other studies
407 (Del Brutto et al., 2016; Hansen et al., 2014). It is worth noting that consumption
408 of oily fish is an important dietary source of long-chain omega-3 polyunsaturated
409 fatty acids, including eicosapentaenoic and docosahexaenoic acids, which play
410 a role in serotonin secretion (Del Brutto et al., 2016). The latter is involved in sleep
411 regulation, especially since serotonin is converted in the body to melatonin, as
412 mentioned above (Burrows et al., 2020).

413

414 In contrast, our results showed that attending a fast-food restaurant more than
415 once a week was associated with poor sleep quality. Keep in mind that fast-food
416 restaurants offer energy-dense foods that are known to be high in fat and sugar,
417 but low in fiber. Therefore, this result would be in line with St-Onge et al (M. St-
418 Onge et al., 2011; Marie-Pierre St-Onge et al., 2016; Zuraikat et al., 2021)
419 findings, which revealed that a higher saturated fat intake and a lower
420 consumption of fiber were associated with more nighttime awakenings and
421 reduced overall sleep quality. However, we cannot ignore that the association
422 between diet and sleep quality is bidirectional (Burrows et al., 2020). Therefore,
423 it is also plausible that poor sleep quality can exacerbate the food-seeking
424 behavior, especially for palatable foods (Blumfield et al., 2018; Greer et al., 2013).

425

426 Our study has certain limitations, starting with the observational nature of the
427 study that prevents us from claiming causation and the use of self-reported
428 data weight and height. In addition, we acknowledge the use of self-reported data
429 to evaluate diet quality and eating behaviors as a limitation of the study. To help
430 avoid underreporting in future studies, we suggest the use of objective data,
431 including actigraphy to assess sleep quality, as well as the measurement of body
432 composition to assess nutritional status. However, our sample size is large
433 enough to provide sufficient strength for the associations of sleep quality with diet
434 quality, eating behaviors, and BMI.

435

436 In summary, our findings indicate that emotional eating and cognitive restraint
437 could underlie the association between poor sleep quality and higher BMI among
438 young adults. In this context, it is likely that young adults with poor sleep quality
439 are more likely to cope with negative emotions with food, which, in turn, could be
440 associated with higher cognitive restraint, becoming a vicious cycle that has a
441 negative impact on body weight. It is also worth noting that eating habits differed
442 as a function of eating behaviors, being emotional and uncontrolled eating
443 associated with a higher consumption of fast-food and sweets (only in the case
444 of emotional eating). Meanwhile, cognitive restraint was related to the
445 consumption of fruits and vegetables, and a lower consumption of commercially
446 baked goods. Although, the latter was not associated with a lower BMI. These
447 findings could open a new framework when designing strategies for obesity
448 prevention in young adults, which according to our results, should consider both
449 sleep and eating behaviors. Finally, our data also revealed that sleep quality has

450 a significant association with the adherence to the Mediterranean diet, suggesting
451 the potential role of nuts and fish as sleep promoting foods in this population.

452

453 **Author Contributions**

454 MFZR, TC and MIP designed the study; MFZR, TC and MIP acquired the data;
455 MFZR and AH analyzed the data; MFZR wrote the first draft; AH, TC, and MIP
456 revised the manuscript. All authors read and approved the final manuscript.

457

458 **Funding**

459 MFZR was supported by a scholarship from the 'Consejo Nacional de Ciencia y
460 Tecnologia' CONACYT from Mexico.

461

462 **Ethical Statement**

463 All study procedures were conducted according to the Declaration of Helsinki and
464 were approved by the Ethics Committee the University of Barcelona
465 (IRB00003099).

466

467 **Conflict of Interests**

468 The authors declare no conflict of interest.

469

470 **References**

471 Al Khatib, H. K., Harding, S. V., Darzi, J., & Pot, G. K. (2017). The effects of
472 partial sleep deprivation on energy balance: A systematic review and meta-
473 analysis. *European Journal of Clinical Nutrition*, 71(5), 614–624.
474 <https://doi.org/10.1038/ejcn.2016.201>

- 475 Barragán, R., Zuraikat, F. M., Tam, V., Scaccia, S., Cochran, J., Li, S., ... St-
476 Onge, M. P. (2021). Actigraphy-derived sleep is associated with eating
477 behavior characteristics. *Nutrients*, 13(3), 1–12.
478 <https://doi.org/10.3390/nu13030852>
- 479 Blumfield, M. L., Bei, B., Zimberg, I. Z., & Cain, S. W. (2018). Dietary
480 disinhibition mediates the relationship between poor sleep quality and body
481 weight. *Appetite*, 120, 602–608. <https://doi.org/10.1016/j.appet.2017.10.022>
- 482 Bryant, E. J., Rehman, J., Pepper, L. B., & Walters, E. R. (2019). Obesity and
483 Eating Disturbance: the Role of TFEQ Restraint and Disinhibition. *Current*
484 *Obesity Reports*, 8(4), 363–372. [https://doi.org/10.1007/s13679-019-](https://doi.org/10.1007/s13679-019-00365-x)
485 00365-x
- 486 Burrows, T., Fenton, S., & Duncan, M. (2020). Diet and sleep health: a scoping
487 review of intervention studies in adults. *Journal of Human Nutrition and*
488 *Dietetics*, 33(3), 308–329. <https://doi.org/10.1111/jhn.12709>
- 489 Cappelleri, J. C., Bushmakin, A. G., Gerber, R. A., Leidy, N. K., Sexton, C. C.,
490 Lowe, M. R., & Karlsson, J. (2009). Psychometric analysis of the Three-
491 Factor Eating Questionnaire-R21: Results from a large diverse sample of
492 obese and non-obese participants. *International Journal of Obesity*, 33(6),
493 611–620. <https://doi.org/10.1038/ijo.2009.74>
- 494 Chaput, J. P., & Dutil, C. (2016). Lack of sleep as a contributor to obesity in
495 adolescents: Impacts on eating and activity behaviors. *International Journal*
496 *of Behavioral Nutrition and Physical Activity*, 13(1), 1–9.
497 <https://doi.org/10.1186/s12966-016-0428-0>
- 498 Chen, W. L., & Chen, J. H. (2019). Consequences of inadequate sleep during

499 the college years: Sleep deprivation, grade point average, and college
500 graduation. *Preventive Medicine*, 124, 23–28.
501 <https://doi.org/10.1016/j.ypped.2019.04.017>

502 Del Brutto, O. H., Mera, R. M., Ha, J. eun, Gillman, J., Zambrano, M., & Castillo,
503 P. R. (2016). Dietary fish intake and sleep quality: A population-based
504 study. *Sleep Medicine*, 17, 126–128.
505 <https://doi.org/10.1016/j.sleep.2015.09.021>

506 Deliens, T., Deforche, B., Chapelle, L., & Clarys, P. (2019). Changes in weight
507 and body composition across five years at university: A prospective
508 observational study. *PLoS ONE*, 14(11), 1–10.
509 <https://doi.org/10.1371/journal.pone.0225187>

510 Dinis, J., & Bragança, M. (2018). Quality of sleep and depression in college
511 students: A systematic review. *Sleep Science*, 11(4), 290–301.
512 <https://doi.org/10.5935/1984-0063.20180045>

513 Dweck, J. S., Jenkins, S. M., & Nolan, L. J. (2014). The role of emotional eating
514 and stress in the influence of short sleep on food consumption. *Appetite*,
515 72, 106–113. <https://doi.org/10.1016/j.appet.2013.10.001>

516 Fedewa, M. V., Das, B. M., Evans, E. M., & Dishman, R. K. (2014). Change in
517 weight and adiposity in college students: A systematic review and meta-
518 analysis. *American Journal of Preventive Medicine*, 47(5), 641–652.
519 <https://doi.org/10.1016/j.amepre.2014.07.035>

520 Ferranti, R., Marventano, S., Castellano, S., Giogianni, G., Nolfo, F., Rametta,
521 S., ... Mistretta, A. (2016). Sleep quality and duration is related with diet
522 and obesity in young adolescent living in Sicily, Southern Italy. *Sleep*

523 *Science*, 9(2), 117–122. <https://doi.org/10.1016/j.slsci.2016.04.003>

524 Gibson, E. (2006). Emotional influences on food choice: Sensory, physiological
525 and psychological pathways. *Physiology and Behavior*, 89(1), 53–61.
526 <https://doi.org/10.1016/j.physbeh.2006.01.024>

527 Godos, J., Ferri, R., Caraci, F., Cosentino, F. I. I., Castellano, S., Galvano, F., &
528 Grosso, G. (2019). Adherence to the Mediterranean Diet is Associated with
529 Better Sleep Quality in Italian Adults. *Nutrients*, 11(5), 976.
530 <https://doi.org/10.3390/nu11050976>

531 Goldstein, A. N., & Walker, M. P. (2014). The role of sleep in emotional brain
532 function. *Annual Review of Clinical Psychology*, 10, 679–708.
533 <https://doi.org/10.1146/annurev-clinpsy-032813-153716>

534 Greer, S. M., Goldstein, A. N., & Walker, M. P. (2013). The impact of sleep
535 deprivation on food desire in the human brain. *Nature Communications*, 4,
536 1–7. <https://doi.org/10.1038/ncomms3259>

537 Hansen, A., Dahl, L., Olson, G., Thornton, D., Graff, I., Frøyland, L., ...
538 Pallesen, S. (2014). Fish Consumption, Sleep, Daily Functioning, and Heart
539 Rate Variability. *Journal of Clinical Sleep Medicine*, 10(5), 567–575.

540 Hayes, A. J. (2018). PROCESS Macro. New York. Retrieved from
541 www.processmacro.org

542 Herman, C. P., & Mack, D. (1975). Restrained and unrestrained eating. *Journal*
543 *of Personality*, 43(4), 647–660. [https://doi.org/10.1111/j.1467-](https://doi.org/10.1111/j.1467-6494.1975.tb00727.x)
544 [6494.1975.tb00727.x](https://doi.org/10.1111/j.1467-6494.1975.tb00727.x)

545 Izydorczyk, B., Sitnik-Warchulska, K., Lizińczyk, S., & Lipiarz, A. (2019).

546 Psychological predictors of unhealthy eating attitudes in young adults.
547 *Frontiers in Psychology*, 10(MAR), 1–14.
548 <https://doi.org/10.3389/fpsyg.2019.00590>

549 Krističević, T., Štefan, L., & Sporiš, G. (2018). The associations between sleep
550 duration and sleep quality with body-mass index in a large sample of young
551 adults. *International Journal of Environmental Research and Public Health*,
552 15(4). <https://doi.org/10.3390/ijerph15040758>

553 Lundahl, A., & Nelson, T. D. (2015). Sleep and food intake: A multisystem
554 review of mechanisms in children and adults. *J Health Psychol*, 20(6), 794–
555 805. <https://doi.org/10.1177/1359105315573427>

556 Martin-Garcia, M., Vila-Maldonado, S., Rodriguez-Gomez, I., Faya, F. M.,
557 Plaza-Carmona, M., Pastor-Vicedo, J. C., & Ara, I. (2016). The Spanish
558 version of the Three Factor Eating Questionnaire-R21 for children and
559 adolescents (TFEQ-R21C): Psychometric analysis and relationships with
560 body composition and fitness variables. *Physiology and Behavior*, 165,
561 350–357. <https://doi.org/10.1016/j.physbeh.2016.08.015>

562 Memon, A. R., Gupta, C. C., Crowther, M. E., Ferguson, S. A., Tuckwell, G. A.,
563 & Vincent, G. E. (2021). Sleep and physical activity in university students: A
564 systematic review and meta-analysis. *Sleep Medicine Reviews*, 58,
565 101482. <https://doi.org/10.1016/j.smrv.2021.101482>

566 Mossavar-Rahmani, Y., Weng, J., Wang, R., Shaw, P. A., Jung, M., Sotres-
567 Alvarez, D., ... Patel, S. R. (2017). Actigraphic sleep measures and diet
568 quality in the Hispanic Community Health Study/Study of Latinos Sueño
569 ancillary study. *Journal of Sleep Research*, 26(6), 739–746.

570 <https://doi.org/10.1111/jsr.12513>

571 Nelson, M. C., Story, M., Larson, N. I., Neumark-Sztainer, D., & Lytle, L. A.
572 (2008). Emerging adulthood and college-aged youth: an overlooked age for
573 weight-related behavior change. *Obesity (Silver Spring, Md.)*, *16*(10),
574 2205–2211. <https://doi.org/10.1038/oby.2008.365>

575 Penaforte, F. R. de O., Minelli, M. C. S., Rezende, L. A., & Japur, C. C. (2019).
576 Anxiety symptoms and emotional eating are independently associated with
577 sweet craving in young adults. *Psychiatry Research*, *271*(December 2017),
578 715–720. <https://doi.org/10.1016/j.psychres.2018.11.070>

579 Pilz, L. K., Keller, L. K., Lenssen, D., & Roenneberg, T. (2018). Time to rethink
580 sleep quality: PSQI scores reflect sleep quality on workdays. *Sleep*, *41*(5),
581 1–8. <https://doi.org/10.1093/sleep/zsy029>

582 Polivy, J., & Herman, C. P. (1985). Dieting and Binging. A Causal Analysis.
583 *American Psychologist*, *40*(2), 193–201. [https://doi.org/10.1037/0003-](https://doi.org/10.1037/0003-066X.40.2.193)
584 [066X.40.2.193](https://doi.org/10.1037/0003-066X.40.2.193)

585 Provencher, V., Polivy, J., & Herman, C. P. (2009). Perceived healthiness of
586 food. If it's healthy, you can eat more! *Appetite*, *52*(2), 340–344.
587 <https://doi.org/10.1016/j.appet.2008.11.005>

588 R Core Team. (2019). R: A language and environment for statistical computing.
589 Vienna, Austria: R Foundation for Statistical Computing. Retrieved from
590 <https://www.r-project.org/>

591 Racine, S. E. (2018). Emotional ratings of high- and low-calorie food are
592 differentially associated with cognitive restraint and dietary restriction.
593 *Appetite*, *121*, 302–308. <https://doi.org/10.1016/j.appet.2017.11.104>

- 594 Ramírez-Contreras, C., Farrán-Codina, A., Izquierdo-Pulido, M., & Zerón-
595 Rugerio, M. F. (2021). A higher dietary restraint is associated with higher
596 BMI : a cross-sectional study in college students. *Physiology & Behavior*,
597 240, 113536.
- 598 Richards, A. L., & Specker, B. (2021). Evaluating hours of sleep and perceived
599 stress on dietary cognitive restraint in a survey of college students. *Journal*
600 *of American College Health*, 68(8), 824–831.
601 <https://doi.org/10.1080/07448481.2019.1618312>
- 602 Roenneberg, T., Wirz-Justice, A., & Mellow, M. (2003). Life between clocks:
603 daily temporal patterns of human chronotypes. *Journal of Biological*
604 *Rhythms*, 18, 80–90. https://doi.org/10.1007/978-1-4419-9893-4_58
- 605 Román-Viñas, B., Ribas-Barba, L., Ngoa, J., & Serra-Majem, L. (2013).
606 Validación en población catalana del cuestionario internacional de
607 actividad física. *Gaceta Sanitaria*, 27(3), 254–257. Retrieved from
608 [http://www.gacetasanitaria.org/es/validacion-poblacion-catalana-del-](http://www.gacetasanitaria.org/es/validacion-poblacion-catalana-del-cuestionario/articulo-resumen/S0213911112002658/)
609 [cuestionario/articulo-resumen/S0213911112002658/](http://www.gacetasanitaria.org/es/validacion-poblacion-catalana-del-cuestionario/articulo-resumen/S0213911112002658/)
- 610 Royuela, A., & Macías, J. . (1997). Propiedades clinimétricas de la versión
611 castellana del cuestionario de Pittsburg. *Vigilia-Sueño*, 9(2), 81–94.
- 612 Saleh-ghadimi, S., Dehghan, P., Farhangi, M. A., & Asghari-jafarabadi, M.
613 (2019). Could emotional eating act as a mediator between sleep quality
614 and food intake in female students ?, 3, 1–9.
- 615 Serra-Majem, L., Ribas, L., Ngo, J., Ortega, R. M., Garcia, A., Perez-Rodrigo,
616 C., & Aranceta, J. (2004). Food, youth and the Mediterranean diet in Spain.
617 Development of KIDMED, Mediterranean Diet Quality Index in children and

618 adolescents. *Public Health Nutr*, 7(7), 931–935.
619 <https://doi.org/10.1079/PHN2004556>

620 Silva, C. M., Mota, M. C., Miranda, M. T., Paim, S. L., Waterhouse, J., Crispim,
621 C. A., ... Chronotype, C. (2016). Chronotype , social jetlag and sleep debt
622 are associated with dietary intake among Brazilian undergraduate students.
623 *Chronobiology International*, 33(6), 740–748.
624 <https://doi.org/10.3109/07420528.2016.1167712>

625 St-Onge, M.-P. (2017). Sleep-obesity relation: underlying mechanisms and
626 consequences for treatment. *Obesity Reviews*, 18(Suppl 1), 34–39.
627 <https://doi.org/10.1111/obr.12499>

628 St-Onge, M. P., McReynolds, A., Trivedi, Z. B., Roberts, A. L., Sy, M., & Hirsch,
629 J. (2012). Sleep restriction leads to increased activation of brain regions
630 sensitive to food stimuli. *American Journal of Clinical Nutrition*, 95(4), 818–
631 824. <https://doi.org/10.3945/ajcn.111.027383>

632 St-Onge, M., Roberts, A., Chen, J., Kelleman, M., O’Keeffe, M., RoyChoudhury,
633 A., & Jones, P. (2011). Short sleep duration increases energy intakes but
634 does not change energy expenditure in normal-weight individuals. *Am J*
635 *Clin Nutr*, 94(2), 410–416.

636 St-Onge, Marie-Pierre, Mikic, A., & Pietrolungo, C. E. (2016). Effects of Diet on
637 Sleep Quality. *Advances in Nutrition*, 7(5), 938–949.
638 <https://doi.org/10.3945/an.116.012336>

639 Wickham, S. R., Amarasekara, N. A., Bartonicek, A., & Conner, T. S. (2020).
640 The Big Three Health Behaviors and Mental Health and Well-Being Among
641 Young Adults: A Cross-Sectional Investigation of Sleep, Exercise, and Diet.

642 *Frontiers in Psychology*, 11(December), 1–10.
643 <https://doi.org/10.3389/fpsyg.2020.579205>

644 Yamamoto, K., Ota, M., Minematsu, A., Motokawa, K., Yokoyama, Y., Yano, T.,
645 ... Yoshizaki, T. (2018). Association between adherence to the Japanese
646 food guide spinning top and sleep quality in college students. *Nutrients*,
647 10(12). <https://doi.org/10.3390/nu10121996>

648 Zerón-Ruggerio, M. F., Cambras, T., & Izquierdo-Pulido, M. (2019). Social Jet
649 Lag Associates Negatively with the Adherence to the Mediterranean Diet
650 and Body Mass Index among Young Adults. *Nutrients*, 11(8), 1756.
651 <https://doi.org/10.3390/nu11081756>

652 Zuraikat, F. M., Wood, R. A., Barragán, R., & St-Onge, M. P. (2021). Sleep and
653 Diet: Mounting Evidence of a Cyclical Relationship. *Annual Review of*
654 *Nutrition*, 41, 309–332. [https://doi.org/10.1146/annurev-nutr-120420-](https://doi.org/10.1146/annurev-nutr-120420-021719)
655 021719

661 **Figure Captions**

662 **Figure 1.** Path diagram for the total effect of sleep quality on the body mass index
663 (BMI) and the indirect effects of sleep quality on the BMI through the potential
664 mediation of emotional eating and cognitive restraint. In the top diagram “c” is the
665 total effect of exposure (X) on outcome (Y) ignoring the mediator (M). In the
666 bottom diagram, the mediation of the effect of sleep quality on the BMI through
667 diet quality and eating behaviors (emotional eating and cognitive restraint) is
668 shown, where “a” is the effect of exposure on mediator and “b” is the effect of
669 mediator on outcome. Effect c’ is the direct effect of exposure on outcome while
670 adjusting for the mediator.

671

672 **Figure 2.** Mediation model highlighting the effect of sleep quality on BMI via
673 emotional eating and cognitive restraint. Unstandardized coefficients [95%
674 confidence interval] are shown. The model was adjusted for age, gender,
675 chronotype, diet quality, and physical activity. Solid lines indicate statistically
676 significant paths, while dotted lines indicate non-significant paths; *** $p < 0.001$.

677

678 **Figure 3.** Associations between Pittsburg Sleep Quality Index and the 16 items
679 of the Mediterranean Diet Quality Index. General linear model adjusted for age,
680 gender, BMI, physical activity, and chronotype were conducted to test these
681 associations. P-values were corrected using the Benjamini–Hochberg method,
682 assuming a False Discovery Rate (FDR) of 5%. * $p < 0.05$

683

Table 1. General characteristics of the population studied.

Total sample, n	925
Age, years	21.4 (2.5)
Gender, %women	77.8
Sleep variables	
Sleep quality, score	5.4 (2.5)
Good sleep quality, %	40.8
Chronotype (MSF), hh:mm	04:47 (1.11)
BMI, kg/m²	21.8 (3.1)
Diet quality, score	
Poor (≤ 3 points), %	5.9
Average (4–7 points), %	47.6
Good (8–12 points), %	46.5
Eating behavior	
Emotional eating, score	1.7 (0.6)
Cognitive restraint, score	2.1 (0.5)
Uncontrolled eating, score	2.0 (0.6)
Physical activity, METs	2 161.1(1 722.6)

BMI, Body mass index; METs, Metabolic Equivalents of Task; MSF, Midpoint of sleep in free-days. Data are expressed as mean and standard deviation for continuous data, and percentages for categorical data.

Table 2. Associations between the Pittsburgh Sleep Quality Index (PSQI) and the body mass index (BMI) with diet quality and eating behaviors (emotional eating, cognitive restraint and uncontrolled eating).

	PSQI, score		BMI, kg/m ²	
	β [95%CI]	<i>p</i> -value	β [95%CI]	<i>p</i> -value
Diet quality, score	-0.14 [-0.19; -0.08]	<0.00001	0.04 [-0.05; 0.13]	0.339
Eating behaviors				
Emotional eating, score	0.04 [0.03; 0.06]	<0.00001	0.89 [0.60; 1.18]	<0.00001
Cognitive restraint, score	0.03 [0.01; 0.04]	<0.001	1.37 [1.02; 1.71]	<0.00001
Uncontrolled eating, score	0.02 [0.01; 0.04]	0.001	0.10 [-0.24; 0.44]	0.556

CI, confidence interval. Associations between sleep quality and BMI were tested using linear regression analyses. All analyses were adjusted for age, gender, BMI, diet quality, physical activity, and chronotype (except when the covariate was the dependent variable of the model). The table shows the unstandardized coefficient (β), CI, and *p*-value associated with each predictor variable. Significant *p*-values are shown in bold.

AUTHOR ACCEPTED VERSION

Table 3. Summary of indirect effects from sleep quality¹ to BMI.

	Coefficient	S.E.	95% CI
Direct effect	0.01	0.04	-0.06; 0.09
Total indirect effect	0.07	0.02	0.04; 0.10*
<i>Indirect effect (via mediators)</i>			
Emotional eating	0.03	0.01	0.02; 0.06*
Cognitive restraint	0.04	0.01	0.02; 0.06*

BMI, Body mass index. ¹Sleep quality was measured with the Pittsburgh Sleep Quality Index.

Pathway analyses were conducted using the PROCESS tool. Analyses were adjusted for age, gender, chronotype, diet quality, and physical activity. *p<0.05

AUTHOR ACCEPTED VERSION

Table 4. Associations between eating behaviors (cognitive restraint, emotional eating and uncontrolled eating) and the 16 items of the Mediterranean Diet Quality Index. General linear model adjusted for age, gender, BMI, physical activity, and chronotype were used to calculate adjusted differences between categories, the reference category was ‘Yes’. P-values were corrected using the Benjamini–Hochberg method, assuming a False Discovery Rate (FDR) of 5%. *p<0.05, ***p<0.001.

	Emotional eating, score	Cognitive restraint, score	Uncontrolled eating, score
	β [95% CI]	β [95% CI]	β [95% CI]
First serving of fruit, daily	0.01 [-0.09; 0.11]	0.06 [-0.02; 0.14]	-0.06 [-0.15; 0.03]
Second serving of fruit, daily	-0.06 [-0.14; 0.02]	0.11 [0.04; 0.18]*	-0.09 [-0.16; -0.01]
Fresh or cooked vegetables, daily	-0.05 [-0.16; 0.06]	0.19 [0.10; 0.28]***	-0.12 [-0.21; -0.02]
Fresh or cooked vegetables, >1 time/day	-0.02 [-0.10; 0.07]	0.13 [0.06; 0.20]***	-0.08 [-0.16; -0.01]
Regular fish consumption, 2-3 times/week	-0.01 [-0.10; 0.07]	0.05 [-0.02; 0.11]	-0.09 [-0.17; -0.02]
Fast-food, >1 time/week	0.30 [0.18; 0.42]***	-0.05 [-0.15; 0.05]	0.26 [0.15; 0.37]***
Legumes, >1 time/week	-0.04 [-0.14; 0.06]	0.02 [-0.06; 0.10]	-0.07 [-0.15; 0.02]
Pasta or rice, \geq 5 times/week	-0.01 [-0.09; 0.07]	-0.06 [-0.13; 0.01]	0.05 [-0.03; 0.12]
Cerales/grains for breakfast	0.00 [-0.12; 0.13]	0.07 [-0.03; 0.17]	0.06 [-0.05; 0.17]
Regular nut consumption, 2-3 times/week	-0.01 [-0.09; 0.08]	0.08 [0.01; 0.15]	-0.07 [-0.15; 0.00]
Use of olive oil at home	-0.22 [-0.54; 0.10]	-0.19 [-0.45; 0.07]	-0.18 [-0.46; 0.11]
Skipping breakfast	0.01 [0.16; -0.14]	0.11 [-0.01; 0.24]	0.08 [-0.05; 0.21]
Dairy product for breakfast	-0.08 [-0.17; 0.01]	-0.06 [-0.13; 0.02]	0.01 [-0.07; 0.10]
Commercially baked goods or pastries for breakfast	0.15 [0.04; 0.25]***	-0.16 [-0.25; -0.08]***	0.11 [0.01; 0.20]
Yogurts or cheese, daily	-0.08 [-0.16; 0.00]	-0.01 [-0.78; 0.06]	-0.03 [-0.10; 0.04]
Sweets and candy, every day	0.29 [0.08; 0.49]***	-0.15 [-0.32; 0.02]	0.24 [0.06; 0.42]*