

7. BIBLIOGRAFIA

A

Acton SL, Scherer PE, Lodish HF, Krieger M. Expression cloning of SR-BI, a CD36-related class B scavenger receptor. *J Biol Chem.* 1994;269:21003-21009.

Al-Haideri M, Goldberg IM, Galeano NF, Gleeson A, Vogel T, Gorecki M, Sturley SL, Deckelbaum RJ. Heparan sulfated proteoglycan-mediated uptake of apolipoprotein E-triglyceride-rich lipoprotein particles: a major pathway at physiological particle concentrations. *Biochemistry.* 1997;36:12766-12772.

Allain CC, Poon LS, Chan CSG, Richmond W, Fu PC. Enzymatic determination of the serum cholesterol. *Clin Chem.* 1974;20:470-475.

Ando J, Tsuboi H, Korenaga R, Takada Y, Toyama-Sorimachi N, Miyasaka M, et al. Shear stress inhibits adhesion of cultured endothelial cells to lymphocytes by downregulating VCAM-1 expression. *Am J Physiol.* 1994;267:C679-687.

Andreeva ER, Pugach IM, Orekhov AN. Collagen-synthesizing cells in initial and advanced atherosclerotic lesions of human aorta. *Atherosclerosis.* 1997;130:133-142.

Aqel NM, Ball RY, Waldmann H, Hutchinson MJ. Identification of macrophages and smooth muscle cells in human atherosclerosis using monoclonal antibodies. *J Pathol.* 1985;146:197-204.

Ardans JA, Economou AP, Martinson JM Jr, Zhou M, Wahl LM. Oxidized low-density and high-density lipoproteins regulate the production of matrix metalloproteinase 1 and 9 by activated monocytes. *J Leukoc Biol.* 2002;71:1012-1008.

Ashcom JD, Tiller SE, Dickerson K et al. The human alpha 2-macroglobulin receptor: identification of a 420-kd cell surface glycoprotein specific for the activated conformation of alpha 2-macroglobulin. *J Cell Biol.* 1990;110:1041-1048.

Aviram M, Maor Y, Keidar S, Hayek T, Oiknine J, Bae EIY, Adler Z, Kertzman V, Milo S. Lesioned low density lipoprotein in atherosclerotic apolipoprotein E-deficient transgenic mice and in humans is oxidized and aggregated. *Biochim Biophys Res Commun.* 1995;216:501-513.

B

Bachem MG, Wendelin D, Schneiderhan W, Haug C, Zorn U, Gross HJ, Schmid-Kotsas A, Grunert A. Depending on their concentration oxidized low density lipoproteins stimulate extracellular matrix synthesis or induce apoptosis in human coronary artery smooth muscle cells. *Clin Chem Lab Med.* 1999;37:319-326.

Badimon L, Chesebro JH, Badimon JJ. Thrombus formation on ruptured atherosclerotic plaques and rethrombosis on evolving thrombi. *Circulation.* 1992;86(SuplIII):74-85.

Bailey AJ, Paul RG, Knott L. Mechanisms of maturation and ageing of collagen. *Mech Ageing Develop.* 1998;106:1-56.

Banerjee SD, Toole BP. Hyaluronan-binding protein in endothelial cell morphogenesis. *J Cell Biol.* 1992;119:643-652.

Barnes MJ, Knight CG, Farndale RW. Collagens and atherosclerosis: cell-collagen interaction. Ed: Jacotot B, Mathè D, Fruchart CC. Elsevier Science Pte Ltd *Atherosclerosis XI*. 1998;299-306.

Barnes MJ, Farndale RW. Collagens and atherosclerosis. *Exp Gerontol.* 1999;34:513-525.

Bayliss High O. Lipids. In Bancroft JD, Stevens A eds. *Theory and practice of histological techniques*. Third edition Churchill Livingstone. Edinburg London Melbourne and New York. 1990:215-244.

Beisiegel U, Krapp A, Weber W, Olivecrona G. The role of alpha 2 M receptor/LRP in chylomicron remnant metabolism. *Ann N Y Acad Sci.* 1994;737:53-69.

Benítez S, Sánchez-Quesada JL, Lucero L, Arcelus R, Ribas V, Jorba O, Castellvi A, Alonso E, Blanco-Vaca F, Ordoñez-Llanos J. Changes in low-density lipoprotein electronegativity and oxidizability after aerobic exercise are related to the increase in associated non-esterified fatty acids. *Atherosclerosis.* 2002;160:223-232.

Bennet MR, Evan GI, Schwartz SM. Apoptosis of human vascular smooth muscle cells derived from normal vessels and coronary atherosclerotic vessels. *J Clin Invest.* 1995;95:2266-2274.

Bennet MR, Boyle JJ. Apoptosis of vascular smooth muscle cells in atherosclerosis. *Atherosclerosis.* 1998;138:3-9.

Bennet MR. Apoptosis of vascular smooth muscle cells in vascular remodelling and atherosclerotic plaque rupture. *Cardiovasc Res.* 1999;41:361-368.

Bhakdi S, Dorweiler B, Kirchmann R, Torzewski J, Weise E, Tranum-Jensen J, Walev I, Wieland E. On the pathogenesis of atherosclerosis: enzymatic transformation of human low density lipoprotein to an atherogenic moiety. *J Exp Med.* 1995;182:1959-1971.

Bihari-Varga M. Proteoglycan-lipoprotein interactions. Ed: Jacotot B, Mathè D, Fruchart CC. Elsevier Science Pte Ltd *Atherosclerosis XI*. 1998:307-310.

Birnboim HC, Doly J. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic acid research.* 1979;7:1513-1522.

Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol.* 1959;37:911-916.

Bo Hu, Dayuan Li, Tatsuya Sawamura, Jawahar L, Mehta JL. Oxidized LDL through LOX-1 modulates LDL-receptor expression in human coronary artery endothelial cells. *Biochem Biophys Res Commun.* 2003;307:1008-1012.

Boucher P, Lorgeil M, Salen P, Crozier P, Delaye J, Valloon JJ, Geyssant A, Dante R. Effect of dietary cholesterol on low density lipoprotein receptor, 3-hydroxy-3-methylglutaryl-CoA reductase and low density lipoprotein receptor-related protein mRNA expression in healthy humans. *Lipids*. 1998;33:1177-1186.

Boucher P, Liu PV, Gotthardt M, Hiesberger T, Anderson RGW, Herz J. Platelet-derived growth factor mediates tyrosine phosphorylation of the cytoplasmic domain of the low density lipoprotein receptor-related protein in caveolae. *J Biol Chem*. 2002;275:15507-15513.

Boucher P, Gotthardt M, Li WP, Anderson RG, Herz J. LRP: role in vascular wall integrity and protection from atherosclerosis. *Science*. 2003;300:329-332.

Boulanger CM, Tanner FC, Bea ML, Hahn AW, Werner A, Luscher F. Oxidized low density lipoproteins induce mRNA expression and release of endothelin from human and porcine endothelium. *Circ Res*. 1992;70:1191-1197.

Bradbury P, Gordon KC. Connective tissue. In Bancroft JD, Stevens A eds. Theory and practice of histological techniques. Third edition Churchill Livingstone. Edinburgh London Melbourne and New York. 1990:119-142.

Breslow JL. Cardiovascular disease burden increases, NIH funding decreases. *Nat Med*. 1997;3:600-601.

Brown MS, Goldstein JL, Krieger M, Ho YK, Anderson RG. Reversible accumulation of cholesteryl esters in macrophages incubated with acetylated lipoproteins. *J Cell Biol*. 1979;82:597-613.

Brown MS, Goldstein JL. Lipoprotein metabolism in the macrophage: implications for cholesterol deposition in atherosclerosis. *Annu Rev Biochem*. 1983;52:223-261.

Brown MS, Goldstein JL. A receptor-mediated pathway for cholesterol homeostasis. *Science*. 1986;232:34-47.

Brown MS, Goldstein JL. The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor. *Cell*. 1997;89:331-340.

Brown CT, Nugent MA, Lau FW, Trinkaus-Randall V. Characterization of proteoglycans synthesized by cultured corneal fibroblasts in response to transforming growth factor β and fetal calf serum. *J Biol Chem*. 1999;274:7111-7119.

C

Calvo D, Gómez-Coronado D, Suárez Y, Lasunción MA, Vega MA. Human CD36 is a high affinity receptor for the native lipoproteins HDL, LDL, and VLDL. *J. Lipid Res*. 1998;39:777-88.

Camejo G, Hurt E, Romano M. Properties of lipoprotein complexes isolated by affinity chromatography from human aorta. *Biomed Biophys Acta*. 1985;44:389-401.

Camejo G, Rosengren B, Olsson U, Lopez F, Olofson SO, Westerlund C, Bondjers G. Molecular basis of the association of arterial proteoglycans with low density lipoproteins: its effects on the structure of the lipoprotein particle. *Eur Heart J.* 1990;11 Suppl E:1644-1673.

^aCamejo G, Hurt E, Wiklund O, Rosengren B, López F, Bondjers G. Modification of low-density lipoproteins induced by arterial proteoglycans and chondroitin-6-sulfate. *Biochem Biophys Acta.* 1991;1096:253-261.

^bCamejo G, Hurt-Camejo E, Rosengren B, Wiklund O, López F, Bondjers G. Modification of copper-catalyzed oxidation of low density lipoprotein by proteoglycans and glycosaminoglycans. *J Lipid Res.* 1991;32:1983-1991.

Camejo G, Fager G, Rosengren B, Hurt-Camejo E, Bondjers G. Binding of low density lipoproteins by proteoglycans synthesized by proliferating and quiescent human arterial smooth muscle cells. *J Biol Chem.* 1993;268:14131-14137.

Camejo G, Hurt-Camejo E, Olsson U, Bondjers G. Proteoglycans and lipoproteins in atherosclerosis. *Curr Opin Lipidol.* 1993;4:385-391.

Camejo G, Hurt-Camejo E, Wiklund O, Bondjers G. Association of apoB lipoproteins with arterial proteoglycans: pathological significance and molecular basis. *Atherosclerosis.* 1998;139:205-222.

Camejo G, Olsson U, Hurt-Camejo E, Baharamian N, Bondjers G. The extracellular matrix on atherogenesis and diabetes-associated vascular disease. *Atherosclerosis Supplements.* 2002;3:3-9.

Cathcart MK, McNally AK, Morel DW, Chisolm GM. Superoxide anion participation in human monocyte-mediated oxidation of low density lipoprotein and conversion of low density lipoprotein to a cytotoxin. *J Immunol.* 1989;142:1963-1969.

Chao FF, Blanchette-Mackie EJ, Tertov VV, Skarlatos SI, Chen YJ, Kruth HS. Hydrolysis of cholesteryl ester in low density lipoprotein converts this lipoprotein into a liposome. *J Biol Chem.* 1992;267:4992-4998.

Chang MY, Potter-Perigo S, Tsui C, Chait A, Wight TN. Oxidized low density lipoproteins regulate synthesis of monkey aortic smooth muscle cell proteoglycans that have enhanced native low density lipoprotein binding properties. *J Biol Chem.* 2000;275:4766-4773.

Chapell DA, Fry GL, Waknitz MA, Muhonen LE, Pladet MW, Iverius PH, Strickland DK. Lipoprotein lipase induces catabolism of normal triglyceride-rich lipoproteins via the low density lipoprotein receptor-related protein/alpha 2-macroglobulin receptor in vitro. A process facilitated by cell-surface proteoglycans. *J Biol Chem.* 1993;268:14168-14175.

Chappey O, Dosquet C, Wautier MP, Wautier JL. Advanced glycation end products, oxidant stress and vascular lesions. *Eur J Clin Invest.* 1997;27:97-108.

Cherchi GM, Coinu R, Demuro P, Formato M. Structural and functional modifications of human aorta proteoglycans in atherosclerosis. *Matrix.* 1990;10:362-372.

Chisolm GM, Penn MS. Oxidized lipoproteins and atherosclerosis. In Fuster V, Ross R, Topol ES, eds. Atherosclerosis and artery disease. Lippincott-Raven Publisher Philadelphia ©. 1996:129-149.

Chomeczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem*. 1987;162:156-159.

Cizmeci-Smith G, Stahl RC, Showalter LJ, Carey DJ. Differential expression of transmembrane proteoglycans in vascular smooth muscle cells. *J Biol Chem*. 1993;268:18740-18747.

Cseh K, Karadi I, Rischak K, Szollar L, Janoki G, Jakab L, Romics L. Binding of fibronectin to human lipoproteins. *Clin Chim Acta*. 1989;182:75-85.

Cushing SD, Berliner JA, Valente AJ, Territo MC, Navab M, Parhami F, Gerrity R, Schwartz CJ, Fogelman AM. Minimally modified low density lipoprotein induces monocyte chemotactic protein 1 in human endothelial cells and smooth muscle cells. *Proc Natl Acad Sci USA*. 1990;87:5134-5138.

Cybulsky MI, Gimbrone MA Jr. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherosclerosis. *Science*. 1991;251:788-791.

D

Daugherty A, Rateri DL. Presence of LDL receptor-related protein/ α_2 -macroglobulin receptors in macrophages of atherosclerotic lesions from cholesterol-fed New Zealand abd heterozygous Watanable heritable hyperlipidemic rabbits. *Arterioscler Thromb*. 1994;14:2017-2024.

Davies PF. Flow-mediated endothelial mechanotransduction. *Physiol Rev*. 1995;75:519-560.

Drake TA, Hannani K, Fei H, Lai S, Berliner JA. Minimally oxidized low density lipoprotein induces tissue factor expression in cultured human endothelial cells. *Am J Pathol*. 1991;138:601-607.

E

Eisenberg S, Sehayek E, Olivecrona T, Vlodavsky i. Lipoprotein lipase enhances binding of lipoproteins to heparan sulfate on cell surfaces and extracellular matrix. *J Clin Invest*. 1992;90:2013-2021.

Endemann G, Stanton LW, Madden KS, Bryant CM, White RT, et al. *J Biol Chem*. 1993;268:11811-11816.

Evanko SP, Raines EW, Ross R, Gold LI, Wight TN. Proteoglycan distribution in lesions of atherosclerosis depends on lesion severity, structural characteristics, and the proximity of platelet-derived growth factor and transforming growth factor-beta. Am J Pathol. 1998;152:533-546.

F

Falk E, Shah PK, Fuster V. Coronary plaque disruption. Circulation. 1995;92:657-671.

Fawcett DW. Sistemas vasculares y linfático. En Fawcet MD, eds. Tratado de histología. Interamericana Mc Graw-Hill. 1989:369-403.

Fernández-Ortiz A, Badimon JJ, Falk E, Fuster V, Meyer B, Maihac A, et al. Characterization of the relative thrombogenicity of atherosclerotic plaque components: implications for consequences of plaque rupture. J Am Coll Cardiol. 1994;23:1562-1569.

Fogelman AM, Shechter I, Seager J, Hokom M, Child JS, Edwards PA. Malondialdehyde alteration of low density lipoproteins leads to cholestryl ester accumulation in hman monocyte-macrophages. Proc Natl Acad Sci USA. 1980;77:2214-2218.

Freeman MW. Macrophage scavenger receptors. Curr Opin Lipid. 1994;5:143-148.

Fujioka Y, Cooper AD, Fong LG. Multiple processes are involved in the uptake of chylomicron remnant by mouse peritoneal macrophages. J Lipid Res. 1998;39:2339-2349.

Fuki IV, Iozzo RV, Williams KJ. Pelecan heparan sulfate proteoglycan. J Biol Chem. 2000;275:25742-25750.

Fuki IV, Kuhn K, Lomazov I, Rothman V, Tuszyński G, Iozzo R, Swenson T, Fisher E, Williams KJ. Perlecan and syndecan proteoglycans: two new classes of lipoprotein receptors. FASEB J. 1996;10:A1140 (Abstract).

Fuki IV, Kuhn K, Lomazov I, Rothman V, Tuszyński G, Iozzo R, Swenson T, Fisher E, Williams KJ. The syndecan family of proteoglycans. Novel receptors mediating internalization of atherogenic lipoproteins in vitro. J Clin Invest. 1997;100:1611-1622.

G

Gaeta BA, Borthwick I, Stanley KK. The 5'-flanking region of the α_2 -MR/LRP genes contains and enhancer-like cluster of Sp1 binding sites. Biochim Biophys Acta. 1994;1219:307-313.

Galeano NF, Milne R, Marcel YL, Walsh MT, Levy E, Ngu'yen TD, Gleeson A, Arad Y, Witte L, al-Haideri M, et al. Apoprotein B structure and receptor recognition of triglyceride-rich low density lipoprotein (LDL) is modified in small LDL but not in triglyceride-rich LDL of normal size. J Biol Chem. 1994;269:511-519.

Galeano NF, Al-Haideri M, Keyserman F, Rumsey SC, Deckelbaum RJ. Small dense low density lipoprotein has increased affinity for LDL receptor-independent cell surface binding sites: a potential mechanism for increased atherogenicity. *J Lipid Res.* 1998;39:1263-1273.

Gal is ZS, Sukhova GK, Lara MW, Lobby P. Increased expression of matrix metalloproteinases and matrix degrading activity in vulnerable regions of human atherosclerotic plaques. *J Clin Invest.* 1994;94:2493-2503.

Garner B, Baoutina A, Dean RT, Jessup W. Regulation of serum-induced lipid accumulation in human monocyte-derived macrophages by interferon-gamma. Correlations with apolipoprotein E production, lipoprotein lipase activity and LDL receptor-related protein expression. *Atherosclerosis.* 1997;128:47-58.

Gardner G, Banka CL, Roberts KA, Mullick AE, Rutledge JC. Modified LDL-mediated increases in endothelial layer permeability are attenuated with 17b-estradiol. *Arterioscler Thromb Vasc Biol.* 1999;19:854-861.

Gewirtz AM, Sokol DL, Ratajczak MZ. Nucleic acid therapeutics: state of the art and future prospects. *Blood.* 1998;92:712-736.

Ghiselli G, Lindahl U, Salmivirta M. Foam cell conversion of macrophages alters the biosynthesis of heparan sulfate. *Biochem Biophys Res Commun.* 1998;247:790-795.

Glass CK, Witztum JL. Atherosclerosis. The Road Ahead. *Cell* 2001;104:503-516.

Godyn a S, Liau G, Popa I, Stefansson S, Argraves WS. Identification of the low density lipoprotein receptor-related protein (LRP) as an endocytic receptor for thrombospondin-1. *J Cell Biol.* 1995;129:1403-1410.

Goldberg IJ, Pillarisetti S, Obunike JC, Blaner WS, Wagner WD, Rutledge JC. Lipoprotein association with cells and matrix: modulation by lipase and proteoglycans. *Atherosclerosis XI* 1998. International Congress Series 1155.

^a**Goldstein JL**, Brown MS. The LDL receptor locus and the genetics of familial hypercholesterolemia. *Annu Rev Genet.* 1979;13:259-289.

^b**Goldstein JL**, HO YK, Basu SK, Brown MS. Binding site on macrophages that mediates uptake and degradation of acetylated low density lipoprotein, producing massive cholesterol deposition. *Proc Natl Acad Sci USA.* 1979;76:333-337.

Gong Q, Pitas RE. Synergistic effects of growth factors on the regulation of smooth muscle cell scavenger receptor activity. *J Biol Chem.* 1995;270:21672-21678.

Grafe M, Auch-Schwelk W, Hertel H, Terbeek D, Steinheider G, Loebe M, et al. Human cardiac microvascular and macrovascular endothelial cells respond differently to oxidatively modified LDL. *Atherosclerosis.* 1998;137:87-95.

Gutierrez P, O'Brian K, Ferguson M, Nikkari ST, Alpers CE, Wight TN. Differences in the distribution of versican, decorin, and biglycan in atherosclerotic human coronary arteries. *Cardiovasc Pathol.* 1997;6:271-278.

Guyton JR, Klemp KF, Mims PM. Altered ultrastructural morphology of self-aggregated low density lipoproteins: coalescence of lipid domains forming droplets and vesicles. *J Lipid Res.* 1991;32:953-961.

Guyton JR, Klemp KF. Transitional features in human atherosclerosis. Intimal thickening cholesterol clefts, and cell loss in human aortic fatty streaks. *Am J Pathol.* 1993;143:1444-1457.

Guyton JR, Llemp KF. Development of the atherosclerotic core region. Chemical and ultrastructural analysis of microdissected atherosclerotic lesions from human aorta. *Arterioscler Thromb.* 1994;14:1305-1314.

Guyton JR, Klemp KF. Development of the lipid-rich core in human atherosclerosis. *Arterioscler Thromb Vasc Biol.* 1996;16:4-11.

H

Haberland ME, Fong L, Cheng L. Malondialdehyde-altered protein occurs in atheroma of Watanabe heritable hyperlipidemic rabbits. *Science.* 1988;241:215-218.

Hahn-Dantona E, Ruiz JF, Bornstein P, Strickland DK. The low density lipoprotein receptor-related protein modulates levels of matrix metalloproteinase 9 (MMP-9) by mediating its cellular catabolism. *J Biol Chem.* 2001;276:15498-15503.

Hakala JK, Öörni K, Pentikäinen MO, Hurt-Camejo E, Kovanen PT. Lipólisis of LDL by human secretory phospholipase A2 induces particle fusion and enhances the retention of LDL to human aortic proteoglycans. *Arterioscler Thromb Vasc Biol.* 2001;21:1053-1061.

Halvorsen B, Aas MA, Kulseth MA, Drevon CA, Christiansen EN, Kolset SO. Proteoglycans in macrophages: characterization and possible role in the cellular uptake of lipoproteins. *Biochem J.* 1998;331:743-752.

Hanahan D. Technique for transformation of *E. coli*. DNA cloning: A practical approach. Glover DM. Ed IRL Press, Oxford. 1985;2:109:135.

Harpel PC, Gordon BR, Parker TS. Plasmin catalyzes binding of lipoprotein (a) to immobilized fibrinogen and fibrin. *Proc Natl Acad Sci USA.* 1989;86:3847-3851.

Havel R, Eder H, Bragdon J. The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J Clin Invest.* 1984;74:1890-1894.

Heinecke JW, Rosen H, Suzuki LA, Chait A. The role of sulfur-containing aminoacids in superoxide production and modification of low density lipoprotein by arterial smooth muscle cells. *J Biol Chem.* 1987;262:10098-10103.

Heinecke JW, Kawamura M, Suzuki LA, Chai A. Oxidation of low density lipoprotein by thiols: superoxide-dependent and independent mechanisms. *J Lipid Res.* 1993;34:2051-2061.

Henriksen T, Mahoney EM, Steinberg D. Enhance macrophage degradation of low density lipoprotein previously incubated with cultured endothelial cells: recognition by receptor for acetylated low density lipoproteins. *Proc Natl Acad Sci USA*. 1981;78:6499-64503.

Herrington GS, McGee JO. Principles and basic methodology of DNA/RNA detection by In situ hibridatin. In Herrington GS, McGee JO. Eds Diagnostic molecular pathology: A practical approach. New York:IRL Press. 1992;1:69-102.

Herz J, Kowal RC, Goldstein JL, Brown MS. Proteolytic processing of the 600 KD low density lipoprotein receptor-related protein (LRP) occurs in a trans-Golgi compartment. *EMBO J*. 1990;9:1769-1776.

Herz J, Clouthier DE, Hammer RE. LDL receptor-related protein internalizes and degrades uPA-PAI-1 complexes and is essential for embryo implantation. [published erratum appeared in Cell. 1993;73:428]. *Cell*. 1992;71:411-421.

Herz J, Strickland DK. LRP: a multifunctional scavenger and signaling receptor. *J Clin Inves*. 2001;108:779-784.

Hevonenja T, Pentikainen MO, Hyvonen MT, Kovanen PT, Ala-Korpela M. Structure of low density lipoprotein (LDL) particles: Basis for understanding molecular changes in modified LDL. *Biochim Biophys Acta*. 2000;1488:189-210.

^aHiltunen TP, Luoma JS, Nikkari T, Yla-Herttuala S. Expression of LDL receptor, VLDL receptor, LDL receptor-related protein, and scavenger receptor in rabbit atherosclerotic lesions: marked induction of scavenger receptor and VLDL receptor expression during lesion development. *Circulation*. 1998;97:1079-1086.

^bHiltunen TP, Ylä-Herttuala S. Expression of lipoprotein receptors in atherosclerotic lesions. *Atherosclerosis*. 1998;137 Supl:S81-S88.

Hoff HF, O'Neil J, Chisolm GM, Cole TB, Quehenberger O, Esterbauer H, Jurgens G. Modification of low density lipoprotein with 4-hydroxynonenal induces uptake by macrophages. *Arteriosclerosis*. 1989;9:538-549.

Hokanson JE, Krauss RM, Albers JJ, Austin MA, Brunzell JD. LDL physical and chemical properties in familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol*. 1995;15:452-459.

Hollmann J, Schmidt A, von Bassewitz D, Buddecke E. Relationship of sulfated glycosaminoglycans and cholesterol content in normal and atherosclerotic human aorta. *Arteriosclerosis*. 1989;9:154-158.

Hunt JV, Smith CCT, Wolff SP. Autoxidative glycosylation and possible involvement of peroxides and free radicals in LDL modification by glucose. *Diabetes*. 1990;39:1420-1424.

Hurt E, Bondjers G, Camejo G. Interaction of LDL with human arterial proteoglycans stimulates its uptake by human monocyte-derived macrophages. *J Lipid Res*. 1990;31:443-454.

Hurt-Camejo E, Camejo G, Rosengren B, López F, Wiklund O, Bondjers G. Differential uptake of proteoglycan-selected subfractions of low density lipoprotein by human macrophages. *J Lipid Res.* 1990;31:1387-1398.

Hurt-Camejo E, Camejo G, Rosengren E, López F, Ahlström C, Pager G, Bondjers G. Effect of arterial proteoglycans and glycosaminoglycans on low density lipoprotein oxidation and its uptake by human macrophages and arterial smooth muscle cells. *Arterioscler Thromb.* 1992;12:569-583.

^aHurt-Camejo E, Andersen S, Sandal R, Rosengren B, Sartipy P, Stadberg E, Johansen B. Localization of non pancreatic secretory phospholipase A2 in normal and atherosclerotic arteries: activity of the isolated enzyme on low density lipoproteins. *Arterioscler Thromb Vasc Biol.* 1997;17:300-309.

^bHurt-Camejo E, Olsson U, Wiklund O, Bondjers G, Camejo G. Cellular consequences of the association of apoB lipoproteins with proteoglycans. *Arterioscler Thromb Vasc Biol.* 1997;17:1011-1017.

Hussaini IM, Sri Kumar K, Quesenberry PJ, Gonias SL. Colony stimulating factor-1 modulates α -macroglobulin receptor expression in murine bone marrow macrophages. *J Biol Chem.* 1990;265:19441-19446.

Hussaini IM, LaMarre J, Lysiak JJ, Karns LR, VandenBerg SR, Gonias SL. Transcriptional regulation of LDL receptor-related protein by IFN- γ and the antagonistic activity of TGF- β_1 IN THE raw 264.7 macrophage-like cell line. *J Leukocyte Biol.* 1996;59:733-739.

I

Iozzo RV, Cohen IR, Grassel S et al The biology of perlecan. *Biochem J.* 1994;302:625-639.

Iozzo RV. The biology of the small leucine-rich proteoglycans. Functional network of interactive proteins. *J Biol Chem.* 1999;274:18843-18846.

Ip JH, Fuster V, Badimon L, Badimon JJ, Taubman MB, Chesebro JH. Syndromes of accelerated atherosclerosis: role of vascular injury and smooth muscle cell proliferation. *J Am Coll Cardiol.* 1990;15:1667-1687.

J

Jang Y, Lincoff AM, Plow EF, Topol EJ. Cell adhesion molecules in coronary artery disease. *J Am Coll Cardiol.* 1994;24:1591-1601.

Jerome WG, Yancey PG. The role of microscopy in understanding atherosclerotic lysosomal lipid metabolism. *Microsc Microanal.* 2003;9:54-67.

Ji ZS, Brecht WJ, Miranda RD, Hussain MM, Innerarity TL, Mahley RW. Role of heparan sulfate proteoglycans in the binding and uptake of apolipoprotein E-enriched remnant lipoproteins by cultured cells. *J Biol Chem.* 1993;268:10160-10167.

Jimi S, Sakata N, Matunaga A, Takebayashi S. Low density lipoproteins bind more to type I and III collagen by negative-charge dependent mechanisms than to type IV and V collagens. *Atherosclerosis.* 1994;107:109-116.

Jimi S, Saku K, Uesugi N, Sakata N, Takebayashi S. Oxidized low density lipoprotein stimulates collagen production in cultured arterial smooth muscle cells. *Atherosclerosis.* 1995;116:15-26.

Joseph J, Ranganathan S, Mehta JL. Low density lipoproteins modulate collagen metabolism in fibroblasts. *J Cardiovasc Pharmacol Ther.* 2003;8:161-6.

Jukema JW, Bruschke AV, van Boven AJ, Reiber JH, Bat ET, Zwinderman AH, Jansen H, Boerma GJ, van Rappard FM, Lie KI, et al. Effects of lipid lowering by pravastatin on progression and regression of coronary artery disease in symptomatic men with normal to moderately elevated serum cholesterol levels. The Regression Growth Evaluation Statin Study (REGRESS). *Circulation.* 1995;91:2528-2540.

Jurgens G, Hoff HF, Chisolm GM, Esterbauer H. Modification of human serum low density lipoprotein by oxidation-characterization and pathophysiological implications. *Chem Phys Lipids.* 1987;45:315-336.

K

Kalela A, Koivu TA, Hoyhtya M, Jaakkola O, Lehtimaki T, Sillanaukee P, Nikkari ST. Association of serum MMP-9 with autoantibodies against oxidized LDL. *Atherosclerosis.* 2002;160:161-165.

Khoo JC, Miller E, Loughlin Mc, Steinberg D. Enhanced macrophage uptake of low density lipoprotein after self-aggregation. *Arteriosclerosis.* 1998;8:348-358.

Kodama T, Reddy P, Kishimoto C, Krieger M. Purification and characterization of a bovine acetyl low density lipoprotein receptor. *Proc Natl Acad Sci USA.* 1988;85:9238-9242.

Komminoth P, Merk FB, Leav I, Wolfe HJ, Roth J. Comparison of 35S- and digoxigenin-labeled RNA and oligonucleotide probes for in situ hybridization. Expression of mRNA of the seminal vesicle secretion protein II and androgen receptor genes in the rat prostate. *Histochemistry.* 1992;98:217-228.

Kounnas MZ, Chappell DA, Wong H, Argraves WS, Strickland DK. The cellular internalization and degradation of hepatic lipase is mediated by low density lipoprotein receptor-related protein and requires cell surface proteoglycans. *J Biol Chem.* 1995;270:9307-9312.

Kowal RC, Herz J, Goldstein JL, Esser V, Brown MS. Low density lipoprotein receptor-related protein mediates uptake of cholesteryl esters derived from apoprotein E-enriched lipoproteins. PNAS. 1989;86:5810-5814.

Kristensen T, Moestrup SK, Gliemann J, Bendtsen L, Sand O, Sottrup-Jensen L. Evidence that the newly cloned low-density-lipoprotein receptor related protein (LRP) is the alpha 2-macroglobulin receptor. FEBS Lett. 1990;276:151-155.

Kubo N, Kikuchi J, Furukawa Y, Sakai T, Ohta H, Iwase S, Yamada H, Sakurabayashi I. Regulatory effects of aggregated LDL on apoptosis during foam cell formation of human peripheral blood monocytes. FEBS letters. 1997;409:177-182.

Kullo IJ, Edwards ED, Schwartz RS. Vulnerable plaque: pathobiology and clinical implications. Ann Intern Med. 1998;129:1050-1060.

Kütt H, Herz J, Stanley KK. Structure of the low density lipoprotein receptor-related protein (LRP) promoter. Biochim Biophys Acta. 1989;1009:229-236.

L

Labat-Robert J, Gruber E, Bihari-Varga M. Interaction between fibronectin, proteoglycans and lipoproteins. Int J Biol Macromol. 1990;12:50-54.

Laemmli UK. Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. Nature. 1970;227:680-685.

Lafont A, Libby P. The smooth muscle cell: sinner or saint in restenosis and the acute coronary syndromes? J Am Coll Cardiol. 1998;32:283-285.

LaMarre J, Wolf BB, Kittler ELW, Quesenberry PJ, Gonias SL. Regulation of macrophage α_2 -macroglobulin/low density lipoprotein receptor-related protein by lipopolysaccharide and interferon- γ . J Clin Invest. 1993;91:1219-1224.

Langeler EG, Snelting-Havinga I, Van Hinsbergh VWM. Passage of low density lipoproteins through monolayers of human arterial endothelial cell: effects of vasoactive substances in an in vitro model. Arteriosclerosis. 1989;9:550-559.

Larsson LI. Immunocytochemistry: Theory and Practice. CRC Press, Inc. Boca Ratón, Florida. 1988.

Latron Y, Chautan M, Anfosso F, Alessi MX, Nalbone G, Lafont H, Juhan-Vague I. Stimulating effect of oxidized low density lipoprotein on plasminogen activator inhibitor-1 synthesis by endothelial cells. Arterioscler Thromb. 1991;11:1821-1829.

Laughton CW, Ruddle DL, Bedford CJ, Alderman EL. Sera containing elevated non-esterified fatty acids from patients with angiographically documented coronary atherosclerosis cause marked lipid accumulation in cultured human arterial smooth muscle-derived cells. Atherosclerosis. 1988;70:233-246.

Lemire JM, Braun KR, Maurel P, Kaplan ED, Schwartz SM, Wight TN. Versican/PG-M isoforms in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol.* 1999;19:1630-1639.

Leskinen MJ, Kovanen PT, Lindstedt KA. Regulation of smooth muscle cell growth, function and death in vitro by activated mast cells a potential mechanism weakening and rupture of atherosclerotic plaques. *Biochem Pharmacol.* 2003;66:1493-1498.

Li DY, Chen HJ, Staples ED, Ozaki K, Annex B, Singh BK, Vermani R, Mehta JL. Oxidized low-density lipoprotein receptor LOX-1 and apoptosis in human atherosclerotic lesions. *J Cardiovasc Pharmacol Ther.* 2002;7:147-153.

Libby P. Molecular bases of acute coronary syndromes. *Circulation.* 1995;91:2844-2850.

Libby P, Geng YJ, Aikawa M, Schoenbeck U, Mach F, Clinton SK, Sukhova GK, Lee RT. Macrophages and atherosclerotic plaque stability. *Curr Opin Lipidol.* 1996;7:330-335.

Libby P, Aikawa M. New insights into plaque stabilisation by lipid lowering. *Drugs.* 1998;56 Suppl1:9-13;discussion 33.

Libby P. Inflammation in atherosclerosis. *Nature.* 2002;420:868-874.

Lindahl U, Lindholt K, Spillman D et al. More to heparin than anticoagulation. *Thrombosis Res.* 1994;75:1-32.

Lipid Research Clinic Program. Manual of Laboratory Operation. Washington, DC: US Government Printing Office. 1974.DHEW publication No.NIH75-628.

Llorente-Cortés V, Martínez-González J, Badimon L. LDL receptor-related protein mediates uptake of aggregated LDL in human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol.* 2000;20:1572-1579.

Löhler J, Timpl R, Jaenisch R. Embryonic lethal mutation in mouse collagen I gene causes rupture of blood vessels and is associated with erythropoietic and mesenchymal cell death. *Cell.* 1984;38:597-607.

Lougheed M, Steinbrecher UP. Mechanism of uptake of copper oxidized low density lipoprotein in macrophages is dependent on its extent of oxidation. *J Biol Chem.* 1996;271:11798-11805.

Loukinova E, Ranganathan S, Kuznetsov S, Gorlatova N, Migliorini MM; Loukinov D, Uery PG, Mikhailenko I, Lawrence DA, Strickland DK. Platelet-derived growth factor (PDGF)-induced tyrosine phosphorylation of the low density lipoprotein receptor-related protein (LRP). *J Biol Chem.* 2002;277:15499-15506.

Lowry O, Rosenbrough N, Farr A, et al. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951;193:265-275.

Lunec J. Free radical activity: Methods of measurement in vivo. Pharmacological methods in the control on inflammation. Modern methods in pharmacology. Joseph Y, Chang y Alan J Lewis eds. New York. 1989;5:59-81.

Luoma J, Hiltunen T, Särkioja T, Moestrup SK, Gliemann J, Kodama T, Nikkari T, Ylä-Herttula S. Expression of α_2 -macroglobulin receptor/low density lipoprotein receptor-related protein and scavenger receptor in human atherosclerotic lesions. *J Clin Invest.* 1994;93:2014-2021.

Lupu F, Heim D, Bachmann F, Kruithof EK. Expression of LDL receptor-related protein/alpha 2-macroglobulin receptor in human normal and atherosclerotic arteries. *Arteriosclerosis Thrombosis.* 1994;14:1438-1444.

Lutz C, Nimpf J, Jenny M, Boecklinger K, Enzinger C, Utermann G, Baier-Bitterlich G, Baier G. Evidence of functional modulation of the MEKK/JNK/cJun signaling cascade by the low density lipoprotein receptor-related protein (LRP). *J Biol Chem.* 2002;277:43143-43151.

M

Mahley RW, Ji ZS, Brecht WJ, Miranda RD, He D. Role of heparan sulfate proteoglycans and the LDL receptor-related protein in remnant lipoprotein metabolism. *Ann N Y Acad Sci.* 1994;737:39-52.

Mao SJ, Yates T, Rechtin E, Jackson RL, van Sickle WA. Antioxidant activity of probucol and its analogues in hypercholesterolemic Watanabe rabbits. *J Med Chem.* 1991;34:298-302.

Maor I, Aviram M. Macrophage released proteoglycans are involved in cell-mediated aggregation of LDL. *Atherosclerosis.* 1999;142:57-66.

Martínez-González J, Llorente-Cortés V, Badimon L. Biología celular y molecular de la lesiones ateroscleróticas. *Rev Esp Cardiol.* 2001;54:218-231.

Mayne R, Brewton RG. New members of collagen superfamily. *Curr Opin Cell Biol.* 1993;5:883-890.

Mehta JL, Li DY. Identification and autoregulation of receptor for ox-LDL in cultured human coronary artery endothelial cells. *Biochem Biophys Res Commun.* 1998;248:511-514.

Meilinger M, Haumer M, Swakamary KA, Steinböck B, Goldenberg H, Huettlinger M. Removal of lactoferrin from plasma is mediated by binding to low density lipoprotein receptor-related protein/ α_2 -macroglobulin receptor and transport to endosomes. *FEBS.* 1995;360:70-74.

Mikhailenko I, Kounnas MZ, Strickland DK. Low density lipoprotein receptor-related protein/alpha 2-macroglobulin receptor mediates the cellular internalization and degradation of thrombospondin. A process facilitated by cell-surface proteoglycans. *J Biol Chem.* 1995;270:9543-9549.

^aMinami M, Kume N, Shimaoka T, Kataika H, Hayashida K, Akiyama Y, Nagata I, Ando K, Nobuyoshi M, Hanyuu M, Komeda M, Yonehara S, Kita T. Expression of SR-PSOX, a novel cell-surface scavenger receptor for phosphatidylserine and oxidized LDL in human atherosclerotic lesions. *Arterioscler Thromb Vasc Biol*. 2001;21:1796-1800.

^bMinami M, Kume N, Shimaoka T, Kataoka H, Hayashida K, Yonehara S, Kita T. Expression of scavenger redceptor for phosphatidylserine and oxidized lipoprotein (SR-PSOX) in human atheroma. *Ann N Y Acad Sci*. 2001;947:373-6.

Misra UK, Gawdi G, González-Gronow M, Pizzo SV. Coordinate regulation of the α_2 -macroglobulin signaling receptor and the low density lipoprotein receptor-related protein/ α_2 -macroglobulin receptor by insulin. *J Biol Chem*. 1999;274:25785-25791.

Mize RR. Quantitative image analysis for immunocytochemistry and in situ hibridization. *J Neurosci Methods*. 1994;54:219-237.

Moestrup SK, Gliemann J, Pallesen G. Distribution of α_2 -macroglobulin receptor/low density lipoprotein receptor-related protein in human tissues. *Cell Tissue Res*. 1992;269:375-382.

Moreno MJ, Falk E, Palacios IF, Newell JB, Fuster V, Fallon JT. Macrophage infiltration in acute coronary syndromes: implications for plaque rupture. *Circulation*. 1994;90:775-778.

Mulder M, Lombardi P, Jansen H, van Berkel TJC, Frants RR, Havekes LM. Low density lipoprotein receptor internalizes low density and very low density lipoproteins that are bound to heparan sulfate proteoglycans via lipoprotein lipase. *J Biol Chem*. 1993;268:9369-9375.

Multhaup HA, Gafvels ME, Kariko K, Jin H, Arenas-Elliott C, Goldman BI, Strauss JF 3rd, Angelin B, Warhol MJ, McCrae KR. Expression of very low density lipoprotein receptor in the vascular wall: analysis of human tissues by in situ hybridization and immunochemistry. *Am J Pathol*. 1996;148:1985-1997.

N

Navab M, Imes SS, Hama Sy. Monocyte transmigration induced by modification of low density lipoprotein in cocultures of human aortic wall cells is due to induction of monocyte chemotactic protein 1 synthesis and is abolished by high density lipoprotein. *J Clin Invest*. 1991;88:2039-2046.

Nermine A, Ismail E, Alavi M.Z., Moore S. Lipoprotein-proteoglycan complexes from injured rabbit aortas accelerate lipoprotein uptake by arterial smooth muscle cells. *Atherosclerosis*. 1994;105:79-87.

Nichols AV, Krauss RM, Musliner TA. Non-denaturing polyacrylamide gradient gel electrophoresis. In: Segrest JP, Albers JJ, eds. *Methods in Enzymology: Plasma Lipoproteins*. New York, NY: Academic Press. 1986;128:417-31.

Nigon F, Lesnik P, Rouis M, Chapman MJ. Discrete subspecies of human low density lipoproteins are heterogeneous in their interaction with the cellular LDL receptor. *J Lipid Res.* 1991;32:1741-1753.

Nordestgaard BG, Nielsen LB. Atherosclerosis and arterial influx of lipoproteins. *Curr Opin Lipidol.* 1994;5:252-257.

Nykjaer A, Petersen CM, Moller B, Jensen PH, Moestrup SK, Holtet TL, Etzerodt M, Thogersen HC, Munch M, Andreasen PA, et al. Purified alpha 2-macroglobulin receptor/LDL receptor-related protein binds urokinase, plasminogen activator inhibitor type-1 complex. Evidence that the alpha 2-macroglobulin receptor mediates cellular degradation of urokinase receptor-bound complexes. *J Biol Chem.* 1992;267:14543-14546.

O

Obunike JC, Edwards IJ, Rumsey SC, Curtiss LK, Wagner WD, Deckelbaum RJ, Goldberg IJ. Cellular differences in lipoprotein lipase-mediated uptake of low density lipoproteins. *J Biol Chem.* 1994;269:13129-13135.

Olin KL, Potter-Perigo S, Barrett PH, Wight TN, Chait A. Lipoprotein lipase enhances the binding of native and oxidized low density lipoproteins to versican and biglycan synthesized by cultured arterial smooth muscle cells. *J. Biol. Chem.* 1999;274:34629-34636.

Olsson U, Bondjers G, Camejo G. Fatty acids modulate the composition of extracellular matrix in cultured human arterial smooth muscle cells by altering the expression of genes for proteoglycan core proteins. *Diabetes.* 1999;48:616-622.

Öörni K, Pentikäinen MO, Ala-Korpela M, Kovanen PT. Aggregation, fusion, and vesicle formation of modified low density lipoprotein particles: molecular mechanisms and effects on matrix interactions. *J Lipid Res.* 2000;41:1703-1714.

Ooshima A. Collagen β chain: Increased proportion in human atherosclerotic. *Science.* 1981;213:666-668.

Orekhov AN, Tertov VV, Mukhin DN. Desialylated low density lipoprotein-naturally occurring lipoprotein with atherogenic potency. *Atherosclerosis.* 1991;86:153-158.

Orth K, Madison EL, Gething MJ, Sambrook JF, Herw J. Complexes of tissue-type plasminogen activator and its serpin inhibitor plasminogen-activator inhibitor type 1 are internalized by means of the low density lipoprotein receptor-related protein/alpha 2-macroglobulin receptor. *PNAS.* 1992;89:7422-7426.

Osborne TF, Goldstein JL, Brown MS. 5' end of HMG-Coa reductase gene contains sequences responsible for cholesterol mediated inhibition of transcription. *Cell.* 1985;42:203-212.

P

Pedreño J, de Castellarnau C, Cullaré C, Ortín R, Sánchez JL, Llopard R, González-Sastre F. Platelet LDL receptor binds with the same apparent affinity both oxidized and native LDL. Evidence that receptor ligand complexes are not internalized. *Arterioscler Thromb*. 1994;14:402-408.

Peet DJ, Turley SD, Ma W, Ma W, Janowski BA, Lobaccaro JM, Hammer RE, Mongeldorf DJ. Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR α . *Cell*. 1998;93:693-704.

Pentikäinen MO, Lehtonen EMP, Kovanen PT. Aggregation and fusion of modified low density lipoprotein. *J Lipid Res*. 1996;17:2638-2649.

Pentikäinen MO, Lehtonen EMP, Öörni K, Lusa S, Somerharju P, Jauhainen M, Kovanen PT. Human arterial proteoglycans increase the rate of proteolytic fusion of low density lipoprotein particles. *J Biol Chem*. 1997;272:25283-25288.

Pillarisetti S, Obunike JC, Goldberg IJ. Lysolecithin induced alterations of subendothelial heparan sulfate proteoglycans increases monocyte binding to matrix. *J Biol Chem*. 1995;270:29760-29765.

Pillarisetti S, Paka S, Obunike J, Berglund L, Goldberg IJ. Subendothelial retention of lipoprotein (a). Evidence that reduced heparan sulfate promotes lipoprotein (a) retention by subendothelial matrix. *J Clin Invest*. 1997;100:867-874.

Pillarisetti S. Lipoprotein modulation of subendothelial heparan sulfate proteoglycans (perlecà) and atherogenicity. *Trends Cardiovasc Med*. 2000;10:60-65.

Pitas RE. Expression of the acetyl low density lipoprotein receptor by rabbit fibroblasts and smooth muscle cells. Up-regulation by phorbol esters. *J Biol Chem*. 1990;265:12722-12727.

Pitt B, Mancini GB, Ellis SG, Rosman HS, Park JS, McGovern ME. Pravastatin limitation of atherosclerosis in the coronary arteries (PLAC I): reduction in atherosclerosis progression and clinical events. PLAC I investigation. *J Am Coll Cardiol*. 1995;26:1133-1139.

Prat WK. Digital image processing. Second edition. A Wiley-Interscience publication. John Wiley & Sons, Inc 1991.

Pritchard KA Jr, Groszek L, Smalley DM, Sessa WC, Wu M, Villalon P, Wolin MS, Stemerman MB. Native low-density lipoprotein increases endothelial cell nitric oxide synthase generation of superoxide anion. *Circ Res*. 1995;77:510-518.

Prockop DJ, Kivirikko KI. Collagens: molecular biology, diseases, and potentials for therapy. *Annu Rev Biochem*. 1995;64:403-434.

Proctor SD, Vine DF, Mamo JCL. Arterial retention of apolipoprotein B₄₈- and B₁₀₀-containing lipoproteins in atherosclerosis. *Curr Opin Lipidol*. 2002;13:461-470.

R

Radhakrishnamurthy B, Srinivasan S, Vijayagopal P, Berenson G. Arterial wall proteoglycans: biological properties related to pathogenesis of atherosclerosis. *Eur Heart J.* 1990;11(Suppl E):148-157.

Ramprasad MP, Terpstra V, Kondratenko N, Quehenberger O, Steinberg D. Cell surface expression of mouse macrosialin and human CD68 and their role as macrophage receptors for oxidized low density lipoprotein. *Proc Natl Acad Sci USA.* 1996;93:14833-14838.

Rangaswamy S, Penn MS, Saidel GM, Chisolm GM. Exogenous oxidized low-density lipoprotein injures and alters the barrier function of endothelium in rats *in vivo*. *Circ Res.* 1997;80:37-44.

Rauterburg J, Jaeger E, Althau M. Collagens in atherosclerotic vessel wall lesion. *Curr Top Pathol.* 1993;87:163-192.

Reaven GM, Chen YK, Jeppesen J, Maheux P, Krauss RM. Insulin resistance and hyperinsulinemia in individuals with small, dense low density lipoprotein particles. *J Clin Invest.* 1993;92:141-146.

Reblin T, Niemeier A, Meyer N, Willnow TE, Kronenberg F, Dieplinger H, Greten H, Beieiegel U. Cellular uptake of lipoprotein (a) by mouse embryonic fibroblasts via the LDL receptor and the LDL receptor-related protein. *J Lipid Res.* 1997;38:2103-2110.

Rekhter MD. Collagen synthesis in atherosclerosis: too much and not enough. *Cardiov Res.* 1999;41:376-384.

Riessen R, Isner JM, Blessing E, Loushin C, Nikol S, Wight TN. Regional differences in the distribution of the proteoglycans biglycan and decorin in the extracellular matrix of atherosclerotic and restenotic human coronary arteries. *Am J Pathol.* 1994;144:962-974.

Rigotti A, Trigatti B, Babitt J, Penman M, Xu S, Krieger M. Scavenger receptor BI—a cell surface receptor for high density lipoprotein. *Curr Opin Lipidol.* 1997;8:181-8.

Rodríguez C, Martínez-González J, Sánchez-Gómez S, Badimon L. LDL down-regulates CYP51 in porcine vascular endothelial cells and in the arterial wall through a sterol regulatory element binding protein-2-dependent mechanism. *Circ Res.* 2001;88:268-274.

Rohlmann A, Gotthardt M, Hammer RE, Herz J. Inducible inactivation of hepatic LRP gene by cre-mediated recombination confirms role of LRP in clearance of chylomicron remnant. *J Clin Invest.* 1998;101:689-695.

Ross R, Wight TN, Strandness E, Thiele B. Human atherosclerosis I. Cell constitution and characteristics of advanced lesions of the superficial femoral artery. *Am J Pathol.* 1984;114:79-93.

Ross R. The pathogenesis of atherosclerosis – an update. *N Engl J Med.* 1986;314:488-500.

Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature.* 1993;362:801-808.

Ross R. Cell biology and atherosclerosis. *Annu Rev Physiol.* 1995;57:791-804.

Ross R. Atherosclerosis—an inflammatory disease. *The New England Journal of Medicine.* 1999;340:115-126.

Rumsey SC, Galeano NF, Lipschitz B, Deckelbaum RJ. Oleate and other long chain fatty acids stimulate low density lipoprotein receptor activity by enhancing acyl coenzyme A: cholesterol acyltransferase activity and altering intracellular regulatory cholesterol pools in cultured cells. *J Biol Chem.* 1995;270:10008-10016.

S

Sakai J, Nohturfft A, Stanley KK. The 5'-flanking region of the α_2 MR/LRP genes contains an enhancer-like cluster of Sp1 binding sites. *Biochim Biophys Acta.* 1994;1219:307-313.

Sakai J, Nohturfft A, Goldstein JL, Brown MS. Cleavage of sterol regulatory element-binding proteins (SREBPs) at site-1 requires interaction with SREBP cleavage-activating protein: evidence from in vivo competition studies. *J Biol Chem.* 1998;273:5785-5793.

Sarafanov AG, Ananyeva NM, Shima M, Saenko EL. Cell surface heparan sulfate proteoglycans participate in factor VIII catabolism mediated by low density lipoprotein receptor-related protein. *J Biol Chem.* 2001;276:11970-11979.

Sartipy P, Johansen B, Camejo G, Rosengren B, Bondjers G, Hurt-Camejo E. Binding of human phospholipase A₂ type II to proteoglycans. *J Biol Chem.* 1996;271:26307-26314.

Sata M, Walsh K. Oxidized LDL activates Fas-mediated endothelial cell apoptosis. *J Clin Invest.* 1991;102:1682-1689.

Sato R, Miyamoto W, Inoue J, Terada T, Imanaka T, Maeda M. Sterol regulatory element-binding protein negatively regulates microsomal triglyceride transfer protein gene transcription. *J Biol Chem.* 1999;274:24714-24720.

Sawamura T, Kume N, Aoyama T, Moriwaki H, Hoshikawa H, Aiba Y, Tanaka T, Miwa S, Katsura Y, Kita T, Mosaki T. An endothelial receptor for oxidized low density lipoprotein. *Nature.* 1997;386:73-7.

Schissel SL, William KJ, Tabas I. Sphingomyelinase (Smase)-induced LDL aggregation: mechanism and relevance. *Circulation.* 1994;90(pt 2):I-403. Abstract.

Schönherr E, Järveläinen HT, Kinsella MG, Sandell LJ, Wight TN. Platelet derived growth factor and transforming growth factor- β 1 differentially affect the synthesis of hyaluronic acid and decorin by monkey arterial smooth muscle cells. *Arterioscler Thromb*. 1993;13:1026-1036.

Schönherr E, Kinsella MG, Wight TN. Genistein selectively inhibits platelet-derived growth factor-stimulated versican biosynthesis in monkey arterial smooth muscle cells. *Arch Biochem Biophys*. 1997;339:353-61.

^aSchwenke DC, Carew TE. Initiation of atherosclerotic lesions in cholesterol-fed rabbits, I: focal increases in arterial LDL concentration precede development of fatty streak lesions. *Arteriosclerosis*. 1989;9:895-907.

^bSchwenke DC, Carew TE. Initiation of atherosclerotic lesions in cholesterol-fed rabbits, II: selective retention of LDL vs. selective increases in LDL permeability in susceptible sites of arteries. *Arteriosclerosis*. 1989;9:908-918.

Sehayek E, Wang XX, Vlodavsky I, Avner R, Levkovitz H, Ilievcrona T, Olivecrona G, Willnow TE, Herz J, Eisenberg S. Heparan sulfate-dependent and low density lipoprotein receptor-related protein independent catabolic pathway for lipoprotein lipase in mouse embryonic fibroblasts. *Isr J Med Sci*. 1996;32:449-454.

Seo T, Clair RW. Heparan sulfate proteoglycans mediate internalization and degradation of β -VLDL and promote cholesterol accumulation by pigeon macrophages. *J Lipid Res*. 1997;38:765-779.

Severs NJ, Shotton DM. Rapid freezing, freeze fracture, and deep etching. Eds. Wiley-Liss. 1995.

Shah PK, Falk E, Badimon JJ, Fernández-Ortiz A, Maihac A, Villareal-Levy G, Fallon JT, Regnstrom J, Fuster V. Human monocyte-derived macrophages induce collagen breakdown in fibrous caps of atherosclerotic plaques: potential role of matrix-degrading metalloproteinases and implications for plaque rupture. *Circulation*. 1995;92:1565-1569.

Skalén K, Gustafsson M, Rydberg EK, Hultén LM, Wiklund O, Innerarity TL, Borén J. Subendothelial retention of atherogenic lipoproteins in early atherosclerosis. *Nature*, 2002;417:750-754.

Simionescu N, Sima A, Dobrian A, Tirziu D, Simionescu M. Pathobiological changes of the arterial wall at the inception of atherosclerosis. *Curr Top Pathol*. 1993;87:1-45.

Sparrow CP, Olszewski J. Cellular oxidation of low density lipoprotein is caused by thiol production in media containing transition metal ions. *J Lipid Res*. 1993;34:1219-1228.

Srinivasan SR, Radhakrishnamurthy B, Vijayagopal P, Berenson GS. Proteoglycans, lipoproteins, and atherosclerosis. *Adv Exp Med Biol*. 1991;285:373-381.

Stanton LW, White RT, Bryant CM, Protter AA, Endemann G. A macrophage Fc receptor for IgG is also a receptor for oxidized low density lipoprotein. *J Biol Chem*. 1992;267:22446-22451.

Stara HC, Chandler AB, Glagov S, Guyton JR, Insull W Jr, Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, Wissler RW. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Arterioscler Thromb*. 1994;14:840-856.

Stary HC, Chandler AB, Glagov S, Guyton JR, Insull W Jr, Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, Wissler RW. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation*. 1994;89:2462-2478.

St. Clair R. Pathogenesis of atherosclerosis. *Cardiol Rev*. 1997;5:14-24.

Steinberg D. Lipoproteins and atherosclerosis: a look back and a look ahead. *Arteriosclerosis*. 1983;3:283-301.

Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witzum JL. Beyond cholesterol. Modifications of low density lipoproteins that increases its atherogenicity. *N Engl J Med*. 1989;320:915-924.

Steinbrecher UP. Role of superoxide in endothelial cell modification of low density lipoprotein. *Biochim Biophys Acta*. 1988;959:20-30.

Strickland DK, Ashcom JD, Williams S et al. Sequence identity between the alpha 2-macroglobulin receptor and low density lipoprotein receptor-related protein suggests that this molecule is a multifunctional receptor. *J Biol Chem*. 1990;265:17401-17404.

T

Tabas I, Li Y, Brocia RW, Xu SW, Swenson TL, William KJ. Lipoprotein lipase and sphingomyelinase synergistically enhance the association of atherogenic lipoproteins with smooth muscle cells and extracellular matrix: a possible mechanism for low density lipoprotein and lipoprotein (a) retention and macrophage foam cell formation. *J Biol Chem*. 1993;268:20419-20432.

Tertov VV, Sobenin IA, Gabbasov ZA, Popov EG, Orekhov AN. Lipoprotein aggregation as an essential step of intracellular lipid accumulation caused by modified low density lipoproteins. *Biochem Biophys Res Commun*. 1989;163:489-494.

Tertov VV, Orekhov AN, Sobenin JA, Gabbasov ZA, Popov EG, Yaroslavov AA, Smirnov VN. Three types of naturally occurring modified lipoproteins induce intracellular lipid accumulation due to lipoprotein aggregation. *Circ Res*. 1992;71:218-228.

Tertov VV, Orekhov AN, Sobenin JA, Morrisett JD, Gotto AM, Guevarra JG. Carbohydrate composition of protein and lipid components in sialic-rich and-poor low density lipoproteins from subjects with and without coronary artery disease. *J Lipid Res*. 1993;34:365-75.

Thyberg J. Differentiated properties and proliferation of arterial smooth muscle cells in culture. *Int Rev Cyt*. 1996;169:183-265.

Tirziu D, Jinga VY, Serban G, Simionescu M. The effects of low density lipoproteins modified by incubation with chondroitin-6-sulfate on human aortic smooth muscle cells. *Atherosclerosis*. 1999;147:155-166.

Tsao PS, Buitrago R, Chan JR, Cooke JP. Fluid flow inhibits endothelial adhesiveness: nitric oxide and transcriptional regulation of VCAM-1. *Circulation*. 1996;94:1682-1689.

Tsukamoto K, Kinoshita M, Kojima K, Mikuni Y, Kudo M, Mori M, Fujita M, Horie E, Shimazu N, Teramoto T. Synergically increased expression of CD36, CLA-1 and CD68, but not of SR-A and LOX-1, with the progression to foam cells from macrophages. *J Atheroscler Thromb*. 2002;9:57-64.

Tunon J, Ruiz-Ortega M, Egido J. Regulation of matrix proteins and impact on vascular structure. *Curr Hypertens Rep*. 2000;2:106-113.

V

Varela O, Martínez-González J, Badimon L. The response of smooth muscle cells to α -thrombin depends on its arterial origin: comparison among different species. *Eur J Clin Invest*. 1998;28:313-323.

Vidal F, Colomé C, Martínez-González J, Badimon L. Atherogenic concentrations of native low density lipoproteins modulates nitric oxide synthase gene expression in endothelial cells. *Eur J Biochem*. 1998;252:378-384.

Vijayagopal P, Srinivasan SR, Jones KM, Radhakrishnamurthy B, Berenson GS. Metabolism of low-density lipoprotein-proteoglycan complex by macrophages: further evidence for a receptor pathway. *Biochim Biophys Acta*. 1988;960:210-219.

^a**Vijayagopal P**. Enhanced synthesis and accumulation of proteoglycans in cholesterol-enriched arterial smooth muscle cells. *Biochem J*. 1993;294:603-611.

^b**Vijayagopal P**, Srinivasan SR, Radhakrishnamurthy B, Berenson GS. Human monocyte-derived macrophages bind low-density-lipoprotein-proteoglycan complexes by a receptor different from the low-density-lipoprotein receptor- *Biochem J*. 1993;289:837-844.

^a**Vijayogopal P**, Figueroa JE, Fontenot JD, Glancy DL. Isolation and characterization of a proteoglycan variant from human aorta exhibiting a marked affinity for low density lipoprotein and demonstration of its enhanced expression in atherosclerotic plaques. *Atherosclerosis*. 1996;127:195-203.

^b**Vijayagopal P**, Figueroa JE, Guo Q, Fontenot JD, Tao Z. Marked alteration of proteoglycan metabolism in cholesterol-enriched human arterial smooth muscle cells. *Biochem J*. 1996;315(Pt3):995-1000.

Völker W, Schmidt A, Buddeke E. Cytochemical changes in a human arterial proteoglycan related to atherosclerosis. *Atherosclerosis*. 1989;77:117-130.

Vora DK, Fang Z, Parhami F, Fogelman A, Territo M, Berliner J. P-selectin induction by mmLDL and its expression in human atherosclerotic lesions. *Circulation*. 1994;90:1-83.

W

Wagner WD, Salisbury BG, Rowe HA. A proposed structure of chondroitin 6-sulfate proteoglycans of human normal and adjacent atherosclerotic plaque. *Arteriosclerosis*. 1988;6:407-417.

Warshawsky I, Herz J, Broze GJ, Schwartz AL. The low density lipoprotein receptor-related protein can function independently from heparan sulfate proteoglycans in tissue factor pathway inhibitor endocytosis. *J Biol Chem*. 1996;271:25873-25879.

Watanabe Y, Inaba T, Shimano H, Gotoda T, Yamamoto K, Mokuno H, Sato H, Yasaki Y, Yamada N. Induction of LDL receptor-related protein during the differentiation of monocyte-macrophages: possible involvement in the atherosclerosis process. *Arterioscler Thromb*. 1994;14:1000-1006.

Watson JD, Gilman M,, Witkowski J, Zoller M. Recombinant DNA. WH Freeman and Co. Eds. 1992.

Weis JR, Pites RE, Wilson BD, Rodgers GM. Oxidized low-density lipoprotein increases cultured human endothelial cell tissue factor activity and reduces protein C activation. *FASEB J*. 1991;5:2459-2465.

Weisgraber KH, Innerarity TL, Mahley RW. Role of lysine residues of plasma lipoproteins in high affinity binding to cell surface receptors on humanfibroblasts. *J Biol Chem*. 1978;253:9053-9062.

Wight TN. Cell biology of arterial proteoglycans. *Arteriosclerosis*. 1989;9:1-20.

Wight TN, Kinsella MG, Qwarnström. The role of proteoglycans in cell adhesion, migration and proliferation. *Curr Opin Cell Biol*. 1992;4:793-801.

Wight TN. The vascular extracellular matrix. In: Fuster V, Ross R, Topol E, editors. *Atherosclerosis and coronary artery disease*. Philadelphia: Lippincot-Raven. 1996;1:421-440.

Williams KJ, Fless GM, Petrie KA, Snyder ML, Brocia RW, Swenson TL. Mechanisms by which lipoprotein lipase alters cellular metabolism of lipoprotein (a), low density lipoprotein, and nascent lipoproteins. Roles for low density lipoprotein receptors and heparan sulfate proteoglycans. *J Biol Chem*. 1992;267:13284-13292.

Williams KJ, Tabas I. The response-to-retention hypothesis of early atherogenesis. *Arterioscler Thromb Vasc Biol*. 1995;15:551-561.

Williams KJ, Fukui IV. Cell-surface heparan sulfate proteoglycans: dynamic molecules mediating ligand catabolism. *Curr Opin Lipidol*. 1997;8:253-262.

Williams KJ. Arterial wall chondroitin sulfate proteoglycans: diverse molecules with distinct roles in lipoprotein retention and atherogenesis. *Curr Opin Lipidol.* 2001;12:477-487.

Wissler RW, Vesselinovitch D, Komatsu A. The contribution of studies of atherosclerotic lesions in young people to future research. *Ann NY Acad Sci.* 1990;598:418-434.

Wissler RW. Morphological characteristics of the developing atherosclerotic plaque: animal studies and studies of lesions from young people. *Atheroscler Rev.* 1991;23:91-103.

Witztum JL, Mahoney EM, Branks MJ, Fisher J, Elam R, Steinberg D. Nonenzymatic glucosylation of low density lipoprotein alters its biologic activity. *Diabetes.* 1982;31:283-91.

Witztum JL. Role of oxidized low density lipoprotein in atherogenesis. *Br Heart J.* 1993;69(suppl):S12-S18.

Wyne KL, Pathak RK, Seabra MC, Hobbs HH. Expression of the VLDL receptor in endothelial cells. *Arterioscler Thromb Vasc Biol.* 1996;16:407-415.

X

Xu C, Zarins CK, Pannaraj PS, Bassiouny HS, Glagov S. Hypercholesterolemia superimposed by experimental hypertension induces differential distribution of collagen and elastin. *Arterioscler Thromb Vasc Biol.* 2000;20:2566-2572.

Y

Yla-Herttula S, Jaakkola O, Solakivi T, Kuivaniemi H, Nikkari T. The effect of proteoglycan, collagen and lysyl oxidase on the metabolism of low density lipoprotein by macrophages. *Atherosclerosis.* 1986;62:73-80.

Yamada Y, Doi T, Hamakubo T, Kodama T. Scavenger receptor family proteins: roles for atherosclerosis, host defence and disorders of the central nervous system. *Cell Mol Life Sci.* 1998;54:628-640.

Z

Zeiher AM, Drexler H, Wollschlager H, Just H. Modulation of coronary vasomotor tone in humans. Progressive endothelial dysfunction with different early stages of coronary atherosclerosis. *Circulation.* 1991;83:391-401.

Zhong-Sheng J, Brecht WJ, Miranda RD, Hussain MM, Innerarity TL, Mahley RW. Role of heparan sulfate proteoglycans in the binding and uptake of apolipoprotein E-enriched remnant lipoproteins by cultured cells. *J Biol Chem.* 1993;268:10160-10167.

Zhong-Sheng J, Dichek HL, Miranda RD, Mahley RW. Heparan sulfate proteoglycans participate in hepatic lipase-and apolipoprotein E-mediated binding and uptake of plasma lipoproteins, including high density lipoproteins. *J Biol Chem.* 1997;272:31285-31292.

Zhong-Sheng J, Pitas RE, Mahley RW. Differential cellular accumulation/retention of apolipoprotein E mediated by cell surface heparan sulfate proteoglycans. *J Biol Chem.* 1998;273:1345213460.

