

RESEARCH PAPER

Carbon and nitrogen partitioning during the post-anthesis period is conditioned by N fertilisation and sink strength in three cereals

I. Aranjuelo^{1,2}, L. Cabrera-Bosquet², J. L. Araus² & S. Nogués²

¹ Instituto de Agrobiotecnología, Universidad Pública de Navarra-CSIC-Gobierno de Navarra, Mutilva Baja, Spain

² Unitat de Fisiologia Vegetal, Facultat de Biologia, Universitat de Barcelona, Barcelona, Spain

Keywords

¹³C and ¹⁵N isotopic labelling; grain filling; N use efficiency; N/C partitioning; wheat.

Correspondence

I. Aranjuelo, Instituto de Agrobiotecnología, Universidad Pública de Navarra-CSIC-Gobierno de Navarra, Campus de Arrosadía, E-31192-Mutilva Baja, Spain.
E-mail: iker.aranjuelo@unavarra.es

Editor

H. Rennenberg

Received: 21 November 2011; Accepted: 27 February 2012

doi:10.1111/j.1438-8677.2012.00593.x

ABSTRACT

Further knowledge of the processes conditioning nitrogen use efficiency (NUE) is of great relevance to crop productivity. The aim of this paper was characterise C and N partitioning during grain filling and their implications for NUE. Cereals such as bread wheat (*Triticum aestivum* L. cv Califa sur), triticale (× *Triticosecale* Wittmack cv. Imperioso) and tritordeum (× *Tritordeum* Asch. & Graebn line HT 621) were grown under low (LN, 5 mm NH₄NO₃) and high (HN, 15 mm NH₄NO₃) N conditions. We conducted simultaneous double labelling (¹²CO₂ and ¹⁵NH₄¹⁵NO₃) in order to characterise C and N partitioning during grain filling. Although triticale plants showed the largest total and ear dry matter values in HN conditions, the large investment in shoot and root biomass negatively affected ear NUE. Tritordeum was the only genotype that increased NUE in both N treatments (NUE_{total}), whereas in wheat, no significant effect was detected. N labelling revealed that N fertilisation during post-anthesis was more relevant for wheat and tritordeum grain filling than for triticale. The study also revealed that the investments of C and N in flag leaves and shoots, together with the 'waste' of photoassimilates in respiration, conditioned the NUE of plants, and especially under LN. These results suggest that C and N use by these plants needs to be improved in order to increase ear C and N sinks, especially under LN. It is also remarkable that even though tritordeum shows the largest increase in NUE, the low yield of this cereal limits its agronomic value.

INTRODUCTION

It is estimated that in the Mediterranean environment, which comprises 10% of the world's total wheat growing area, N availability is the main environmental factor, alongside drought stress, that limits productivity (Passioura 2002; Cossani *et al.* 2007). The fact that cereals require large amounts of N to increase maximum yield has led to a worldwide increase in fertiliser application from 1.3 million tonnes in 1930 to 85–90 million tonnes by 2004. Furthermore, it is predicted that by 2050, fertiliser application will reach 240 million tonnes (Masclaux-Daubresse *et al.* 2008). However, in developed economies, nitrogen use efficiency (NUE) is very low and estimated to be ca. 33% (Raun & Johnson 1999). The abuse of N fertilisers can lead to contamination of ground and surface water (Matson *et al.* 2002). To cope with this problem, European policies encourage an increase in the NUE of agriculture through crop management and plant breeding in order to reduce the excessive input of fertilisers, while maintaining acceptable yields (Lea & Azevedo 2006, 2007).

Improving wheat performance and NUE under low N availability conditions requires improved knowledge of yield determining factors (Masclaux-Daubresse *et al.* 2008, 2010). Grain filling, and by extension grain yield, are mainly sus-

tained by assimilation and management of C and N compounds. In wheat, grain N content is the result of the N taken up after anthesis, together with the amount of remobilised N originating from pre-anthesis uptake (Dupont & Altenbach 2003). On the other hand, C required for grain filling is mostly provided by flag leaf photosynthesis (Evans *et al.* 1975), translocation of C assimilated before anthesis (mainly stored in the internodes; Gebbing *et al.* 1999) and ear photosynthesis (Tambussi *et al.* 2007). Contrasting results concerning the origin of C and N contributing to grain filling are reported in the literature (Palta & Fillery 1995; Ercoli *et al.* 2008) because the relative contribution of those sources varies with growth conditions and the cereal studied (Dupont & Altenbach 2003; Aranjuelo *et al.* 2009). The relative contribution of remobilisation to grain yield depends mainly on C/N source/sink interaction during grain filling (Asseng & Van Herwaarden 2003). These authors observed that when crops are exposed to water stress during grain filling, pre-anthesis reserves of carbohydrate and N contribute to grain filling to a greater extent under high N status than under low N. Among other factors, grain filling has been described as being affected by the rate and timing of N application and on the level and form of soil N (Spiertz & De Vos 1983; Borghi *et al.* 1997). Furthermore, as observed by Papakosta & Gagianas (1991), high soil N availability during grain filling

favours post-anthesis N uptake and reduces the remobilisation of pre-anthesis N.

Improvements conducted in plant breeding programmes derived from the application of genetic studies appear to be reaching a plateau. Approaches that complement traditional breeding methodologies with new approaches for more efficient phenotyping, along with a deeper understanding of the metabolic mechanisms that modulate plant responses to environmental conditions, are a central goal of breeding programmes (Araus *et al.* 2008). In this context, the use of stable isotopes has provided key information for the identification of new traits to assess plant performance (Avice *et al.* 1996; Aranjuelo *et al.* 2009, 2011a,b; Masclaux-Daubresse & Chardon 2011). Previous studies conducted in wheat have revealed that the 'classical' assumptions concerning C partitioning during grain filling were, in some cases, not totally accurate. The low number of studies analysing the loss of photoassimilates by respiration during wheat grain filling underscores the relevancy of this process, a topic that has been observed to require a large amount of carbohydrate (Aranjuelo *et al.* 2009, 2011b). These studies on canopy-level C labelling conducted with wheat plants revealed that 2 weeks after anthesis, nearly half of the CO₂ respired by the plants came from C fixed in the previous week. These studies also revealed that at the same phenological stage, 21% of C present in root total organic matter (TOM) was labelled (*i.e.* C assimilated in the previous 7 days). Overall, these results suggest that a substantial part of the C assimilated during grain filling effectively does not contribute to this important process. It appears reasonable to speculate that this may be due to low sink strength.

Traditional breeding strategies are limited by the low genetic variability of yield components under stress conditions (Cook 1999). Plant genetic resources for agriculture might be one of the biological bases of future agriculture. *Tritordeum* is a doubled haploid hexaploid (\times *Tritordeum* Asch. & Graebn) that resulted from an interspecific cross between the diploid *Hordeum chilensis* and the tetraploid wheat, *Triticum turgidum* L. ssp. *durum*. *Tritordeum* is considered as a new crop having great potential (Martin 1988) as well as a bridge species for transferring useful barley traits, *e.g.* adaptation to Mediterranean conditions, to wheat (Martin *et al.* 1999; Ballesteros *et al.* 2005). Studies carried out in *2* *tritordeum* (Barro *et al.* 1991, 1996) have highlighted an increased capacity for N absorption. Maximising the NUE of N fertilisers is a matter of great concern, especially in the context of sustainable agriculture.

Ensuring an adequate C and N balance in the sink and source tissues is an essential target point for maximising the response of cereals to growth under low N availability conditions. The main goal of this paper was to determine the C and N partitioning during grain filling and its implications in the NUE of two contrasting genotypes (wheat and triticale) together with a new crop (*tritordeum*) with potential interest for introgression (*i.e.* bridging traits) in wheat breeding programmes for Mediterranean conditions. Moreover, these three cereals show a clear range in harvest index (the ratio of grain weight to total shoot biomass), with bread wheat and *tritordeum* exhibiting the highest and the lowest values, respectively (Villegas *et al.* 2010). Therefore, the three cereals show different constitutive (*i.e.* in the absence of any stress)

sink strengths. ¹²CO₂ labelling was conducted at the whole plant level to better understand C partitioning of these plants during the post-anthesis period.

Preliminary studies carried out previously in *tritordeum* plants suggested that they have higher NUE, which means that their performance should be better in low N conditions than other cereals. This hypothesis was tested in low N and compared with high yielding cultivars. The sampling and subsequent analyses of respired CO₂ isotopic composition through GC-C-IRMS enabled the characterisation of labelled C invested in respiration processes. Respiration represents an important C sink that has not been considered in classical approaches studying the contribution of C assimilated during pre/post-anthesis in grain filling. In parallel to C labelling, N labelling was applied to the plants in a ¹⁵NH₄¹⁵NO₃ enriched solution in order to test N uptake and partitioning to different plant organs.

MATERIAL AND METHODS

Experimental design

Seeds of bread wheat (*Triticum aestivum* L. cv. Califa Sur) and its two amphiploids, triticale (\times *Triticosecale* Wittm. cv. Imperioso) and *tritordeum* (\times *Tritordeum* Asch. & Graeb, HT 621) were sown (eight plants per pot) in 4-l pots filled with sand. Being amphidiploids, the latter two genotypes contain the diploid chromosome sets of both parents. Triticale is the result of a cross between wheat (*Triticum turgidum*) and rye (*Secale cereale*), whereas *tritordeum* is a fertile amphiploid between a wild barley (*Hordeum chilense*) and a durum wheat (*Triticum turgidum*). The plants were grown in two controlled environment chambers (Conviron E15; Controlled Environments Ltd, Winnipeg, MB, Canada) throughout the experiment. Inside the growth chambers, the growth conditions were 400 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ photosynthetic photon flux density (PPFD), 70% relative humidity, 25/15 °C day/night temperature and a 14-h photoperiod. The air of each module was sampled every 5 min using an industrial programmer (PLS C20K; Omron, Kyoto, Japan) and the CO₂ concentration was analysed through an infrared gas analyser (IRGA; Lira 3600; MSA, Barcelona, Spain). Air probes were placed at the centre of each module. The data were continuously recorded with a computer through analogue-digital converters (Microlink 751; Biodata Ltd, Manchester, UK) and Windmill software with a TEST-SEQ programming tool (Biodata Ltd).

The plants were watered with a modified Hoagland nutrient solution (Arnon & Hoagland 1939). A random selection of half of the pots was watered with 5 mM NH₄NO₃ [low N treatment (LN)] and the other half with 15 mM NH₄NO₃ [high N treatment (HN)]. The two levels of N were selected according to observations of previous studies conducted by our group with wheat (Lopes *et al.* 2006; Cabrera-Bosquet *et al.* 2009) and barley (Yousfi *et al.* 2010) plants exposed to different nutritional levels.

For each species we carried out double (C and N) labelling over 1 week, starting 1 week after anthesis (defined when at least 50% of the spikes in a pot showed protruding anthers). This period was selected because growth conditions between 20 days pre- and 10 post-anthesis are considered a critical

period for grain yield (Schnyder 1992). The labelling was conducted in a random selection of half of the pot plants that were placed in separated growth chambers. The pots were rotated regularly within the corresponding chamber to avoid chamber effects. All the determinations were conducted in samples collected immediately after the end of the labelling period.

N and C isotope labelling procedures

One week after anthesis, coinciding with the period of largest photoassimilate contribution to grain filling, we undertook simultaneous N and C labelling (for 1 week). The N labelling was conducted by replacing the $^{14}\text{NH}_4^{14}\text{NO}_3$ in the Hoagland solution with $^{15}\text{NH}_4^{15}\text{NO}_3$ that had a $\delta^{15}\text{N}$ of ca. 12.60‰. For C labelling, $^{12}\text{CO}_2$ -enriched gas was used in one of the growth chambers, whereas the other chamber was maintained at ambient conditions. Bottles of CO_2 required for the C labelling were provided by Carbueros Metálicos S.A. (Barcelona, Spain). Mixing the commercial CO_2 ($\delta^{13}\text{C}$ ca. $-32.0 \pm 0.1\text{‰}$) with the ambient air ($\delta^{13}\text{C}$ ca. -11.3‰) resulted in a $^{13}\text{CO}_2$ isotopic composition of $\delta^{13}\text{C} -23.3\text{‰}$ in the labelling chamber. Air isotopic composition inside the corresponding growth chamber was collected daily using 50-ml syringes (SGE International PTY Ltd, Ringwood, Vic., Australia) and kept in 10-ml vacutainers (BD Vacutainers, Plymouth, UK). To avoid contamination with the air present in the syringe and the needle, both were purged with N_2 prior to each sampling. The vacutainers were also over-pressurised with the same N_2 gas so that the pressure inside the vacutainer was above ambient.

Plant growth

Plant biomass was determined by harvesting just after the end of labelling (14 days after anthesis). Ears, flag leaves, shoots (including the rest of the leaves) and roots were collected separately. After drying at 60 °C for 48 h, dry mass (DM) was determined. Total nitrogen use efficiency ($\text{NUE}_{\text{total}}$) was determined as the ratio between total dry matter and total N content (Aranjuelo *et al.* 2005). Ear nitrogen use efficiency (NUE_{ear}) was calculated as the ratio between ear DM and total N content. The shoot/root ratio was determined as the total shoot DM (sum of ear, stem, including leaf sheaths, and leaf blades DM) divided by the root DM.

$\delta^{13}\text{C}$ of dark-respired CO_2 determination

Prior to the dark respiration determinations, the plants were dark-adapted for 45 min in a dark room. Flag leaf blades, ears and roots were placed separately in a plastic gas analysis chamber to determine the $\delta^{13}\text{C}$ of dark respired CO_2 ($\delta^{13}\text{C}_{\text{DR}}$). In the case of root respiration analysis, root samples were gently cleaned and immediately placed on damp paper inside the chamber; determinations were conducted during the first 4 min. Due to methodological constraints associated with shoot structure, shoot respiration was not determined. The plastic chamber ($20 \times 12 \times 6 \times 10^{-6} \text{ m}^3$) included two fans and was connected to a LI-6400 photosynthesis system (Li-Cor, Lincoln, NE, USA). Temperature in the chamber was maintained at ca. 25 °C by cool-watering

the jacket around the leaf chamber. Ingoing air was passed through the chamber at a rate of $1 \text{ l} \cdot \text{min}^{-1}$, monitored by the LI-6400.

Molar fractions of CO_2 and humidity were measured with the IRGA of the LI-6400. The PPFD inside the chamber was maintained at $0 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ by covering the entire system with a black cover. The gas analysis chamber, included in the closed system, was first flushed with CO_2 -free air to ensure that only the CO_2 respired in the chamber was accumulated. The CO_2 concentration inside the chamber was measured by the LI-6400. When the CO_2 inside reached $300 \mu\text{mol} \cdot \text{mol}^{-1}$, CO_2 samples were collected and analysed as described in more detail below.

N and C isotope composition ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) and total C and N content

Ear, flag leaf blade, shoot (including the rest of leaves) and root samples were collected on the last day of labelling (*i.e.* last day of the experiment), dried at 60 °C for 48 h, and analysed for the C isotopic composition of total organic matter ($\delta^{13}\text{C}$; TOM). A ground sample of 1 mg was used for each determination. The $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ ratios (R) of plant material were determined using an elemental analyser (EA1108, Series 1; Carlo Erba Instrumentazione, Milan, Italy) coupled to an isotope ratio mass spectrometer (Delta C; Finnigan, Mat., Bremen, Germany) operating in continuous flow mode.

The $^{13}\text{C}/^{12}\text{C}$ ratios (R) of air samples were determined by gas chromatography-combustion-isotope ratio mass spectrometry (GC-C-IRMS) as previously described (Nogués *et al.* 2008). Briefly, water vapour and oxygen from gas samples were removed and the CO_2 , argon, and N gases were separated by gas chromatography (Agilent 6890 Gas Chromatograph) coupled to an isotope ratio mass spectrometer Delta^{plus} via a GC-C Combustion III interphase (Thermo-Finnigan). The column used was 30 m \times 0.32 mm i.d. GS-GASPRO (J. and W. Scientific, USA). The carrier gas was helium at a flow rate of $1.2 \text{ ml} \cdot \text{min}^{-1}$. The injection port temperature was 220 °C. The oven temperature was kept at 60 °C during the entire run. Injection was conducted in the split mode (injected volume 0.3 ml, split flow $20 \text{ ml} \cdot \text{min}^{-1}$).

The $^{13}\text{C}/^{12}\text{C}$ ratios (R) of plant material and air samples were expressed as $\delta^{13}\text{C}$ values using international secondary standards of known $^{13}\text{C}/^{12}\text{C}$ ratios (IAEA CH7 polyethylene foil, IAEA CH6 sucrose and USGS 40 L-glutamic acid) calibrated against Vienna Pee Dee Belemnite calcium carbonate (VPDB) with an analytical precision of 0.1‰:

$$\delta^{13}\text{C} = \left(\frac{R_{\text{sample}}}{R_{\text{standard}}} \right) - 1$$

The $^{15}\text{N}/^{14}\text{N}$ ratios of plant material were also expressed in δ notation ($\delta^{15}\text{N}$) using international secondary standards of known $^{15}\text{N}/^{14}\text{N}$ ratios (IAEA N₁ and IAEA N₂ ammonium sulphate and IAEA NO₃ potassium nitrate) referred to N_2 in air, with analytical precision at about 0.2‰:

$$\delta^{15}\text{N} = \left(\frac{R_{\text{sample}}}{R_{\text{standard}}} \right) - 1$$

The N isotope discrimination ($\Delta^{15}\text{N}$, ‰), calculated in non-labelled plants was expressed following Evans (2001):

$$\Delta^{15}\text{N} = (\delta^{15}\text{N}_f - \delta^{15}\text{N}_p) / [1 + (\delta^{15}\text{N}_p/1000)]$$

where $\delta^{15}\text{N}_f$ and $\delta^{15}\text{N}_p$ refer to the fertiliser and plant N isotope composition, respectively.

The N and C isotopic composition analyses of TOM and air samples (in case of C) were conducted at the Serveis Científic-Tècnics of the University of Barcelona.

Statistical analyses

The effect of N fertilisation and plant species was tested by two-way analysis of variance (ANOVA). The analysis was conducted with the SPSS 12.0 statistical software package (SPSS Inc., Chicago, IL, USA). When there were differences between means, we made comparisons using a Tukey b multiple comparison test. The results were accepted as significant at $P < 0.05$.

RESULTS

The N treatment ($F = 26.41$, $P = 0.00$) had more effect on total DM than on analysed genotype ($F = 4.38$, $P = 0.03$). Triticale was the genotype that produced the most biomass under HN conditions, while tritordeum was the least productive, with wheat in between (Table 1). Low N availability decreased biomass production by 42% (wheat), 58% (triticale) and 25% (tritordeum), and no differences among cereals were observed (Tables 1 and S1). The more specific analyses of growth parameters revealed that the larger DM of triticale at HN was related to its larger ears ($F = 8.32$, $P = 0.00$) and shoots ($F = 5.38$, $P = 0.02$) DM. The effect of LN availability on plant growth was highest on the DM of

shoots ($F = 48.56$, $P = 0.00$) and ears ($F = 26.24$, $P = 0.00$; Table S1). LN had no effect on wheat root DM, whereas root DM increased in triticale and tritordeum. Low N availability reduced ($F = 98.52$, $P = 0.00$) the shoot/root ratio (Table 1). Under HN conditions, the largest shoot/root ratio was observed in triticale plants and the lowest in wheat. The ear DM/total DM ratio (Table 1) highlighted that LN negatively affected DM values in triticale and wheat, whereas no effect was observed in tritordeum.

Low N availability (LN) decreased the plant N content in all three cereals by 43% (wheat), 61% (triticale) and 56% (tritordeum) ($F = 52.75$, $P = 0.00$; Table 2). LN affected ear N content (N_{ear}) negatively in the three genotypes ($F = 17.44$, $P = 0.00$) but this decrease was more marked in tritordeum, where its content decreased by 50% (Table S2). Flag and shoot N content (N_{flag} and N_{shoot} , respectively) highlighted that these values were negatively affected by LN in wheat (79% and 32%, respectively) tritordeum (78% and 70%, respectively) and in triticale (74% and 83%, respectively). The effect of N availability on total and ear N use efficiencies (NUE_{total} and NUE_{ear} , respectively) was different depending on the analysed genotype ($F = 27.74$, $P = 0.00$ and $F = 5.82$, $P = 0.01$, respectively; Table S2). Although no effect was observed in wheat, NUE_{total} increased in triticale (13%) and especially in tritordeum (73%) plants. LN negatively affected NUE_{ear} in wheat and triticale (24% and 25%, respectively) but in the case of tritordeum plants, NUE_{ear} increased by 146%.

The $\delta^{13}\text{C}_{\text{TOM}}$ of labelled plants was more ^{13}C -depleted than the corresponding non-labelled plants in wheat ($F = 58.92$, $P = 0.00$), triticale ($F = 395.85$, $P = 0.00$) and tritordeum ($F = 138.75$, $P = 0.00$; Fig. 1). Under HN conditions, the ears were the main C sink in wheat and triticale

Table 1. Low (LN) versus high (HN) N fertilisation effect on total dry matter (DM, g·plant⁻¹), ear DM (g·plant⁻¹), flag DM (g·plant⁻¹), shoot DM, (g·plant⁻¹), root (g·plant⁻¹), and ear DM/total DM and shoot/root of wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants.

ecotype	treatment	total DM	ear DM	flag DM	shoot DM	root DM	ear DM/total DM	shoot/root
wheat	HN	6.78 ± 0.04	3.23 ± 0.45	0.34 ± 0.03	2.00 ± 0.16	1.21 ± 0.30	0.48 ± 0.07	4.85 ± 1.42
	LN	3.91 ± 0.83	1.38 ± 0.20	0.08 ± 0.02	1.56 ± 0.58	0.98 ± 0.34	0.37 ± 0.13	3.25 ± 0.95
triticale	HN	9.23 ± 1.25	4.20 ± 0.65	0.26 ± 0.03	4.07 ± 0.78	0.69 ± 0.20	0.46 ± 0.04	12.92 ± 3.24
	LN	3.90 ± 1.26	1.18 ± 0.41	0.08 ± 0.03	1.23 ± 0.09	1.42 ± 0.37	0.30 ± 0.01	2.12 ± 1.06
tritordeum	HN	4.94 ± 1.99	1.06 ± 0.45	0.20 ± 0.11	3.20 ± 0.71	0.48 ± 0.12	0.19 ± 0.05	9.31 ± 0.52
	LN	3.71 ± 0.88	1.19 ± 0.13	0.06 ± 0.03	1.56 ± 0.46	0.91 ± 0.35	0.28 ± 0.12	3.35 ± 1.40

Each value represents the mean of four replicates ± SD.

Table 2. Low (LN) versus high (HN) N fertilisation effect on total N (N_{total} , mg·plant⁻¹), ear N (N_{ear} , mg·plant⁻¹), flag leaf N (N_{flagleaf} , mg·plant⁻¹), shoot N (N_{shoot} , mg·plant⁻¹), root N (N_{root} , mg·plant⁻¹), total N use efficiency (NUE_{total} , mg·g⁻¹ plant) and ear N use efficiency (NUE_{ear} , mg·g⁻¹ plant) of wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants.

ecotype	treatment	N_{total}	N_{ear}	N_{flagleaf}	N_{shoot}	N_{root}	NUE_{total}	NUE_{ear}
wheat	HN	162.24 ± 2.20	76.12 ± 10.48	16.97 ± 1.64	51.34 ± 4.17	17.79 ± 4.33	41.83 ± 0.52	19.92 ± 2.62
	LN	91.89 ± 11.03	43.14 ± 6.30	3.41 ± 0.99	34.52 ± 12.95	10.81 ± 8.83	41.42 ± 3.03	15.20 ± 4.33
triticale	HN	188.24 ± 29.01	81.29 ± 15.01	9.81 ± 1.48	82.01 ± 19.07	15.11 ± 1.78	48.11 ± 0.38	21.85 ± 2.44
	LN	72.61 ± 16.70	27.36 ± 9.48	2.51 ± 0.88	13.51 ± 1.01	19.22 ± 5.85	54.57 ± 3.32	16.33 ± 0.47
tritordeum	HN	152.27 ± 45.95	24.38 ± 11.92	10.79 ± 4.16	108.23 ± 23.89	8.85 ± 3.97	32.24 ± 1.21	6.21 ± 2.96
	LN	66.96 ± 24.49	24.39 ± 13.12	2.34 ± 1.12	32.04 ± 9.56	8.19 ± 4.04	55.93 ± 4.55	15.29 ± 6.47

Each value represents the mean of four replicates ± SD.




Fig. 1. ^{13}C isotopic composition ($\delta^{13}\text{C}$) of non-labelled (Control) and labelled total organic matter (TOM) of ears, flag leaf, shoot and roots in wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants. A, C and E are high N treatments, whereas B, D and F are low N treatments. Each value represents the mean of three replicates \pm SD.

plants, whereas in tritordeum plants more labelled C was detected in flag leaves and shoots. However, in LN, the $\delta^{13}\text{C}_{\text{TOM}}$ was more homogeneous among the different organs (Fig. 1). By contrast, in unlabelled plants the flag leaf had the lowest $\delta^{13}\text{C}$ (Fig. 1). On the other hand, the $\delta^{13}\text{C}_{\text{DR}}$ also revealed significant differences (Fig. 2, Table S4) between organs for wheat ($F = 543.72$, $P = 0.00$), triticale ($F = 443.17$, $P = 0.00$) and tritordeum ($F = 779.37$, $P = 0.00$). Regardless of the genotype analysed and the N treatment, the ears respired the largest amount of labelled C (Fig. 2). In triticale and tritordeum plants, higher $\delta^{13}\text{C}$ values (*i.e.* ^{13}C -enriched) for respired CO_2 were observed in LN plants. In the case of wheat, although labelled C decreased in flag leaves in LN treatments, in ears and roots no significant differences were detected (Fig. 2). The most depleted $^{13}\text{CO}_2$ was respired by ears. Interestingly, the $\delta^{13}\text{C}_{\text{DR}}$ was similar in ears of the three genotypes exposed to HN. Although the ear $\delta^{13}\text{C}_{\text{DR}}$ values were similar in LN and HN plants of wheat and triticale, in tritordeum plants the $\delta^{13}\text{C}_{\text{DR}}$ was lower (-34% versus -32%).

Although no N treatment effect was observed on the $\delta^{15}\text{N}$ of wheat ($F = 0.82$, $P = 0.38$), triticale ($F = 2.78$, $P = 0.11$)




Fig. 2. ^{13}C isotopic composition ($\delta^{13}\text{C}$) of non-labelled (Control) and labelled dark respired CO_2 (DR) of ears, flag leaf, shoot and roots of wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants. A, C and E are high N treatments whereas B, D and F are low N treatments. Each value represents the mean of three replicates \pm SD.

and tritordeum ($F = 0.80$, $P = 0.38$), the ^{15}N labelling showed that ears were the main N sink in wheat and tritordeum plants subjected to HN, whereas in triticale, more labelled N was detected in flag leaves and shoots (Fig. 3). In LN treatments, with the exception of wheat (where the $\delta^{15}\text{N}$ was lower), the $\delta^{15}\text{N}$ of ears was similar to the values detected in flag leaves, shoots and roots.

In non-labelled plants, with the exception of wheat flag leaves, triticale ears and tritordeum roots, the $\delta^{15}\text{N}$ values obtained under LN were lower than those recorded in the corresponding HN plants (Fig. 3). As a consequence, N isotope discrimination ($\Delta^{15}\text{N}$) values recorded in LN treatments were higher for wheat ($F = 144.35$, $P = 0.00$), triticale ($F = 92.52$, $P = 0.00$) and tritordeum ($F = 31.77$, $P = 0.00$) than those in the corresponding HN treatments (Tables 3 and S3).

The negative correlation across organs and species ($R^2 = 0.58$) observed between TOM $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ revealed that, in general terms, the sink strength of the organs was similar for the C and N assimilated during the previous 7 days of the post-anthesis period (Fig. 4). The organ correlation analyses (Fig. 5) highlighted that ears ($R^2 = 0.69$), fol-

LOW RESOLUTION FIG




Fig. 3. ^{15}N isotopic composition ($\delta^{15}\text{N}$) of non-labelled (Control) and labelled total organic matter (TOM) of ears, flag leaf, shoot and roots of wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants. A, C and E are high N treatments whereas B, D and F correspond to low N treatments. Each value represents the mean of three replicates \pm SD.

lowed by shoots ($R^2 = 0.55$) and flag leaves ($R^2 = 0.54$) had similar C and N sinks. However, in the case of roots, no correlation ($R^2 = 0.06$) was detected.

DISCUSSION

In this study the NUE, together with C and N partitioning, of three cereals fertilised with HN and LN solutions were characterised. Similar to observations in a previous study (Aranjuelo *et al.* 2009), triticale, followed by wheat, was the

ecotype	treatment	ear	flag leaf	shoot	root
wheat	HN	2.52 \pm 0.03	2.83 \pm 0.14	2.09 \pm 0.01	2.85 \pm 0.31
	LN	-2.57 \pm 0.44	-2.67 \pm 0.23	-3.72 \pm 0.31	-0.21 \pm 0.11
triticale	HN	1.71 \pm 0.19	1.95 \pm 0.01	0.76 \pm 0.13	0.18 \pm 0.15
	LN	-2.23 \pm 0.09	-2.87 \pm 0.53	-3.33 \pm 0.35	-1.99 \pm 0.80
tritordeum	HN	2.29 \pm 0.18	1.90 \pm 0.01	2.47 \pm 0.21	2.89 \pm 0.05
	LN	-1.45 \pm 0.38	-2.67 \pm 0.13	-1.82 \pm 0.14	-1.38 \pm 0.85

Each value represents the mean of three replicates \pm SD.




Fig. 4. The relationship between ^{15}N isotopic composition ($\delta^{15}\text{N}$) and ^{13}C isotopic composition ($\delta^{13}\text{C}$) wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants.

genotype that produced the most biomass under optimal growth conditions. Similar results were described in Ehdaie & Waines (2001), who also observed that fertilisation with low N affected wheat and triticale production similarly. However, in this study, the data obtained showed that the production of triticale was the most N availability-dependent of the three genotypes (Table S1). Total DM and ear DM decreased in triticale by 58% and 72%, respectively, under LN and revealed that cultivation of the selected triticale cultivar (Imperioso) in soils with low N availability might be of little interest. Apparent discrepancies between our findings and the results obtained by Ehdaie & Waines (2001) could be explained by the fact the cultivars selected in these studies were derived from different breeding programmes. On the other hand, although total DM decreased by 25% and ear production was not affected by low N availability in tritordeum, the low DM (75% and 52% lower than triticale and wheat respectively) in HN conditions constrains its agromomic interest.

The negative effect of LN treatment (compared with the corresponding HN treatment) on harvest index (expressed as ear DM per total DM; Table 1) observed in wheat and triticale, revealed that these species had lower ear sink strength as N availability decreased. This result is in agreement with previous studies (Ehdaie & Waines 2001; Mergoum *et al.* 2004)

Table 3. Low (LN) versus high (HN) N fertilisation effect on the ear, flag leaf, shoot and root ^{15}N isotopic discrimination ($\Delta^{15}\text{N}$, ‰) of non-labelled wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants.




Fig. 5. The relationship between $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ of ears (Δ), flag leaf (\bullet), shoot (\circ) and roots (\blacktriangledown) of wheat (Califa sur), triticale (Imperioso) and tritordeum (HT 621) plants.

that describe how triticale yield is strongly conditioned by N fertilisation levels. It can also be highlighted that under HN conditions, tritordeum had an ear DM/total DM ratio lower than the other two cereals. Similar results have been described in a multi-local field study conducted with the same genotypes (Villegas *et al.* 2010). According to the observations of these authors, tritordeum had fewer grains per spike and the kernel weight was also the lowest compared with wheat and triticale plants. The fact that in tritordeum plants the relative investment in the shoots was higher than in the ears (Aranjuelo *et al.* 2009) reflects a lack of agronomic improvement of this amphiploid compared with wheat and triticale. Since tritordeum is a quite a new species, its present genotypes have not been subjected to extensive plant breeding programmes with the consequent limitations on yield of those plants. Our data highlight the need to increase breeding efforts directed towards increasing ear sink strength in tritordeum.

Grain filling is conditioned by the size and availability of C and N pools (Fuentes-Mendizábal *et al.* 2010). As mentioned before, the main C sources for grain filling are the flag leaf, the stem internodes and the C assimilated by the ears. The fact that the flag leaves had the lowest labelled C in wheat and tritordeum is indicative that these organs acted as a C source for grain filling, with grain having more labelled C. Interestingly, the large labelled C values of flag leaves and shoots in triticale suggest two possibilities: (i) shoots represented major labelled C sinks that competed with the ears as C sinks; or (ii) shoots acted as major labelled C sinks due to the poor sink strength of triticale ears. In fact, triticale is characterised by constitutively (*i.e.* in the absence of stress) wrinkled grains. Despite the large investment of C compounds in respiration, little attention has been given to this point in previous studies in cereals (Bort *et al.* 1996; Dupont & Altenbach 2003; Lopes *et al.* 2006; Aranjuelo *et al.* 2009). The fact that in wheat and tritordeum the $\delta^{13}\text{C}_{\text{DR}}$ was less

labelled than the corresponding TOM revealed that those plants used respiratory substrates formed before the period when labelling was performed (*i.e.* pre-labelling C), which means that more labelled C was available for grain filling. However, in the case of triticale plants, $\delta^{13}\text{C}_{\text{DR}}$ respired by the ears was composed of more labelled C that could not be directed toward grain filling. Since, under LN conditions, a similar $\delta^{13}\text{C}$ was observed in TOM and respired CO_2 , it can be expected that triticale plants invested more labelled C in respiration processes to the detriment of grain filling. This study highlights that in spite of species and N treatment, the plants 'invested' a large amount of labelled C in respiratory processes, which means that less labelled C was available for grain filling.

The stable N isotope signature ($\delta^{15}\text{N}$) is a useful screening tool for plant N metabolism (Coque *et al.* 2006; Yousfi *et al.* 2010). $\delta^{15}\text{N}$ is conditioned by the N uptake from the medium into the root cells or by the subsequent enzymatic assimilation into other N forms (Kolb & Evans 2003; Tcherkez & Hodges 2008) and by the discrimination of enzymes involved in N uptake (Tcherkez & Hodges 2008). Furthermore, soil N concentration levels have been described (Serret *et al.* 2008; Yousfi *et al.* 2010) to condition ^{15}N discrimination ($\Delta^{15}\text{N}$). In agreement with these authors, our data revealed that $\Delta^{15}\text{N}$ is conditioned by the plant N demand/availability ratio. The fact that (under non-labelled conditions) LN plants had higher $\delta^{15}\text{N}$ values highlighted that when N availability is limiting, the $\Delta^{15}\text{N}$ in LN plants was reduced.

Our study showed different NUE response patterns depending on the genotype analysed and the N treatment. Under HN conditions, tritordeum plants had the lowest total and ear N use efficiency ($\text{NUE}_{\text{total}}$ and NUE_{ear}). The fact that N_{shoot} represented 71% of the total N, whereas N_{ear} only represented 16% of it, highlights the low ear N sink strength, and this explained the low NUE in tritordeum. However, when N availability was limiting, tritordeum was the only genotype that increased $\text{NUE}_{\text{total}}$ and NUE_{ear} . Compared with the corresponding HN, under LN tritordeum increased in ear sink strength (ear contained 36% of total N and shoots 48%; Tables 1 and 2). However, in the case of triticale, $\text{NUE}_{\text{total}}$ increased under HN, and the large shoot and root biomass in LN treatments decreased ear sink strength and, consequently, grain filling. In this study, wheat plants proved to be the only genotype where N treatment had no effect on $\text{NUE}_{\text{total}}$ and NUE_{ear} . The stability of NUE under both N fertilisation regimes is explained by the fact that the present wheat cultivars are the consequence of improved plant breeding programmes.

The ^{15}N labelling enabled the further characterisation of N assimilation and partitioning during this post-anthesis phenological stage. During grain filling, shoot proteins are hydrolysed to amino acids to enable the remobilisation of N content toward grains (Gebbing *et al.* 1998, 1999; Masclaux-Daubresse *et al.* 2008). As described in previous studies (Palta & Fillery 1995; Lopes *et al.* 2006), 60–95% of N present in grains comes from the remobilisation of N stored in shoots and roots before anthesis. The lower amount of labelled N present in triticale ears (compared with wheat and tritordeum) highlights that pre-anthesis N remobilisation contributed toward grain filling in this genotype (Table S4). In tritordeum and wheat plants the redistribution of labelled N from

source organs (flag leaves and shoots) toward sink organs (ears) was more marked in HN, especially in HN tritordeum (Gebbing *et al.* 1999). Such differences between HN and LN were caused by the fact that N remobilisation starts earlier in plants under LN fertilisation conditions compared to HN fertilisation conditions (Ta & Weiland 1992). These results suggest that N fertilisation during this period was more relevant for grain filling of wheat and tritordeum plants compared with triticale.

The double ^{13}C and ^{15}N labelling also highlighted that C and N assimilated post-anthesis were similarly partitioned (Fig. 3), regardless of the genotype and N treatment (Gebbing & Schnyder 1999). The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ correlation analyses (Fig. 4) confirmed that the distribution among organs of the labelled C and N was similar. The more specific analyses at the level of the different organs (Fig. 5) revealed that the ears showed the highest correlation between $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ ($R^2 = 0.69$), indicating their major roles as C and N sinks. Negative correlations between $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ were also found in shoots ($R^2 = 0.55$) and to a lesser extent in flag leaves ($R^2 = 0.54$). These results are in agreement with the finding of Gebbing *et al.* (1998), who observed that post-anthesis C was mainly accumulated in glumes and to a lesser extent in the stem.

Although the role of ears as a major sink has been widely described (Schnyder 1992; Gebbing *et al.* 1998, 1999; Gebbing & Schnyder 1999), the C and N sink strength during post-anthesis has been related to the synthesis of structural compounds and elongation of the peduncle (Borrell *et al.* 1991; Bonnett & Incoll 1992). Such expansion is associated with the synthesis of cell wall material, which requires the utilisation of post-anthesis photosynthate (Gebbing *et al.* 1998). On the other hand, no such correlation was observed in roots (Fig. 5). The use of N and C compounds by plants involves, in part, resource remobilisation (Masclaux-Daubresse *et al.* 2008). Furthermore, as observed by Kichey *et al.* (2007), grain yield is based on N uptake before flowering and on the remobilisation of N during grain filling. The redistribution of labelled N from source organs (flag leaves and shoots) toward sink organs (ears) of N assimilated during the post-anthesis period was more marked in HN triticale and in HN tritordeum (Gebbing *et al.* 1999). Such differences between HN and LN were caused by the fact that N remobilisation starts earlier when plants are under low N fertilisation conditions compared to high N fertilisation conditions (Ta & Weiland 1992).

CONCLUSIONS

As mentioned above, the challenge for breeders is to select plants with high yields and high NUE. This study revealed that even if triticale had the largest total and ear DM values in HN conditions, it proved to be the genotype that was the most agronomically negatively affected by LN. The largest allocation of labelled C and N in shoots was involved in the decreased ear N use efficiency (NUE_{ear}) of triticale. Furthermore, N fertilisation during post-anthesis was least relevant in triticale. Although wheat NUE was similar to values in triticale, it did not improve under LN conditions. Tritordeum plants were the only ones where NUE ($\text{NUE}_{\text{total}}$ and NUE_{ear}) values increased under LN treatment. The remobilisation of

pre-labelled N in these plants was shown to be more relevant to the increase in grain filling and NUE in LN.

In summary, this study has shown that ear sink and NUE strength need to be improved in wheat and triticale LN plants, whereas in the case of tritordeum, although NUE was enhanced, the low yield in HN conditions reduced the potential of this genotype. The 'competition' for photoassimilates between ear DM and respiration processes, together with the investment of C and N resources in leaves and shoots also limited the NUE of plants. The study also highlights the potential of breeding for improved NUE in LN, especially in triticale plants that were revealed to be more sensitive to fertiliser levels. Furthermore, the promising results for tritordeum plants highlight the need to improve their yield under LN conditions and more especially under HN.

ACKNOWLEDGEMENTS

The authors would like to thank Pilar Teixidor and Pilar Rubio for technical support with the isotopic analyses. We also thank Ricardo Simoneau, Josep Matas and Toni Pardo for technical assistance. The authors would also like to thank the two anonymous reviewers who contributed to improvement of the manuscript. This work was funded by the Spanish National Research and Development Programme–European Regional Development Fund ERDF (project nos. AGF2005-07257-C04-01, CGL2009-13079-CO2 and AGL2010-20180). Iker Aranjuelo was the recipient of a Ramón y Cajal research grant (Ministerio de Ciencia e Innovación). Finally, we would like to thank the private company AGRASYS and Antonio Martín for providing HT 621 seeds.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Table S1. Mean squares explained by ANOVA for the main effects (genotype and N treatments and their corresponding interactions) for total dry matter (DM), ear DM, flag leaf DM, shoot DM, root DM, ear DM/total DM and shoot/root. Percentage of the sum of squares is expressed in brackets. The different symbols indicate significant differences at 5% (*) and 1% (**), whereas n.s. refers to no significant difference.

Table S2. Mean squares explained by ANOVA for the main effects (genotype and N treatment and their corresponding interactions) for total N (N_{total}), ear N (N_{ear}), flag leaf N ($\text{N}_{\text{flagleaf}}$), shoot N (N_{shoot}), root N (N_{root}), total N use efficiency ($\text{NUE}_{\text{total}}$) and ear N use efficiency (NUE_{ear}). Percentage of the sum of squares is in brackets. The different symbols indicate significant differences at 5% (*) and 1% (**), whereas n.s. refers to no significant difference.

Table S3. Mean squares explained by ANOVA for the main effects (genotype and N treatments and their corresponding interactions) for ^{15}N isotopic discrimination ($\Delta^{15}\text{N}$, ‰). Percentage of the sum of squares is in brackets. The different symbols indicate significant differences at 5% (*) and 1% (**), whereas n.s. refers to no significant difference.

Table S4. Mean squares explained by ANOVA for the main effects (labelling, N treatments, organ and their corresponding interactions) for total organic matter (TOM) ^{13}C and ^{15}N

isotopic composition and dark respired (DR) CO₂ δ¹³C. Percentage of the sum of squares is in brackets. The different symbols indicate significant differences at 5% (*) and 1% (**), whereas n.s. refers to no significant difference.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

REFERENCES

- Aranjuelo I., Pérez P., Hernández L., Irigoyen J.J., Zita G., Martínez-Carrasco R., Sánchez-Díaz M. (2005) The response of nodulated alfalfa to water supply, temperature and elevated CO₂: photosynthetic downregulation. *Physiologia Plantarum*, **123**, 348–358.
- Aranjuelo I., Cabrera-Bosquet L., Mottaleb S.A., Arous J.L., Nogués S. (2009) ¹³C/₁₂ C isotope labelling to study carbon partitioning and dark respiration in cereals subjected to water stress. *Rapid Communications in Mass Spectrometry*, **23**, 2819–2828.
- Aranjuelo I., Cabrera-Bosquet L., Morcuende R., Avicé J.C., Nogués S., Arous J.L., Martínez-Carrasco R., Pérez P. (2011a) Does ear C sink strength contribute to overcoming photosynthetic acclimation of wheat plants exposed to elevated CO₂? *Journal of Experimental Botany*, **62**, 3957–3969.
- Aranjuelo I., Ebberts A.L., Evans D.R., Tissue D.T., Nogués S., van Gestel N., Payton P., Ebbert V., Adams W.W. III, Nowak R.S., Smith S.D. (2011b) Maintenance of C sinks sustains enhanced C assimilation during long-term exposure to elevated [CO₂] in Mojave Desert shrubs. *Oecologia*, **167**, 339–354.
- Arous J.L., Slafer G.A., Royo C., Serret M.D. (2008) Breeding for yield potential and stress adaptation in cereals. *Critical Reviews in Plant Sciences*, **27**, 377–412.
- Arnon D.I., Hoagland D.R. (1939) A comparison of water culture and soil as media for crop production. *Science*, **89**, 512–514.
- Asseng S., Van Herwaarden A.F. (2003) Analysis of the benefits to wheat yield from assimilates stored prior to grain filling in a range of environments. *Plant and Soil*, **256**, 217–229.
- Avicé J.C., Oury A., Lemaire G., Boucaud J. (1996) Nitrogen and carbon flows estimated by ¹⁵N and ¹³C pulse-chase labeling during regrowth of Alfalfa. *Plant Physiology*, **112**, 281–290.
- Ballesteros J., Ramírez M.C., Martínez C., Atienza S.G., Martín A. (2005) Registration of HT621, a high carotenoid content tritordeum germplasm line. *Crop Science*, **45**, 2662–2663.
- Barro F., Fontes A.G., Maldonado J.M. (1991) Organic nitrogen content and nitrate and nitrite reductase activities in tritordeum and wheat grown under nitrate or ammonium. *Plant and Soil*, **135**, 251–256.
- Barro F., González-Fontes A., Maldonado J.M. (1996) Relation between photosynthesis and dark respiration in cereal leaves. *Journal of Plant Physiology*, **149**, 64–68.
- Bonnett G.D., Incoll L.D. (1992) The potential pre-anthesis and post-anthesis contributions of stem internodes to grain yield in crops of winter barley. *Annals of Botany*, **69**, 219–225.
- Borghesi B., Corbellini M., Minoia C., Palumbo M., Di Fonzo N., Perenzin M. (1997) Effects of Mediterranean climate on wheat bread-making quality. *European Journal of Agronomy*, **6**, 145–154.
- Borrell A.K., Incoll L.D., Dalling M.J. (1991) The influence of the Rht1 and Rht2 alleles on the growth of wheat stems and ears. *Annals of Botany*, **67**, 103–110.
- Bort J., Brown R.H., Arous J.L. (1996) Refixation of respiratory CO₂ in the ears of C₃ cereals. *Journal of Experimental Botany*, **47**, 1567–1575.
- Cabrera-Bosquet L., Molero G., Nogués S., Arous J.L. (2009) Water and nitrogen conditions affect the relationships of Δ¹³C and Δ¹⁸O to gas exchange and growth in durum wheat. *Journal of Experimental Botany*, **60**, 1633–1644.
- Cook D.R. (1999) *Medicago truncatula* – a model in the making! *Current Opinion in Plant Biology*, **2**, 301–304.
- Coque M., Bertin P., Hirel B., Gallais A. (2006) Genetic variation and QTLs for ¹⁵N natural abundance in a set of maize recombinant inbred lines. *Field Crops Research*, **97**, 310–321.
- Cossani C.M., Savin R., Slafer G.A. (2007) Contrasting performance of barley and wheat in a wide range of conditions in Mediterranean Catalonia (Spain). *Annals of Applied Biology*, **151**, 167–173.
- Dupont F.M., Altenbach S.B. (2003) Molecular and biochemical impacts of environmental factors on wheat grain development and protein synthesis. *Journal of Cereal Science*, **38**, 133–146.
- Ehdaie B., Waines J.G. (2001) Sowing date and nitrogen rate effects on dry matter and nitrogen partitioning in bread and durum wheat. *Field Crop Research*, **73**, 47–61.
- Ercoli L., Lulli L., Mariotti M., Masoni A., Arduini I. (2008) Post-anthesis dry matter and nitrogen dynamics in durum wheat as affected by nitrogen supply and soil water availability. *European Journal of Agronomy*, **28**, 138–147.
- Evans R.D. (2001) Physiological mechanism influencing plant nitrogen isotope composition. *Trends in Plant Science*, **6**, 121–126.
- Evans L.T., Wardlaw I.F., Fischer R.A. (1975) *Crop physiology: some case histories*. Cambridge University Press, Cambridge, 101–150.
- Fuertes-Mendizábal T., Setián I., Estavillo J.M., Begoña González-Moro M. (2010) Late nitrogen fertilization affects carbohydrate mobilization in wheat. *Journal of Plant Nutrition and Soil Science*, **173**, 907–919.
- Gebbing T., Schnyder H. (1999) Pre-anthesis reserve utilization for protein and carbohydrate synthesis in grains of wheat. *Plant Physiology*, **121**, 871–878.
- Gebbing T., Schnyder H., Kühbauch W. (1998) Carbon mobilization in shoot parts and roots of wheat during grain filling: assessment by ¹³C/¹²C steady-state labeling, growth analysis and balance sheets of reserves. *Plant, Cell and Environment*, **21**, 301–313.
- Gebbing T., Schnyder H., Kühbauch W. (1999) The utilization of pre-anthesis reserves in grain filling of wheat. Assessment by steady-state ¹³CO₂/¹²CO₂ labeling. *Plant, Cell and Environment*, **22**, 851–858.
- Kichey T., Hirel B., Heumez E., Dubois F., Le Gouis J. (2007) In winter wheat (*Triticum aestivum* L.), post-anthesis nitrogen uptake and remobilisation to the grain correlates with agronomic traits and nitrogen physiological markers. *Field Crop Research*, **102**, 22–32.
- Kolb K.J., Evans R.D. (2003) Influence of nitrogen source and concentration on nitrogen isotopic discrimination in two barley genotypes (*Hordeum vulgare* L.). *Plant, Cell and Environment*, **26**, 1431–1440.
- Lea P.J., Azevedo R.A. (2006) Nitrogen use efficiency. 1. Uptake of nitrogen from the soil. *Annals of Applied Biology*, **149**, 243–247.
- Lea P.J., Azevedo R.A. (2007) Nitrogen use efficiency. 2. Amino acid metabolism. *Annals of Applied Biology*, **151**, 269–275.
- Lopes M.S., Cortadellas N., Kichey T., Dubois F., Habash D.Z., Arous J.L. (2006) Wheat nitrogen metabolism during grain filling: comparative role of glumes and the flag leaf. *Planta*, **225**, 165–181.
- Martin A. (1988) Tritordeum: the first ten years. *Rachis*, **7**, 12–15.
- Martin A., Alvarez J.B., Martín L.M., Barro F., Ballesteros J. (1999) The development of tritordeum: a novel cereal for food processing. *Journal of Cereals Science*, **30**, 85–95.
- Masclaux-Daubresse C., Reisdorf-Cren M., Orsel M. (2008) Leaf nitrogen remobilisation for plant development and grain filling. *Plant Biology*, **10**, 23–36.
- Masclaux-Daubresse C., Daniel-Vedele F., Dechorgnat J., Chardon F., Gaufichon L., Suzuki A. (2010) Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. *Annals of Botany*, **105**, 1141–1157.
- Masclaux-Daubresse C., Chardon F. (2011) Exploring nitrogen remobilization for seed filling using natural variation in *Arabidopsis thaliana*. *Journal of Experimental Botany*, **62**, 2131–2142.
- Matson P., Lohse K.A., Hall S.J. (2002) The globalization of nitrogen deposition: consequences for terrestrial ecosystems. *Ambio*, **31**, 113–119.
- Mergoum M., Pfeiffer W.H., Peña R.J., Ammar K., Rajaram S. (2004) Triticale crop improvement: the CIMMYT programme. In: Mergoum M., Gómez-Macpherson H. (Eds), *Triticale improvement and production*. FAO, Rome, Italy, pp 11–26.
- Nogués S., Aranjuelo I., Pardo A., Azcón-Bieto J. (2008) Assessing the stable carbon isotopic composition of intercellular CO₂ in a CAM plant using gas chromatography-combustion-isotope ratio mass spectrometry. *Rapid Communications in Mass Spectrometry*, **22**, 1017–1022.
- Palta J.A., Fillery I.R. (1995) N application increases pre-anthesis contribution of dry matter to grain yield in wheat grown on a duplex soil. *Australian Journal of Agricultural Research*, **46**, 507–518.
- Papakosta D.K., Gagianas A.A. (1991) Nitrogen and dry matter accumulation, remobilisation, and losses for Mediterranean wheat during grain filling. *Agronomy Journal*, **83**, 864.
- Passioura J.B. (2002) Environmental biology and crop improvement. *Functional Plant Biology*, **29**, 537–546.
- Raun W.R., Johnson G.V. (1999) Improving nitrogen use efficiency for cereal production. *Agronomy Journal*, **91**, 357–363.
- Schnyder H. (1992) Long-term steady-state labeling of wheat plants by use of natural ¹³CO₂/¹²CO₂ mixtures in an open, rapidly turned-over system. *Planta*, **187**, 128–135.

- Serret M.D., Ortiz-Monasterio I., Pardo A., Araus J.L. (2008) The effects of urea fertilisation and genotype on yield, nitrogen use efficiency, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in wheat. *Annals of Applied Biology*, **153**, 243–257.
- Spieritz J.H.J., De Vos N.M. (1983) Agronomical and physiological aspects of the role of nitrogen in yield formation of cereals. *Plant and Soil*, **75**, 379–391.
- Ta C.T., Weiland R.T. (1992) Nitrogen partitioning in maize during ear development. *Crop Science*, **32**, 443–451.
- Tambussi E.A., Bort J., Guiamet J.J., Nogués S., Araus J.L. (2007) The photosynthetic role of ears in C_3 cereals: metabolism, water use efficiency and contribution to grain yield. *Critical Reviews in Plant Sciences*, **26**, 1–16.
- Tcherkez G., Hodges M. (2008) How stable isotopes may help to elucidate primary nitrogen metabolism and its interaction with (photo)respiration in C_3 leaves. *Journal of Experimental Botany*, **59**, 1685–1693.
- Villegas D., Casadesús J., Atienza S.G., Martos V., Maalouf F., Karam F., Aranjuelo I., Nogués S. (2010) Tritordeum, wheat and triticale yield components under multi-local Mediterranean drought conditions. *Field Crops Research*, **116**, 68–74.
- Yousfi S., Serret M.D., Voltas J., Araus J.L. (2010) Effect of salinity and water stress during the reproductive stage on growth, ion concentrations, $\Delta^{13}\text{C}$, and $\delta^{15}\text{N}$ of durum wheat and related amphiploids. *Journal of Experimental Botany*, **61**, 3529–3542.

Author Query Form

Journal: PLB

Article: 593

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	Author: Starting in 2011, research articles will be subject to a page charge at a rate of GBP 100 per page above the 8-page limit (the first 8 pages are free of charge). Your article exceeds the page limit. Please either make the necessary changes for the article to fit into 8 pages or go to the journal website, download the Page Charges Form, fill it in and send it to the production editor at Wiley-Blackwell (please, click on the following link to download the form: http://www.blackwellpublishing.com/pdf/PLB_Page_Charge_Form.pdf)	
2	AUTHOR: Barro <i>et al.</i> 1993, has been changed to Barro <i>et al.</i> 1991, so that this citation matches the Reference List. Please confirm that this is correct.	
3	AUTHOR: Alphabets have been changed to symbols to match with physical Figure 5. Please check.	
4	AUTHOR: Figure 1 has been saved at a low resolution of 213 dpi. Please resupply at 600 dpi. Check required artwork specifications at http://authorservices.wiley.com/submit_illust.asp?site=1	
5	AUTHOR: Figure 2 has been saved at a low resolution of 200 dpi. Please resupply at 600 dpi. Check required artwork specifications at http://authorservices.wiley.com/submit_illust.asp?site=1	
6	AUTHOR: Figure 3 has been saved at a low resolution of 202 dpi. Please resupply at 600 dpi. Check required artwork specifications at http://authorservices.wiley.com/submit_illust.asp?site=1	
7	AUTHOR: Figure 4 has been saved at a low resolution of 226 dpi. Please resupply at 600 dpi. Check required artwork specifications at http://authorservices.wiley.com/submit_illust.asp?site=1	
8	AUTHOR: Figure 5 has been saved at a low resolution of 178 dpi. Please resupply at 600 dpi. Check required artwork specifications at http://authorservices.wiley.com/submit_illust.asp?site=1	

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	⧵	New matter followed by ⧵ or ⧵ [Ⓢ]
Delete	/ through single character, rule or underline or ⎓ through all characters to be deleted	⧻ or ⧻ [Ⓢ]
Substitute character or substitute part of one or more word(s)	/ through letter or ⎓ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↵
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⧵
Change bold to non-bold type	(As above)	⧵
Insert 'superior' character	/ through character or ⧵ where required	Y or Y under character e.g. Y or Y
Insert 'inferior' character	(As above)	⧵ over character e.g. ⧵
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	Y or Y and/or Y or Y
Insert double quotation marks	(As above)	Y or Y and/or Y or Y
Insert hyphen	(As above)	⎓
Start new paragraph	⌞	⌞
No new paragraph	⌞	⌞
Transpose	⌞	⌞
Close up	linking ○ characters	○
Insert or substitute space between characters or words	/ through character or ⧵ where required	Y
Reduce space between characters or words		↑