

CONCISE REPORT

Identification of the *PTPN22* functional variant R620W as susceptibility genetic factor for giant cell arteritis

A Serrano,¹ A Márquez,¹ S L Mackie,² F D Carmona,¹ R Solans,³ J A Miranda-Filloy,⁴ J Hernández-Rodríguez,⁵ M C Cid,⁵ S Castañeda,⁶ IC Morado,⁷ J Narváez,⁸ R Blanco,⁹ B Sopeña,¹⁰ M J García-Villanueva,¹¹ J Monfort,¹² N Ortego-Centeno,¹³ A Unzurrunzaga,¹⁴ B Marí-Alfonso,¹⁵ J Sánchez-Martín,¹⁶ E de Miguel,¹⁷ C Magro,¹⁸ E Raya,¹⁸ N Braun,¹⁹ J Latus,¹⁹ O Molberg,²⁰ B A Lie,^{21,22} F Moosig,²³ T Witte,²⁴ A W Morgan,² M A González-Gay,⁹ J Martín,¹ UK GCA Consortium Spanish GCA Consortium

Handling editor Tore K Kvien

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2013-203641>).

For numbered affiliations see end of article.

Correspondence to

Dr Ana Márquez, Instituto de Parasitología y Biomedicina López-Neyra. Consejo Superior de Investigaciones Científicas. Parque Tecnológico Ciencias de la Salud. Avenida del Conocimiento s/n 18016-Armiña (Granada), Spain; anamaort@ipb.csic.es

AS, AM and SLM contributed equally to this work. MAGG and JM share senior authorship.

Received 18 March 2013

Revised 2 July 2013

Accepted 9 July 2013

Published Online First

14 August 2013

ABSTRACT

Objective To analyse the role of the *PTPN22* and *CSK* genes, previously associated with autoimmunity, in the predisposition and clinical phenotypes of giant cell arteritis (GCA).

Methods Our study population was composed of 911 patients diagnosed with biopsy-proven GCA and 8136 unaffected controls from a Spanish discovery cohort and three additional independent replication cohorts from Germany, Norway and the UK. Two functional *PTPN22* polymorphisms (rs2476601/R620W and rs33996649/R263Q) and two variants of the *CSK* gene (rs1378942 and rs34933034) were genotyped using predesigned TaqMan assays.

Results The analysis of the discovery cohort provided evidence of association of *PTPN22* rs2476601/R620W with GCA ($P_{FDR}=1.06E-04$, OR=1.62, CI 95% 1.29 to 2.04). The association did not appear to follow a specific GCA subphenotype. No statistically significant differences between allele frequencies for the other *PTPN22* and *CSK* genetic variants were evident either in the case/control or in stratified case analysis. To confirm the detected *PTPN22* association, three replication cohorts were genotyped, and a consistent association between the *PTPN22* rs2476601/R620W variant and GCA was evident in the overall meta-analysis ($P_{MH}=2.00E-06$, OR=1.51, CI 95% 1.28 to 1.79).

Conclusions Our results suggest that the *PTPN22* polymorphism rs2476601/R620W plays an important role in the genetic risk to GCA.

INTRODUCTION

Giant cell arteritis (GCA) is a chronic vasculitis that shows a complex aetiology derived from the interaction between both genetic and environmental factors.¹ Similar to most immune-related disorders, the highest susceptibility signals belong to the human leukocyte antigen region. However, different studies have highlighted that genes involved in inflammation pathways may also be implicated in GCA susceptibility.² In spite of these findings, the genetic background of this condition is still poorly understood.

Although the aetiology of GCA remains unclear, it is well known that innate and adaptive immune responses are involved in its pathogenesis. Several lines of evidence indicate that this vasculitis is a T cell-mediated disease with both Th17 and Th1 cells contributing to inflammation. While Th1 response is associated with chronically persistent vascular lesions, Th17 immunity appears to be more important for acute manifestations, both systemically and in the blood vessels.^{3 4}

The *PTPN22*/*CSK* pathway is a master regulator of autoimmunity, with a key role in the negative control of the signalling mediated by the T cell receptor (TCR).⁵ Interestingly, several single-nucleotide polymorphisms (SNPs) located within these two genes have been associated with autoimmunity,^{6–11} suggesting that this is one of the molecular pathways shared by different autoimmune disorders.

Regarding *PTPN22*, it has been reported that two autoimmune disease-associated variants, rs2476601 (R620W) and rs33996649 (R263Q) influence the function of the protein.^{12 13} On the other hand, two *CSK* polymorphisms, rs34933034 and rs1378942, were recently identified as susceptibility factors for systemic sclerosis¹¹ and systemic lupus erythematosus (SLE),¹⁰ respectively. A functional role for the *CSK* genetic variant rs34933034*A in SLE patients has been proposed in a recent study.¹⁰

Based on this, we decided to assess the role of the disease-associated *PTPN22* and *CSK* polymorphisms in both predisposition to and the clinical phenotypes of GCA.

METHODS

Study population

A total of 911 GCA patients and 8136 unrelated healthy controls were included in this study. First, we analysed a discovery cohort of 623 GCA patients and 1729 healthy controls of Spanish Caucasian ancestry. Subsequently, three independent replication cohorts were analysed (72 GCA and 937 controls from Germany; 60 GCA and 271

To cite: Serrano A, Márquez A, Mackie SL, et al. *Ann Rheum Dis* 2013;**72**:1882–1886.

controls from Norway; 156 GCA and 5199 controls from the UK). Case and control sets were matched by geographical origin and ethnicity, but not by age, which may represent a limitation of the study. *PTPN22* rs2476601 genotype data from the control population of Germany were obtained from Hüffmeier *et al.*,¹⁴ since this set matched geographically and ethnically our German GCA cohort. More detailed information about the UK controls can be obtained from Morgan *et al.*¹⁵ Informed written consent from all participants and approval from the local ethical committees were obtained in accordance with the tenets of the Declaration of Helsinki. All patients had a positive temporal artery biopsy (disruption of the internal elastic laminae with infiltration of mononuclear cells into the arterial wall with or without multinucleated giant cells) and fulfilled the 1990 American College of Rheumatology classification criteria for GCA.¹⁶ In the subphenotype analysis, the patients were stratified according to manifestations of polymyalgia rheumatica (PMR) and the presence or absence of visual ischaemic manifestations (VIM; if they experienced transient visual loss including amaurosis fugax, permanent visual loss or diplopia) and irreversible occlusive disease (IOD; if they had at least one of the following features: permanent visual loss, stroke or occlusive disease in the upper extremities or lower extremities).

Genotyping methods

Genomic DNA was extracted from peripheral white blood cells using standard procedures. Two SNPs located within *PTPN22*, rs2476601/R620W and rs33996649/R263Q, and two SNPs located within *CSK*, rs1378942 and rs34933034, were genotyped using the TaqMan allelic discrimination assay technology

on a 7900HT Fast Real-Time PCR System, both from Applied Biosystems (Foster City, California, USA). For the UK samples, rs2476601/R620W was genotyped by direct sequencing.

Statistical analysis

The overall statistical power of the analysis, according to Power Calculator for Genetic Studies 2006 software (<http://www.sph.umich.edu/csg/abecasis/CaTS/>), is shown in online supplementary table S1. Plink (v1.07) (<http://pngu.mgh.harvard.edu/purcell/plink/>) and StatsDirect V2.6.6 (StatsDirect Ltd, Cheshire, UK) were used to perform 2×2 contingency tables and χ^2 test and/or Fisher's exact test. ORs and 95% CIs were obtained according to Woolf's method. The Benjamini and Hochberg (1995) step-up false discovery rate (FDR) control correction for multiple testing¹⁷ was applied to the p values of the discovery cohort. After correction, p values lower than 0.05 were considered statistically significant. The allelic combinations were tested using Plink and Haploview (V4.2). The analysis of the combined data from all populations was performed using Plink and StatsDirect. Breslow–Day (BD) test method was used to estimate the homogeneity among populations. Pooled analyses were performed by Mantel–Haenszel test under fixed effects.

RESULTS

After genotyping, no divergence from Hardy–Weinberg equilibrium was observed either in controls or cases ($p > 0.01$), and control allelic frequencies were similar to those previously reported in equivalent European Caucasian populations.^{8 10 11}

First, we conducted an association study in a case–control set of Spanish Caucasian origin. As shown in table 1, when allelic frequencies were compared between cases and controls, a clear

Table 1 Genotype and allele distribution of *PTPN22* rs2476601, rs33996649 and *CSK* rs1378942, rs34933034 in Spanish biopsy-proven GCA patients and healthy controls

SNP	Locus	1/2	Subgroup (N)	Genotype, N (%)			MAF (%)	Allele test		
				1/1	1/2	2/2		p Value*	P _{FDR} †	OR (CI 95%)‡
rs2476601	<i>PTPN22</i>	A/G	Controls (n=1729)	13 (0.75)	200 (11.57)	1516 (87.68)	6.54			
			GCA (n=623)	4 (0.64)	119 (19.10)	500 (80.26)	10.19	2.66E-05	1.06E-04	1.62 (1.29 to 2.04)
			PMR+ (n=259)	2 (0.77)	53 (20.46)	204 (78.76)	11.00	2.26E-04	9.02E-04	1.77 (1.30 to 2.40)
			VIM+ (n=168)	1 (0.60)	36 (21.43)	131 (77.98)	11.31	1.03E-03	4.10E-03	1.82 (1.27 to 2.62)
			IOD+ (n=96)	2 (2.08)	21 (21.88)	73 (76.04)	13.02	5.47E-04	2.19E-03	2.14 (1.38 to 3.33)
rs33996649	<i>PTPN22</i>	T/C	Controls (n=1729)	4 (0.23)	110 (6.36)	1615 (93.41)	3.41			
			GCA (n=623)	1 (0.16)	39 (6.26)	583 (93.58)	3.29	0.838	0.919	0.96 (0.67 to 1.38)
			PMR+ (n=259)	0 (0.00)	12 (4.63)	247 (95.37)	2.32	0.191	0.382	0.67 (0.37 to 1.23)
			VIM+ (n=168)	0 (0.00)	12 (7.14)	156 (92.86)	3.57	0.878	0.878	1.05 (0.57 to 1.92)
			IOD+ (n=96)	0 (0.00)	6 (6.25)	90 (93.75)	3.13	0.831	0.831	0.91 (0.40 to 2.10)
rs1378942	<i>CSK</i>	C/A	Controls (n=1729)	281 (16.25)	798 (46.15)	650 (37.59)	39.33			
			GCA (n=623)	129 (20.71)	230 (36.92)	264 (42.38)	39.17	0.919	0.919	0.99 (0.87 to 1.13)
			PMR+ (n=259)	55 (21.24)	99 (38.22)	105 (40.54)	40.35	0.658	0.756	1.04 (0.86 to 1.26)
			VIM+ (n=168)	34 (20.24)	69 (41.07)	65 (38.69)	40.77	0.605	0.807	1.06 (0.85 to 1.33)
			IOD+ (n=96)	20 (20.83)	39 (40.63)	37 (38.54)	41.15	0.616	0.822	1.08 (0.80 to 1.45)
rs34933034	<i>CSK</i>	A/G	Controls (n=1729)	72 (4.16)	537 (31.06)	1120 (64.78)	19.69			
			GCA (n=623)	21 (3.37)	176 (28.25)	426 (68.38)	17.50	0.091	0.182	0.86 (0.73 to 1.02)
			PMR+ (n=259)	10 (3.86)	79 (30.50)	170 (65.64)	19.11	0.756	0.756	0.96 (0.76 to 1.22)
			VIM+ (n=168)	7 (4.17)	47 (27.98)	114 (67.86)	18.15	0.497	0.807	0.90 (0.68 to 1.21)
			IOD+ (n=96)	5 (5.21)	24 (25.00)	67 (69.79)	17.71	0.500	0.822	0.88 (0.60 to 1.28)

*All p values have been calculated for the allelic model.

†Benjamini and Hochberg step-up FDR control.

‡OR for the minor allele.

FDR, false discovery rate; GCA, giant cell arteritis; IOD, irreversible occlusive disease; MAF, minor allele frequency; PMR, polymyalgia rheumatica; SNP, single-nucleotide polymorphism; VIM, visual ischaemic manifestations.

Basic and translational research

Table 2 Replication and pooled analysis of the *PTPN22* rs2476601 variant in Caucasian biopsy-proven GCA patients and controls

Population	Subgroup (N)	Genotype, N (%)			MAF (%)	Allele test	
		1/1	1/2	2/2		p Value*	OR (CI 95%)†
Germany	Controls (n=937)	9 (0.98)	164 (17.94)	741 (81.07)	9.96	0.9280	0.97 (0.54 to 1.73)
	GCA (n=72)	0 (0.00)	14 (19.44)	58 (80.56)	9.72		
Norway	Controls (n=271)	1 (0.37)	56 (20.66)	214 (78.97)	10.70	0.0376	1.77 (1.03 to 3.05)
	GCA (n=60)	1 (1.67)	19 (31.67)	40 (66.67)	17.50		
UK	Controls (n=5199)	42 (0.80)	933 (17.95)	4224 (81.25)	9.78	0.0319	1.44 (1.03 to 2.00)
	GCA (n=156)	3 (1.92)	36 (23.08)	117 (75.00)	13.46		
Replication meta-analysis‡	Controls (n=6384)	52 (0.81)	1153 (18.06)	5179 (81.12)	9.84	0.0154	1.38 (1.07 to 1.77)
	GCA (n=288)	4 (1.39)	69 (23.96)	215 (74.65)	13.37		
Overall meta-analysis§	Controls (n=8113)	65 (0.80)	1353 (16.68)	6695 (82.52)	9.14	2.00E-06	1.51 (1.28 to 1.79)
	GCA (n=911)	8 (0.88)	188 (20.64)	715 (78.49)	11.20		

*All p values have been calculated for the allelic model.

†OR for the minor allele.

‡Including independent cohorts from Germany, Norway and UK.

§Including independent cohorts from Spain, Germany, Norway and UK.

GCA, giant cell arteritis.

association of the *PTPN22* rs2476601/R620W*A allele with GCA was observed ($P_{FDR}=1.06E-04$, $OR=1.62$, $CI\ 95\% 1.29$ to 2.04). Subsequently, to examine whether *PTPN22* and *CSK* polymorphisms might influence the clinical manifestations of the disease, GCA patients were stratified according to the presence of PMR, VIM and IOD (table 1). Consistently, the subphenotype analysis also reached statistical significance for the rs2476601 polymorphism (PMR+ vs controls: $P_{FDR}=9.02E-04$, $OR=1.77$, $CI\ 95\% 1.30$ to 2.40 ; VIM+ vs controls: $P_{FDR}=4.10E-03$, $OR=1.82$, $CI\ 95\% 1.27$ to 2.62 ; IOD+ vs controls: $P_{FDR}=2.19E-03$, $OR=2.14$, $CI\ 95\% 1.38$ to 3.33). However, no statistically significant differences between GCA patients with and without these clinical characteristics were observed (data not shown). No association with any other *PTPN22* and *CSK* genetic variants was evident either in the case/control or subphenotype analysis (table 1).

To follow-up the positive finding of an association between *PTPN22* rs2476601/R620W and GCA in the Spanish

population, we attempted to confirm the detected association in a replication set of three independent cohorts of Caucasian ancestry. No heterogeneity between the ORs from the three replication cohorts was evident by BD test ($p=0.05$), and therefore a combined meta-analysis was performed (table 2 and online supplementary table S2). Statistically significant differences were observed for the *PTPN22* rs2476601*A allele in the pooled analysis ($P_{MH}=0.0154$, $OR=1.38$, $CI\ 95\% 1.07$ to 1.77) (table 2). Subsequently, the overall meta-analysis including both the discovery and the three replication cohorts showed a consistent association between the *PTPN22* rs2476601*A variant and GCA ($P_{MH}=2.00E-06$, $OR=1.51$, $CI\ 95\% 1.28$ to 1.79 ; figure 1). Again, no significant differences were found when GCA patients with and without specific clinical features were compared (data not shown).

The comparisons of the different detected allelic combinations between cases and controls did not yield additional information (data not shown).




Figure 1 Forest plot showing the ORs and CIs of the *PTPN22* rs2476601 association in the discovery and replication cohorts. OR and CI were calculated under the fixed effect model.

DISCUSSION

Our data indicate, for the first time, an important role for *PTPN22* in the genetic susceptibility of GCA. The combined analysis of the four independent cohorts showed a strong association between the *PTPN22* rs2476601/R620W variant and this disease. The effect size detected in our study (OR=1.51) is similar to that described for other autoimmune conditions, such as rheumatoid arthritis (OR=1.45), SLE (OR=1.45) or type 1 diabetes mellitus (OR>1.80)^{6,7} and, interestingly, for other vasculitides, such as Behçet's disease (OR>2.0) or antineutrophil cytoplasmic antibodies-associated vasculitis (OR>1.90).^{18,19} Despite this, a previous study failed to show association between *PTPN22* rs2476601 and GCA²⁰; however, it should be noted that the statistical power of this study was compromised because of the small sample size included in this report (96 GCA cases and 229 controls). In the subphenotype analysis, no specific association with any analysed clinical feature was observed, indicating that this variant may represent a risk factor for the global disease. Nevertheless, this should be taken with caution because of the low statistical power, which was a limitation of this stratified analysis.

Regarding *CSK*, our analysis had enough statistical power to detect a possible weak signal (power > 80% to detect an OR>1.25 in the discovery cohort); therefore, it is unlikely that *CSK* may play an important role in GCA susceptibility. Since an association between *PTPN22* and GCA was observed, it makes sense that its interacting partner, *CSK*, may also play a role in this pathology, but in most of the diseases in which an involvement of *PTPN22* has been described, an association with *CSK* has not been reported. Nevertheless, an effect of other *CSK* polymorphisms, showing low linkage disequilibrium with those analysed in our study, in GCA susceptibility cannot be discarded.

Initially, the *PTPN22* allele rs2476601*A, located within a protein-protein interaction domain, was reported as a gain-of-function allele that causes a decrease in TCR signalling.¹² However, a recent study has reported that this variant is a loss-of-function allele, leading to an accelerated degradation of lyp that results in enhanced signalling in several immune cell types.¹³ Although the mechanisms underlying the role of the *PTPN22* rs2476601 genetic variant in autoimmunity remain unclear, the association of this SNP with GCA suggests that a deregulation of TCR signalling is involved in the pathophysiological mechanisms of this vasculitis.

Author affiliations

- ¹Instituto de Parasitología y Biomedicina López-Neyra, CSIC, Granada, Spain
- ²NIHR-Leeds Musculoskeletal Biomedical Research Unit, Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, West Yorkshire, UK
- ³Department of Internal Medicine, Hospital Vall d'Hebron, Barcelona, Spain
- ⁴Department of Rheumatology, Hospital Xeral-Calde, Lugo, Spain
- ⁵Department of Autoimmune and Systemic Diseases, Vasculitis Research Unit, Hospital Clínic, University of Barcelona, Centre de Recerca Biomèdica Cellex (IDIBAPS), Barcelona, Spain
- ⁶Department of Rheumatology, Hospital de la Princesa, IIS-Princesa, Madrid, Spain
- ⁷Department of Rheumatology, Hospital Clínic San Carlos, Madrid, Spain
- ⁸Department of Rheumatology, Hospital Universitario de Bellvitge-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain
- ⁹Department of Rheumatology, Hospital Universitario Marqués de Valdecilla, IFIMAV, Santander, Spain
- ¹⁰Department of Internal Medicine, Complejo Hospitalario Universitario de Vigo, Spain
- ¹¹Department of Rheumatology, Hospital Ramón y Cajal, Madrid, Spain
- ¹²Department of Rheumatology, Grup de recerca cel·lular en inflamació i cartilag. IMIM (Institut de Recerca Hospital del Mar), Barcelona, Spain
- ¹³Department of Internal Medicine, Hospital Clínic San Cecilio, Granada, Spain
- ¹⁴Department of Internal Medicine, Hospital de Galdakano, Vizcaya, Spain
- ¹⁵Department of Internal Medicine, Corporació Sanitària Parc Taulí, Institut Universitari Parc Taulí, UAB, Sabadell, Barcelona, Spain
- ¹⁶Department of Rheumatology, Hospital Universitario 12 de Octubre, Madrid, Spain
- ¹⁷Department of Rheumatology, Hospital Universitario de La Paz, Madrid, Spain

¹⁸Department of Rheumatology, Hospital Clínic Universitario San Cecilio, Granada, Spain

¹⁹Department of Internal Medicine, Division of Nephrology, Robert-Bosch-Hospital, Stuttgart, Germany

²⁰Department of Rheumatology, Oslo University Hospital, Oslo, Norway and Institute of Clinical Medicine, University of Oslo, Oslo, Norway

²¹Department of Medical Genetics, University of Oslo and Oslo University Hospital, Oslo, Norway

²²Department of Immunology, Oslo University Hospital and University of Oslo, Oslo, Norway

²³Department of Clinical Immunology and Rheumatology, University of Luebeck, Bad Bramstedt, Germany

²⁴Hannover Medical School, Hannover, Germany

Acknowledgements The authors thank Sofia Vargas, Sonia García and Gema Robledo for their excellent technical assistance, and all the patients and healthy controls for kindly accepting their essential collaboration. Banco Nacional de ADN (University of Salamanca, Spain) is thanked for supplying part of the control material. The Norwegian Systemic Vasculitis and Connective Tissue Disease Registry (NOSVAR) at Oslo University Hospital is acknowledged for providing data on the Norwegian patients and the Norwegian Bone Marrow Donor Registry is acknowledged for providing the Norwegian controls.

Contributors FDC, AWM, MAG-G and JM were involved in the conception and design of the study. AS, AM, SLM and FDC contributed in the analysis and interpretation of data. AM and AS drafted the manuscript. RS, JAM-F, JH-R, MCC, SC, ICM, JN, RB, BS, MJG-V, JM, NO-C, AU, BM-A, JSM, EdM, CM, ER, NB, JL, OM, BAL, FM, TW, AWM and MAG-G collected samples and participated in analysis and interpretation of data. FDC, SLM, JM, MAG-G, RS, JAM-F, JH-R, MCC, SC, ICM, JN, RB, BS, MJG-V, JM, NO-C, AU, BM-A, JSM, EDM, CM, ER, NB, JL, OM, BAL, FM, TW and AWM revised critically the manuscript draft. All authors approved the final version of the manuscript.

Funding This study was supported by 'Fondo de Investigaciones Sanitarias' through grants PI06-0024 and PS09/00748 (Spain), and partially supported by RETICS Program RD08/0075 (RIER) from 'Instituto de Salud Carlos III' (ISCIII). FDC was supported by Consejo Superior de Investigaciones Científicas (CSIC) through the program JAE-DOC. MCC and JH-R were granted by SAF 11/30073. TW was granted by DFG WI 1031/6.1. SLM was supported by a Clinical Lectureship from the National Institute for Health Research, UK, and by a grant from the Wellcome Trust/Academy of Medical Sciences.

Competing interests None.

Patient consent Obtained.

Ethics approval The Ethical Board of each participating centre.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Gonzalez-Gay MA, Vazquez-Rodriguez TR, Lopez-Diaz MJ, *et al*. Epidemiology of giant cell arteritis and polymyalgia rheumatica. *Arthritis Rheum* 2009;61:1454–61.
- 2 Ly KH, Regent A, Tamby MC, *et al*. Pathogenesis of giant cell arteritis: More than just an inflammatory condition? *Autoimmun Rev* 2010;9:635–45.
- 3 Deng J, Younge BR, Olshen RA, *et al*. Th17 and Th1 T-cell responses in giant cell arteritis. *Circulation* 2010;121:906–15.
- 4 Espigol-Frigolè G, Corbera-Bellalta M, Planas-Rigol E, *et al*. Increased IL-17A expression in temporal artery lesions is a predictor of sustained response to glucocorticoid treatment in patients with giant-cell arteritis. *Ann Rheum Dis* 2013;72:1481–7.
- 5 Levinson NM, Seeliger MA, Cole PA, *et al*. Structural basis for the recognition of c-Src by its inactivator Csk. *Cell* 2008;134:124–34.
- 6 Bottini N, Musumeci L, Alonso A, *et al*. A functional variant of lymphoid tyrosine phosphatase is associated with type I diabetes. *Nat Genet* 2004;36:337–8.
- 7 Orozco G, Sanchez E, Gonzalez-Gay MA, *et al*. Association of a functional single-nucleotide polymorphism of *PTPN22*, encoding lymphoid protein phosphatase, with rheumatoid arthritis and systemic lupus erythematosus. *Arthritis Rheum* 2005;52:219–24.
- 8 Diaz-Gallo LM, Gourh P, Broen J, *et al*. Analysis of the influence of *PTPN22* gene polymorphisms in systemic sclerosis. *Ann Rheum Dis* 2011;70:454–62.
- 9 Rodriguez-Rodriguez L, Taib WR, Topless R, *et al*. The *PTPN22* R263Q polymorphism is a risk factor for rheumatoid arthritis in Caucasian case-control samples. *Arthritis Rheum* 2011;63:365–72.
- 10 Manjarrez-Orduno N, Marasco E, Chung SA, *et al*. *CSK* regulatory polymorphism is associated with systemic lupus erythematosus and influences B-cell signaling and activation. *Nat Genet* 2012;44:1227–30.
- 11 Martin JE, Broen JC, Carmona FD, *et al*. Identification of *CSK* as a systemic sclerosis genetic risk factor through Genome Wide Association Study follow-up. *Hum Mol Genet* 2012;21:2825–35.

Basic and translational research

- 12 Vang T, Congia M, Macis MD, *et al.* Autoimmune-associated lymphoid tyrosine phosphatase is a gain-of-function variant. *Nat Genet* 2005;37:1317–19.
- 13 Zhang J, Zahir N, Jiang Q, *et al.* The autoimmune disease-associated PTPN22 variant promotes calpain-mediated Lyp/Pep degradation associated with lymphocyte and dendritic cell hyperresponsiveness. *Nat Genet* 2011;43:902–7.
- 14 Huffmeier U, Steffens M, Burkhardt H, *et al.* Evidence for susceptibility determinant(s) to psoriasis vulgaris in or near PTPN22 in German patients. *J Med Genet* 2006;43: 517–22.
- 15 Morgan AW, Thomson W, Martin SG, *et al.* Reevaluation of the interaction between HLA-DRB1 shared epitope alleles, PTPN22, and smoking in determining susceptibility to autoantibody-positive and autoantibody-negative rheumatoid arthritis in a large UK Caucasian population. *Arthritis Rheum* 2009;60:2565–76.
- 16 Hunder GG, Bloch DA, Michel BA, *et al.* The American College of Rheumatology 1990 criteria for the classification of giant cell arteritis. *Arthritis Rheum* 1990;33:1122–8.
- 17 Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* 1995;57:289–300.
- 18 Baranathan V, Stanford MR, Vaughan RW, *et al.* The association of the PTPN22 620W polymorphism with Behcet's disease. *Ann Rheum Dis* 2007;66:1531–3.
- 19 Martorana D, Maritati F, Malerba G, *et al.* PTPN22 R620W polymorphism in the ANCA-associated vasculitides. *Rheumatology (Oxford)* 2012;51:805–12.
- 20 Gonzalez-Gay MA, Oliver J, Orozco G, *et al.* Lack of association of a functional single nucleotide polymorphism of PTPN22, encoding lymphoid protein phosphatase, with susceptibility to biopsy-proven giant cell arteritis. *J Rheumatol* 2005;32:1510–12.



Identification of the *PTPN22* functional variant R620W as susceptibility genetic factor for giant cell arteritis

A Serrano, A Márquez, S L Mackie, F D Carmona, R Solans, J A Miranda-Filloo, J Hernández-Rodríguez, M C Cid, S Castañeda, IC Morado, J Narváez, R Blanco, B Sopeña, M J García-Villanueva, J Monfort, N Ortego-Centeno, A Unzurrunzaga, B Marí-Alfonso, J Sánchez-Martín, E de Miguel, C Magro, E Raya, N Braun, J Latus, O Molberg, B A Lie, F Moosig, T Witte, A W Morgan, M A González-Gay, J Martín and UK GCA Consortium Spanish GCA Consortium

Ann Rheum Dis 2013 72: 1882-1886 originally published online August 14, 2013

doi: 10.1136/annrheumdis-2013-203641

Updated information and services can be found at:
<http://ard.bmj.com/content/72/11/1882>

These include:

Supplementary Material

Supplementary material can be found at:
<http://ard.bmj.com/content/suppl/2013/08/14/annrheumdis-2013-203641.DC1.html>

References

This article cites 20 articles, 8 of which you can access for free at:
<http://ard.bmj.com/content/72/11/1882#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Immunology \(including allergy\)](#) (4772)
[Vasculitis](#) (279)
[Genetics](#) (912)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>