



Killing Vector Fields and Holonomy Algebras

Author(s): Carlos Currás-Bosch

Source: *Proceedings of the American Mathematical Society*, Vol. 90, No. 1 (Jan., 1984), pp. 97-102

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/2044677>

Accessed: 06/02/2009 04:48

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

KILLING VECTOR FIELDS AND HOLONOMY ALGEBRAS

CARLOS CURRÁS-BOSCH

ABSTRACT. We prove that for each Killing vector field X on a complete Riemannian manifold, whose orthogonal distribution is involutive, the $(1,1)$ skew-symmetric operator A_X associated to X by $A_X = L_X - \nabla_X$ lies in the holonomy algebra at each point. By using the same techniques, we also study when that operator lies in the infinitesimal and local holonomy algebras respectively.

Introduction. Kostant (see [2 or 3]) proved that if X is a Killing vector field on a compact Riemannian manifold M , for each point x of M , $(A_X)_x$ lies in the holonomy algebra $\mathfrak{G}(x)$, A_X is associated to X by $A_X = L_X - \nabla_X$ (L is the Lie derivative and ∇ the Riemannian connection). The proof is based on a decomposition, $A_X = S_X + B_X$, where $(S_X)_x \in \mathfrak{G}(x)$ and $(B_X)_x \in \mathfrak{G}(x)^\perp$, where $\mathfrak{G}(x)^\perp$ is the orthogonal complement of $\mathfrak{G}(x)$ in the algebra $E(x)$ of skew-symmetric endomorphisms of $T_x(M)$, with respect to the inner product given by $(A, B) = -\text{trace}(A \circ B)$. One observes that B_X is parallel and as $\text{div}(B_X X) = -\text{trace}(B_X^2)$, if M is compact it results in $B_X = 0$.

In this paper we are concerned with noncompact and complete Riemannian manifolds and a Killing vector field X whose orthogonal distribution is involutive. We prove in §2 that $(A_X)_x \in \mathfrak{G}(x)$, $\forall x \in M$. We apply the techniques used in this proof to study in §3 when $(A_X)_x$ lies in $\mathfrak{G}'(x)$ and $\mathfrak{G}^*(x)$ (infinitesimal and local holonomy algebras respectively).

1. Properties of some distributions. Throughout this paper we suppose that M is a complete, connected Riemannian manifold of dimension $n + 1$, with metric g and Riemannian connection ∇ . We set $w = i_X g$, where X is a Killing vector field.

LEMMA 1.1. $dw(Y, Z) = -g(A_X Y, Z)$.

PROOF.

$$\begin{aligned} 2dw(Y, Z) &= Y(w(Z)) - Z(w(Y)) - w([Y, Z]) \\ &= Y(g(X, Z)) - Z(g(X, Y)) - g(X, [Y, Z]) \\ &= g(\nabla_Y X, Z) + g(X, \nabla_Y Z) - g(\nabla_Z X, Y) \\ &\quad - g(X, \nabla_Z Y) - g(X, \nabla_Y Z - \nabla_Z Y) \\ &= g(\nabla_Y X, Z) - g(\nabla_Z X, Y). \end{aligned}$$

Received by the editors July 2, 1982 and, in revised form, April 6, 1983. Partially presented in September 1981 to "VI Congrès du Groupement des Mathématiciens d'Expression Latine".

1980 *Mathematics Subject Classification*. Primary 53C20.

Key words and phrases. Complete Riemannian manifolds, Killing vector fields, holonomy algebras.

©1984 American Mathematical Society

0002-9939/84 \$1.00 + \$.25 per page

As $A_X U = -\nabla_U X$ for each vector field U , and A_X is skew-symmetric, we have $2dw(Y, Z) = -2g(A_X Y, Z)$.

We assume that the orthogonal distribution to X is involutive; i.e., there exists a 1-form ϕ such that $dw = \phi \wedge w$.

We can assume that dw is nowhere zero because if $(A_X)_x = 0$, at some point x of M , $(B_X)_x = 0$ and as B_X is parallel, $B_X = 0$.

We also assume that X is nowhere zero, because we consider its orthogonal distribution, but as is well known, if a vector bundle has a nowhere zero cross-section then its Euler class must be zero (see [1, Theorem 8.3, p. 242]). So from now on, the manifolds to consider must have Euler class zero.

Let \mathcal{P} be the involutive orthogonal distribution to X , and let $\{Z_1, \dots, Z_n\}$ be a local orthonormal basis of \mathcal{P} near x , so that throughout that neighborhood $dw = \lambda z^1 \wedge w$, where $\{(1/\|X\|)w; z^1, \dots, z^n\}$ is the dual basis of $\{(1/\|X\|)X; Z_1, \dots, Z_n\}$.

In such a basis the matrix of A_X takes the form

$$\begin{bmatrix} 0 & -\lambda\|X\| & 0 & \dots & 0 \\ \lambda\|X\| & 0 & \cdot & \dots & 0 \\ 0 & 0 & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \\ 0 & 0 & \cdot & \dots & 0 \end{bmatrix}.$$

From A_X we know all the curvature transformations $R(X, Z_i)$ ($1 \leq i \leq n$) because $R(X, Z_i) = \nabla_{Z_i} A_X$.

We can observe

LEMMA 1.2. *The integral manifolds of \mathcal{P} are all totally geodesic.*

PROOF. $g(\nabla_{Z_i} Z_j, X) = -g(Z_j, \nabla_{Z_i} X) = g(Z_j, A_X Z_i) = 0$ ($1 \leq i, j \leq n$).

LEMMA 1.3. *$\{Z_2, \dots, Z_n\}$ and $\{X, Z_1\}$ are involutive distributions.*

PROOF. Consider the indices α, β ; $2 \leq \alpha, \beta \leq n$. We already know that

$$g(X, \nabla_{Z_\alpha} Z_\beta) = 0.$$

Now,

$$\begin{aligned} \lambda\|X\|^2 g(Z_1, \nabla_{Z_\alpha} Z_\beta) &= g(A_X X, \nabla_{Z_\alpha} Z_\beta) = -g(\nabla_{Z_\alpha}(A_X X), Z_\beta) \\ &= -g((\nabla_{Z_\alpha} A_X)(X), Z_\beta) - g(A_X(\nabla_{Z_\alpha} X), Z_\beta) \\ &= -g(R(X, Z_\alpha)X, Z_\beta). \end{aligned}$$

So we have

(1) $g(A_X X, \nabla_{Z_\alpha} Z_\beta) = -g(R(X, Z_\alpha)X, Z_\beta)$ and

(2) $g(A_X X, \nabla_{Z_\beta} Z_\alpha) = -g(R(X, Z_\beta)X, Z_\alpha)$.

Subtracting (2) from (1) gives us $g(A_X X, [Z_\alpha, Z_\beta]) = 0$.

To prove that $\{X, Z_1\}$ is involutive, we consider

$$L_X(A_X X) = L_X(\lambda\|X\|^2 Z_1) = X(\lambda\|X\|^2)Z_1 + \lambda\|X\|^2 L_X Z_1 = 0$$

because $L_X(A_X X) = L_X(-\nabla_X X) = 0$, so $L_X Z_1 = 0$.

It is well known (see [2, Corollary 4.3, p. 246]) that if X is an infinitesimal affine transformation, then at each $x \in M$, $(A_X)_x$ belongs to the normaliser of $\mathfrak{G}(x)$, $N(\mathfrak{G}(x))$. The same result is true for $\mathfrak{G}'(x)$ as can be proved easily, using that $\mathfrak{G}'(x)$ is spanned by all linear endomorphisms of $T_x(M)$ of the form $(\nabla^k R)(Y, Z; V_1, \dots, V_k)$, where $Y, Z; V_1, \dots, V_k \in T_x(M)$ and $0 \leq k < \infty$.

It is not difficult to give some examples of Riemannian manifolds with a Killing vector field X , such that the condition $d\omega = \phi \wedge \omega$ is verified. We give here three examples such that:

- (i) the orbits of X are all diffeomorphic to R ;
- (ii) the orbits of X are all diffeomorphic to S^1 ;
- (iii) the orbits of X are solenoids on a torus.

(i) Let (N, g') be a complete Riemannian manifold. We consider $R \times N$. Let t be the parameter in R given by the identity map and $X = d/dt$.

Take in $R \times N$ the Riemannian metric g such that its restriction to $T(N)$ is g' , X is orthogonal to $T(N)$ at all points and $\|X\|$ is given by a definite positive function on N . Take $w_X = i_X g$. It is verified that $d\omega_X = \phi \wedge w_X$ and the vector field Z_1 considered above is $\gamma \cdot \text{grad}(\|X\|)$.

(ii) Taking S^1 instead of R , by the same argument, we obtain a Killing vector field with $d\omega = \phi \wedge \omega$, and now the orbits are all diffeomorphic to S^1 .

(iii) Taking $S^1 \times S^1 \times N$, we associate Y to the first S^1 factor and Z to the second S^1 factor. We define g such that $g(Y, Z) = 0$ and $g(Y, Y) = g(Z, Z)$ is given by a definite positive function on N ; Y and Z are orthogonal to $T(N)$, and g restricted to $T(N)$ is g' . Now it is verified that $d\omega_Y = \phi \wedge w_Y$, $d\omega_Z = \phi \wedge w_Z$, ϕ is the same for w_Y and w_Z ; so the same condition is verified considering $X = \alpha Y + \beta Z$, α and β real numbers; then taking α and β such that $\alpha/\beta \notin Q$, it is verified that $d\omega_X = \phi \wedge w_X$ and the orbits of X are solenoids on a torus.

It should be pointed out that these three examples give us the three possible types of orbit for Killing vector fields on noncompact manifolds (take N noncompact in (ii) and (iii)).

2. $(A_X)_x$ belongs to $\mathfrak{G}(x)$. We calculate $R(X, Z_i)$ ($1 \leq i \leq n$). We begin by the following observations.

$$A_X X = \lambda \|X\|^2 Z_1, \text{ as we proved in §1, so } \nabla_X X = -\lambda \|X\|^2 Z_1, \text{ and } \nabla_X((1/\|X\|)X) = (1/\|X\|)\nabla_X X = -\lambda \|X\| Z_1.$$

As the integral manifolds of \mathfrak{P} are totally geodesic, $\nabla_{Z_i}((1/\|X\|)X) = 0$, so for $i = 1$,

$$\begin{aligned} \nabla_{Z_1}((1/\|X\|)X) &= Z_1(1/\|X\|)X + (1/\|X\|)\nabla_{Z_1}X \\ &= Z_1(1/\|X\|)X - (1/\|X\|)A_X Z_1 \\ &= Z_1(1/\|X\|)X + (1/\|X\|)\lambda X = 0. \end{aligned}$$

For $Z_\alpha, 2 \leq \alpha \leq n$, we have

$$\nabla_{Z_\alpha}((1/\|X\|)X) = 0, \text{ so } Z_\alpha(\|X\|) = 0.$$

As $R(X, Z_i) = \nabla_{Z_i} A_X$, in the basis $\{(1/\|X\|)X, Z_1, \dots, Z_n\}$, we obtain

$$R(X, Z_i) = \begin{bmatrix} 0 & -Z_i(\lambda\|X\|) & -\lambda\|X\|\Gamma_{i1}^2 & \cdots & -\lambda\|X\|\Gamma_{i1}^n \\ Z_i(\lambda\|X\|) & 0 & 0 & \cdots & 0 \\ \lambda\|X\|\Gamma_{i1}^2 & 0 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \lambda\|X\|\Gamma_{i1}^n & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

In order to simplify the notations, we let

$$\eta = \begin{bmatrix} 0 & -a & -a_2 & \cdots & -a_n \\ a & 0 & 0 & \cdots & 0 \\ a_2 & 0 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ a_n & 0 & 0 & \cdots & 0 \end{bmatrix}, \quad \kappa = \begin{bmatrix} 0 & -b & 0 & \cdots & 0 \\ b & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

$R(X, Z_i)$ is of the form η and belongs to $\mathfrak{G}(x)$, while A_X is of the form κ and belongs to $N(\mathfrak{G}(x))$.

We calculate the elements of $\mathfrak{G}(x)$, $[\kappa, \eta]$ and $[\kappa, [\kappa, \eta]]$.

$$[\kappa, \eta] = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & -ba_2 & \cdots & -ba_n \\ 0 & ba_2 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ 0 & ba_n & 0 & \cdots & 0 \end{bmatrix},$$

$$[\kappa, [\kappa, \eta]] = \begin{bmatrix} 0 & 0 & b^2a_2 & \cdots & b^2a_n \\ 0 & 0 & 0 & \cdots & 0 \\ -b^2a_2 & 0 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ -b^2a_n & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

If $a \neq 0$, there exist real numbers A, B such that $\eta = A\kappa + (1/B^2)[\kappa, [\kappa, \eta]]$, so $\kappa = (1/A)(\eta - (1/B^2)[\kappa, [\kappa, \eta]])$, whence the right side clearly belongs in $\mathfrak{G}(x)$.

If $a = 0$ and at least one of a_2, \dots, a_n is $\neq 0$, we calculate $[\eta, [\kappa, \eta]] \in \mathfrak{G}(x)$ and obtain

$$[\eta, [\kappa, \eta]] = \begin{bmatrix} 0 & -b(a_2^2 + \dots + a_n^2) & 0 & \dots & 0 \\ b(a_2^2 + \dots + a_n^2) & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \\ 0 & 0 & 0 & \dots & 0 \end{bmatrix}.$$

Therefore, there exists a real number C such that $\kappa = C[\eta, [\kappa, \eta]]$.

If $a = a_2 = \dots = a_n = 0$, then this means that at all points of M , $i_x R = 0$. Hence $\Gamma_{11}^\alpha = \Gamma_{\alpha\beta}^1 = 0$, for all $\alpha, \beta \in \{2, \dots, n\}$. In this case the distributions $\{X, Z_1\}$ and $\{Z_2, \dots, Z_n\}$ are parallel, so that the elements of $\mathfrak{G}(x)$ are of the form

$$\left[\begin{array}{cc|ccc} \Lambda_1 & & 0 & \dots & 0 \\ & & 0 & \dots & 0 \\ \hline 0 & 0 & & & \\ \cdot & \cdot & & & \\ \cdot & \cdot & & \Lambda_2 & \\ \cdot & \cdot & & & \\ 0 & 0 & & & \end{array} \right],$$

where Λ_1 is a 2×2 matrix and Λ_2 is $(n - 1) \times (n - 1)$.

At each point x , $R(X, Z_1) = 0$, $\Lambda_1 = 0$; so, $(A_X)_x \in (\mathfrak{G}(x))^\perp$ and $A_X = B_X$.

In this case ψ_x^0 (the restricted holonomy group) is reducible and $T_x(M)$ has an invariant subspace on which ψ_x^0 acts trivially, of dimension ≥ 2 , which contains X at all points. Let \tilde{M} be the universal covering manifold of M with the induced Riemannian metric. The holonomy group of \tilde{M} at \tilde{x} ($\pi(\tilde{x}) = x$) is ψ_x^0 and by the De Rham Decomposition Theorem, we have $\tilde{M} = R^{2+q} \times N$, where in the R^{2+q} factor we have the Killing vector field induced by X , which we also represent by X , such that $dw = \phi \wedge w$. But this cannot occur in R^{2+q} , unless X is parallel and $A_X = 0$.

So we have

THEOREM 2.1. *Let (M, g) be a complete and connected Riemannian manifold. Let X be a Killing vector field on M , such that $dw = \phi \wedge w$, for $w = i_x g$. Then, $(A_X)_x \in \mathfrak{G}(x)$, for all x in M .*

3. Local and infinitesimal holonomy algebras. Throughout this section we continue to assume that $dw = \phi \wedge w$.

LEMMA 3.1. *If $(i_x R)_x \neq 0$, $(A_X)_x \in \mathfrak{G}'(x)$.*

PROOF. There exists some i , $1 \leq i \leq n$, such that $R(X, Z_i) \neq 0$, so in $\mathfrak{G}'(x)$ there is some element of the form η , and as $(A_X)_x$ is of the form $\kappa \in N(\mathfrak{G}'(x))$, by using the same method as in §2, we obtain $(A_X)_x \in \mathfrak{G}'(x)$.

We study which of the above mentioned holonomy algebras contains A_X .

We can assume that $A_X \neq 0$, so $X \neq 0$, because we have supposed $dw = \phi \wedge w$.

Let $U = \{x \in M \mid (i_X R)_x \neq 0\}$. U is an open set and if $x \in M \overset{\circ}{=} U$, there exists an open neighborhood of x , V_x , where $i_X R \neq 0$. In V_x , $\{X, Z_1\}$ and $\{Z_2, \dots, Z_n\}$ are parallel distributions and, in the same way as in §2, we can prove that $(A_X)_x \in (\mathfrak{G}^*(x))^\perp$, $\psi^*(x)$ (the local holonomy group) is reducible and $T_x(M)$ has an invariant subspace on which $\psi^*(x)$ acts trivially, of dimension ≥ 2 .

If $x \in \partial(U)$, the boundary of U , we consider an open neighborhood of x , W_x , such that for each $y \in W_x$ and the parallel transport τ along any curve in W_x from x to y , we have $\tau^{-1}\mathfrak{G}^*(y)\tau \subset \mathfrak{G}^*(x)$. Such W_x exists by Proposition 10.1, Chapter II of [2].

Let $\{y_n\}$ be a sequence $\{y_n\} \rightarrow x$ and $y_n \in W_x \cap U$, $\forall n$; we have $(A_X)_{y_n} \in \mathfrak{G}'(y_n) \subset \mathfrak{G}^*(y_n)$. Let τ_n be the parallel transport along the minimizing geodesic from x to y_n (we can take W_x convex), then $\tau_n^{-1}\mathfrak{G}^*(y_n)\tau_n \subset \mathfrak{G}^*(x)$, and as $(A_X)_{y_n} \in \mathfrak{G}'(y_n) \subset \mathfrak{G}^*(y_n)$, we obtain $\tau_n^{-1}(A_X)_{y_n}\tau_n \in \mathfrak{G}^*(x)$; so $(A_X)_x \in \mathfrak{G}^*(x)$.

We have proved

THEOREM 3.1. *There exists an open set U , defined by $U = \{x \in M \mid (i_X R)_x \neq 0\}$, such that:*

- (a) if $x \in U$, $(A_X)_x \in \mathfrak{G}'(x)$,
- (b) if $x \in M \overset{\circ}{=} U$, $(A_X)_x \in (\mathfrak{G}^*(x))^\perp$,
- (c) if $x \in \partial(U)$, $(A_X)_x \in \mathfrak{G}^*(x)$.

REFERENCES

1. D. Husemoller, *Fibre bundles*, 2nd ed., Graduate Texts in Math., Vol. 20, Springer-Verlag, Berlin and New York, 1975.
2. S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Vol. I, Interscience, New York, 1963.
3. B. Kostant, *The holonomy and the Lie algebra of infinitesimal motions of a Riemannian manifold*, Trans. Amer. Math. Soc. **80** (1955), 528–542.

DEPARTAMENTO GEOMETRÍA Y TOPOLOGÍA, FACULTAD MATEMÁTICAS, UNIVERSIDAD DE BARCELONA, BARCELONA, SPAIN