

Manuscript Number: TIV-D-15-00229R3

Title: Protection against Oxidative Damage in Human Erythrocytes and Preliminary Photosafety Assessment of Punica granatum Seed Oil Nanoemulsions Entrapping Polyphenol-rich Ethyl Acetate Fraction

Article Type: Research Paper

Section/Category: Mechanisms

Keywords: Punica granatum; seed oil; polyphenol-rich ethyl acetate fraction; fluorescent probes, haemolysis; photohaemolysis.

Corresponding Author: Prof. Maria Pilar Vinardell,

Corresponding Author's Institution: Universitat de Barcelona

First Author: Thaisa Baccarin

Order of Authors: Thaisa Baccarin; Montserrat Mitjans; Elenara Lemos-Senna Lemos-Senna ; Maria Pilar Vinardell

Abstract: The main purpose of the present study is to evaluate the ability of nanoemulsion entrapping pomegranate peel polyphenol-rich ethyl acetate fraction (EAF) prepared from pomegranate seed oil and medium chain triglyceride to protect human erythrocyte membrane from oxidative damage and to assess preliminary in vitro photosafety. In order to evaluate the phototoxic effect of nanoemulsions, human red blood cells (RBCs) are used as a biological model and the rate of haemolysis and photohaemolysis (5 J cm⁻² UVA) is assessed in vitro. The level of protection against oxidative damage caused by the peroxy radical generator AAPH in human red blood cells (RBCs) as well as its effects on bilayer membrane characteristics such as fluidity, protein profile and RBCs morphology are determined. EAF-loaded nanoemulsions do not promote haemolysis or photohaemolysis. Anisotropy measurements show that nanoemulsions significantly restrain the increase in membrane fluidity caused by AAPH. SDS-PAGE analysis reveal that AAPH induced degradation of membrane proteins, but that nanoemulsions reduce the extend of degradation. Scanning electron microscopy examinations corroborate the interaction between AAPH, nanoemulsions and the RBCs membrane bilayer. Our work demonstrates that P. granatum nanoemulsions are photosafe and protect RBCs against oxidative damage and possible disturbance of the lipid bilayer of biomembranes. Moreover it suggests that these nanoemulsions could be promising new topical products to reduce the effects of sunlight on skin.

Suggested Reviewers: Qasim Chaudhry
Qasim.Chaudhry@fera.gsi.gov.uk
Expertise in nanomaterials

Adela Lopez de Cerain
acerain@unav.es
expertise in nanoformulations

Maria Dusinska
maria.dusinska@nilu.no
Expertise in nanomaterials

Björn Dahlbäck
bjorn.dahlback@med.lu.se

Reviewer #2:

We thank the authors for the response. I still think that the chemical characterization about the extract fraction should be included. The % distribution of the different polyphenolic compounds is important.

We added the information requested by the reviewer at section 4 (discussion) - page 24. We also added at section 2.2.1 (methods) that the EAF was characterized by HPLC and Folin-Ciocalteu methods. For more detailed information about the characterization methods the reader must look for our previously published work.

1 **Protection against Oxidative Damage in Human Erythrocytes**
2
3 **and Preliminary Photosafety Assessment of *Punica granatum* Seed**
4
5 **Oil Nanoemulsions Entrapping Polyphenol-rich Ethyl Acetate Fraction**
6
7
8
9

10 *Thaisa Baccarin*^{a,b}, *Montserrat Mitjans*^a, *Elenara Lemos-Senna*^b, *Maria Pilar*
11 *Vinardell*^{a*}
12
13
14
15

16 ^a Departament de Fisiologia, Facultat de Farmàcia, Universitat de Barcelona, Av.
17 Joan XXIII s/n, E-08028 Barcelona, Spain.
18
19

20
21
22 ^b Programa de Pós-Graduação em Farmácia, Centro de Ciências da Saúde,
23 Universidade Federal de Santa Catarina, Campus Universitário Trindade, 88040-970
24 Florianópolis, SC, Brazil.
25
26
27

28
29
30
31 Corresponding author email: mpvinardellmh@ub.edu. Departament de Fisiologia,
32 Facultat de Farmàcia, Universitat de Barcelona, Av. Joan XXIII s/n, E-08028
33 Barcelona, Spain. Phone number: +34 934 024505; Fax number: +34 934035901.
34
35
36
37
38
39

40 **Abstract**
41

42 The main purpose of the present study is to evaluate the ability of nanoemulsion
43 entrapping pomegranate peel polyphenol-rich ethyl acetate fraction (EAF) prepared
44 from pomegranate seed oil and medium chain triglyceride to protect human
45 erythrocyte membrane from oxidative damage and to assess preliminary *in vitro*
46 photosafety. In order to evaluate the phototoxic effect of nanoemulsions, human red
47 blood cells (RBCs) are used as a biological model and the rate of haemolysis and
48 photohaemolysis (5 J cm⁻² UVA) is assessed *in vitro*. The level of protection against
49 oxidative damage caused by the peroxy radical generator AAPH in human RBCs as
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 well as its effects on bilayer membrane characteristics such as fluidity, protein profile
2 and RBCs morphology are determined. EAF-loaded nanoemulsions do not promote
3 haemolysis or photohaemolysis. Anisotropy measurements show that nanoemulsions
4 significantly retrain the increase in membrane fluidity caused by AAPH. SDS-PAGE
5 analysis reveal that AAPH induced degradation of membrane proteins, but that
6 nanoemulsions reduce the extend of degradation. Scanning electron microscopy
7 examinations corroborate the interaction between AAPH, nanoemulsions and the
8 RBCs membrane bilayer. Our work demonstrates that *P. granatum* nanoemulsions
9 are photosafe and protect RBCs against oxidative damage and possible disturbance
10 of the lipid bilayer of biomembranes. Moreover it suggests that these nanoemulsions
11 could be promising new topical products to reduce the effects of sunlight on skin.

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 **Keywords:** *Punica granatum*, seed oil, polyphenol-rich ethyl acetate fraction,
29 fluorescent probes, haemolysis, photohaemolysis.
30
31
32
33
34

35 1. Introduction

36
37
38
39
40 The peel extract of *Punica granatum*, popularly known as pomegranate,
41 contains a large quantity of phenolic compounds such as ellagic acid, gallic acid,
42 punicalagin, punicalin, luteolin and others (Amakura et al., 2000; Seeram et al., 2005;
43 Van Elswijk et al., 2004). Pomegranate seed oil (PSO) contains conjugated fatty
44 acids, in which the main constituent is punიცic acid (El-Nemr et al., 2006; Fadavi et
45 al., 2006), that it has been reported to have antioxidant properties (Qu et al., 2010).
46 Phenolic compounds are natural antioxidants and are candidates for the ultraviolet
47 photoprotection and prevention of ultraviolet radiation-induced oxidative stress and
48 skin alteration, especially given their capacity to reduce the production of radicals,
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 and their ability to stabilize reactions induced by oxygen and its radical species
2
3 (González et al., 2013); promising applications of polyphenols exist in the cosmetic
4
5 and pharmaceutical industries.
6
7

8 In a previous study, we developed PSO and medium chain triglyceride (MCT)
9
10 nanoemulsion entrapping a pomegranate polyphenol-rich ethyl acetate fraction (EAF)
11
12 for UV photoprotection application. Nanoemulsions (NEs) have attracted
13
14 considerable attention for cosmetic and personal care as potential vehicles for
15
16 releasing and improving the permeation of active compounds through the skin
17
18 (Guglielmini, 2008). The *in vitro* antioxidant activity of free EAF and NE loaded with
19
20 EAF (EAF-PSO-NE and EAF-MCT-NE) was confirmed using the ferric reducing
21
22 antioxidant power (FRAP) and 2,2-diphenyl-1-picrylhydrazyl (DPPH) methods
23
24 (Baccarin and Lemos-Senna, 2014).
25
26
27
28
29

30 The assessment of the toxicity potential of new molecules and formulations for
31
32 human use is fundamental and it has become imperative for the nonclinical
33
34 toxicologist to establish new approaches for early screening of potential cosmetic and
35
36 pharmaceutical products candidates (Ahuja and Sharma, 2014). In particular,
37
38 phototoxicity is of increasing concern in dermatology because of the increased level
39
40 of ultraviolet radiation reaching the earth from the sun (Onoue et al., 2009).
41
42 Phototoxicity can be caused by several classes of pharmaceuticals and cosmetics,
43
44 which have the potential to provoke photoirritant, photoallergic and photogenotoxic
45
46 events in light-exposed tissues through oxidation and chemical modification under
47
48 exposure to sunlight (Stein et al., 2007; Elkeeb et al. 2012).
49
50
51
52
53

54 Although there is no single method that provides all the toxicity information
55
56 required, and since different nanosized systems elicit different biological responses,
57
58 during the process of determining and studying the toxicological behaviour of a
59
60
61
62
63
64
65

1 sample a combination of assays is often required. *In vitro* haemolysis is one of these
2
3 assays (Doktorovova et al., 2014) and evaluates the biocompatibility of nanodroplets
4
5 and formulation components (*i.e.* surfactants, lipid) (Schubert and Müller-Goymann,
6
7 2005; Söderlind and Karlsson, 2006) via the impact of their physico-chemical
8
9 characteristics on human red blood cells (RBCs), which is evaluated by quantifying
10
11 the release of haemoglobin (Arora et al., 2012).
12
13
14
15

16 Therefore, the aim of this work was to evaluate the potential protective effect
17
18 of free EAF, unloaded and EAF-loaded NEs against oxidative damage caused by the
19
20 peroxy radical generator 2,2'-azobis (amidinopropane dihydrochloride) (AAPH) in
21
22 human erythrocytes and also to identify any possible irritant reactions; the potential
23
24 irritancy and photo irritancy were assessed by *in vitro* haemolysis and
25
26 photohaemolysis, respectively. The erythrocyte was chosen as an *in vitro* model
27
28 since its membrane is rich in polyunsaturated fatty acids, which are extremely
29
30 susceptible to free radical-mediated peroxidation; besides, erythrocytes have no
31
32 internal organelles and because they are the simplest cell model available, they are
33
34 the most popular cell membrane system for verifying possible membrane interactions
35
36 and are considered to be representative of the plasma membrane in general
37
38 (Magalhães et al., 2009; Svetina et al., 2004). Oxidative damage of the erythrocyte
39
40 membrane may cause its malfunctioning by altering its fluidity and protein profile,
41
42 manifested by decreased reduction in cytoskeletal protein content (low-molecular-
43
44 weight, LMW) and the production of high-molecular-weight (HMW) proteins (Flynn et
45
46 al., 1983; Snyder et al., 1985) and abnormalities in erythrocyte shape (Yang et al.,
47
48 2006). Since these events have been proposed as a general mechanism leading to
49
50 the haemolysis involved in cell injury and death (Alvarez-Suarez et al., 2012), we
51
52 also evaluated them after RBCs were incubated with treatments.
53
54
55
56
57
58
59
60
61
62
63
64
65

1 To the best of our knowledge, this is the first time that the protective effect
2
3 against oxidative damage in human erythrocytes of NE entrapping *P. granatum* peel
4
5 polyphenol-rich EAF prepared using PSO or MCT as oil phases and their preliminary
6
7 photosafety have been assessed using the above mentioned assays.
8
9

10 11 12 13 **2. Materials and Methods**

14 15 16 17 **2.1. Materials**

18
19 Polysorbate 80 (Tween 80[®]), 2,2'-Azobis(2-methyl-propionamide) dihydro-
20 chloride (AAPH), triethanolamine, hydrogen peroxide 30% (w/w), sodium azide,
21 sodium dodecyl sulfate (SDS), Triton X-100 and glycine were purchased from Sigma-
22 Aldrich (St. Louis, MO, USA). NaCl, Na₂HPO₄ and KH₂PO₄ were purchased from
23 Merck (Darmstadt, Germany). Ethyl acetate, dichloromethane, acetic acid and
24 methanol were obtained from Vetec[®] (Rio de Janeiro, Brazil). Pomegranate seed oil
25 and pomegranate fruit peel dry extract were purchased from Via Farma (Sao Paulo,
26 Brazil). Soy lecithin (Lipoid[®] S100) was from Lipoid AG (Steinhausen, Switzerland).
27 Medium chain triglyceride was from Brasquim (Porto Alegre, Brazil). Fluorescent
28 probes DPH (1,6-diphenyl-1,3,5-hexatriene) and TMA-DPH (1-(4-trimethylammonium
29 phenyl)-6-phenyl-1,3,4-hexatriene *p*-toluenesulfonate) were purchased from
30 Molecular Probes (Eugene, OR). Acrylamide PAGE 40%, methylenebisacrylamide
31 2%, TEMED, mercaptoethanol, ammonium persulfate and bromophenol blue used
32 for SDS-PAGE were obtained from GE Healthcare Amersham Biosciences (Uppsala,
33 Sweden). Finally, water was purified in a Milli-Q system (Millipore, Bedford, MA).
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.2. Methods

2.2.1. Ethyl acetate fraction (EAF)

The pomegranate polyphenol-rich ethyl acetate fraction (EAF) was obtained from a commercial pomegranate peel dried extract and characterized through a high performance liquid chromatography (HPLC) method and spectrophotometric method (Folin-Ciocalteu) as described in our previously work (Baccarin and Lemos-Senna, 2014). Briefly, the dried powder was extracted by 24 hour dynamic maceration with a 90:10 (v/v) methanol:water mixture. The extractive solution was evaporated under reduced pressure for solvent removal, and the precipitate was suspended with a 2% aqueous acetic acid solution. The resulting mixture was then partitioned with dichloromethane and ethyl acetate. After that, the pooled ethyl acetate fraction was evaporated under reduced pressure to dryness.

2.2.2. Preparation of nanoemulsions

EAF-loaded pomegranate seed oil nanoemulsion (EAF-PSO-NE) or EAF-loaded medium chain triglyceride nanoemulsion (EAF-MCT-NE) were prepared using an ultrasonic emulsification method followed by solvent evaporation or spontaneous emulsification method, respectively (Baccarin and Lemos-Senna, 2014). Briefly, for the EAF-PSO-NE, the ethyl acetate fraction (EAF) (0.5%; w/v), soy lecithin (0.4%; w/v) and PSO (2%; w/v) were dissolved in 10 mL of ethyl acetate. This ethyl acetate solution was slowly poured into 40 mL of a polysorbate 80 (2.1%; w/v) aqueous solution. The oil in water dispersion was sonicated for 3 minutes using an Ultrasonic Processor UP200S (Hielscher, Germany), adjusted to pH 5.0-6.5 with triethanolamine (2%; w/v) and then it was and kept under magnetic stirring for 24

1 hours. The resulting nanoemulsion was evaporated under reduced pressure up to
2
3 volume of 15 mL.
4

5 For the EAF-MCT-NE, 10 mL of an ethanolic solution containing EAF (0.5%;
6 w/v), soy lecithin (0.4%; w/v), and MCT (1.8%, w/v) was poured into a 2.1% (w/v)
7 polysorbate 80 aqueous solution under magnetic stirring and adjusted to pH 5.0-6.5
8 with triethanolamine (2%; w/v). The nanoemulsion was then evaporated under
9 reduce pressure to eliminate the organic solvent and concentrated up to volume of
10 15 mL. All formulations were filtered through 8 μm quantitative filter paper. Unloaded
11 PSO-NE and MCT-NE were prepared in the same manner.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 *2.2.3. Physicochemical characterization of nanoemulsions*

26
27 The mean droplet size and zeta potential were determined by dynamic light
28 scattering (DLS) and laser-Doppler anemometry, respectively, using a Zetasizer
29 Nano Series (Malvern Instruments, Worcestershire, UK). The size analyses were
30 performed at a scattering angle of 173° , after appropriated dilution of the samples in
31 ultrapure water (Milli-Q®, Millipore, USA). The size analyses were
32 performed at a scattering angle of 173° , after appropriated dilution of the samples in
33 ultrapure water (Milli-Q®, Millipore, USA). For zeta potential analysis, the samples
34 were diluted in Milli-Q® water and placed in electrophoretic cells where a potential of
35 ± 150 mV was applied. The zeta potential values were calculated as mean
36 electrophoretic mobility values by using Smoluchowski's equation.
37
38
39
40
41
42
43
44
45
46
47
48

49 *2.2.4. Preparation of red blood cell suspensions*

50
51 Blood samples were obtained from anesthetized rats by cardiac puncture or
52 from healthy volunteers from the Blood Bank of "Hospital Clinic" (Barcelona, Spain).
53 All procedures followed the ethical guidelines of the University of Barcelona. Red
54 blood cells (RBCs) were isolated by centrifugation at 3000 rpm at 4 °C for 10
55
56
57
58
59
60
61
62
63
64
65

1 minutes, and washed three times in isotonic PBS containing 123.3 mM NaCl, 22.2
2
3 mM Na₂HPO₄ and 5.6 mM KH₂PO₄ in distilled water (pH 7.4).
4

5 6 2.2.5. *Haemolysis assay*

7
8 The membrane-lytic activity of EAF, NE and formulation components
9 (polysorbate 80, lecithin, MCT and PSO) was determined through haemolysis assay.
10 A 25 µL aliquot of rat or human erythrocyte suspension was exposed to each sample
11 at various concentrations: EAF (10-1000 µg mL⁻¹), polysorbate 80 (0.5-80 mg mL⁻¹),
12 lecithin (0.1-10 mg mL⁻¹), MCT (1-55 mg mL⁻¹), PSO (1-60 mg mL⁻¹), unloaded (MCT-
13 NE and PSO-NE) and EAF-loaded nanoemulsions (EAF-MCT-NE and EAF-PSO-NE)
14 (5-500 µg mL⁻¹) diluted in PBS solution in a total volume of 1 mL. Negative and
15 positive controls were prepared by resuspending erythrocyte in buffer alone or
16 distilled water, respectively. The samples were incubated at room temperature under
17 constant shaking for 10 min, and then centrifuged at 10,000 rpm for 5 min. The
18 absorbance of the haemoglobin released in the supernatant was measured at 540
19 nm using a Shimadzu UV-160A spectrophotometer (Shimadzu, Kyoto, Japan), and
20 the percentages of haemolysis were obtained by comparing the sample absorbance
21 with the positive control (totally haemolysed). Dose-response curves were plotted
22 from haemolysis results and the concentrations inducing 50% haemolysis (HC₅₀)
23 were calculated. (Nogueira et al., 2012a)
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 2.2.6. *Haemolysis induced by H₂O₂*

51 The haemolysis induced by H₂O₂ was measured as reported by Ugartondo
52 and collaborators (Ugartondo et al., 2009) with modifications. First, 250 µL of the
53 human erythrocyte suspension, previously incubated with sodium azide (2 mM), was
54 mixed with various concentrations of EAF (25-150 µg mL⁻¹), EAF-MCT-NE (25-100
55
56
57
58
59
60
61
62
63
64
65

1 $\mu\text{g mL}^{-1}$), EAF-PSO-NE (50-200 $\mu\text{g mL}^{-1}$). Then H_2O_2 was added and the mixture
 2
 3 incubated for 90 min at 37 °C under stirring. The volume of each sample was
 4
 5 adjusted to 1 mL with PBS. After final incubation, the samples were centrifuged at
 6
 7 10,000 rpm for 5 min, the supernatant was diluted 1:10 in PBS and the absorbance
 8
 9 of the haemoglobin released was measured at 540 nm using a Shimadzu UV-160A
 10
 11 spectrophotometer (Shimadzu, Kyoto, Japan). The percentages of haemolysis were
 12
 13 obtained by comparing the sample absorbance with the positive control (totally
 14
 15 haemolysed). The concentration inhibiting 50% of the haemolysis caused by H_2O_2
 16
 17 (IC_{50}) was determined for each sample. The experiment was carried out three times
 18
 19 using three replicate samples for each sample concentration tested.
 20
 21
 22
 23
 24
 25
 26

27 2.2.7. Photohaemolysis assay

28
 29 Twenty-five microliter of rat or human red blood cell suspension were added to
 30
 31 a 24-well plate containing 500 $\mu\text{g mL}^{-1}$ of EAF, EAF-MCT-NE, EAF-PSO-NE, MCT-
 32
 33 NE or PSO-NE. Then, one plate was exposed to UVA (TL-D 15 W/10 UVA lamp,
 34
 35 Royal Philips Eletronics-The Netherlands) and the other was kept in the dark.
 36
 37 Irradiance was measured using a Delta OHM photoradiometer (HD2302-Italy) to
 38
 39 determine the time of exposure, using the formula:
 40
 41
 42
 43
 44
 45
 46

$$47 \quad E (\text{J}/\text{cm}^2) = t(\text{s}) \times P(\text{W}/\text{cm}^2) \quad \text{Eq. 1}$$

48
 49 where E is UV dose, t is the time expressed in seconds and, finally, P is the lamp
 50
 51 potency. Cells were irradiated with 1.6-2.1 mW/cm^2 to give a final exposure of 5 J
 52
 53 cm^2 .
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

1 After irradiation the contents of each well were transferred to a tube and
2
3 centrifuged at 1000 rpm for 5 minutes. The absorbance of the supernatant was
4
5 measured at 525 nm using a Shimadzu UV-160A spectrophotometer (Shimadzu,
6
7 Kyoto, Japan). The HC₅₀ was determined for plates exposed and non-exposed to the
8
9 UVA irradiation. The photohaemolysis factor (PHF) was calculated by dividing the
10
11 HC₅₀ of the non-irradiated cells by the HC₅₀ of the irradiated cells. The haemoglobin
12
13 oxidation assay measured the formation of intracellular and extracellular
14
15 methaemoglobin. The same assay conditions as those of the photohaemolysis assay
16
17 were used, except that 100 µL of Triton X-100 1% (v/v) solution was added to each
18
19 irradiated and non-irradiated well and the absorbance of methaemoglobin was
20
21 measured at 630 nm. The amount of methaemoglobin in the sample was determined
22
23 by the difference in optical density (Δ OD) between the irradiated and non-irradiated
24
25 plate. A product is considered photoirritant if the PHF is ≥ 3.0 or if the Δ OD is ≥ 0.05 .
26
27 (Pape et al., 2001)
28
29
30
31
32
33
34
35
36

37 2.2.8. Haemolysis mediated by AAPH

38 The haemolysis mediated by AAPH followed a method previously described in
39
40 the literature, with some modifications (Miki et al., 1987). The addition of AAPH (a
41
42 peroxy radical initiator) to the human erythrocyte suspension induces the oxidation of
43
44 cell membrane lipids and proteins, resulting in haemolysis. Aliquots of 83 µL of the
45
46 human erythrocyte suspension were incubated with AAPH at a final concentration of
47
48 50 mM for 2.5 h in a shaker at 37 °C to achieve 100% haemolysis. The
49
50 antihaemolytic activity of EAF (0.5-50 µg mL⁻¹), MCT-NE (5-100 µg mL⁻¹), EAF-MCT-
51
52 NE (5-100 µg mL⁻¹), PSO-NE (5-100 µg mL⁻¹) and EAF-PSO-NE (5-100 µg mL⁻¹) was
53
54 tested by adding several concentrations of each sample to the erythrocyte
55
56
57
58
59
60
61
62
63
64
65

1 suspension in the presence of AAPH. A negative control (erythrocyte suspension
2 with PBS alone) was included in all experiments to monitor spontaneous haemolysis.
3
4 Samples plus erythrocyte suspension were also incubated without AAPH to
5
6 determine whether the sample alone could cause haemolysis during 2.5 h of
7
8 incubation. After incubation, the samples were centrifuged at 10,000 rpm for 5 min
9
10 and the absorbance of the supernatant was measured at 540 nm in a Shimadzu
11
12 spectrophotometer. The concentration inhibiting 50% of the haemolysis (IC₅₀) was
13
14 determined for each sample.
15
16
17
18
19
20
21
22

23 *2.2.9. Scanning electron microscopy - SEM*

24
25 After 2.5 h of incubation, unwashed samples were fixed by adding 1 mL of
26
27 5.0% glutaraldehyde in PBS solution and incubated at 4 °C for 2 h. The samples
28
29 were then centrifuged (1500 rpm for 10 min), the supernatant was discarded, and 1
30
31 mL of 2.5% glutaraldehyde in PBS was added. Fixed samples were washed with
32
33 PBS solution, post-fixed with 1% osmium tetroxide, placed over a glass coverslip,
34
35 dehydrated in an ascending series of ethyl alcohol (50–100%), air-dried using the
36
37 critical point drying method in a CPD 7501 apparatus (Polaron, Watford, UK), and
38
39 finally mounted on an aluminium stub and gold-coated in an SEM coating system SC
40
41 510 (Fisons Instruments, East Grinstead, UK). The resulting specimens were
42
43 examined in a Zeiss DSM 940A scanning electron microscope (Carl Zeiss SMT AG,
44
45 Jena, Germany).
46
47
48
49
50
51
52
53

54 *2.2.10. Erythrocyte ghost preparation*

55
56 After treatments and incubation for 2.5 hours, human erythrocyte ghost
57
58 membranes were prepared following the procedure proposed by Fairbanks
59
60
61
62
63
64
65

1 (Fairbanks et al., 1971). Samples were centrifuged and the erythrocyte pellet was
2
3 resuspended and washed with buffer (Na_2HPO_4 5mM, pH 8.0) several times until
4
5 white ghost membranes were obtained. The protein content of erythrocytes ghost cell
6
7 membranes was measured using the Bio-Rad assay (Bio-Rad, Hercules, CA), which
8
9 is based on the dye-binding procedure of Bradford (Bradford, 1976), using bovine
10
11 serum albumin (BSA) as a protein standard.
12
13
14
15
16
17

18 2.2.11. SDS-PAGE

19
20 Sodium dodecyl sulfate-polyacrylamine gel electrophoresis of membrane
21
22 proteins was performed according to Miki and collaborators (Miki et al., 1987) using a
23
24 Mini-PROTEAN Tetra Cell unit (Bio-Rad, Hercules, CA). Erythrocyte ghost
25
26 membranes were mixed with SDS sample buffer, heated at 95 °C for 5 min and then
27
28 a total of 25 µg of protein was electrophoresed in parallel into 12.5% SDS-
29
30 polyacrylamide gel under reducing conditions (slab gel consisted of 7.5%
31
32 polyacrylamide resolving gel and 5% polyacrilamide stacking gel). Electrophoresis
33
34 was carried out at 60V for 10 min followed by 35 min at 200V. Protein bands were
35
36 viewed by staining with Coomassie brilliant blue R-250 for 50 min under gentle
37
38 shaking and destained with a mixture of 7.5% methanol and 10% acetic acid. The
39
40 molecular weight of the membrane proteins was estimated from the molecular size
41
42 marker (Bio-Rad Precision Plus Unstained Standard), ranging from 10 to 250 kDa.
43
44 Electrophoresis gel was digitized on a computer through a video camera. Thereafter,
45
46 ImageJ software was used to calculate the average pixel intensity of bands 1, 2, 3,
47
48 4.1, 4.2 and 5 within regions of interest (ROI) drawn on the collected gel images.
49
50 Bands colour intensity was measured in triplicate for each treatment. The results
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 were expressed as percentage of band colour intensity compared with control
 2
 3 (100%) (Nogueira et al., 2012b).
 4
 5
 6
 7
 8
 9

10 2.2.12. Cell membrane fluidity

11 To determine cell membrane fluidity, fluorescence anisotropy measurements
 12 were carried out using DPH and TMA-DPH fluorescent probes.(Nogueira et al.,
 13 2012b) Treated and non-treated human red blood cells suspensions (haematocrit of
 14 0.01%) in PBS were labelled with the fluorescent probes (final concentration in
 15 samples 10^{-6} M) at room temperature for 1 hour in dark conditions. Changes in
 16 membrane fluidity were also evaluated in samples incubated without AAPH to
 17 determine whether treatment alone can interfere in cell membrane fluidity. Steady-
 18 state anisotropy was measured using an SLM-Amino AB-2 spectrofluorometer, with
 19 polarizers in the L configuration in a quartz cuvette at room temperature. Samples
 20 were illuminated with linearly (vertically or horizontally) polarized monochromatic light
 21 ($\lambda_{\text{ex}} = 365$ nm), and the fluorescence intensities ($\lambda_{\text{em}} = 425$ nm), emitted parallel or
 22 perpendicular to the direction of the excitation beam (slit widths of 8 nm), were
 23 recorded. Fluorescence anisotropy (r) was calculated automatically by software
 24 provided with the instrument, according to the following equation:
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49

$$50 \quad r = \frac{(I_{vv} - I_{vh} \times G)}{(I_{vv} + 2I_{vh} \times G)} \quad \text{Eq. 2}$$

51 where I_{vv} and I_{vh} represent the components of the light intensity emitted in parallel
 52 and perpendicular, respectively, to the direction of the vertically polarized excitation
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

1 light. The factor $G = I_{hv}/I_{hh}$ was used to correct the inequality of the detection beam to
2
3 horizontally and vertically polarized emission.
4
5
6
7
8
9

10 11 12 13 2.2.13. *Statistical analysis* 14

15 Each assay was carried out on three independent experiments using three
16 replicate samples for each sample concentration tested. Statistical analyses were
17 performed using one-way analysis of variance (ANOVA) followed by Dunnett's or
18 Tukey's post-hoc test for multiple comparisons using GraphPad Prism 5.0 software.
19 Differences were considered significant for $p < 0.05$ and $p < 0.001$.
20
21
22
23
24
25
26
27
28
29

30 **3. Results** 31

32 33 34 35 **3.1 Physicochemical characterization of the nanoemulsions** 36

37 It is important to determine the physicochemical characteristics of NE since
38 these characteristics may not only alter the biological but also the toxicological
39 response (Rivera Gil et al., 2010). Table 1 summarizes the physicochemical
40 characteristics of unloaded and EAF-loaded NEs. The average size of all
41 formulations, were about 200 nm, with the unloaded NE smaller than the loaded
42 ones. The polydispersity indexes (PI) were < 0.2 , indicating that monodisperse
43 colloidal dispersions were obtained. Zeta potential, which describes the surface
44 charge of the NE droplets, varied from -20.1 to -24.7 mV, giving an indication of the
45 physical stability of the colloidal dispersions (Mishra et al., 2009; Mitri et al., 2011).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Physicochemical characterization of unloaded and EAF-loaded nanoemulsions.

<i>Samples</i>	<i>Size (nm)</i>	<i>Zeta potential (mV)</i>	<i>Polydispersity Index</i>
EAF-PSO-NE	201.2 ± 3.8	-21.9 ± 2.2	0.17 ± 0.02
PSO-NE	180.7 ± 4.0	-20.1 ± 1.8	0.17 ± 0.01
EAF-MCT-NE	202.5 ± 1.3	-25.7 ± 4.0	0.11 ± 0.03
MCT-NE	186.5 ± 3.3	-22.3 ± 0.3	0.18 ± 0.01

*Results are expressed as mean ± SD of at least 3 independent analyses.

3.2 Haemocompatibility and photohaemolysis studies

In the haemolysis assay the HC_{50} for the isolated formulations components was $7.25 \pm 0.02 \text{ mg mL}^{-1}$, $1790 \pm 1.83 \text{ mg mL}^{-1}$, $45.9 \pm 0.07 \text{ mg mL}^{-1}$, $15.37 \pm 0.54 \text{ mg mL}^{-1}$ and $> 1 \text{ mg mL}^{-1}$, for soy lecithin, polysorbate 80, PSO, MCT and free EAF, respectively; for EAF-loaded nanoemulsions (EAF-PSO-NE; EAF-MCT-NE) and unloaded nanoemulsions (PSO-NE; MCT-NE) was $> 0.5 \text{ mg mL}^{-1}$. When the RBCs were irradiated, the EAF-loaded NEs (EAF-PSO-NE; EAF-MCT-NE) did not demonstrate a photohaemolytic behaviour.

3.3 Haemolysis mediated by H_2O_2 and AAPH

After verifying the haemolytic capacity of EAF and EAF-loaded NEs, their capacity to protect human RBCs against haemolysis induced by H₂O₂ and AAPH was evaluated (Figure. 1).




Figure 1. Protective effect of *P. granatum* EAF and EAF-loaded nanoemulsions against oxidative haemolysis induced by H₂O₂ and AAPH. Results are expressed as mean ± SEM of at least 3 independent assays. Statistical analyses were performed by ANOVA followed by Tukey's multiple comparison test ($p < 0.05$). Same letters mean no statistical difference.

Oxygen free radicals are implicated in many biological processes and for the most part, the endogenous antioxidant system can manage their detoxification. However, certain external events can trigger the production of these damaging free radicals. Hydroxyl (OH[•]) and superoxide (O₂^{•-}) free radical are two oxygen free radical species in biological systems. Normally, the enzymes catalase, superoxide dismutase, and glutathione peroxidase maintain these free radicals at low levels, but

1 under certain circumstances, higher levels of these compounds may be available,
2
3 thus leading to oxidative damage (Floyd and Lewis, 1983).
4

5
6 Hydrogen peroxide is generated by the dismutation of superoxide radical,
7
8 catalysed by superoxide dismutase enzymes and directly produced by oxidase
9
10 enzymes. It can lead the production of hydroxyl free radicals either by exposure to
11
12 ultraviolet light or by interaction with transition metal ions, mostly iron (Fenton-type
13
14 reaction) (Halliwell et al., 2000). Also, hydrogen peroxide may enter the human body
15
16 through skin contact; it crosses the RBC lipid membrane and acts on the intracellular
17
18 moiety, forming hydroxyl radicals that can initiate lipid peroxidation and cause cell
19
20 damage (Alam et al., 2013; Blasa et al., 2007). In the case of haemolysis induced by
21
22 H_2O_2 the IC_{50} of EAF and EAF-MCT-NE did not differ significantly, being 31.62 and
23
24 $27.16 \mu g mL^{-1}$, respectively; EAF-PSO-NE presented a slightly higher IC_{50} , at 76.05
25
26 $\mu g mL^{-1}$. During AAPH-induced haemolysis the EAF showed an IC_{50} of $1.20 \mu g mL^{-1}$
27
28 which places it at the same level of two well-known antioxidants, ascorbic acid IC_{50}
29
30 $7.5 \mu g mL^{-1}$ (Aman et al., 2013) and curcumin IC_{50} $2.94 \mu g mL^{-1}$ (Banerjee et al.,
31
32 2008). In the absence of AAPH, RBCs were stable and haemolysis was negligible
33
34 (data not shown). The percentage of haemolysis in RBCs incubated only with EAF or
35
36 EAF-loaded NEs was almost identical to that of the control sample (RBCs + PBS)
37
38 indicating that EAF and EAF-loaded NEs themselves do not induce haemolysis.
39
40
41
42
43
44
45
46
47
48
49

50 **3.4 Erythrocyte membrane fluidity**

51

52
53
54 Table 2 shows the anisotropy values for both probes after AAPH assay. An
55
56 increase in the r values (anisotropy measurement) of a probe is indicative of a
57
58 decrease in the fluidity of the membrane. The baseline fluorescence for DPH and
59
60
61
62
63
64
65

TMA-DPH was 0.2404 ± 0.0059 and 0.2456 ± 0.0065 , respectively. Free EAF and EAF-loaded NEs containing PSO (EAF-PSO-NE) or MCT (EAF-MCT-NE) maintained the fluidity in the outer and inner parts of erythrocytes membranes at around baseline levels (control PBS). The presence of AAPH alone increased anisotropy values, indicating a decrease in membrane fluidity. In AAPH-induced peroxidation, free radicals are formed in the solution and attack the membranes from the external medium forming lipid hydroperoxides (Arora et al., 2012). Co-incubated free EAF or EAF-loaded NEs managed to restrain the fluidity diminution caused by AAPH, suggesting that the polyphenols present in the EAF stop the free radicals attacking and accessing the bilayer.

Table 2. Steady-state Fluorescence Anisotropy of the probes DPH and TMA-DPH incorporated into erythrocyte membrane after AAPH assay.

<i>Samples ($\mu\text{g mL}^{-1}$)</i>	<i>(r^a) DPH (mean \pm SE)</i>	<i>(r^a) TMA-DPH (mean \pm SE)</i>
Control	0.2404 ± 0.0059	0.2456 ± 0.0065
AAPH (50 mM)	0.3427 ± 0.0069^b	0.3315 ± 0.0053^b
EAF (1)	0.2488 ± 0.0060^c	0.2465 ± 0.0064^c
*EAF (1)	0.2351 ± 0.0073^c	0.2534 ± 0.0066^c
EAF-MCT-NE (50)	0.2498 ± 0.0060^c	0.2489 ± 0.0059^c
*EAF-MCT-NE (50)	0.2465 ± 0.0058^c	0.2443 ± 0.0060^c
EAF-PSO-NE (50)	0.2329 ± 0.0058^c	0.2382 ± 0.0052^c
*EAF-PSO-NE (50)	0.2449 ± 0.0079^c	0.2496 ± 0.0076^c

*Samples incubated without AAPH.

^a*r* values = anisotropy measurements. ^b*p* < 0.01 when compared to control PBS (Dunnett's post hoc test). ^c*p* < 0.01 when compared to AAPH (Dunnett's post hoc

test). Data represent the means of three independent experiments \pm SE, each conducted in triplicate.

3.5 SDS-PAGE

Given the changes in membrane fluidity, the next step was to run an SDS-PAGE experiment to assess possible changes in membrane proteins in human erythrocytes. Oxidants basically alter the erythrocyte membrane by decreasing the LMW proteins and producing HMW proteins (Flynn et al., 1983; Snyder et al., 1985).

Figure 2 shows the electrophoretic profile of the erythrocyte membrane proteins.




Figure 2. Protective effect of *P. granatum* EAF and EAF-loaded nanoemulsions for human erythrocyte membrane skeletal proteins after incubation with AAPH, determined by SDS-PAGE. Following preincubation of the erythrocytes for 2.5 hours

1 in the absence or presence of AAPH and treatments, membranes were separated
2
3 and washed as described in section 2.12. The major cytoskeletal protein bands are
4
5 identified following the classification of Fairbanks *et al.* (Fairbanks *et al.*, 1971) and
6
7 are given on the right of the gel. Each lane corresponds to a different treatment: (1)
8
9 protein standard; (2) control PBS (untreated membrane); (3) AAPH control; treatment
10
11 with ethyl acetate fraction and nanoemulsions in the presence of AAPH (4) EAF (1 μg
12
13 mL^{-1}), (5) EAF-PSO-NE (50 $\mu\text{g mL}^{-1}$), (6) EAF-MCT-NE (50 $\mu\text{g mL}^{-1}$).
14
15
16
17
18
19
20

21 When RBCs were incubated with EAF or NE alone, the level of in erythrocyte-
22
23 membrane proteins was maintained at a background level similar to that in the
24
25 untreated samples (control) (data not shown). The well-established normal
26
27 distribution of the major membrane cytoskeletal proteins is shown in lane 2, which
28
29 contains untreated erythrocytes ghosts. AAPH treatment produced changes in the
30
31 protein pattern, leading to a remarkable loss of spectrins (band 1 and 2), and band 3,
32
33 4.1, 4.2 and 5, as seen in lane 3, confirming previous results from other authors
34
35 (Hseu *et al.*, 2002; Martínez *et al.*, 2012; Yang *et al.*, 2006). As shown in lanes 4, 5
36
37 and 6, the analysis of LMW proteins revealed that EAF and EAF-loaded NEs
38
39 protected against AAPH-induced changes in the amount of erythrocytes membrane
40
41 proteins. Table 3 shows the results expressed as the percentage of band colour
42
43 intensity compared with the control (100%).
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 3. Effect of AAPH and EAF and EAF-loaded nanoemulsions (EAF-PSO-NE and EAF-MCT-NE) on protein bands of human erythrocyte membrane.

<i>Treatment</i>	<i>Band 1 and 2</i>	<i>Band 3</i>	<i>Band 4.1 and 4.2</i>	<i>Band 5</i>
AAPH 50 Mm	37.42 ± 1.45 [*]	39.65 ± 2.42 [*]	4.96 ± 4.97 [*]	22.94 ± 1.06 [*]
AAPH + EAF 1 µg mL ⁻¹	59.28 ± 2.05 ^{*#a}	70.03 ± 4.32 ^{*#b}	32.63 ± 3.49 ^{*#}	50.43 ± 1.01 ^{*#}
AAPH + EAF-PSO-NE 50 µg mL ⁻¹	81.08 ± 1.24 ^{*#}	88.02 ± 4.47 ^{*#c}	61.10 ± 3.40 ^{*#d}	69.08 ± 0.85 ^{*#e}
AAPH + EAF-MCT-NE 50 µg mL ⁻¹	60.35 ± 1.90 ^{*#a}	75.29 ± 4.18 ^{*#b,c}	59.49 ± 2.42 ^{*#d}	65.80 ± 0.45 ^{*#e}

The results are expressed as percentage of control taking control as 100%. Values are mean ± SEM of three experiments. See the Materials and Methods section for details of the analysis performed. Statistical analyses were performed by ANOVA followed by Tukey's multiple comparison tests (same letter means no statistical difference).

^{*}Significant difference from control group ($p < 0.01$).

[#]Significant difference from AAPH group ($p < 0.01$).

1 After AAPH treatment, the amount of band 1, 2, 3, 4 and 5 proteins dropped to
2
3 37%, 40%, 5% and 23 %, respectively, in comparison with untreated erythrocytes.
4
5 On the other hand, the co-treatment of erythrocytes with AAPH and EAF or EAF-
6
7 loaded NEs containing PSO (EAF-PSO-NE) or MCT (EAF-MCT-NE) as the oil phase,
8
9 reduced the effects of the oxidant agent on those proteins. This reduction was better
10
11 and more evident when the EAF was nanoemulsified (EAF-PSO-NE or EAF-MCT-
12
13 NE).
14
15
16
17
18
19

20 **3.6 SEM analysis**

21
22
23
24

25 The protein organization network found mainly in the inner membrane layer is
26
27 responsible for the maintenance of shape and stability; once the protein network is
28
29 compromised the shape of the erythrocyte may change (Vittori et al., 2002). As
30
31 mentioned previously, oxidative damage in cell membranes leads to changes in the
32
33 protein profile. The effect of AAPH alone, EAF and EAF-loaded NEs co-incubated
34
35 with AAPH on erythrocyte morphology was studied by SEM analysis (Figure. 3).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 3. Scanning electron micrograph of normal human erythrocyte and protective effects of ethyl acetate fraction (EAF) and nanoemulsions of *P. granatum* peel extract against AAPH induced oxidative damage on RBC. A) AAPH at 50 mM; B-1) Control in PBS and B-2) Control in PBS after 2.5 h incubation; C) EAF ($1.5 \mu\text{g mL}^{-1}$) with AAPH at 50 mM; D) EAF-loaded NEs ($50 \mu\text{g mL}^{-1}$) with AAPH at 50 mM. Scale bars correspond to $10 \mu\text{m}$.

SEM examinations corroborated that AAPH, nanoemulsions and the EAF interacted with the lipid bilayer by altering the normal morphology of the RBCs. Treatment of erythrocytes with AAPH alone resulted in a significant change in shape to echinocytic or acanthocytic (Figure. 3A). In our study the control erythrocytes incubated with PBS solution for 2.5 hours presented a leptocyte-type shape (Fig. 4B-1), *i.e.* a flattened cell with reduced volume; assuming that the long incubation time was somehow deforming the RBCs, SEM preparation was then conducted immediately after adding RBCs to the PBS solution, but the findings were the same (Figure. 3B-2); it is thus likely that this was a blood sample characteristic. EAF (Figure. 3C) and EAF-loaded NEs (Figure. 3D) incubated with AAPH showed the

1 knizocyte and stomatocyte-type deformation; knizocyte is a cell with two or three
2 concavities while a stomatocyte is a cell with one concavity in the shape of a mouth.
3
4 The EAF and EAF-loaded NEs incubated with RBCs alone did not alter the
5 morphology of the erythrocytes that remained as the control ones (data not shown).
6
7
8
9

10 11 12 13 **4. Discussion** 14 15 16 17

18 *Punica granatum* is well known for its considerable polyphenolics content; they
19 are bioactive secondary metabolites widely distributed in plants (Harborne, 1989) and
20 contain one or more aromatic hydroxyl groups that actively scavenge free radicals
21 and are responsible for antioxidant activity. It is essential to identify and quantify the
22 polyphenolic compounds in both free EAF and EAF-loaded NEs, given that their
23 capacity to protect erythrocytes against oxidation is largely related to the
24 characteristics of these compounds. EAF contains at least 600 mg g⁻¹ GAE (gallic
25 acid equivalents) of total phenolic content, among which ellagic acid, gallic acid, and
26 punicalagin are the major ones (8.47%, 23.86% and 29.67% w/w, respectively). The
27 nanoemulsions presented entrapment efficiency near or above 50% depending on
28 the chemical compound lipophilicity. Details of the polyphenol identification and
29 quantification through HPLC method are described in a previous study (Baccarin and
30 Lemos-Senna, 2014).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Haemocompatibility testing, a major part of biocompatibility testing, comprises
50 the evaluation of interactions of foreign material such as active compounds,
51 excipients and/or formulations with human red blood cells and to explore possible
52 damaging effects arising from these combined exposure (Szebeni, 2012).
53
54
55
56
57
58
59
60
61
62
63
64
65

1 induced oxidative damage to membranes and to evaluate the antioxidant activity of
2 new compound or formulation, but also to determine the topical irritation and/or
3 phototoxic potential (photohaemolysis) of these substances (Pape and Hoppe, 1991;
4 Pape et al., 1987). With respect to the potential topical application of EAF-loaded
5 NEs as a photoprotector and according to the Safety Evaluation Guide for Cosmetic
6 Products (Brasil, 2003) it is extremely important to evaluate their potential topical
7 irritation and photoirritation. Some studies have reported the photohaemolytic
8 potential of hydroalcoholic extracts (Madariaga et al., 2010; Toledo et al., 2012).
9 Nevertheless, there are few studies in the literature related to the effects of
10 nanodispersions (*i.e.* nanoemulsion) and even fewer related to nanoemulsions
11 loaded with plant extracts or extract fractions.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 Several studies support the use of RBCs as a model of oxidative damage
28 induced by AAPH (Ajila and Prasada Rao, 2008; Alvarez-Suarez et al., 2012;
29 Banerjee et al., 2008; Blasa et al., 2007; Botta et al., 2014; Chirinos et al., 2008;
30 Hseu et al., 2002; Magalhães et al., 2009; Martínez et al., 2012; Yang et al., 2006).
31 When AAPH is added as a radical initiator, it decomposes at physiological
32 temperature in aqueous solution to generate alkyl radicals, which in the presence of
33 oxygen are converted to the corresponding peroxy radicals; in turn, these peroxy
34 radicals induce oxidation of polyunsaturated lipids in the RBCs membrane, causing a
35 chain reaction known as lipid peroxidation (Banerjee et al., 2008).
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Both EAF and EAF-loaded NEs demonstrated antioxidant properties against
50 oxidative damage in the erythrocyte membrane induced by H₂O₂ or AAPH. The
51 different IC₅₀ values obtained for the H₂O₂ and AAPH assays only emphasizes the
52 fact that conclusions about antioxidant activity should not be based on a single test
53 model, because each assay represents a different type of oxidative damage and
54
55
56
57
58
59
60
61
62
63
64
65

1 varies in different respects; therefore samples may also behave differently. With
2
3 regard to AAPH assay, it was observed that the NE presented a higher IC₅₀ than the
4
5 free EAF, which can be explained by the fact that as the EAF is encapsulated it is not
6
7 totally available in the assay medium in order to directly neutralize the peroxy
8
9 radicals generated from AAPH.
10
11

12
13 The incorporation of EAF and NE into RBCs could cause conformational
14
15 alterations in membrane cytoskeletal proteins and changes in the internal viscosity of
16
17 the cells. To determine whether free EAF or EAF-loaded NEs disturb the
18
19 phospholipid bilayer across its thickness, steady-state fluorescence anisotropy (*r*)
20
21 was measured to test membrane fluidity, as this is a sensitive indicator for monitoring
22
23 fluorophore binding to regions of biomembranes with constrained movement (Hou et
24
25 al., 2011; Nogueira et al., 2012b). This is an important parameter related to the
26
27 structure and functional state of the cell membrane (Marczak, 2009).
28
29
30
31

32
33 The fluorescent probes, DPH and TMA-DPH, were incorporated into the
34
35 human erythrocyte membranes to detect possible changes in membrane fluidity.
36
37 DPH is incorporated within the hydrophobic region of the bilayer membrane, while
38
39 TMA-DPH is incorporated near the surface of the cell membrane (Marczak, 2009).
40
41 The fluorescence anisotropy values are inversely proportional to cell membrane
42
43 fluidity. A high degree of fluorescence anisotropy, represents a high structural order
44
45 and/or low cell membrane fluidity. Therefore, it is possible to assess the arrangement
46
47 and mobility of membrane components in different regions of the bilayer (Marczak,
48
49
50
51
52 2009).
53

54
55 Our findings suggest that the possible localization of phenolic compounds of
56
57 EAF-loaded NEs on the erythrocyte membrane is not restricted to one determined
58
59 zone since EAF-loaded NEs when together with AAPH were able to elevate the
60
61
62
63
64
65

1 fluidity of both the inner and outer layer, allowing better interaction between the
2
3 antioxidant compounds and lipophilic radicals.
4

5
6 The degree of protection afforded by free EAF and EAF-loaded NEs against
7
8 human erythrocyte oxidation might be attributed to different features. First, the
9
10 partition coefficient or degree of lipophilicity of phenolic compounds determines their
11
12 interaction with biomembranes and influences their antioxidant capacity and/or ability
13
14 to interact with free radicals present in the aqueous medium of the oxidation system
15
16 (Liao and Yin, 2000). Second, the chemical structure of the phenolic compounds
17
18 determines their ability to react with free radicals; for phenolic acids, the free radical-
19
20 scavenging properties depend mainly on their number of hydroxyl groups that have
21
22 the ability to donate hydrogen, thereby forming a stable end-product, which does not
23
24 initiate or propagate further oxidation of the lipid (Hsieh et al., 2005). Third, in
25
26 phenolic mixtures *i.e* extract and/or extract fraction, the presence of synergism may
27
28 modify the characteristics of oxidation inhibition (Chirinos et al., 2008).
29
30
31
32
33
34

35 Regarding the membrane protein profile, when the RBCs were treated with the
36
37 EAF-loaded NEs (EAF-PSO-NE or EAF-MCT-NE) a more efficient protection against
38
39 the oxidant agent on the membrane proteins was observed. This is probably due to
40
41 the fact that one of the advantages of NE is that they better solubilize lipophilic
42
43 compounds and enhance their transportation through lipid layers. In this way, it is not
44
45 only the water-soluble radical scavengers present in the EAF that efficiently
46
47 scavenge the peroxy radicals in the medium before they attack the erythrocytes, but
48
49 also the lipid-soluble ones which are located in lipophilic regions of the membrane
50
51 (Blasa et al., 2007). Furthermore, EAF-PSO-NE appears to deliver an even better cell
52
53 membrane protection, since PSO also presents antioxidants compounds that may act
54
55 in a synergistic manner with EAF.
56
57
58
59
60
61
62
63
64
65

1 SEM studies corroborated the interaction of AAPH, EAF-PSO-NE and EAF-
2 MCT-NE and free EAF with the lipid bilayer, as demonstrated by the changes in cell
3 shape. According to the bilayer couple hypothesis, the shape changes induced in
4 erythrocytes by foreign molecules are due to differential expansion of the two
5 monolayers of the RBCs membrane. Thus, stomatocytes are formed when the
6 compound inserts into the inner monolayer, whereas speculated-shaped echinocytes
7 are produced when it inserts into the outer moiety, causing surface expansion (Lim
8 H. W. et al., 2002; Sheetz and Singer, 1974).
9

10 Finally, nanoemulsions entrapping pomegranate peel polyphenol-rich EAF
11 have a great potential for cosmetic application as antioxidant and photoprotector. The
12 present study indicates that EAF-PSO-NE and EAF-MCT-NE were photosafe in a
13 preliminary *in vitro* test and protected the RBC membrane against oxidative damage.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **Conflict of Interest**

32 The authors declare no competing financial interest.
33
34
35
36
37
38
39

40 **Acknowledgments**

41 The authors acknowledge financial support from CAPES (CAPES/PDSE
42 Project No. BEX 5613/13-2), the Ministerio de Economía y Competitividad - Spain
43 (Project MAT2012-38047-C02-01) and European Union (FEDER). The authors would
44 like to thank Professor Dr. Lourdes Pérez for Malvern Zetasizer Nano ZS use and
45 Professor Dr. Maria Antonia Busquets for SLM-Amino AB-2 Spectrofluorometer use.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Ajila, C.M., Prasada Rao, U.J.S., 2008. Protection against hydrogen peroxide induced oxidative damage in rat erythrocytes by *Mangifera indica* L. peel extract. *Food Chem. Toxicol.* 46, 303-309.
- Alam, M.N., Bristi, N.J., Rafiquzzaman, M., 2013. Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharm. J.* 21, 143-152.
- Alvarez-Suarez, J.M., Giampieri, F., González-Paramás, A.M., Damiani, E., Astolfi, P., Martínez-Sánchez, G., Bompadre, S., Quiles, J.L., Santos-Buelga, C., Battino, M., 2012. Phenolics from monofloral honeys protect human erythrocyte membranes against oxidative damage. *Food Chem. Toxicol.* 50, 1508-1516.
- Amakura, Y., Okada, M., Tsuji, S., Tonogai, Y., 2000. High-performance liquid chromatographic determination with photodiode array detection of ellagic acid in fresh and processed fruits. *J. Chromatogr. A* 896, 87-93.
- Aman, S., Moin, S., Owais, M., Siddiqui, M.U., 2013. Antioxidant activity of thymol: protective role in AAPH-induced hemolysis in diabetic erythrocytes. *Int J Pharm Sci Invent* 2, 55-60.
- Arora, S., Rajwade, J.M., Paknikar, K.M., 2012. Nanotoxicology and in vitro studies: The need of the hour. *Toxicol. Appl. Pharmacol.* 258, 151-165.
- Baccarin, T., Lemos-Senna, E., 2014. Pomegranate Seed Oil Nanoemulsions Encapsulating Pomegranate Peel Polyphenol-Rich Ethyl Acetate Fraction: Development and Antioxidant Assessment. *Journal of Nanopharmaceutics and Drug Delivery* 2, 333-343.
- Banerjee, A., Kunwar, A., Mishra, B., Priyadarsini, K.I., 2008. Concentration dependent antioxidant/pro-oxidant activity of curcumin: Studies from AAPH induced hemolysis of RBCs. *Chem.-Biol. Interact.* 174, 134-139.
- Blasa, M., Candiracci, M., Accorsi, A., Piacentini, M.P., Piatti, E., 2007. Honey flavonoids as protection agents against oxidative damage to human red blood cells. *Food Chem.* 104, 1635-1640.
- Botta, A., Martínez, V., Mitjans, M., Balboa, E., Conde, E., Vinardell, M.P., 2014. Erythrocytes and cell line-based assays to evaluate the cytoprotective activity of antioxidant components obtained from natural sources. *Toxicol. In Vitro* 28, 120-124.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254.
- Brasil, 2003. Guia para avaliação de segurança de produtos cosméticos. Agência Nacional de Vigilância Sanitária, 1-47.
- Chirinos, R., Campos, D., Warnier, M., Pedreschi, R., Rees, J.-F., Larondelle, Y., 2008. Antioxidant properties of mashua (*Tropaeolum tuberosum*) phenolic extracts against oxidative damage using biological in vitro assays. *Food Chem.* 111, 98-105.
- Doktorovova, S., Souto, E.B., Silva, A.M., 2014. Nanotoxicology applied to solid lipid nanoparticles and nanostructured lipid carriers – A systematic review of in vitro data. *Eur. J. Pharm. Biopharm.* 87, 1-18.
- El-Nemr, S.E., Ismail, I.A., Ragab, M., 2006. Chemical composition of juice and seeds of pomegranate fruit. *Die Nahrung* 34, 601-606.
- Fadavi, A., Barzegar, M., Azizi, H.M., 2006. Determination of fatty acids and total lipid content in oilseed of 25 pomegranates varieties grown in Iran. *J Food Compost Anal* 19, 676-680.
- Fairbanks, G., Steck, T.L., Wallach, D.F.H., 1971. Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochemistry* 10, 2606-2617.

- 1 Floyd, R.A., Lewis, A.C., 1983. Hydroxyl Free Radical Formation from Hydrogen Peroxide
2 by Ferrous Iron-Nucleotide Complexes. *Biochem* 22, 2645-2649.
- 3 Flynn, T.P., Allen, D.W., Johnson, G.J., White, J.G., 1983. Oxidant damage of the lipids and
4 proteins of the erythrocyte membranes in unstable hemoglobin disease. Evidence for the role
5 of lipid peroxidation. *J. Clin. Invest.* 71, 1215-1223.
- 6 González, N., Ribeiro, D., Fernandes, E., Nogueira, D.R., Conde, E., Moure, A., Vinardell,
7 M.P., Mitjans, M., Domínguez, H., 2013. Potential use of *Cytisus scoparius* extracts in topical
8 applications for skin protection against oxidative damage. *J. Photochem. Photobiol.*, B 125,
9 83-89.
- 10 Guglielmini, G., 2008. Nanostructured novel carrier for topical application. *Clinics in*
11 *Dermatology* 26, 341-346.
- 12 Halliwell, B., Clement, M.V., Long, L.H., 2000. Hydrogen peroxide in the human body.
13 *FEBS Lett* 486, 10-13.
- 14 Harborne, J.B., 1989. General procedures and measurements of total phenolics, In: Harborne,
15 J.B. (Ed.), *Methods in plant biochemistry. Plant phenolics.* Academic Press, London.
- 16 Hou, S.-Z., Su, Z.-R., Chen, S.-X., Ye, M.-R., Huang, S., Liu, L., Zhou, H., Lai, X.-P., 2011.
17 Role of the interaction between puerarin and the erythrocyte membrane in puerarin-induced
18 hemolysis. *Chem.-Biol. Interact.* 192, 184-192.
- 19 Hseu, Y.-C., Chang, W.-C., Hseu, Y.-T., Lee, C.-Y., Yech, Y.-J., Chen, P.-C., Chen, J.-Y.,
20 Yang, H.-L., 2002. Protection of oxidative damage by aqueous extract from *Antrodia*
21 *camphorata* mycelia in normal human erythrocytes. *Life Sciences* 71, 469-482.
- 22 Hsieh, C.-L., Yen, G.-C., Chen, H.-Y., 2005. Antioxidant Activities of Phenolic Acids on
23 Ultraviolet Radiation-Induced Erythrocyte and Low Density Lipoprotein Oxidation. *J. Agric.*
24 *Food Chem.* 53, 6151-6155.
- 25 Liao, K.-I., Yin, M.-c., 2000. Individual and Combined Antioxidant Effects of Seven Phenolic
26 Agents in Human Erythrocyte Membrane Ghosts and Phosphatidylcholine Liposome
27 Systems: Importance of the Partition Coefficient. *J. Agric. Food Chem.* 48, 2266-2270.
- 28 Lim H. W., G., Wortis, M., Mukhopadhyay, R., 2002. Stomatocyte–discocyte–echinocyte
29 sequence of the human red blood cell: Evidence for the bilayer– couple hypothesis from
30 membrane mechanics. *Proc. Natl. Acad. Sci. U. S. A.* 99, 16766-16769.
- 31 Madariaga, Y.G., Cárdenas, M.B., Toledo, D.B., Alfonso, O.C., Montalbán, C.M.M., Bernal,
32 Y.M., 2010. In vitro photohemolytic assessment of *Cissus sicyoides* L. and *Achyranthes*
33 *aspera*. *Rev Cubana Plant Med.* 15, 126-132.
- 34 Magalhães, A.S., Silva, B.M., Pereira, J.A., Andrade, P.B., Valentão, P., Carvalho, M., 2009.
35 Protective effect of quince (*Cydonia oblonga* Miller) fruit against oxidative hemolysis of
36 human erythrocytes. *Food Chem. Toxicol.* 47, 1372-1377.
- 37 Marczak, A., 2009. Fluorescence anisotropy of membrane fluidity probes in human
38 erythrocytes incubated with anthracyclines and glutaraldehyde. *Bioelectrochemistry* 74, 236-
39 239.
- 40 Martínez, V., Ugartondo, V., Vinardell, M.P., Torres, J.L., Mitjans, M., 2012. Grape
41 Epicatechin Conjugates Prevent Erythrocyte Membrane Protein Oxidation. *J. Agric. Food*
42 *Chem.* 60, 4090-4095.
- 43 Miki, M., Tamai, H., Mino, M., Yamamoto, Y., Niki, E., 1987. Free-radical chain oxidation
44 of rat red blood cells by molecular oxygen and its inhibition by α -tocopherol. *Arch. Biochem.*
45 *Biophys.* 258, 373-380.
- 46 Mishra, A., Ram, S., Ghosh, G., 2009. Dynamic Light Scattering and Optical Absorption in
47 Biological Nanofluids of Gold Nanoparticles in Poly(vinyl pyrrolidone) Molecules. *J. Phys.*
48 *Chem. C* 113, 6976-6982.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Mitri, K., Shegokar, R., Gohla, S., Anselmi, C., Müller, R.H., 2011. Lipid nanocarriers for
2 dermal delivery of lutein: preparation, characterization, stability and performance. *Int. J.*
3 *Pharm.* 414, 267-275.
- 4 Nogueira, D., Mitjans, M., Morán, M.C., Pérez, L., Vinardell, M.P., 2012a. Membrane-
5 destabilizing activity of pH-responsive cationic lysine-based surfactants: role of charge
6 position and alkyl chain length. *Amino Acids* 43, 1203-1215.
- 7 Nogueira, D.R., Mitjans, M., Busquets, M.A., Pérez, L., Vinardell, M.P., 2012b. Phospholipid
8 Bilayer-Perturbing Properties Underlying Lysis Induced by pH-Sensitive Cationic Lysine-
9 Based Surfactants in Biomembranes. *Langmuir* 28, 11687-11698.
- 10 Onoue, S., Seto, Y., Gandy, G., Yamada, S., 2009. Drug-induced phototoxicity; an early in
11 vitro identification of phototoxic potential of new drug entities in drug discovery and
12 development. *Curr. Drug Saf.* 4, 123-136.
- 13 Pape, W.J.W., Hoppe, U., 1991. In vitro methods for the assessment of primary local effects
14 of topically applied preparations. *Skin Pharmacol.* 4, 205-212.
- 15 Pape, W.J.W., Maurer, T., Pfannenbecker, U., Steiling, W., 2001. The red blood cell
16 phototoxicity test (photohaemolysis and haemoglobin oxidation): EU/COLIPA validation
17 programme on phototoxicity (phase II). *ATLA, Altern. Lab. Anim.* 29, 145-162.
- 18 Pape, W.J.W., Pfannenbecker, U., Hoppe, U., 1987. Validation of the red blood cell system as
19 in vitro assay for rapid screening of irritation potential of surfactants. *Mol. Toxicol.* 1, 525-
20 536.
- 21 Qu, W., Pan, Z., Ma, H., 2010. Extraction modeling and activities of antioxidants from
22 pomegranate marc. *J Food Eng* 99, 16-23.
- 23 Rivera Gil, P., Oberdörster, G., Elder, A., Puentes, V., Parak, W.J., 2010. Correlating Physico-
24 Chemical with Toxicological Properties of Nanoparticles: The Present and the Future. *ACS*
25 *Nano* 4, 5527-5531.
- 26 Schubert, M.A., Müller-Goymann, C.C., 2005. Characterisation of surface-modified solid
27 lipid nanoparticles (SLN): Influence of lecithin and nonionic emulsifier. *Eur. J. Pharm.*
28 *Biopharm.* 61, 77-86.
- 29 Seeram, N.P., Adams, L.S., Henning, S.M., Niu, Y., Zhang, Y., Nair, M.G., Heber, D., 2005.
30 In vitro antiproliferative, apoptotic and antioxidant activities of punicalagin, ellagic acid and a
31 total pomegranate tannin extract are enhanced in combination with other polyphenols as
32 found in pomegranate juice. *J. Nutr. Biochem.* 16, 360-367.
- 33 Sheetz, M.P., Singer, S., 1974. Biological membranes as bilayer couples. A molecular
34 mechanism of drug-erythrocyte interactions. *Proc. Natl. Acad. Sci. U. S. A.* 71, 4457-4461.
- 35 Snyder, L.M., Fortier, N.L., Trainor, J., Jacobs, J., Leb, L., Lubin, B., Chiu, D., Shohet, S.,
36 Mohandas, N., 1985. Effect of hydrogen peroxide exposure on normal human erythrocyte
37 deformability, morphology, surface characteristics, and spectrin-hemoglobin cross-linking. *J.*
38 *Clin. Invest.* 76, 1971-1977.
- 39 Söderlind, E., Karlsson, L., 2006. Haemolytic activity of maltopyranoside surfactants. *Eur. J.*
40 *Pharm. Biopharm.* 62, 254-259.
- 41 Svetina, S., Kuzman, D., Waugh, R.E., Zihlerl, P., Žekš, B., 2004. The cooperative role of
42 membrane skeleton and bilayer in the mechanical behaviour of red blood cells.
43 *Bioelectrochemistry* 62, 107-113.
- 44 Szebeni, J., 2012. Hemocompatibility testing for nanomedicines and biologicals: predictive
45 assays for complement mediated infusion reactions. *Eur J Nanomed.* 4, 33-53.
- 46 Toledo, D.B., Cárdenas, M.B., Díaz, A.V., Montalbán, C.M.M., Rodriguez, N.I., 2012.
47 Evaluación fotohemolítica in vitro de *Parthenium hysterophorus* L. *Revista Científica Villa*
48 *Clara* 16, 43-48.
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Ugartondo, V., Mitjans, M., Vinardell, M.P., 2009. Applicability of lignins from different
2 sources as antioxidants based on the protective effects on lipid peroxidation induced by
3 oxygen radicals. *Ind. Crops Prod.* 30, 184-187.

4 Van Elswijk, D.A., Schobel, U.P., Lansky, E.P., Irth, H., Van der Greef, J., 2004. Rapid
5 dereplication of estrogenic compounds in pomegranate (*Punica granatum*) using on-line
6 biochemical detection coupled to mass spectrometry. *Phytochemistry* 65, 233-241.

7 Vittori, D., Garbossa, G., Lafourcade, C., Pérez, G., Nesse, A., 2002. Human erythroid cells
8 are affected by aluminium. Alteration of membrane band 3 protein. *Biochim. Biophys. Acta,*
9 *Biomembr.* 1558, 142-150.

10 Yang, H.-L., Chen, S.-C., Chang, N.-W., Chang, J.-M., Lee, M.-L., Tsai, P.-C., Fu, H.-H.,
11 Kao, W.-W., Chiang, H.-C., Wang, H.-H., Hseu, Y.-C., 2006. Protection from oxidative
12 damage using *Bidens pilosa* extracts in normal human erythrocytes. *Food Chem. Toxicol.* 44,
13 1513-1521.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figura 1




Figure 2






Figure 3



Graphical Abstract

Title: Protection against Oxidative Damage in Human Erythrocytes and Preliminary Photosafety Assessment of *Punica granatum* Seed Oil Nanoemulsions Entrapping Polyphenol-rich Ethyl Acetate Fraction

T. Baccarin, M. Mitjans, E. Lemos-Senna, M. P. Vinardell*



Highlights

Title: Protection against Oxidative Damage in Human Erythrocytes and Preliminary Photosafety Assessment of *Punica granatum* Seed Oil Nanoemulsions Entrapping Polyphenol-rich Ethyl Acetate Fraction

T. Baccarin, M. Mitjans, E. Lemos-Senna, M. P. Vinardell*

- EAF-loaded NEs were able to elevate the fluidity of both the inner and outer layer of RBCs.
- EAF-loaded NEs managed to reduce the RBC membrane proteins changes.
- EAF-PSO-NE appears to deliver better RBC membrane protection.
- SEM studies corroborated the interaction of EAF-loaded NEs with the lipid bilayer.
- EAF-loaded NEs were photosafe in a preliminary *in vitro* test and protected the RBC membrane against oxidative damage.

***Conflict of Interest**

[Click here to download Conflict of Interest: coi_disclosure.pdf](#)

***Conflict of Interest**

[Click here to download Conflict of Interest: coi_disclosure Elenara.pdf](#)

***Conflict of Interest**

[Click here to download Conflict of Interest: coi_disclosure Thaisa.PDF](#)

***Conflict of Interest**

[Click here to download Conflict of Interest: coi_disclosure_Mitjans.pdf](#)