

1 **Differential requirements for Tausled like kinases 1 and 2 in mammalian**
2 **development**

3 Sandra Segura-Bayona^{1,*}, Philip A. Knobel^{1,*}, Helena González Burón^{1,*}, Sameh A. Youssef^{2,3},
4 Aida Peña-Blanco¹, Étienne Coyaude^{4,5}, Teresa López-Rovira¹, Katrin Rein¹, Lluís Palenzuela¹,
5 Julien Colombelli¹, Stephen Forrow¹, Brian Raught^{4,5}, Anja Groth⁶, Alain de Bruin^{2,7} and
6 Travis H. Stracker^{1,**}

7 ¹Institute for Research in Biomedicine (IRB Barcelona), The Barcelona Institute of Science
8 and Technology, Barcelona 08028, Spain

9 ²Dutch Molecular Pathology Center, Department of Pathobiology, Faculty of Veterinary
10 Medicine, Utrecht University, Utrecht 3512 JE, The Netherlands

11 ³Department of Pathology, Faculty of Veterinary Medicine, Alexandria University, Edfina,
12 Egypt.

13 ⁴Princess Margaret Cancer Centre, University Health Network, Canada

14 ⁵Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada

15 ⁶Biotech Research and Innovation Centre (BRIC) and Centre for Epigenetics, Faculty of
16 Health and Medical Sciences, University of Copenhagen, Copenhagen, DK-2200, Denmark.

17 ⁷University Medical Center Groningen, Department of Pediatrics, University of Groningen, NL-
18 9713 Groningen, The Netherlands

19 *These authors contributed equally to this work

20 **To whom correspondence should be addressed. Tel: +34 93 403 11 83; Fax: +34 93 403 71
21 14; Email: travis.stracker@irbbarcelona.org. Institute for Research in Biomedicine (IRB)
22 Barcelona, Oncology Programme, Baldori Reixac, 10, 08028 Barcelona, Spain

23 **Running title:** Tausled like kinases in development

24 **Abstract**

25 The regulation of chromatin structure is critical for a wide range of essential cellular processes.
26 The Tausled like kinases, TLK1 and TLK2, regulate ASF1, a histone H3/H4 chaperone, and
27 likely other substrates, and their activity has been implicated in transcription, DNA replication,
28 DNA repair, RNA interference, cell cycle progression, viral latency, chromosome segregation
29 and mitosis. However, little is known about the functions of TLK activity *in vivo* or the relative
30 functions of the highly similar TLK1 and TLK2 in any cell type. To begin to address this, we
31 have generated *Tlk1* and *Tlk2* deficient mice. We found that while TLK1 was dispensable for
32 murine viability, TLK2 loss led to late embryonic lethality due to placental failure. TLK2 was
33 required for normal trophoblast differentiation and the phosphorylation of ASF1 was reduced
34 in placentas lacking TLK2. Conditional bypass of the placental phenotype allowed the
35 generation of apparently healthy TLK2 deficient mice, while only the depletion of both TLK1
36 and TLK2 led to extensive genomic instability, indicating that both activities contribute to
37 genome maintenance. Our data identifies a specific role for TLK2 in placental function during
38 mammalian development and suggests that TLK1 and TLK2 play largely redundant roles in
39 genome maintenance.

Introduction

The Tousled kinase (TSL) and Tousled-like kinases (TLKs) are serine-threonine kinases whose activity has been linked to DNA replication, DNA repair, transcription, chromatin structure, chromosome segregation, RNA interference, viral latency and cell cycle checkpoint control in various organisms(1-9). TSL was identified in *Arabidopsis* where mutations in the gene led to pleiotropic defects, including delays in flowering time and leaf development(1). In *Drosophila* and *C.elegans*, deficiency in the single TLK gene led to severe mitotic defects and lethality during development(8, 10). In *Drosophila*, these effects were proposed to result from defective chromatin maintenance and could be rescued by the overexpression of the histone H3-H4 chaperone ASF1, a substrate of TLK(3, 10). In *C.elegans*, the major defects observed following ceTLK-1 downregulation were transcriptional, reflecting reduced phosphorylation of RNA polymerase II, and mitotic, through interactions with Aurora B(7, 8).

Vertebrates encode 2 distinct TLK homologues, Tousled like kinases 1 and 2 (TLK1 and TLK2), that interact with each other and both ASF1a and ASF1b (3). In *Drosophila*, TLK phosphorylation of ASF1 controls its stability, while in vertebrates, TLK1 mediated phosphorylation of several sites on the C-terminal tail of ASF1 promote its binding affinity for the H3-H4 heterodimer(11, 12). Following DNA damage, TLK1 is phosphorylated by the checkpoint kinase CHK1, inhibiting its activity, potentially to coordinate global ASF1 histone binding capacity with the checkpoint response(13, 14). TLK activity has also been linked to cell cycle checkpoint recovery through the phosphorylation of RAD9, a component of the 9-1-1 (RAD9/HUS1/RAD1) alternative clamp loader that plays multiple roles in the response to DNA damage, including the regulation of CHK1 activation(15-17), and through TLK2-ASF1a mediated transcriptional events(18). Although TLK2 is less well characterized, the amplification of its activity has been specifically linked to breast cancer progression in recent work(19, 20).

Despite their identification nearly 15 years ago, the roles of the ubiquitously expressed TLK1 and TLK2 in mammalian physiology remain unexplored. In particular, whether they have distinct, overlapping or entirely redundant roles in particular tissues or cell types, remains

unknown. Here we describe the generation and characterization of both *Tlk1* and *Tlk2* deficient mice and cells that reveal distinct and redundant roles for TLK1 and TLK2 activities in development and genome maintenance.

Results

TLK1 is dispensable for embryogenesis and murine viability

To determine if TLK1 and TLK2 had distinct or redundant roles during development or aging, we generated mice deficient for either *Tlk1* or *Tlk2*. Mice homozygous for a gene-trapped allele of *Tlk1* (*Tlk1^T*) (Figure 1A and B) showed efficient reduction of *Tlk1* expression (Figure 1C). In addition, no TLK1 protein was detectable in Westerns from embryonic fibroblast (MEF) lysates (Figure 1D) and no compensatory changes in *Tlk2* mRNA or protein levels were evident (Figure 1C and D). Despite the implication of TLK1 in many critical cellular processes, *Tlk1^{T/T}* mice were born at normal Mendelian ratios (Figure 1E) and showed no obvious morphological or size differences at birth or during the first months of development (Figure 1F)(4, 13-16, 21-23). In addition, *Tlk1^{T/T}* mice did not exhibit any obvious developmental phenotypes or accelerated morbidity over 18 months (Figure 1G). Both male and female *Tlk1^{T/T}* mice were fertile, indicating that meiotic recombination was largely functional, and *Tlk1^{T/T}* females produced litter sizes similar to that of *Tlk1^{+/+}* and *Tlk1^{+T}* littermates (Figure 1H). Additionally, we examined T-cell development that requires the repair of DNA double-strand breaks (DSBs) induced by the RAG recombinase, as previous work has implicated TLK1 in DNA repair(16, 21-24). Similar to what was observed in MEFs, no TLK1 protein was detectable in T-cells from *Tlk1^{T/T}* mice (Figure 1I) and the distribution of T-cells in *Tlk1^{+/+}* and *Tlk1^{T/T}* deficient mice was indistinguishable (Figure 1J), indicating that non-homologous end-joining (NHEJ) mediated DNA repair in this tissue operates efficiently in the absence of TLK1. While we cannot rule out more subtle effects on repair processes or developmental programs, these data indicated that TLK1 was dispensable for murine development, aging and multiple physiological DNA repair events.

TLK2 is an essential gene

Based on previous data from lower organisms, we expected that loss of TLK1 activity would result in severe phenotypes in the mouse (8, 10). The lack of any clear phenotypes in *Tlk1*^{TT} animals suggested potentially redundant activities, with TLK2 being the most likely candidate. To address this, we generated a conditional, gene-trapped allele of *Tlk2* (Figure 2A, 2B and Supplementary Figure S1) and interbred heterozygous mice to generate *Tlk2* deficient animals. In sharp contrast to *Tlk1*, we did not observe any live births of *Tlk2*^{TT} mice, indicating embryonic lethality (Figure 2C).

To understand the cause of death, we examined embryos at different developmental stages. The number of *Tlk2*^{TT} embryos observed from E10.5 to E13.5 was consistent with expected Mendelian ratios (Figure 2D), although embryos were smaller in all cases (Figure 2E to H). *Tlk2*^{TT} embryos appeared anemic (Figure 2F and G) and perished by E15.5, as no heartbeat was detectable. By E16.5 extensive tissue autolysis was evident (Figure 2H). Fibroblast cultures from E14.5 embryos confirmed that *Tlk2* mRNA and protein levels were reduced and that there was no compensatory increase of TLK1 (Figure 2I and J). The further examination of *Tlk2*^{TT} embryos by conventional immunohistochemistry (IHC) (Figure 2K) and lightsheet-based fluorescence microscopy (Supplementary Figure S2) did not reveal any consistent defects and suggested that embryos were developmentally delayed, but in the majority of cases morphologically normal. Collectively, our data suggested that TLK2, but not TLK1, was essential for embryonic viability.

TLK2 is essential for placental development

As consistent defects in embryonic morphology were not observed, we examined placental function. At E14.5 *Tlk2*^{TT} placentas were smaller and there was a lack of blood circulation in the umbilical cord (Supplementary Figures S3A and S3B). To address this phenotype further without potential complications from the genetrapped allele, we excised it by crossing to FlpO expressing mice, thus rendering Exon 4 floxed (*Tlk2*^F) (Figure 2A). Crosses with mice expressing *Sox2-Cre*, that is active in the embryonic epiblast cells, effectively deleted Exon 4 and yielded *Tlk2*^{+/-} animals that were then interbred to generate *Tlk2*^{-/-} embryos (Supplementary Figure S3C).

We histologically examined *Tlk2*^{-/-} placentas from embryonic days E10.5 to E16.5 and at all stages, *Tlk2*^{-/-} placentas were moderately to severely smaller, thinner and less cellular compared to *Tlk2*^{+/+} or *Tlk2*^{+/-} littermates, and poorly vascularized (Figure 3A to 3E and Supplementary Figure S3). At E10.5 through E12.5 the labyrinths of *Tlk2*^{-/-} placentas were disorganized and composed of less differentiated and less mature trophoblasts (Figure 3D and Supplementary Figure S3). *Tlk2*^{-/-} placentas had moderately to severely reduced numbers of labyrinth trophoblast, syncytiotrophoblast, spongiotrophoblast (including glycogen cells) and mildly to moderately reduced numbers of trophoblastic giant cells and vasculature (Figure 3C, 3D, 3E and Supplementary Figure S3). These data indicated that TLK2 was required for normal placental development and function, and that its loss led to late embryonic lethality

The influence of TLK2 deficiency on proliferation and gene expression in the placenta

As the regulation of ASF1 by TLK activity has been linked to ongoing DNA replication, we examined trophoblast proliferation by IHC using the marker Ki67 (2). By E12.5 most cells of the trophoblastic lineage are differentiated and their rate of proliferation decreases (25, 26). Regardless of genotype, we observed similar percentages of Ki67 positive cells at each stage of development suggesting that the overall number of proliferating cells was similar (Figure 4A). In addition, we did not observe any significant increase in either the DNA double strand break marker γ H2AX or the apoptosis marker cleaved caspase 3 (CC3) (Figure 4B and Supplementary Figure S4A), suggesting that overall levels of proliferation, DNA damage, and cell death were similar.

In addition to DNA replication, TLK2 has been implicated in transcriptional regulation through the phosphorylation of ASF1a (18, 27). To address the possibility that transcription may be affected, we examined the mRNA expression levels of several genes required for trophoblast differentiation; *Cdx2*, *Tpbpa*, *Pl1* and *Pl2* (28). We observed lower levels of *Cdx2*, *Tpbpa*, and *Pl2* in the *Tlk2*^{-/-} placentas, reflecting reduced numbers and/or defective differentiation of trophoblast stem cells, spongiotrophoblasts and canal or sinusoidal trophoblast giant cells, respectively (Figure 4C). In contrast, the expression of *Pl1*, that is a marker for parietal

trophoblast giant cells and is normally reduced by E11.5, was slightly upregulated compared to wild type placenta, although this was more variable amongst individual samples (Figure 4C)(28). In addition, the expression of many imprinted genes that influence placental development, including *Igf2*, *H19*, *Peg10* and *Grb10*, was reduced in *Tlk2*^{-/-} placentas, while others, such as *IgfR2* and *Phlda2*, were expressed at similar levels to that of wild type (Figure 4D)(28, 29). The reduced size of the *Tlk2*^{-/-} labyrinth and reduced area of Tpbpa positive cells was further confirmed with IHC (Figure 4E). Together, these data suggested that the loss of TLK2 did not cause clear defects in trophoblast proliferation or survival, but reduced the expression of numerous genes important for their normal function and led to impaired differentiation and placental function.

TLK2 interacts with ASF1 and influences its phosphorylation in the placenta

To understand why placental function was dependent on TLK2 and not TLK1, we examined their relative mRNA expression and protein levels. We found similar mRNA expression patterns of both genes in the wild type placenta at E13.5, E15.5 and E17.5 (Figure 5A). The protein levels of TLK2 were similar in placental tissue compared to tissue from the embryo (heart and fetal liver), while TLK1 protein levels were strongly increased in the embryonic tissues compared to the placenta, although TLK1 was detectable in placentas (Figure 5B and Supplementary Figure S4B). This suggested that TLK1 was significantly less abundant in the placenta than the embryo, providing a potential explanation as to why placental development was particularly sensitive to *Tlk2* deficiency.

ASF1a and ASF1b are the most clearly defined targets of TLK1 and TLK2 but other substrates, such as RAD9 and NEK1, have been reported for TLK1(15, 23, 30). To address the potential substrate spectrum of TLK2, we carried out immunoprecipitations of tagged TLK2 or kinase dead TLK2 (TLK2-KD) followed by quantitative mass spectrometry (IP-MS, Supplementary Figure S5A). In addition, we overexpressed TLK2 fused to the BirA* biotin ligase to identify the proximity interactome of TLK2 using BioID (Supplementary Figure S5B). Only a small number of proteins were consistently identified by both approaches, including ASF1a and ASF1b, TLK1 and LC8 (DYNLL1) (Figure 5C, Supplementary Figure S5 and

Supplementary Tables S1-S6). Additional proteins known to be involved in a variety of cellular processes, including DNA replication and chromatin assembly (ex. MCM helicase components, RIF1 and HIRA/UBN1-2), transcription (ex. GATAD2A, JUN) and translation (ex. EIF3F, EIF2S1) were identified in one or both approaches below our statistical thresholds (Figure 5C). We focused on the most high confidence interactions and found that while we could confirm the LC8 interaction, it did not appear to be a substrate of TLK2 activity (Supplementary Figure S5E-H). We therefore focused on determining the level and phosphorylation status of ASF1 in normal and *Tlk2* deficient placentas. In *Tlk2*^{-/-} placentas total ASF1 protein levels were similar to wild type controls but the phosphorylation of S166 (p-ASF1), a target site of TLKs, was reduced, suggesting reduced H3/H4 binding potential in the absence of TLK2 (Figure 5D and Supplementary Figure S6A)(11). To confirm the interaction of TLK2 with ASF1 in the placenta, we performed co-immunoprecipitations (IPs) of endogenous TLK2 or ASF1 from placental lysates. Following the IP of TLK2, we readily observed both ASF1 and TLK1 and reciprocal IPs with anti-ASF1 antibodies efficiently co-IPd TLK2 (Figure 5E and Supplementary Figure S6B). Together, these results indicated that despite some levels of detectable TLK1, TLK2 is required for full ASF1 phosphorylation in the placenta, potentially influencing the regulation of genes required for trophoblast differentiation and placental development.

TLK2 is dispensable for viability in embryonic and adult tissues

The differential protein levels of TLK1 between the placenta and embryonic tissue provided a potential explanation for placental dependence on TLK2 (Figure 5B). We considered that a similar situation in other cell types in the embryo could also contribute to the embryonic death of *Tlk2*^{-/-} animals. To address this, we crossed female *Tlk2*^{+/-} mice carrying the Sox2-Cre transgene, that expresses Cre in the epiblast, with males carrying the floxed conditional allele (*Tlk2*^{+F}) to allow the generation of *Tlk2*^{-/-} embryos, while maintaining at least 1 functional copy of *Tlk2* in the placenta (Figure 6A)(31). This strategy bypassed the embryonic lethality and we observed the birth of live, morphologically normal, *Tlk2*^{-/-} animals (Figure 6A). These animals, although sometimes smaller than littermates, showed a similar growth capacity (Figure 6B) and no increased morbidity has been observed up to 18 months (Figure 6C), although we

cannot yet rule out tissue specific or aging related phenotypes that are not overtly evident. Importantly, we could not detect TLK2 protein in any tissues examined by Western blotting, indicating that Sox2-Cre was effective at eliminating *Tlk2* expression (Figure 6D). These data demonstrate that, like TLK1, TLK2 is not required for organismal viability, despite its critical role in the development and function of the placenta.

As relative TLK1 and TLK2 protein levels were different between embryonic and placental tissues, we sought to determine if a similar situation existed in other adult tissues. We freshly isolated mRNA and protein from a panel of wild type tissues at 2 months of age. Using quantitative real time PCR we determined the expression levels of *Tlk1* and *Tlk2* in different tissues (Figure 6E). In general, the expression of *Tlk1* and *Tlk2* followed a similar pattern across tissues, although the expression of *Tlk2* was particularly high in the testes, where the human *TLK2* gene was originally identified (6). To determine if the mRNA levels reflected relative protein levels, we performed Western blotting from lysates prepared from some of the same tissues of both male and female mice (Figure 6F). These results were generally consistent with the mRNA expression levels and the patterns of TLK1 and TLK2 protein levels were similar relative to tissue type. One exception was the kidney, where both TLK1 and TLK2 protein were undetectable, despite similar levels of mRNA expression as other tissues (Figure 6E and 6F). These results suggested that the lack of clear phenotypes in adult animals deficient for either TLK1 or TLK2 could be due to their redundancy in most tissues, in contrast to the placenta where TLK2 is required due to the low levels of TLK1. Supporting this possibility, we found that *Tlk2* is haploinsufficient on a *Tlk1* deficient background, as all *Tlk1*^{T/T} *Tlk2*^{+T} pups we observed displayed severe birth defects (Figure 6G and Supplementary Figure S7).

TLK1 and TLK2 cooperate to maintain cell viability and chromosomal stability

As TLK1 or TLK2 deficient animals appeared phenotypically normal, we wanted to investigate the effects of depleting both activities. To accomplish this, we generated single (*Tlk1*^{+/-}*Tlk2*^{F/-}) and double conditional (*Tlk1*^{C/C}*Tlk2*^{F/-}) transformed MEF lines that allow the 4-hydroxytamoxifen (4OHT) inducible trapping and deletion of *Tlk1* and *Tlk2*, respectively.

Depletion of TLK1 and TLK2 protein levels following 4OHT induced Cre recombinase induction was effective (Figure 7A) and led to reduced colony formation compared to wild type or single mutants for *Tlk2* (Figure 7B).

The simultaneous depletion of TLK1 and TLK2 resulted in a marked increase in DNA double-strand breaks, indicated by increased intensity of γ H2AX foci in the nucleus (Figure 7C and 7D). These cells also displayed mitotic defects, such as chromosome bridges (Figure 7E), as well as increased levels of chromosome aberrations scored in metaphase spreads, particularly chromosome fragments and fusions (Figure 7F). This was accompanied by a reduction in total ASF1 phosphorylation, similar to what was observed in the placenta (Figure 7G). Collectively, our *in vivo* and *in vitro* results indicated that TLK1 and TLK2 activities are largely redundant for cell viability and chromosome maintenance, but that TLK2 is critical for mammalian development through its role in supporting placental function (Figure 7H).

Discussion

Here we describe for the first time, the relative contributions of TLK1 and TLK2 to mammalian embryogenesis. Of the 2 proteins, TLK1 has been the most studied and has been implicated in many critical cellular processes, such as transcription, DNA repair, DNA replication and mitosis(2, 4, 7, 8, 10, 11, 13-16, 21-23, 32-36). While it may play a role directly or indirectly in these processes, our results indicate that TLK1 is not required for essential cellular functions when TLK2 is expressed normally. We did however observe that TLK1 becomes essential in *Tlk2*^{+/-} animals, (Figure 6G) and we believe that this likely reflects a need for TLK1 in supporting placental, or possibly embryonic function, when *Tlk2* is haploinsufficient, but have not investigated this further. Regardless, the loss of TLK1 does not impair the generation of viable, apparently healthy animals, notably impact several physiological repair processes or detectably influence genomic stability.

In contrast to TLK1, TLK2 was strictly required for embryogenesis through its role in supporting placental development. As both *Tlk1* and *Tlk2* mRNAs are readily detectable in every tissue that we (Figure 6E), and others (37), have examined, this essential role for TLK2 was unexpected. We observed reduced ASF1 phosphorylation, impaired gene expression

and histological evidence that indicated a requirement for TLK2 in the proper differentiation of the trophoblast lineages. As we did not observe any indications that proliferation was strongly affected, we believe this is likely through a role in transcriptional regulation(18). We however, cannot rule out the possibility that loss of TLK2 impairs DNA replication to some extent, leading to either cellular attrition or indirect defects on transcription and that TLK2 may influence cell migration(19) or other cellular functions that have yet to be uncovered. A similar differential requirement for ASF1a and ASF1b in development has been previously reported. The depletion of murine ASF1a leads to lethality by E9.5(38), notably earlier than we observed with TLK2 deletion, potentially consistent with the incomplete effect we observed on total ASF1 phosphorylation. In contrast, ASF1b is dispensable for viability, although it affects fertility(39). While mice lacking either TLK1 or TLK2 were fertile, we cannot rule out subtle or age related differences in fertility from existing data. Future analysis of mice lacking combinations of both TLK and ASF1 alleles may be informative for identifying more specific functions and determining how much of the TLK deficient phenotype relies on ASF1 regulation.

Although mRNA levels of *Tlk1* and *Tlk2* were similar from E13.5 to 17.5 of placental development, TLK1 protein levels were much higher in embryonic tissues compared to placenta, while TLK2 protein levels were similar regardless of the tissue. We believe this provides the most parsimonious explanation for the essential nature of TLK2 in the placenta, although we cannot rule out that TLK2 is uniquely capable of modifying a cell type specific substrate or carrying out another unknown, essential role during development. Whether similar unique dependencies of either TLK1 or TLK2 occur in other cell types of embryonic or adult animals remains to be determined, as do the mechanisms that regulate TLK1 or TLK2 protein levels, activity and substrate engagement.

TLK1 activity appears to buffer *Tlk2* haploinsufficiency (Figure 6G), but it is insufficient to fully support placental development in the absence of TLK2. This may be due to its lower levels, and presumably lower activity, relative to embryonic tissue, reduced affinity for ASF1a(27), or due to specific interactions that have yet to be identified. However, in other tissues of the embryo and adult animals, the protein levels of TLK1 and TLK2 appear to be sufficient to act

interchangeably to support cellular functions, evidenced by the viability of both *Tlk1*^{+/+} and *Tlk2*^{-/-} cell cultures and animals (Figures 1, 6 and 7).

While our results suggest that there may be a high level of redundancy between TLK1 and TLK2 in many cell types, recent work has specifically connected *TLK2* mutations to patients with intellectual disabilities(40) and it is tempting to speculate that this could reflect placental defects, although we would expect them to be less dramatic than what we have observed in mice with a full TLK2 knockout. Moreover, TLK2 has been identified as a regulator of starvation induced autophagy(41) and gammaherpesvirus latency(9), amplifications and hyperphosphorylation of specifically TLK2 have been identified in subtypes of breast cancer and shown to influence cancer progression(19, 20, 42, 43) and TLK2 has been shown to play a critical role in cell cycle checkpoint recovery that was not observed following TLK1 depletion(18). These apparently specific roles may reflect differences in their transcriptional or translational regulation and relative ratios, differential control of their activity or distinct substrate specificity. Recent work identified around 150 putative TLK1 interactors(30) and we found only 7 proteins in common using different experimental approaches for TLK2, including ABLIM1, DNAJC7, EMG1, PDCD6, SERBP1, SNRPC and SRSF1, although these fell below the stringent thresholds we used for selecting interactions to validate (Figure 5C and Supplementary Tables S1-S6). Further investigation in a wider array of cell types will be required to understand the unique and overlapping substrates and functions of TLK1 and TLK2 and their specific roles in development, genome maintenance and other cellular processes.

Materials and Methods

Generation of *Tlk1* and *Tlk2* deficient mice

Embryonic stem (ES) cells with a gene-trapped *Tlk1* allele were purchased from the German Genetrap Consortium (clone ID E067A02). A single insertion was confirmed and cells injected into 3.5-day mouse C57B6/j blastocysts (Fig. S1). 12-15 ES cells were injected into each blastocyst and re-implanted into the oviduct of 2.5-day pseudo-pregnant foster mice (CD1s). Chimeras were scored and mated with C57B6/j wild-type mice. Agouti offspring were

321 screened by PCR for the *Tlk1* allele and pups were screened for zygosity using PCR or by
322 quantitative real time PCR (Transnetyx). Primer details available upon request.

323
324 The knockout-first *Tlk2* targeting construct was designed and constructed in the IRB Mutant
325 Mouse facility using standard recombineering methods(44). The targeting vector was
326 confirmed by sequencing and linearized vector transfected into ES cells for targeting. Cells
327 were selected with neomycin and digested genomic DNA analyzed by Southern blot
328 (performed by service in the Centro Ricerche Biotecnologiche of the Università Cattolica del
329 Sacro Cuore) for correct targeting (Fig. S1). For Southern analysis, 3 probes were used and
330 correctly targeted clones injected as described for *Tlk1*. We analyzed the 5' (5P) and 3' (3P)
331 integration sites following KpnI digestion (probe primers in Table S7). *Pgk1-FlpO* (011065),
332 *Sox2-Cre* (008454) were purchased from Jackson laboratories and *ERT2-Cre* mice were
333 previously described (31, 45, 46). All animals were maintained in strict accordance with the
334 European Community (86/609/EEC) guidelines at the animal facilities in the Barcelona
335 Science Park (PCB). Protocols (CEAA13/0008) were approved by the Animal Care and Use
336 Committee of the PCB (IACUC; CEEA-PCB) in accordance with applicable legislation (Law
337 5/1995/GC; Order 214/1997/GC; Law 1201/2005/SG). All efforts were made to minimize use
338 and suffering.

339 **Generation of mouse embryonic fibroblasts (MEFs)**

340 Mouse embryonic fibroblasts (MEFs) cultures were established from E14.5 embryos that were
341 dissected into fresh PBS and kept overnight at 4°C in trypsin/EDTA (Life Technologies)
342 solution. Following incubation at 37°C for 20 minutes, cells were disaggregated with a
343 serological pipette and supernatant plated for culture in DMEM supplemented with 10% (v/v)
344 fetal bovine serum (FBS, Sigma), 2mM L-glutamine and 100 U/ml penicillin. Transformed
345 MEFs were generated by transfecting a linearized p129 plasmid, containing an origin-less
346 SV40 genome, using an Amaxa nucleofector with MEF reagent 2 (Amaxa). For the induction
347 of Cre recombinase in *ERT2-Cre* expressing cell lines, cells were exposed to 4-
348 hydroxytamoxifen (Sigma H7904, 1uM) for 72h, washed, and plated for experiments.

349 **Immunological analysis**

Analysis of thymocytes was performed as described previously(47). Thymi were dissected from 8-week old animals and a single-cell suspension was obtained using a 45 µm nylon mesh (Falcon). Apoptosis of thymocytes was assessed by flow cytometry after double staining with FITC-conjugated Annexin V (BD Biosciences) and propidium iodide. Thymic subpopulations were quantified by staining with fluorescence-conjugated antibodies for CD3e, CD4 and CD8 (BD Biosciences) and analysis by flow cytometry using FlowJo software (Tree Star).

Immunohistochemistry (IHC)

Paraffin-embedded tissue sections (3 µm) were air dried and further dried at 60°C over-night. IHC was performed using an Autostainer Plus (Dako-Agilent) or manually (Tpbpa). Prior to IHC, sections were dewaxed using the low pH EnVision™ FLEX Target Retrieval Solutions for caspase 3 detection (Dako, Burlington) for 20 min at 97°C using a PT Link (Dako-Agilent), or dewaxed and rehydrated and thereafter antigen retrieval was performed using a PT link (Dako-Agilent) for 20 min at 97°C with citrate pH 6 buffer. Quenching of endogenous peroxidase was performed by 10 min with Peroxidase-Blocking Solution (Dako REAL S2023). Primary antibodies rabbit polyclonal anti-Ki67 (Thermo scientific; 1:75), rabbit polyclonal anti-CD31 (Abcam ab28364; 1:500), mouse monoclonal anti-phospho-Histone H2A.X-pSer139 (clone JBW301, Millipore 05-636; 1:600), rabbit polyclonal anti-Cleaved Caspase-3 (Asp175) (Cell Signalling #9661; 1:300) and rabbit polyclonal anti-trophoblast specific protein alpha (Abcam ab104401; 1:1500) were incubated for 120 min at room temperature. BrightVision Poly-HRP-Anti Rabbit IgG Biotin-free (Immunologic, DPVR-110HRP) was used as a secondary antibody. Antigen-antibody complexes were revealed with 3-3'-diaminobenzidine (K3468, Dako), with the same time exposure per antibody (5 min). Sections were counterstained with hematoxylin (Dako, S202084) and mounted with Mounting Medium, Toluene-Free (CS705, Dako) using a Dako CoverStainer. Specificity of staining was confirmed by omission of the primary antibody. Slides were imaged using Olympus DP25 camera mounted on Olympus BX45 microscope or using a E600 Nikon epifluorescence microscope equipped with 40x NA 0.75 objective lens and a charge-coupled Olympus DP72 device camera.

Histopathology

Embryos and placentas were harvested and fixed in 4% paraformaldehyde overnight at room temperature. Representative sections were trimmed, processed, embedded in paraffin, blocked, sectioned, and stained with H&E and periodic acid/Schiff reagent (PAS). Stained slides were examined microscopically, unbiased by 2 experienced board-certified pathologists (S.A.Y., A.d.B). The different types of trophoblastic cell lineage (i.e. labyrinth trophoblast, syncytiotrophoblast, giant cell trophoblast, and trophospongium) were identified based on the characteristic histologic features of every cell. The numbers and/or size of a particular cell from the trophoblastic cell lineage were compared between different genotypes and scored by an ordinal semi-quantitative system as described (48) where normal = no change in the numbers/size from control tissue, mild = decreased between 10-25% , moderate = decreased between 25-50%, and severe = the numbers and/or size of a particular cell type is decreased >50% when compared to control tissue.

Isolation of RNA and quantitative real time PCR (qRT-PCR)

Dissected tissue or primary MEFs were collected on ice, washed in PBS and snap frozen on dry ice. Tissues were disrupted in Tri-Reagent (Sigma) by zirconium beads in a mechanical tissue disruptor (Precellys 24, Bertin technologies). RNA was isolated by centrifugation followed by chloroform extraction, isopropanol precipitation, washing twice in 75% ethanol and resuspension in DEPC treated water. Nucleic acid quantification was performed with a Nanodrop 8000 instrument (ThermoScientific). Reverse transcription reaction was carried out using High Capacity RNA-to-cDNA Kit (ABI), following manufacturers instructions, in a reaction volume of 20ul and with the primers contained in the 2X RT buffer mix, which is composed of dNTPs, random octamers, and oligo(dT)16 and cDNA was stored at -20°C. qPCR was performed using the comparative CT method and a Step-One-Plus real-time PCR Instrument (Applied Biosystems). SYBR Green and Taqman reactions were done in technical triplicate in a final volume of 10 µl. For SYBR Green 1X SYBR Green PCR Master Mix (#4364344, ABI), Forward and Reverse primer (Sigma; 100nM each) and 25 ng of template were used. For TaqMan reactions 1X TaqMan Universal PCR Master Mix (#4324018, ABI), 0.5 µl of probe and 25 ng of template were used. Thermocycling parameters used were: for SYBR Green reactions (95°C 20 sec; 40 cycles 95°C 3 sec, 60°C 30 sec; melting curve) and for TaqMan reactions (95°C 20 sec; 40 cycles 95°C 1 sec, 60°C 20 sec). For TaqMan assays,

mACTB probe was used as an endogenous control for normalization and for SYBR Green assays, primers for 18S were used as endogenous controls for normalization. The 2^{-ddCT} method was used for the analysis of the amplification products. Primer pairs used are indicated in Supplementary Table S7. Data are presented as the means $\pm$ standard deviation of at least 3 experiments, with at least 3 animals within each experimental group. Statistical significance was determined using an unpaired t-test (**p=<0.0001, *p=<0.001, * p=<0.05, n.s.= non significant).

Affinity purification of TLK2 and quantitative mass spectrometry

The human TLK2 cDNA was obtained in a Gateway compatible pENTR223 vector (Sigma Mission cDNA library) and introduced into the N-SF-TAP-DEST vector (a kind gift from J. Gloeckner(49)) by recombination reaction (300 ng TLK2-pENTR223, 300 ng N-SF-TAP DEST, 4 μ l Gateway® buffer (Invitrogen) and 4 μ l clonase LR (Invitrogen)) and incubated 1 h at 25°C. Proteinase K (2 μ l) (Sigma) was added and incubated 10 min at 37°C to stop the reaction. Competent *E. coli* DH5 α (Sigma) were transformed with the resulting recombination reaction and selected with kanamycin. Plasmids were checked by restriction digests and sequencing (Macrogen) and DNA prepared with MaxiPreps (Promega). The kinase dead mutation (D592V) was introduced using QuickChange Lightning (Agilent).

HEK293T cells were grown in DMEM (Life Technologies) supplemented with 10% fetal bovine serum (FBS) and penicillin/streptomycin at 37°C in a 5% CO₂ incubator. Transient transfections of 15 μ g plasmid per 15 cm plate were carried out using polyethylenimine (PEI) (Polysciences Inc.). Cells were harvested 48 h after transfection and collected by centrifugation. Pellets were lysed in 1 ml lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% Tween-20, 0.5% NP-40 with 1x protease inhibitor cocktail (Roche) and 1x phosphatase inhibitor cocktail (Sigma)) on ice for 20 minutes. Cells were freeze-thawed 3 times and lysates were cleared by centrifugation at 16K g for 20 min at 4°C. Proteins were quantified using the DC Protein Assay (BioRad) and 100 μ l was retained for input. Protein extracts were incubated with 100 μ l pre-washed streptactin superflow resin (IBA) for 2 h at 4°C using an overhead tumbler. Resin was spun down (7000xg for 30 s) and transferred to microspin columns (GE Healthcare). Resin was washed 3 times with 500 μ l washing buffer (30 mM Tris, pH 7.4, 150

mM NaCl, 0.1% NP-40, protease and phosphatase inhibitors). Fusion proteins were eluted from the Strep-Tactin matrix with 500 µl of desthiobiotin elution buffer (IBA) (30 mM Tris, pH 7.4, 150 mM NaCl, 1mM EDTA, 2 mM desthiobiotin, protease and phosphatase inhibitors) for 10 min on ice. 150 µl was retained for Westerns and remaining protein was concentrated in centrifugal filter units and taken to a final concentration of 6M Urea, 100mM ammonium bicarbonate. Then they were reduced, alkylated, and digested to peptides using sequence-grade trypsin at 1:10 ratio (w:w; enzyme:substrate) at 37°C overnight. Tryptic peptide mixtures were desalted using a C18 UltraMicroSpin (ThermoFisher Scientific) column(50).

Samples were analyzed in a LTQ-Orbitrap Velos Pro mass spectrometer (ThermoFisher Scientific) coupled to nano-LC (Proxeon) equipped with a reversed-phase chromatography 12-cm column with an inner diameter of 75 µm, packed with 5 µm C18 particles (Nikkyo Technos Co., Ltd.). Chromatographic gradients from 93% buffer A, 7% buffer B to 65% buffer A, 35% buffer B in 60 min at a flow rate of 300 nl/min, in which buffer A: 0.1% formic acid in water and buffer B: 0.1% formic acid in acetonitrile. The instrument was operated in DDA mode and full MS scans with 1 micro scans at resolution of 60,000 were used over a mass range of m/z 250-2,000 with detection in the Orbitrap. Following each survey scan the top twenty most intense ions with multiple charged ions above a threshold ion count of 5000 were selected for fragmentation at normalized collision energy of 35%. Fragment ion spectra produced via collision-induced dissociation (CID) were acquired in the linear ion trap. All data were acquired with Xcalibur software v2.2.

Acquired data were analyzed using the Proteome Discoverer software suite (v1.3.0.339, ThermoFisher Scientific) and the Mascot search engine (v2.3, Matrix Science) were used for peptide identification. Data were searched against an in-house generated database containing all proteins corresponding to human in the SwissProt database plus the most common contaminants, as previously described(51). A precursor ion mass tolerance of 7 ppm at the MS1 level was used, and up to three miscleavages for trypsin were allowed. The fragment ion mass tolerance was set to 0.5 Da. Oxidation of methionine and protein acetylation at the N-terminal were defined as variable modification. Carbamidomethylation on

cysteines was set as a fix modification. The identified peptides were filtered using a FDR < 1%. For protein quantitation, the top3 method was used(52). In brief, the average of the peptide areas for the three most intense peptides per protein was used a measure of protein abundance. Additional data is presented in Supplementary Tables S1 to S4.

BioID analysis of proximity interactions

Human TLK2 was amplified with forward primer containing 5' Ascl (TLK2-Ascl-F:AAAAAAGGCGCGCCTATG-GAAGAATTGCATAGCC) or reverse primer containing 3' NotI restriction sites (TLK2-NotI-R: AAAAAAGCGGCCGCTTA-ATTAGAAGAACTGTT) using KOD Hot Start DNA Polymerase (Millipore) and cycling conditions recommended from the manufacturer (Polymerase activation at 95°C for 2 minutes, denaturation at 95°C for 20 seconds, annealing at 55°C for 10 seconds and extension at 70°C for 50 seconds, repeated for 40 cycles). PCR products were purified using the PureLink Quick Gel Extraction Kit (Invitrogen) and cloned into pCR2.1-TOPO vector (Invitrogen). Top10 competent *E. coli* cells (Invitrogen) were transformed with pCR2.1-TLK2 and colonies were selected in carbenicillin. Constructs were verified by restriction digestion and sequencing (Macrogen) with primers for the TOPO vector (T7 Promoter-F and M13-R). Afterwards, TLK2 was cut from the pCR2.1-TOPO vector by restriction digest with Ascl (NEB) and NotI-HF (NEB), purified using the PureLink Quick Gel Extraction Kit (Invitrogen) and ligated into pcDNA5/FRT/TO-N-FLAG-hBirA* (TLK2) using Quick Ligation Kit (BioLabs). Top10 competent *E. coli* cells (Invitrogen) were transformed with pcDNA5/FRT/TO-N-Flag-hBirA*-TLK2 vector and carbenicillin selected. The constructs were confirmed by restriction digestion with Ascl (NEB) and NotI-HF (NEB) and sequencing (Macrogen).

AD293 cells were seeded and transfected the next day with either pcDNA5/FRT/TO-N-FLAG-hBirA*-TLK2 or pcDNA5/FRT/TO-N-FLAG-hBirA* using PEI (Sigma) +/- 50µM biotin (IBIANTechnology 2-1016-002). For mass spectrometry, 5x15cm plates per condition were harvested 24 hours post-transfection by scraping cells into PBS, washing 2X in PBS and snap freezing on dry ice. Cell pellets were lysed in 5 ml modified RIPA buffer (1% TX-100, 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 0.1%SDS, 0.5% Sodium deoxycholate and protease inhibitors) on ice, treated with 250 U benzonase (EMD Millipore)

and biotinylated proteins were isolated using streptavidin-sepharose beads (GE). Proteins were washed in Ammonium bicarbonate and digested with trypsin. Mass spectrometry was performed as previously described(53). Additional data is presented in Supplementary Tables S5 and S6.

Endogenous co-immunoprecipitation

Dissected placentas were collected on ice and disrupted in Extraction Buffer (300 mM NaCl, 0.5% NP-40, 50 mM Tris pH 7.8, 0.2 mM EDTA, 5% glycerol, 1x protease (Roche) and phosphatase inhibitors (Sigma)) by zirconium beads in a mechanical tissue disruptor (Precellys 24, Bertin technologies). After two centrifugation steps (10 min, max speed, 4 °C and 5 min, max speed, 4°C), supernatants were collected, precleared, and immunoprecipitated (overnight, 4 °C, on rotation) with 10 µg of anti-TLK2 antibody (Bethyl Laboratories A301-257A), 10 µl of polyclonal ASF1 serum (generated in the Groth laboratory) or 10 µg of rabbit IgG (Sigma) bound to 50 µl of Dynabeads Protein G (Invitrogen). Beads were washed five times with cold Wash Buffer (150mM NaCl, 0.2% NP-40, 50mM Tris pH 7.8, 0.2 mM EDTA, 5% glycerol, 1x protease (Roche) and phosphatase inhibitors (Sigma)), resuspended in 30 µl of Laemmli Sample Buffer and boiled for 20 min.

Protein extraction for Western blotting

Dissected tissues or cells were collected on ice, washed in PBS and resuspended in TNG-150 buffer (50 mM Tris/Hcl, 150 mM NaCl, 1% Tween-20, 0.5% NP-40, 1x protease (Roche) and phosphatase inhibitors (Sigma)). Tissues were disrupted by zirconium beads in a mechanical tissue disruptor (Precellys 24, Bertin technologies). Protein concentration was quantified using the DC Protein Assay (Biorad), separated by SDS-PAGE and transferred to PVDF membrane (Millipore). Membranes were probed with antibodies for TLK1 (Cell Signaling 4125, 1:1000), TLK2 (Bethyl Laboratories A301-257A, 1:1000), ASF1 (Sigma A5236, 1:800, for HEK-293T human cells; Groth laboratory rabbit polyclonal, 1:2000, for mouse tissue and MEFs)(14), Actin (Sigma A4700, 1:3000), FLAG (Sigma F3165, 1:1000), LC8 (Abcam ab51603, 1:1000), GAPDH (Millipore MAB374, 1:10000) and p-ASF1 S166 (Groth laboratory rabbit polyclonal, 1:30000)(11). Primary antibodies were detected with appropriate secondary antibodies conjugated to HRP and visualized by ECL-Plus (GE).

Analysis of genomic instability

For colony formation assays, cells were seeded in duplicate on 6-cm plates and the number of seeded cells adjusted to the plating efficiency. Cells were incubated at 37°C for 10-14 days and colonies stained with crystal violet (Sigma-Aldrich) and counted using ImageJ-plugin Trainable Weka Segmentation. For immunofluorescence and high-throughput microscopy (HTM), MEFs were grown in LabTek II Chamber slides (Labclinics). Cells were pre-extracted for 5 min in 0.2% TX-100/PBS on ice and fixed for 10 min in 4% paraformaldehyde at room temperature. Fixed cells were blocked in 3% BSA with 0.1% Tween/PBS for 1 h and stained using standard procedures. For HTM, 48 images were automatically acquired per well with a Scan[^]R (Olympus) at 40× magnification and non-saturating conditions. Images were segmented using the DAPI staining to generate nuclei masks from which the corresponding signals were calculated using an in-house—developed package based on Cell Profiler. For immunofluorescence analysis the antibody anti- γ H2AX (sc-101696, Santa Cruz) and the corresponding secondary antibody conjugated with Alexa Fluor 568 were used. For metaphase spread preparations, cells were treated with colcemid (0.1 g/ml for 2 h). Cells were trypsinized, hypotonically swollen in 0.075 M KCl for 20 min at 37°C and fixed (75% MeOH and 25% acetic acid, ice cold). Metaphase preparations were spread on glass slides and stained with DAPI in Vectashield mounting medium. Images were taken using a Leica DM6000 microscope and analyzed using Fiji software.

Acknowledgements

We are grateful to the Stracker lab, I.del Barco, A.Igea, J.Roig and C.Gekas for helpful input, A.Nebreda for critical reading of the manuscript, R.Medema for sharing unpublished results, S.Aivio for animal maintenance, D.Reverter and P.Gallego Alonso for purified LC8 protein, J. Gloeckner for SF-TAP vectors, the IRB Histopathology facility and N.Prats, M.Aguilera, B.Domínguez, E.Llonch and N.Vasconcelos for assistance, The IRB Advanced Digital Microscopy Core Facility for use of the Lightsheet Microscope, L.Bardia, M.Marsal and S.Tosi for sample prep and image analysis and E.Sabido for technical advice and mass spectrometry measurements and analysis performed in the CRG/UPF Proteomics Unit, part of the “Plataforma de Recursos Biomoleculares y Bioinformàtics (ProteoRed-Instituto de Salud Carlos III, PT13/0001). T.H.S. was a Ramon y Cajal scholar and supported by the

Ministerio de Economía y Competitividad (MINECO) (BFU2012-39521, BFU2015-68354, Ayudas para incentivar la incorporación estable de doctores (IED) 2015) and institutional funding from MINECO through the Centres of Excellence Severo Ochoa award and from the CERCA Programme of the Catalan Government. A.G. is an EMBO Young Investigator, and supported by the Danish National Research Foundation to the Center for Epigenetics [DNRF82], the European Commission ITN FP7 'aDDress', a European Research Council Starting Grant [ERC2011StG, no. 281,765], the Danish Cancer Society, the Danish Medical Research Council, the Novo Nordisk Foundation and the Lundbeck Foundation, S.S.B. and H.G. were supported by PhD fellowships from Fundacio La Caixa, P.A.K. was supported by an Advanced Postdoc Mobility fellowship from the Swiss National Science Foundation and the Kurt und Senta Herrmann Stiftung, and K.R. was supported by an FPI fellowship (MINECO).

Conflict of interest

The authors declare no conflict of interest.

Supplementary Information

Supplementary information is available at Cell Death & Differentiation's website.

Author Contributions

SSB, PAK and HG designed, performed and analyzed most of the experiments, HG and AP carried out IP-MS experiments, PAK, EC and BR carried out BioID experiments, SAY and AdB assessed placental pathology, KR performed experiments, LP and PAK maintained mouse cohorts, JC performed light-sheet imaging, AG, TLR and SF generated and provided critical reagents and experimental advice and THS designed and analyzed experiments and wrote the manuscript.

References

1. Roe JL, Rivin CJ, Sessions RA, Feldmann KA, Zambryski PC. The *Tousled* gene in *A. thaliana* encodes a protein kinase homolog that is required for leaf and flower development. *Cell*. 1993;75(5):939-50.
2. Sillje HH, Takahashi K, Tanaka K, Van Houwe G, Nigg EA. Mammalian homologues of the plant *Tousled* gene code for cell-cycle-regulated kinases with maximal activities linked to ongoing DNA replication. *EMBO J*. 1999;18(20):5691-702.
3. Sillje HH, Nigg EA. Identification of human Asf1 chromatin assembly factors as substrates of *Tousled*-like kinases. *Curr Biol*. 2001;11(13):1068-73.
4. Li Y, DeFatta R, Anthony C, Sunavala G, De Benedetti A. A translationally regulated *Tousled* kinase phosphorylates histone H3 and confers radioresistance when overexpressed. *Oncogene*. 2001;20(6):726-38.
5. Delacroix S, Wagner JM, Kobayashi M, Yamamoto K, Karnitz LM. The Rad9-Hus1-Rad1 (9-1-1) clamp activates checkpoint signaling via TopBP1. *Genes Dev*. 2007;21(12):1472-7.
6. Shalom S, Don J. Tlk, a novel evolutionarily conserved murine serine threonine kinase, encodes multiple testis transcripts. *Mol Reprod Dev*. 1999;52(4):392-405.
7. Han Z, Riefler GM, Saam JR, Mango SE, Schumacher JM. The *C. elegans* *Tousled*-like kinase contributes to chromosome segregation as a substrate and regulator of the Aurora B kinase. *Curr Biol*. 2005;15(10):894-904.
8. Han Z, Saam JR, Adams HP, Mango SE, Schumacher JM. The *C. elegans* *Tousled*-like kinase (TLK-1) has an essential role in transcription. *Curr Biol*. 2003;13(22):1921-9.
9. Dillon PJ, Gregory SM, Tamburro K, Sanders MK, Johnson GL, Raab-Traub N, et al. *Tousled*-like Kinases Modulate Reactivation of Gammaherpesviruses from Latency. *Cell Host Microbe*. 2013;13(2):204-14.
10. Carrera P, Moshkin YM, Gronke S, Sillje HH, Nigg EA, Jackle H, et al. *Tousled*-like kinase functions with the chromatin assembly pathway regulating nuclear divisions. *Genes Dev*. 2003;17(20):2578-90.
11. Klimovskaia IM, Young C, Stromme CB, Menard P, Jasencakova Z, Mejlvang J, et al. *Tousled*-like kinases phosphorylate Asf1 to promote histone supply during DNA replication. *Nature communications*. 2014;5:3394.

614 12. Pilyugin M, Demmers J, Verrijzer CP, Karch F, Moshkin YM. Phosphorylation-
615 mediated control of histone chaperone ASF1 levels by Tousled-like kinases. *PLoS One*.
616 2009;4(12):e8328.

617 13. Groth A, Lukas J, Nigg EA, Sillje HH, Wernstedt C, Bartek J, et al. Human Tousled
618 like kinases are targeted by an ATM- and Chk1-dependent DNA damage checkpoint. *EMBO J*.
619 2003;22(7):1676-87.

620 14. Krause DR, Jonnalagadda JC, Gatei MH, Sillje HH, Zhou BB, Nigg EA, et al.
621 Suppression of Tousled-like kinase activity after DNA damage or replication block requires
622 ATM, NBS1 and Chk1. *Oncogene*. 2003;22(38):5927-37.

623 15. Kelly R, Davey SK. Tousled-like kinase-dependent phosphorylation of Rad9 plays a
624 role in cell cycle progression and G2/M checkpoint exit. *PLoS One*. 2013;8(12):e85859.

625 16. Sunavala-Dossabhoy G, De Benedetti A. Tousled homolog, TLK1, binds and
626 phosphorylates Rad9; TLK1 acts as a molecular chaperone in DNA repair. *DNA Repair*
627 (Amst). 2009;8(1):87-102.

628 17. Abreu CM, Kumar R, Hamilton D, Dawdy AW, Creavin K, Eivers S, et al. Site-specific
629 phosphorylation of the DNA damage response mediator rad9 by cyclin-dependent kinases
630 regulates activation of checkpoint kinase 1. *PLoS Genet*. 2013;9(4):e1003310.

631 18. Bruinsma W, van den Berg J, Aprelia M, Medema RH. Tousled-like kinase 2
632 regulates recovery from a DNA damage-induced G2 arrest. *EMBO Rep*. 2016;17(5):659-70.

633 19. Kim JA, Tan Y, Wang X, Cao X, Veeraraghavan J, Liang Y, et al. Comprehensive
634 functional analysis of the tousled-like kinase 2 frequently amplified in aggressive luminal
635 breast cancers. *Nature communications*. 2016;7:12991.

636 20. Kim JA, Anurag M, Veeraraghavan J, Schiff R, Li K, Wang X. Amplification of TLK2
637 Induces Genomic Instability via Impairing the G2/M Checkpoint. *Mol Cancer Res*. 2016.

638 21. Sunavala-Dossabhoy G, Balakrishnan SK, Sen S, Nuthalapaty S, De Benedetti A.
639 The radioresistance kinase TLK1B protects the cells by promoting repair of double strand
640 breaks. *BMC Mol Biol*. 2005;6:19.

641 22. Sen SP, De Benedetti A. TLK1B promotes repair of UV-damaged DNA through
642 chromatin remodeling by Asf1. *BMC Mol Biol*. 2006;7:37.

643 23. Canfield C, Rains J, De Benedetti A. TLK1B promotes repair of DSBs via its
644 interaction with Rad9 and Asf1. *BMC Mol Biol.* 2009;10:110.

645 24. Ronald S, Sunavala-Dossabhoy G, Adams L, Williams B, De Benedetti A. The
646 expression of Tousled kinases in CaP cell lines and its relation to radiation response and DSB
647 repair. *Prostate.* 2011;71(13):1367-73.

648 25. Cross JC. Genetic insights into trophoblast differentiation and placental
649 morphogenesis. *Semin Cell Dev Biol.* 2000;11(2):105-13.

650 26. Simmons DG, Cross JC. Determinants of trophoblast lineage and cell subtype
651 specification in the mouse placenta. *Dev Biol.* 2005;284(1):12-24.

652 27. Abascal F, Corpet A, Gurard-Levin ZA, Juan D, Ochsenbein F, Rico D, et al.
653 Subfunctionalization via adaptive evolution influenced by genomic context: the case of histone
654 chaperones ASF1a and ASF1b. *Molecular biology and evolution.* 2013;30(8):1853-66.

655 28. Simmons DG, Fortier AL, Cross JC. Diverse subtypes and developmental origins of
656 trophoblast giant cells in the mouse placenta. *Dev Biol.* 2007;304(2):567-78.

657 29. Cleaton MA, Edwards CA, Ferguson-Smith AC. Phenotypic outcomes of imprinted
658 gene models in mice: elucidation of pre- and postnatal functions of imprinted genes. *Annual*
659 *review of genomics and human genetics.* 2014;15:93-126.

660 30. Singh V, Connelly ZM, Shen X, De Benedetti A. Identification of the proteome
661 complement of humanTLK1 reveals it binds and phosphorylates NEK1 regulating its activity.
662 *Cell Cycle.* 2017:0.

663 31. Hayashi S, Lewis P, Pevny L, McMahon AP. Efficient gene modulation in mouse
664 epiblast using a Sox2Cre transgenic mouse strain. *Mech Dev.* 2002;119 Suppl 1:S97-S101.

665 32. Kodym R, Mayerhofer T, Ortmann E. Purification and identification of a protein kinase
666 activity modulated by ionizing radiation. *Biochem Biophys Res Commun.* 2004;313(1):97-103.

667 33. Sunavala-Dossabhoy G, Li Y, Williams B, De Benedetti A. A dominant negative
668 mutant of TLK1 causes chromosome missegregation and aneuploidy in normal breast
669 epithelial cells. *BMC Cell Biol.* 2003;4:16.

670 34. Sunavala-Dossabhoy G, Fowler M, De Benedetti A. Translation of the radioresistance
671 kinase TLK1B is induced by gamma-irradiation through activation of mTOR and
672 phosphorylation of 4E-BP1. *BMC Mol Biol.* 2004;5:1.

673 35. De Benedetti A. Tousled kinase TLK1B mediates chromatin assembly in conjunction
674 with Asf1 regardless of its kinase activity. *BMC Res Notes*. 2010;3:68.

675 36. Li Z, Gourguechon S, Wang CC. Tousled-like kinase in a microbial eukaryote
676 regulates spindle assembly and S-phase progression by interacting with Aurora kinase and
677 chromatin assembly factors. *J Cell Sci*. 2007;120(Pt 21):3883-94.

678 37. Petryszak R, Keays M, Tang YA, Fonseca NA, Barrera E, Burdett T, et al. Expression
679 Atlas update--an integrated database of gene and protein expression in humans, animals and
680 plants. *Nucleic Acids Res*. 2016;44(D1):D746-52.

681 38. Hartford SA, Luo Y, Southard TL, Min IM, Lis JT, Schimenti JC. Minichromosome
682 maintenance helicase paralog MCM9 is dispensible for DNA replication but functions in germ-
683 line stem cells and tumor suppression. *Proc Natl Acad Sci U S A*. 2011;108(43):17702-7.

684 39. Messiaen S, Guiard J, Aigueperse C, Fliniaux I, Tourpin S, Barroca V, et al. Loss of
685 the histone chaperone ASF1B reduces female reproductive capacity in mice. *Reproduction*.
686 2016;151(5):477-89.

687 40. Lelieveld SH, Reijnders MR, Pfundt R, Yntema HG, Kamsteeg EJ, de Vries P, et al.
688 Meta-analysis of 2,104 trios provides support for 10 new genes for intellectual disability.
689 *Nature neuroscience*. 2016;19(9):1194-6.

690 41. McKnight NC, Jefferies HB, Alemu EA, Saunders RE, Howell M, Johansen T, et al.
691 Genome-wide siRNA screen reveals amino acid starvation-induced autophagy requires
692 SCOC and WAC. *EMBO J*. 2012;31(8):1931-46.

693 42. Mertins P, Mani DR, Ruggles KV, Gillette MA, Clauser KR, Wang P, et al.
694 Proteogenomics connects somatic mutations to signalling in breast cancer. *Nature*.
695 2016;534(7605):55-62.

696 43. Srihari S, Singla J, Wong L, Ragan MA. Inferring synthetic lethal interactions from
697 mutual exclusivity of genetic events in cancer. *Biology direct*. 2015;10(1):57.

698 44. Copeland NG, Jenkins NA, Court DL. Recombineering: a powerful new tool for
699 mouse functional genomics. *Nat Rev Genet*. 2001;2(10):769-79.

700 45. Wu Y, Wang C, Sun H, LeRoith D, Yakar S. High-efficient FLPO deleter mice in
701 C57BL/6J background. *PLoS One*. 2009;4(11):e8054.

46. Indra AK, Warot X, Brocard J, Bornert JM, Xiao JH, Chambon P, et al. Temporally-controlled site-specific mutagenesis in the basal layer of the epidermis: comparison of the recombinase activity of the tamoxifen-inducible Cre-ER(T) and Cre-ER(T2) recombinases. *Nucleic Acids Res.* 1999;27(22):4324-7.
47. Knobel PA, Belotserkovskaya R, Galanty Y, Schmidt CK, Jackson SP, Stracker TH. USP28 is recruited to sites of DNA damage by the tandem BRCT domains of 53BP1 but plays a minor role in double-strand break metabolism. *Mol Cell Biol.* 2014;34(11):2062-74.
48. Gibson-Corley KN, Olivier AK, Meyerholz DK. Principles for valid histopathologic scoring in research. *Vet Pathol.* 2013;50(6):1007-15.
49. Gloeckner CJ, Boldt K, Schumacher A, Roepman R, Ueffing M. A novel tandem affinity purification strategy for the efficient isolation and characterisation of native protein complexes. *Proteomics.* 2007;7(23):4228-34.
50. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat Protoc.* 2007;2(8):1896-906.
51. Bunkenborg J, Garcia GE, Paz MI, Andersen JS, Molina H. The minotaur proteome: avoiding cross-species identifications deriving from bovine serum in cell culture models. *Proteomics.* 2010;10(16):3040-4.
52. Silva JC, Gorenstein MV, Li GZ, Vissers JP, Geromanos SJ. Absolute quantification of proteins by LCMSE: a virtue of parallel MS acquisition. *Mol Cell Proteomics.* 2006;5(1):144-56.
53. Coyaud E, Mis M, Laurent EM, Dunham WH, Couzens AL, Robitaille M, et al. BioID-based Identification of Skp Cullin F-box (SCF) β -TrCP1/2 E3 Ligase Substrates. *Mol Cell Proteomics.* 2015;14(7):1781-95.

FIGURE LEGENDS

Figure 1. TLK1 is dispensable for murine development. (A) Schematic of the *Tlk1* genetrap allele (*Tlk1*^T). The position of the Flex-vector genetrap cassette in the *Tlk1* locus and approximate positions of primers used for quantitative real time PCR are shown. (B) The genetrap clone was screened for single copy insertion using a probe for the neomycin cassette in a Southern blot of EcoRV digested genomic DNA. A single band of the expected size was observed (red arrow). GTF2 is an unrelated line with a neomycin cassette (positive control) and C57BL6 is wild type genomic DNA (negative control). (C) Quantitative real time PCR analysis of *Tlk1* and *Tlk2* levels in mouse embryonic fibroblast (MEF) cultures derived from littermates of the indicated genotype. The mean (red bar) and standard deviation (SD) of 3 replicates are plotted. (D) Western blotting of TLK1 and TLK2 protein levels in MEF cultures of the indicated genotype. Ponceau red stained blot showed equal loading. (E) *Tlk1*^{T/T} mice are born at the expected (exp.) Mendelian ratios. Numbers of pups of the indicated genotype observed (obs.) from 32 litters of *Tlk1*^{+T} mice (110 pups total) are indicated. Percentage observed is indicated above the bar graphs. (F) Normal weight and growth of *Tlk1*^{T/T} mice. Weights of littermate animals at 18 and 30 days post partum are plotted with the mean (red bars) and SD indicated (n=16 for *Tlk1*^{+/+} and 15 for *Tlk1*^{T/T} respectively). (G) Normal survival of *Tlk1*^{T/T} mice. Kaplan Meier plot of animal survival over 18 months (n=25, 41 and 26 for *Tlk1*^{+/+}, *Tlk1*^{+T} and *Tlk1*^{T/T} respectively). (H) Normal litter sizes are observed in *Tlk1*^{T/T} female mice of 2-5 months of age compared to wild type of heterozygous littermates (n=20, 11 and 11 for *Tlk1*^{+/+}, *Tlk1*^{+T} and *Tlk1*^{T/T} respectively). The mean (red bars) and SD are plotted and statistical analysis using the Wilcoxon Rank Sum test indicated no significant differences (p=0.65; *Tlk1*^{+/+} vs. *Tlk1*^{+T}, 0.75; *Tlk1*^{+/+} vs. *Tlk1*^{T/T} and 0.52; *Tlk1*^{+T} vs. *Tlk1*^{T/T}). (I) TLK1 protein levels in thymocytes of the indicated genotype are shown. Actin serves as a loading control. (J) Analysis of T-cell subsets in 2-month old animals of the indicated genotype (n=3 per genotype, each in triplicate). The mean (red bars) and SD are shown and representative flow cytometry data showing CD4 and CD8 staining of CD3 positive T-cells is shown in the right panel.

Figure 2. TLK2 is an essential gene. (A) Schematic of the *Tlk2* knockout first allele (*Tlk2*^T). The position of the genetrap cassette, the allelic configurations and their designations following FLP and CRE expression are shown. (B) Southern blot of *SpeI* digested genomic DNA with a probe for Neomycin to identify clones with single insertions. 6 targeted clones and a negative control (C57BL/6) are shown and the expected size indicated. Additional Southern blots confirming correct targeting are shown in Supplementary Figure S1. (C) *Tlk2* deficiency is embryonic lethal. The number of animals expected (exp.) from 27 *Tlk2*^{+T} breedings (112 pups total) assuming normal Mendelian inheritance is shown compared to the number of observed pups (obs.). Percentage observed is indicated above the bar graphs. Based on the genotypes of a total of 112 pups from 14 independent litters, a p-value of <0.0001 was determined using binomial distribution and indicated by ***. (D) Normal Mendelian distribution of genotypes was observed amongst 41 E10.5 to E13.5 embryos from 6 individual *Tlk2*^{+T} breedings. Percentage observed is indicated above the bar graphs. (E) Examples of littermate E10.5 embryos (scale bar = 1mm) (F) E12.5 embryos (scale bar = 1mm) (G) E15.5 embryos (scale bar = 2mm) and (H) E16.5 embryos (scale bar = 2mm). (I) Quantitative real time PCR analysis of *Tlk1* and *Tlk2* levels in MEFs from littermates of the indicated genotype. The mean (red bars) and standard deviation (SD) of 3 replicates are plotted. (J) Western blotting of TLK1 and TLK2 protein levels in MEFs of the indicated genotype. Histone H3 is shown as a loading control and a non-specific (NS) band recognized by the TLK2 antibody is indicated. (K) H&E (left panels) or Ki67 immunohistochemical staining (right panels) of E12.5 embryo sections of the indicated genotype (scale bar = 1 mm).

Figure 3. TLK2 is required for placental development. (A) Semi-quantitative analysis of histological phenotypes in *Tlk2*^{-/-} placentas. In each case, *Tlk2*^{-/-} placentas (n=10) at the indicated age were compared to wild type littermates. Phenotypes are abbreviated as follows: Placental thickness (Pl.Th), numbers/size of Labyrinth trophoblasts (LT), Syncytiotrophoblasts (Syn.T), Spongiotrophoblasts including glycogen cells (TS), Giant cell trophoblasts (Gi.T) and Vasculature (Vas). Summary of the grading system used (48) provided in the Materials and Methods. (B) Sections from E11.5 and E12.5 littermate placentas stained with H&E. *Tlk2*^{-/-} placentas are thinner, less cellular and poorly vascularized compared to *Tlk2*^{+/+}. Scale bars =

500 microns. (C) CD31 staining reveals that the vasculature of *Tlk2*^{-/-} placentas is miniaturized, collapsed and appears slit-like (red arrowheads) compared to the well-developed, widely opened vasculature of *Tlk2*^{+/+} (blue arrowheads). Scale bar = 20 microns. (D) H&E staining from the indicated age and genotype showing fewer and smaller trophoblasts (arrows) and Syncytiotrophoblasts (blue arrowheads) with an absence of fetal vasculature (red arrowhead) in *Tlk2*^{-/-} placentas. Scale bar = 20 microns. (E) H&E staining of placentas showing a thinner Trophospongium (Spongiotrophoblasts and glycogen cells) in *Tlk2*^{-/-} placentas. The Trophospongium (TS), labyrinth (L) and decidua (D) are indicated. Scale bar = 50 microns.

Figure 4. Proliferation and gene expression in placentas lacking TLK2. (A) Immunohistochemical staining of the proliferative marker Ki67 or (B) the DNA double-strand break marker γ H2AX in E10.5, E11.5 and E12.5 placentas of the indicated genotype. Similar numbers of proliferative cells are visible in the labyrinth of *Tlk2*^{-/-} placentas compared to wild type at each stage. Mean percent positive cells (red bars) and SD is graphed in right panels. n=3 and 2 at E10.5, n=3 and 3 at E11.5 and n=5 and 6 at E12.5 for *Tlk2*^{+/+} and *Tlk2*^{-/-}, respectively. At least 200 total cells were counted for each individual and scale bar in (A) = 10 microns and in (B) = 100 microns. (C) and (D) Quantitative real time PCR analysis of genes expressed in trophoblast lineages from E12.5 placenta samples. Combined data represents the mean of triplicate samples from 3 individual animals of each genotype. Statistical significance was determined using an unpaired t-test (***p<0.0001, **p<0.001, * p<0.05). (E) Immunohistochemical staining for Tpbpa reveals a thinner trophospongium (TS) in the absence of TLK2 at both E10.5 and E12.5. The orientation of the labyrinth (L) is also indicated. Quantification of signal intensity distribution in the TS is shown in the right panel, normalized to area. Scale bar = 500 microns.

Figure 5. TLK2 interacts with ASF1 and influences its phosphorylation in the placenta. (A) Quantitative real time PCR of *Tlk1* and *Tlk2* in wild type placentas of the indicated age. Mean (red bars) and SD of triplicate samples is graphed (n=1, 3 and 1 for each age indicated). (B) Western blotting of TLK1, TLK2 and ASF1 in lysates from E13.5 placenta (Pon. = Ponceau

red stained membrane). TLK2 levels are similar in either tissue (placenta = Pla. and heart/fetal liver = h-l) while TLK1 levels are much higher in embryonic tissue relative to placenta. Samples from 2 littermate wild type placentas are shown. Additional examples are shown in Supplementary Figure S4B. (C) Graphic depiction of the combined results from multiple proteomics approaches to identify TLK2 interacting proteins from HEK293T or AD293 cells (IP-MS and BioID approaches and data are described in further detail in Supplementary Figure S5 and Supplementary Tables S1-S6). TLK1, ASF1a, ASF1b and DYNLL1 (LC8) are the only high confidence interacting proteins identified. LC8 does not appear to be a substrate of TLK2 (Supplementary Figure S5G and S5H). (D) Reduced ASF1-S166 phosphorylation (p-ASF1) in the E12.5 and E13.5 *Tlk2*^{-/-} placentas. Western blotting of lysates for TLK2, ASF1 and p-ASF1 from littermates of the indicated age. In each case, blots shown are from the same membrane (Pon. = Ponceau red stained membrane). Quantification of p-ASF1/ASF1 levels in multiple samples of the indicated genotype is shown in the right panel (additional Western examples are shown in Supplementary Figure S6A). Statistical significance (p=0.0064) was determined using an unpaired t-test and the mean (red bars) and SD are indicated (n=14 and 11, respectively). (E) Co-immunoprecipitation of endogenous TLK2 and ASF1 from wild type E13.5 placental lysates using antibodies for ASF1 (top panels) or TLK2 (bottom panels). (Pon. = Ponceau red stained membrane). Additional examples are shown in Figure S6B.

Figure 6. TLK2 is dispensable in embryonic and adult tissues. (A) *Tlk2*^{+/+} and *Tlk2*^{-/-} littermates are shown from breedings of female *Tlk2*^{+/+}, Sox2-Cre⁺ and male *Tlk2*^{+/F} mice. (B) Weight of mice lacking TLK2 is slightly reduced but growth is similar to wild type. Mean (red bars) and SD are indicated (n=20 and 9 for 1 mo., n=10 and 7 for 2 mo., n=6 and 5 for 3 mo., n=5 and 4 for 4 mo. for *Tlk2*^{+/+} and *Tlk2*^{-/-}, respectively) (C) Kaplan Meier survival curve of mice with the indicated genotypes (*Tlk2*^{+/+} n=42, *Tlk2*^{-/-} n=26). (D) Western blotting of TLK1 and TLK2 using lysates from selected tissues of *Tlk2*^{+/+} and *Tlk2*^{-/-} mice. NS refers to a non-specific band observed with the TLK2 antibody in many tissues and cell lines. (E) Similar expression patterns of *Tlk1* and *Tlk2* in adult mice. Taqman real time PCR analysis of *Tlk1* and *Tlk2* mRNA levels (Log2 transformed) relative to *ActB*. Mean (blue bars for *Tlk1* and red

bars for *Tlk2*) and SD of triplicate reactions from at least 2 animals are plotted. In each case, levels are normalized to that of the liver (set to 1). (F) Western blotting of TLK1/2 protein levels in 2-month old male and female wild type tissues. GAPDH is shown as a loading control. (G) Examples of littermate *Tlk1^{+/+} Tlk2^{+/+}* and *Tlk1^{+/+} Tlk2^{+/-}* animals at different ages. All *Tlk1^{+/+} Tlk2^{+/-}* animals (n=4) observed were severely runted and 1 lacked limbs on the left torso. All mice were sacrificed prior to 1 month following veterinary advice due to lack of mobility and trembling. A Kaplan Meier survival plot for all related genotypes is presented in Supplementary Figure S7.

Figure 7. TLK1 and TLK2 cooperate to maintain cell viability and chromosome stability. (A) Western Blot with antibodies against TLK1 and TLK2 in whole cell extracts of transformed MEFs of the indicated genotypes mock treated or treated with 4OHT to induce trapping of TLK1 and/or deletion of TLK2 (Pon. = Ponceau red stained membrane). (B) Deletion of TLKs reduces colony formation. MEFs were mock treated or treated with 4OHT for 72h, washed, plated and cultured for 10-14 days. Relative colony number to mock treated cells is displayed. *Tlk1^{+/+} Tlk2^{+/+} Cre^{+/-}* n= 3; *Tlk1^{C/C} Tlk2^{F/-} Cre^{+/-}* n=6; *Tlk2^{F/-} Cre^{+/-}* n=6. Results are shown as mean $\pm$ SEM. Experiment was performed in technical duplicates, at least in biological triplicate. Statistical significance was determined using an unpaired t-test (****p<0.0001, ***p<0.001). (C) Representative images of IF experiments staining for γ H2AX are shown from *Tlk1^{+/+} Tlk2^{F/-}* cells that were mock treated or treated with 4OHT. Nuclei were counterstained with 4',6-diamidino- 2-phenylindole (DAPI). (D) Increased DNA double-strand break formation following the depletion of TLK1 and TLK2. High throughput microscopy of γ H2AX levels per individual nucleus in response to TLK depletion in MEFs mock treated or treated with 4OHT. At least 800 nuclei were quantified per condition. Mean is shown as a red line; a.u., arbitrary units. Statistical significance was determined using Ordinary one-way ANOVA Tukey's multiple comparisons test (****p<0.0001). (E) Increased chromosome bridges are evident in 4OHT induced double TLK deficient cultures relative to mock treated or *Tlk2* deletion alone. A minimum of 200 cells was scored in each independent experiment. *Tlk1^{+/+} Tlk2^{+/+} Cre^{+/-}* n= 2; *Tlk1^{C/C} Tlk2^{F/-} Cre^{+/-}* n=5; *Tlk2^{F/-} Cre^{+/-}* n=4. Results are shown as mean $\pm$ SD. Statistical significance was determined using an unpaired t-test (*p<0.05). (F) Plot

of chromosomal aberration types scored from multiple transformed MEF cultures depicted as percent aberrations per chromosome. Results are shown as mean $\pm$ SEM. Statistical significance was determined using Fisher's exact test (****=p<0.0001). (G) Western blotting of *Tlk1*^{C/C} *Tlk2*^{F/-} MEF lysates mock treated or treated with 4OHT for TLK1, TLK2, ASF1 and p-ASF1. Experiment performed in biological triplicate and a representative example shown. (H) Proposed model for the contribution of TLK1 and TLK2 activity to the placenta and adult tissues. In the placenta, and possibly other cell types or tissues, TLK2 is the prevalent activity and is required to maintain a threshold of signalling required for viability. This likely involves ASF1a/b and potentially other substrates. In most adult tissues, TLK1 and TLK2 are largely redundant and the presence of either is sufficient to provide the necessary activity to support cellular functions required for the maintenance of genome integrity and viability.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6







Figure 7




Supplementary Figure S1



Supplementary Figure S2




Supplementary Figure S3




Supplementary Figure S4

A




B




Supplementary Figure S5


A IP-MS




B BioID




D




E




G




C





F




H



Supplementary Figure S6



Supplementary Figure S7



Differential requirements for Tlk2 like kinases 1 and 2 in mammalian development

Sandra Segura-Bayona, Philip A. Knobel, Helena González Burón, Aida Peña-Blanco, Sameh A. Youssef, Étienne Coyaoud, Teresa López-Rovira, Katrin Rein, Lluís Palenzuela, Julien Colombelli, Stephen Forrow, Brian Raught, Anja Groth, Alain de Bruin and Travis H. Stracker

Supplementary Figure Legends

Figure S1. Generation of *Tlk2* conditional knockout mice. (A) Following transfection and neomycin selection, ES clones were screened by long range PCR for correct targeting. Clone IDs are indicated below and the expected band for the targeted allele indicated in red. C57BL/6 refers to wild type genomic DNA (negative control). (B) Southern blotting of KpnI digested genomic DNA from targeted ES cell clones selected by long range PCR. The 5P probe recognizes a 12.5 kb fragment and the 3P probe recognizes a 14 kb fragment in the correctly targeted locus. The untargeted locus gives a 26 kb band in both cases. For the Neo probe, we digested with SspI to recognize a 10.5 kb fragment (Figure 2B). Expected size of recombined products with probes for the 5' and 3' homology arms are indicated (red arrowhead). Clone IDs are indicated above the lanes. Additional details regarding the generation and screening of targeted ES cells are available upon request.

Figure S2. Growth retardation of *Tlk2*^{+/+} E12.5 embryos. PFA fixed embryos were embedded in a 0.8% agarose block before optical clearing, dehydrated in 100% methanol for 24 hours and transferred into Murray's Clear (BABB: Benzyl Alcohol Benzyl Benzoate) for at least 2 days before imaging. Whole mount imaging of specimens was performed with a custom-built lightsheet system, at 1.32X with a macroscope detection unit AZ100M (Nikon, Japan) for autofluorescence with a 488nm laser forming a lightsheet of about 12 microns thickness, incident from the left of the image and flat across the whole field of view (zoom: 1.32x, colored in 'Orange Hot' Look-up-Table). The images were recorded with an ORCA R2 CCD camera (Hamamatsu, Japan). Image stacks were stitched vertically with ImageJ. The 4 planes (from left to right) show different depths through the whole embryos (2, 4, 6 and 7.2 mm respectively) from the dorsal surface. The embryos are mounted to face the camera dorsally.

Figure S3. Analysis of TLK2 deficient placentas. (A) Reduced size and vascularization of *Tlk2*^{+/+} placenta compared to a wild type littermate. (B) Lack of blood circulation in the umbilical cord (red arrowhead) of an E14.5 *Tlk2*^{+/+} embryo. (C) Representative PCR analysis of the deletion of *Tlk2* Exon 4 following Sox2-Cre expression, genotypes are indicated and shown schematically to the right. (D) CD31 staining of E11.5 and E12.5 placentas of the indicated genotypes. Areas cropped and presented in Figure 3C are indicated. Scale bar is 20 microns. (E) Sections from E10.5 placentas stained with H&E. *Tlk2*^{-/-} placentas are thinner,

less cellular and poorly vascularized compared to *Tlk2*^{+/+}. Fewer numbers of labyrinth trophoblast (black arrows) and syncytiotrophoblasts (blue arrows) are present in *Tlk2*^{-/-} placentas. L=labyrinth, TS=trophospongium and D=decidua in all panels. Scale bar in 10X=100 microns and in 20X=50 microns. (F) The E11.5 *Tlk2*^{+/+} labyrinth (L) is well vascularized, has widely opened sinusoids and is composed of many well-developed trophoblasts (black arrows) and syncytiotrophoblasts (blue arrows). The *Tlk2*^{-/-} labyrinth is poorly vascularized, has collapsed or has very narrow sinusoids and is composed of fewer and smaller trophoblasts (black arrows) and syncytiotrophoblasts (blue arrows). Also the *Tlk2*^{-/-} trophospongium (TS) is significantly thinner compared to *Tlk2*^{+/+} and lacks glycogen trophoblasts. Scale bar in 10X=100 microns and in 20X=50 microns. (G) Lower magnification images of E11.5 placentas presented in Figure 3D from the indicated genotypes. Boxed regions indicate cropped regions in main figure. Scale bars are 20 microns. (H) Sections from E12.5 placentas stained with H&E. The *Tlk2*^{+/+} placenta has a histologically normal labyrinth (L) that is well vascularized and is composed of many mature trophoblasts discerned by their large spherical nuclei (black arrows). In contrast, the *Tlk2*^{-/-} labyrinth is composed of clusters of small, undifferentiated hyperchromatic trophoblasts that tend to arrange in clusters or aggregates (arrowheads). Also the trophospongium (TS) zone of the *Tlk2*^{-/-} placenta appears thinner and lacks glycogen trophoblasts. Scale bar in 10X=100 microns and in 20X=50 microns. (I) Lower magnification images of E12.5 placentas presented in Figure 3D from the indicated genotypes. Boxed regions indicate cropped regions in main figure. Scale bars are 20 microns. (J) The E15.5 *Tlk2*^{-/-} labyrinth (L) is collapsed, lacks vasculature, has fewer trophoblasts (black arrows), and exhibits increased deposition of extracellular matrix (asterisks) when compared to the highly vascularized *Tlk2*^{+/+} labyrinth. Scale bar in 10X=100 microns and in 20X=50 microns.

Figure S4. Histochemical and protein analysis. (A) Immunohistochemical staining of the cleaved caspase-3 apoptosis marker in littermate E10.5, E11.5 and E12.5 placentas of the indicated genotype (n=3 and 2 at E10.5, n=3 and 3 at E11.5 and n=5 and 6 at E12.5 for *Tlk2*^{+/+} and *Tlk2*^{-/-}, respectively. At least 200 total cells were counted for each individual). Few positive cells were observed regardless of genotype or developmental stage. Quantification of percent positive cells is graphed in the right panels with mean and standard deviation indicated. (B) Western blotting of lysates from embryonic tissue (fetal heart and liver = L) and placenta (P) from littermate animals of the indicated genotypes. While TLK2 protein levels are similar between tissues, TLK1 protein levels are much higher in the embryonic tissues relative to placenta.

Figure S5. Proteomic approaches to identify TLK2 interacting proteins by mass spectrometry. (A) Schematic of affinity purification strategy (IP-MS) for Strep-FLAG tagged TLK2 or a kinase dead allele, TLK2-KD (D592V). Replicates of the affinity purifications were digested with trypsin and subjected to LC-MS/MS to quantitatively identify interacting proteins. (B)

Schematic of the BioID approach using TLK2 fused to the promiscuous biotin ligase BirA*. (C) Validation of TLK2 or TLK2 KD expression and affinity purification of TLK2 by Western blotting following transfection of AD293 cells (bottom panel). The effectiveness of the kinase dead mutation was also validated in IP-kinase assays (top panel). The TLK2 KD mutation abolished autophosphorylation and substrate phosphorylation (Myelin Basic Protein: MBP). (D) Validation of BirA*-TLK2 expression and activity in AD293 cells supplemented with biotin. Note that Streptavidin signal that was detected in the nucleus and cytoplasm for the BirA*-FLAG control, was exclusively nuclear in the case of BirA*-FLAG-TLK2. Biotinylated peptides were purified from cells expressing the BirA* constructs and identified by LC-MS/MS. (E) Validation of high confidence interacting proteins by IP-Western. Data is summarized in Figure 5C and presented in full in Supplementary Tables S1 to S4. The only proteins identified that were not detected in any controls and present in all 6 TLK2/TLK2-KD replicates were TLK1, ASF1a, ASF1b, LC8 (DYNLL1) and RPS11 (Supplementary Table S2). Of these, all but RPS11 were also found highly enriched in the BioID experiments (Supplementary Table S5 and S6). We could readily validate the interactions between TLK2 and endogenous TLK1 and ASF1 in HEK293Ts. In addition, we could detect binding to endogenous LC8. (F) Validation of ASF1a as a TLK2 substrate. IP-kinase assays were performed with TLK2 or TLK2 KD affinity purified from HEK293T cells using GST-ASF1a as a substrate. (G) Addition of LC8 did not influence the kinase activity of TLK2 (based on autophosphorylation) and recombinant LC8 was not phosphorylated by TLK2, suggesting that it is unlikely to be a substrate. (H) Kinetic analysis of TLK2 or TLK KD activity on MBP and LC8. MBP phosphorylation is clearly increased over time while no signal is observed for LC8 in the phosphorimager, despite similar levels of protein loaded.

Figure S6. Analysis of ASF1 levels and interaction with TLK2. (A) Analysis of total ASF1 and ASF1-pS166 (p-ASF1) levels in placentas from the indicated genotypes and age. Western blotting results from 3 different gels/transfers are shown. Compiled quantification of all samples is shown in Figure 5D. ID codes indicate the litter number and embryo number, *Tlk2*^{-/-} placentas are indicated in green font. (B) Additional example of ASF1 Westerns following IP of endogenous TLK2 from placental extracts (related to main Figure 5E). Both TLK1 and ASF1 consistently interact with TLK2 in IPs. A non-specific band is indicated by “ns”.

Figure S7. Kaplan Meier survival curve of all genetrap allelic configurations observed to date. The number of animals represented is shown in parentheses.