



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Differences and similarities in endothelial and angiogenic profiles of preeclampsia and COVID-19 in pregnancy

Marta PALOMO, Msc, PhD, Lina YOUSSEF, MD, PhD, Alex RAMOS, Mr., Sergi TORRAMADE-MOIX, Msc, Ana Belen MORENO-CASTAÑO, MD, Julia MARTINEZ-SANCHEZ, Msc, Laura BONASTRE, Ms., Marc PINO, Mr., Pilar GOMEZ-RAMIREZ, Ms., Lidia MARTIN, Ms., Estefania GARCIA. MATEOS, Ms., Pablo SANCHEZ, PhD, Sara FERNANDEZ, MD, Francesca CROVETTO, MD, PhD, Ginés ESCOLAR, MD, PhD, Enric CARRERAS, MD, PhD, Pedro CASTRO, PhD, Eduard GRATACOS, MD, PhD, Fàtima CRISPI, MD, PhD, Maribel DIAZ-RICART, PhD

PII: S0002-9378(22)00227-7

DOI: <https://doi.org/10.1016/j.ajog.2022.03.048>

Reference: YMOB 14392

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 6 August 2021

Revised Date: 22 March 2022

Accepted Date: 23 March 2022

Please cite this article as: PALOMO M, YOUSSEF L, RAMOS A, TORRAMADE-MOIX S, Belen MORENO-CASTAÑO A, MARTINEZ-SANCHEZ J, BONASTRE L, PINO M, GOMEZ-RAMIREZ P, MARTIN L, MATEOS EG, SANCHEZ P, FERNANDEZ S, CROVETTO F, ESCOLAR G, CARRERAS E, CASTRO P, GRATACOS E, CRISPI F, DIAZ-RICART M, Differences and similarities in endothelial and angiogenic profiles of preeclampsia and COVID-19 in pregnancy, *American Journal of Obstetrics and Gynecology* (2022), doi: <https://doi.org/10.1016/j.ajog.2022.03.048>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Differences and similarities in endothelial and angiogenic profiles of preeclampsia and COVID-19 in pregnancy

Marta PALOMO, Msc, PhD^{1,2,3*}; Lina YOUSSEF, MD, PhD^{4*}; Alex RAMOS, Mr.^{1,2,3}; Sergi TORRAMADE-MOIX, Msc²; Ana Belen MORENO-CASTAÑO, MD^{2,3,5}; Julia MARTINEZ-SANCHEZ, Msc^{1,2,3}; Laura BONASTRE, Ms.²; Marc PINO, Mr.²; Pilar GOMEZ-RAMIREZ, Ms.²; Lidia MARTIN, Ms.²; Estefania GARCIA MATEOS, Ms.²; Pablo SANCHEZ, PhD⁶; Sara FERNANDEZ, MD⁷; Francesca CROVETTO, MD, PhD^{4,8}; Ginés ESCOLAR, MD, PhD^{2,6}; Enric CARRERAS, MD, PhD^{1,3}; Pedro CASTRO, PhD^{5,6}; Eduard GRATACOS, MD, PhD^{4,5,8}; Fàtima CRISPI, MD, PhD^{4,5,8#}; Maribel DIAZ-RICART, PhD^{2,3,5#}.

¹Josep Carreras Leukaemia Research Institute, Hospital Clinic, University of Barcelona Campus, Barcelona, Spain.

²Laboratory of Hemostasis and Eritropathology, Hematopathology, Pathology Department, Centre Diagnòstic Biomèdic (CDB), Hospital Clinic, University of Barcelona, Barcelona, Spain.

³Barcelona Endothelium Team (BET), Barcelona, Spain.

⁴BCNatal (Hospital Clínic and Hospital Sant Joan de Déu), Barcelona, Spain.

⁵Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, Barcelona, Spain.

⁶Department of Marine Biology and Oceanography, Institut de Ciències del Mar (ICM), CSIC, Barcelona, Spain.

⁷Medical Intensive Care Unit, Hospital Clinic, School of Medicine, Barcelona, Spain.

⁸Centre for Biomedical Research on Rare Diseases (CIBER-ER), Madrid, Spain.

*contributed equally as first authors

contributed equally as last authors

Corresponding author: Marta Palomo de Udaeta, PhD
Josep Carreras Leukaemia Research Institute
Hospital Clinic/University of Barcelona Campus
C/ Villarroel 170, 08036 Barcelona, Spain
Phone: 34-93-227 54 00 Ext. 2034 FAX: 34-93-227 93 69
e-mail: mpalomo@carrerasresearch.org

Disclosure of interests: The authors report no conflict of interest.

Word count

Abstract: 500

Main text: 3420

40 **Funding sources**

41 We would like to acknowledge support from Fundació Clínic, Barcelona (HCB/2020/0401),
42 Fundació La Marató de TV3 (202026-10), Jazz Pharmaceuticals Plc (IST-16-10355), German
43 José Carreras Leukaemia Foundation (03R/ 2019), Instituto de Salud Carlos III from Spanish
44 Government (PI19/00888), Bristol Myers-Squibb & Pfizer (ERISTA 15), “La Caixa”
45 Foundation, the Kids Corona Child and Mother COVID-19 Open Data and Biobank Initiative
46 from Hospital Sant Joan de Déu (Stavros Niarchos Foundation, Santander Foundation and
47 others), Fundació Privada Daniel Bravo Andreu, Generalitat de Catalunya (2017-SGR675 and
48 CERCA Programme), Fundació Catalana de Trasplantament (FCT 2021) and by ad hoc
49 patronage funds for research on COVID-19 from donations from citizens and organizations to
50 the Hospital Clínic de Barcelona-Fundació Clínic per a la Recerca Biomèdica, Barcelona,
51 Spain. All funders had no role in the study design, data collection, data analysis, data
52 interpretation and writing the manuscript.

Condensation

Preeclampsia and COVID-19 exhibit a differential profile of circulating biomarkers with a similar end-stage *in vitro* induced endothelial dysfunction.

Short version of the title

Endothelial dysfunction in preeclampsia vs. COVID-19 in pregnancy

AJOG at a glance

Why was this study conducted?

We conducted this study to characterize the profile of endothelial damage, coagulation, innate immune response and angiogenesis in preeclampsia and COVID-19 in pregnancy which are both considered disorders associated with endothelial dysfunction.

What are the key findings?

Severe COVID-19 in pregnancy and preeclampsia share a similar end-stage *in vitro* induced p38MAPK phosphorylation in endothelial cells but a differential profile of circulating endothelial and angiogenic biomarkers. Severe COVID-19 is characterized by higher VCAM-1, sTNFR-I, HS, VWF antigen and NETS and reduced PlGF while preeclampsia is marked by increased VCAM-1, sTNFR-I, sFlt-1, Ang2, C5b9 and NETS and a reduction in VWF antigen, VWF activity, α 2AP and PlGF.

What does this study add to what is already known?

Soluble biomarkers of coagulopathy [VWF], endothelial inflammation [sTNFR-I], barrier damage [HS] and angiogenesis [sFlt1] seem to be highly specific to differentiate preeclampsia from severe COVID-19 in pregnancy. These findings improve our understanding of the

- 78 pathophysiological pathways in preeclampsia and COVID-19 and may help in the differential
79 diagnosis of these disorders during pregnancy.

ABSTRACT

Background: COVID-19 presents a spectrum of signs and symptoms in pregnant women that might resemble preeclampsia. Differentiation between severe COVID-19 and preeclampsia is difficult in some cases.

Objective: To study biomarkers of endothelial damage, coagulation, innate immune response and angiogenesis in preeclampsia and COVID-19 in pregnancy in addition to *in vitro* alterations in endothelial cells exposed to sera from pregnant women with preeclampsia and COVID-19.

Methods: Plasma and sera samples were obtained from pregnant women with COVID-19 infection classified into mild (n=10) or severe (n=9) in addition to normotensive pregnancies as controls (n=10) and patients with preeclampsia (n=13). A panel of plasmatic biomarkers was assessed including vascular cell adhesion molecule-1 (VCAM-1), soluble TNF-receptor I (sTNFR-I), heparan sulfate (HS), von Willebrand factor (VWF) antigen, activity and multimeric pattern, α 2-antiplasmin (α 2AP), C5b9, neutrophil extracellular traps (NETS), placental growth factor (PlGF), fms-like tyrosine kinase-1 (sFlt-1) and angiopoietin 2 (Ang2). Additionally, microvascular endothelial cells were exposed patient's serum, and changes in the cell expression of intercellular adhesion molecule 1 (ICAM-1) on cell membrane and VWF release to the extracellular matrix were evaluated through immunofluorescence. Changes in inflammation cell signaling pathways were also assessed by of P38MAPK phosphorylation. Statistical analysis included univariate and multivariate methods.

Results: Biomarker profiles in mild COVID-19 were similar to controls. Both preeclampsia and severe COVID-19 showed significant alterations in the majority of circulating biomarkers with distinctive profiles. While severe COVID-19 exhibited higher concentrations of VCAM-1, sTNFR-I, HS, VWF antigen and NETS with a significant reduction of PlGF as compared to controls; preeclampsia presented a marked increase in VCAM-1, sTNFR-I (significantly increased compared to controls and to severe COVID-19) with a striking reduction in VWF

antigen, VWF activity and α 2AP. As expected, reduced PlGF, increased sFlt-1 and Ang2 and a very high sFlt-1/PlGF ratio were also observed in preeclampsia. In addition, a significant increase in C5b9 and NETS was also detected in preeclampsia compared to controls. The principal component analysis demonstrated a clear separation between preeclampsia and the rest of groups (first and second components explained 42.2% and 13.5% of the variance), mainly differentiated by variables related to VWF, sTNFRI, HS and sFlt-1. VWF multimeric analysis revealed the absence of VWF high-molecular-weight multimers in preeclampsia (similar profile to von Willebrand disease type 2A) whereas in healthy pregnancies and COVID-19 patients, VWF multimeric pattern was normal.

Sera from both preeclampsia and severe COVID-19 patients induced an overexpression of ICAM-1 and VWF in endothelial cells in culture compared to controls. However, the effect of preeclampsia was less pronounced than the one triggered by severe COVID-19. Immunoblots of lysates from endothelial cells exposed to mild and severe COVID-19, and preeclampsia sera showed an increase in p38MAPK phosphorylation. Severe COVID-19 and preeclampsia were statistically different from controls, suggesting that both severe COVID-19 and preeclampsia sera can activate inflammatory signaling pathways.

Conclusion: While similar *in vitro* endothelial dysfunction, preeclampsia and severe COVID-19 exhibit distinctive profiles of circulating biomarkers related to endothelial damage, coagulopathy and angiogenic imbalance that could aid in the differential diagnosis of these entities.

Keywords: angiogenic factors, angiopoietin, C5b9, COVID-19, endothelial dysfunction, heparan sulfate, hypertensive disorders of pregnancy, NETS, PlGF, preeclampsia, SARS-CoV-2, sFlt-1, sTNFRI, Von Willebrand factor.

INTRODUCTION

Preeclampsia is a pregnancy complication and a leading cause of maternal and perinatal morbidity and iatrogenic prematurity¹⁻³. Although its etiology is not completely understood^{4,5}, it is accepted that this condition relies on placental insufficiency and maternal cardiovascular maladaptation underlined by angiogenic imbalance, endothelial dysfunction, coagulopathy and complement dysregulation⁶⁻⁸, which lead clinically to hypertension and proteinuria that can progress to multi-organ dysfunction during pregnancy. The multifactorial nature of preeclampsia explains a variable clinical/laboratory presentation, mainly determined by gestational age at onset: early vs late.

Clinical and analytical data from patients infected by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), which causes coronavirus disease 2019 (COVID-19), suggest that endothelial dysfunction plays an important role in the pathophysiology of this condition⁹⁻¹² involving extrapulmonary manifestations of COVID-19 like hypertension, kidney disease, thrombocytopenia, and liver injury. Some of these clinical features overlap with those observed in preeclampsia. In addition, an increased incidence of preeclampsia has been reported in association with COVID-19¹³⁻¹⁵. Despite their clinical resemblance, the mechanisms underlying endothelial dysfunction might differ between COVID-19 and preeclampsia. Understanding endothelial and angiogenic profiles could enlighten the pathophysiological basis of these two entities.

The endothelium is a monolayer of cells that lines the interior of blood vessels acting as a protective layer between circulating blood and other tissues. The endothelium is crucial for the regulation of vascular homeostasis, coagulation cascade, immune response and angiogenesis. Circulating biomarkers related to endothelial activation and loss of barrier integrity seem to be associated to disease severity in COVID-19^{12,16}. Inflammatory effects on these damaged

endothelial cells activates the innate immune response and induces a hypercoagulable state with impaired fibrinolysis and angiogenic imbalance^{17,18}. On the other hand, angiogenesis dysregulation has emerged as one of the main pathophysiological features in the development of preeclampsia^{19,20}. Finally, *in vitro* studies enabled us to describe the endothelial cell proinflammatory and thrombogenic response in COVID-19²¹⁻²³.

The aim of the present study was to comprehensively investigate the endothelial and angiogenic profiles in preeclampsia and SARS-CoV-2 infection in pregnancy using circulating biomarkers and *in vitro* studies.

MATERIALS AND METHODS

Study populations and design

Pregnant women with laboratory confirmed SARS-CoV-2 infection were selected from a large multicenter prospective population-based cohort study conducted from March 15 to May 31, 2020, in Barcelona, Spain including consecutive cases detected during the study period²⁴. SARS-CoV-2 infection was confirmed by a positive real time polymerase chain reaction (RT-PCR) on nasopharyngeal swab or a positive serological result. SARS-CoV-2 positive pregnancies were subdivided into mild (n=9) and severe disease (n=8) according to the presence of pneumonia or coexistence of fever, dry cough and dyspnea. In addition, we also included SARS-CoV-2 negative pregnant women including preeclampsia (n=13) and normotensive pregnancies as controls (n=10) who were matched to COVID-19 cases by gestational age at blood sampling. preeclampsia was defined as high blood pressure (systolic blood pressure $\geq$ 140 mmHg and/or diastolic blood pressure $\geq$ 90 mmHg on two occasions, at least four hours apart) developed after 20 weeks of gestation with proteinuria ($\geq$ 300 mg/24 h or protein/creatinine ratio $\geq$ 0.3), thrombocytopenia (platelet count $< 100 \times 10^9/l$, renal insufficiency (serum creatinine concentrations > 1.1 mg/dl), impaired liver function (elevated blood concentrations of liver transaminases to twice normal concentration), pulmonary edema or a new-onset headache unresponsive to medication and not accounted for by alternative diagnoses or visual symptoms²⁵. Early-onset preeclampsia was defined by gestational age at delivery before 34 weeks of gestation²⁶. Baseline and perinatal data were obtained by interviews and from electronic medical records. Gestational age was calculated based on the crown-rump length at first trimester ultrasound²⁷. Birthweight centiles were assigned according to local standards²⁸. Pregnancies with chromosomal/structural anomalies or intrauterine infection were excluded. Endothelial and angiogenic profiles were studied in all participants by analyzing

circulating molecules in maternal peripheral blood and by *in vitro* study of endothelial cells exposed to patient's sera. Details of the laboratory methodology used is detailed below and in the Supplementary material.

This study was approved by the ethics committee of Hospital Clinic (HCB/2020/0401) and conformed to the ethical guidelines of the Helsinki Declaration. All participants provided informed written consent before sample collection.

Maternal blood sample collection

Peripheral maternal blood was obtained by venipuncture within 24-48 hours of the onset of symptoms and before starting any treatment. Plasma and sera samples were obtained by centrifugation of blood anticoagulated with EDTA and by incubation for 30 min at room temperature to allow clotting and subsequently centrifuged at 1500×g for 10 min at 4 °C to separate the serum from clots, respectively. All samples were aliquoted and stored at -80°C until used.

Assessment of circulating biomarkers

Endothelial damage was assessed by measuring plasmatic concentrations of vascular cell adhesion molecule-1 (VCAM-1), soluble TNF- α receptor I (sTNFR1) and heparan sulfate (HS) by ELISA (R&D systems, MN, USA; Biomatik Corporation, DE, USA and AttendBio Research, Spain, respectively). The kit used for the detection of HS do not show any significant cross-reactivity or interference between HS and analogs according to the manufacturer's instructions.

Biomarkers for *coagulation/fibrinolysis* included von Willebrand antigen (VWF:Ag) and activity (VWF:GPIbM) and α 2-antiplasmin (α 2AP) evaluated by immunoturbidimetry (Atellica 180 360 COAG, Siemens Healthineers, Germany). Visualization of VWF multimers

was achieved using a commercially available enhanced chemiluminescence kit for detecting HRP-labeled antibodies on Western blots.²⁹ In addition, VWF factor-cleaving protease (ADAMTS-13) activity was assessed by fluorescence resonance energy transfer (Fluoroskan Ascent FL; Thermolab Systems, MA, USA). Plasminogen activator inhibitor antigen (PAI) and thrombomodulin (TM) were measured by ELISA (Imubind, Toronto, Canada; and Biomatik Corporation, MN, USA, respectively).

Activation of *innate immune response* was determined by circulating terminal complement complex (C5b9) and dsDNA for neutrophil extracellular traps (NETS) quantified by QuantiTTM PicoGreenTM dsDNA Assay Kit (Invitrogen, Thermo Fisher, MA, USA) on a fluorescence reader.

Angiogenic profile was assessed by sera concentrations of free Placental Growth Factor (PlGF) and soluble fms-like tyrosine kinase-1 (sFlt1) by ELISA (R&D Systems Europe Ltd, Abingdon, UK) and Angiopoietin-2 (Ang2) (R&D systems, MN, USA). The ratio sFlt1/PlGF was calculated as previously described³⁰

***In vitro* studies**

For the *in vitro* studies, human dermal microvascular endothelial cells (ATCC, CRL-3243, Lot:62630587) in culture were exposed to patient's sera in order to study cell response to: a) the expression of adhesion receptors at the cell surface (InterCellular Adhesion Molecule 1, ICAM-1), as an indicator of a proinflammatory response of cells; b) the presence or the adhesive protein VWF, involved in thrombogenicity, on the extracellular matrix generated by these cells; and c) the activation of the endothelial intracellular signaling pathway related to inflammation p38MAPK. Details of the laboratory methodology used is detailed in the Supplementary material.

Statistical analysis

Baseline and perinatal data were analyzed with the statistical software STATA 14.2 (StataCorp LLC, Texas, USA) and results are expressed as median and interquartile range or percentage as appropriate. Statistical analysis comprised the comparison of each group of complicated pregnancies vs. controls. Soluble markers are expressed as median (interquartile range). Further statistical analyses were performed in R (version 4.0.0) using Student's t-test with the Benjamini-Hochberg correction for multiple comparisons after checking data normality and homoscedasticity. Results were considered statistically significant when adjusted P value was < 0.05 . Data were ordinated and plotted using principal component analysis. An additional unsupervised hierarchical clustering was performed based on the univariate results comparing severe COVID-19 vs. preeclampsia. A sub analysis comparing early- vs late-onset preeclampsia was performed using Student's t-test and Benjamini-Hochberg procedure for multiple pairwise comparisons and included in the Supplementary material.

RESULTS

Baseline and perinatal characteristics of the study populations

Baseline characteristics of the study populations are summarized in Table 1. Study groups were mainly similar in terms of maternal and perinatal characteristics. However, preeclamptic patients had higher rates of Asian ethnicity and a tendency to younger age. Chronic hypertension was present in three preeclamptic patients and systemic lupus erythematosus in one control. None of the patients included in this study had pregestational diabetes or previous respiratory disorders. All the pregnancies complicated by preeclampsia were proteinuric, four were early-onset cases that were treated with corticosteroids for fetal lung maturity and five preeclamptic patients had preeclampsia with severe features that was treated with magnesium sulfate. Preeclamptic patients showed an earlier gestational age at delivery with a trend to higher rates of small-for-gestational-age fetus and admissions to neonatal intensive care unit. Three cases of preeclampsia were complicated by peripartum hemorrhage. Severe COVID-19 cases were all detected by RT-PCR. Among the mild cases, two were detected by RT-PCR and the rest by positive serology. Since this study has been conducted at the beginning of the pandemic, convalescent subjects should have been infected during the four weeks preceding the blood analysis. Two cases of mild COVID-19 had hypertension and one of them had associated proteinuria. None of the COVID-19 patients (mild or severe) had thrombocytopenia, elevated liver enzymes or elevated creatinine. All COVID-19 cases were followed up to 40 days postpartum to exclude the diagnosis of evolving preeclampsia. The diagnosis of atypical preeclampsia in COVID-19 cases was excluded as none of them presented signs of placental insufficiency nor abnormal sFlt1/PlGF ratio (according to our institutional protocol for the differential diagnosis of hypertensive disorders in pregnancy). Severe COVID-19 cases were not critically ill (no mortality and only one case required invasive mechanical ventilation). Six

patients with severe COVID-19 were treated with low-molecular-weight heparin, three of them were additionally treated with hydroxychloroquine and azithromycin and one of these three has been also given lopinavir / ritonavir and corticosteroids. As mentioned earlier, maternal blood samples were obtained before starting any treatment. Gestational age at sampling was similar between the study groups at median (interquartile range) of 40.2 (38.9 -41) weeks in controls, 39.1 (38.7 – 39.6) weeks in mild COVID-19, 39.3 (34.9 – 41.1) weeks in severe COVID-19 and 39.1 (35.1 – 39.6) weeks in preeclampsia. No cases of perinatal mortality were observed in the study population.

Endothelial and angiogenic circulating biomarkers are differentially altered in COVID-19 vs. preeclampsia

Results on soluble biomarkers in the study populations are displayed in Figure 1 and Table S1. Most soluble biomarkers were similar in **mild COVID-19** and controls with the exception of a significant increase in VWF:Ag. In contrast, profound alterations in endothelial, coagulation, immune and angiogenic biomarkers were detected in **severe COVID-19** including significantly higher concentrations of VCAM-1, sTNFR-I, HS, VWF:Ag and NETS with a significant reduction of PlGF as compared to controls. No differences were observed in Ang2, sFlt1, C5b9, ADAMTS13, PAI nor TM between severe COVID-19 and controls. Pregnant women with preeclampsia exhibited also remarkable alterations in soluble biomarkers in a distinct profile from the one observed in COVID-19. Cases of **preeclampsia** showed a marked increase in VCAM-1, sTNFR-I (significantly increased compared to controls and to severe COVID-19) with a striking reduction in VWF:Ag, VWF:GPIbM, VWF:Ag/ VWF:GPIbM and α 2AP. As expected, reduced PlGF, increased sFlt-1 and Ang2 and a very high sFlt-1/PlGF ratio were also observed in preeclampsia. In addition, a significant increase in C5b9 and NETS was also

detected in preeclampsia compared to controls. HS, ADAMTS13, PAI and TM remained unchanged in preeclampsia.

Principal component analysis demonstrated a clear separation between preeclampsia and the other study populations (controls and mild/severe COVID-19) (Figure 2A). The first and second components explained 42.2% and 13.5% of the variance between groups. Unsupervised hierarchical clustering also showed a complete separation between severe COVID-19 cases and preeclampsia (Figure 2B) with the most remarkable differences observed in VWF:GPIbM, VWF:Ag and VWF:Ag/ VWF:GPIbM followed by HS (significantly lower in preeclampsia) and sTNFRI, sFlt1 and sFlt-1/PlGF ratio (significantly higher in preeclampsia).

VWF multimeric analysis revealed the absence of VWF high-molecular-weight multimers in preeclampsia, comparable to a diagnosis of von Willebrand disease type 2A, with an accumulation of low-molecular-weight multimers (Figure 3). In healthy pregnancies and SARS-CoV-2 positive patients, VWF multimeric pattern was normal.

A sub analysis revealed a similar pattern of endothelial damage, coagulopathy and angiogenic imbalance in early- vs late-onset preeclampsia (Supplementary Table S2), with much remarkable changes in early-onset cases. In contrast, C5b9 and NETS were more altered in late-onset preeclampsia.

Severe COVID-19 and preeclampsia sera induce similar endothelial damage and inflammation *in vitro*

Endothelial cells incubation with sera from mild and severe COVID-19 patients induced a significant overexpression of ICAM-1 and VWF compared with controls (Figure 4). Cells exposed to preeclampsia sera showed also significantly increased ICAM-1 and VWF

314 expression although preeclampsia effect was less pronounced than the one caused by severe
315 COVID-19 ($p < 0.05$).

316 Immunoblots of lysates from endothelial cells exposed to mild and severe COVID-19, and
317 preeclampsia sera showed an increase in p38MAPK phosphorylation. Severe COVID-19 and
318 preeclampsia were statistically different from controls (Figure 5), suggesting that both severe
319 COVID-19 and preeclampsia sera can activate inflammatory signaling pathways.

COMMENT

Principal findings of the study

A comprehensive *ex vivo* and *in vitro* study revealed distinct endothelial and angiogenic profiles of severe COVID-19 *versus* preeclampsia. While severe COVID-19 exhibited alterations in heparan sulfate (HS), neutrophil extracellular traps (NETS) and placental growth factor (PlGF), preeclampsia presented abnormal levels of vascular adhesion molecule-1 (VCAM-1), soluble TNF receptor type I (sTNFRI), von Willebrand factor (VWF), complement C5b9, angiopoietin 2 (Ang2) and soluble fms-like tyrosine kinase-1 (sFlt-1). Sera from both severe COVID-19 patients and preeclampsia induced an overexpression of intercellular adhesion molecule-1 (ICAM-1) and VWF and activation of p38MAPK phosphorylation in endothelial cells in culture even though the effect of preeclampsia was less pronounced than the one triggered by severe COVID-19.

Preeclampsia vs COVID-19: a distinct profile of circulating endothelial damage biomarkers

Both preeclampsia and severe COVID-19 showed signs of endothelial damage, but with a differential pattern. Preeclamptic patients presented a very significant increase in VCAM-1 and sTNFRI with preserved HS. These results are consistent with previous reports demonstrating elevated VCAM-1^{31,32}. The presence of sTNFRI has been only anecdotally described³³. sTNFRI is the soluble receptor of tumor necrosis factor alpha, a pro-inflammatory cytokine that triggers the expression of inflammatory molecules, including cell adhesion molecules, such as VCAM-1 and ICAM-1³⁴ resulting in inflammation, apoptosis, reactive oxygen species generation, cell proliferation, and cell survival. In contrast, severe COVID-19 cases showed a milder increase

in VCAM-1 and sTNFRI with a significant alteration of HS. These data is in line with previous reports on non-pregnant COVID-19 patients showing a good correlation of VCAM-1 and sTNFRI with disease severity^{12,16}. The increased levels of HS suggest endothelial glycocalyx barrier disruption and degradation. This finding is consistent with previous reports in critically ill non-pregnant COVID-19 patients as HS is used by SARS-CoV-2 to interact with endothelial cells through its receptor-binding domain, leading to a damaged endothelial barrier³⁵.

Preeclampsia is associated with remarkable alterations in VWF antigen and functionality

Interestingly, the most remarkable differences between preeclampsia and COVID-19 were observed in VWF concentrations and activity. Our data in COVID-19 pregnancies are consistent with the previously described positive correlation of VWF with disease severity¹². Conversely, in preeclampsia, we observed a striking decrease in VWF levels contrary to the increase reported formerly in the literature³⁶. Interestingly, these changes were more pronounced in more severe early-onset cases. A potential explanation to this observation is acute VWF consumption due to endothelial cell exhaustion³⁷ in preeclampsia, as indeed, the *in vitro* exposure of endothelial cells to preeclampsia sera resulted in a relevant increase in VWF release. Other potential explanations could be bleeding or drug interaction (with corticosteroids given to ensure lung maturity). Moreover, our results suggest a qualitative VWF defect in preeclampsia manifested by low VWF:GPIbM/VWF:Ag ratio and confirmed by the multimeric analysis of VWF. Since ADAMTS-13 activity was similar in preeclampsia and the other study groups, the loss of high-molecular-weight multimers might be due to the lysis by other proteases such as plasmin. In fact, the degradation of VWF by plasmin has been described in hyperfibrinolytic states³⁸ and preeclampsia is known to be a hypercoagulable and hyperfibrinolytic state³⁹. Thus, it is plausible that a fibrinolytic imbalance might be underlying

VWF proteolysis, specifically an imbalance in plasmin regulation since $\alpha 2AP$ was significantly reduced in preeclampsia compared to the other groups.

Innate immune dysregulation in preeclampsia vs. COVID-19 in pregnancy

Our data confirm the previously reported increase in the soluble C5b9 in preeclampsia.^{40,41} Damaged endothelial cells in preeclampsia seem to activate the innate immune response including the complement system. In addition, we also report the formation of NETS both in preeclampsia and severe COVID-19 in pregnancy. NETS are large structures of chromatin and antimicrobial proteins released by dying neutrophils in order to capture extracellular pathogens, limit the spread of infections and directly activate alternative complement pathway. Our results are consistent with the previously reported activation of NETS directly by SARS-CoV-2 in non-pregnant individuals⁴². Hyperactivation of NETS formation in preeclampsia has been proposed to be induced by placental derived factors.⁴³ Interestingly, these changes were more remarkable in cases of late-onset preeclampsia. Overall, dysregulation of innate immune response seems to play a role in the complex pathological cascade leading to endothelial damage in both SARS-CoV-2 infection and preeclampsia^{43,44}. Interestingly, certain aspects of the complement cascade and NETS facilitate coagulation and interfere with anticoagulation⁴⁵. Therefore, the crosstalk between the complement and coagulation cascades, along with endothelial damage, may create the prothrombotic environment associated with adverse outcomes in COVID-19 and preeclampsia^{46,47}.

Preferential angiogenic imbalance in preeclampsia vs COVID-19

Finally, our results show a profound disruption of the angiogenic balance in preeclampsia as compared to controls and COVID-19 with very high levels of sFlt-1 and Ang2 together with reduced PlGF⁴⁸. As previously described, angiogenic was more severely altered in early-onset preeclampsia⁴⁹. Interestingly, COVID-19 cases showed also significantly low PlGF but normal concentrations of sFlt-1 and therefore preserved sFlt-1/PlGF ratio. PlGF is mainly synthesized in the endothelium, which might explain a reduction in any case of endothelial damage. In contrast, sFlt-1 and Ang2 seems to be distinctive of preeclampsia. These findings are consistent with angiogenesis dysregulation being proposed one of the main pathophysiological features in the development of preeclampsia.^{19,30} These results are also in line with previous reports⁵⁰ proposing sFlt1/PlGF ratio for the differential diagnosis of preeclampsia and COVID-19 in pregnancy.

Similar *in vitro* induced endotheliopathy in preeclampsia and SARS-CoV-2 infection

Our *in vitro* results demonstrate a strong activation of p38MAPK induced by both severe COVID-19 and preeclampsia sera, together with a potent induction of ICAM-1 and VWF expression. This functional approach reflects the direct deleterious effect of both sera inducing microvascular endothelial damage *in vivo*. The slightly superior effect of severe COVID-19 sera could be attributed not only to the soluble factors present in the sera but to a direct viral infection. The observed activation of ICAM-1 and VWF is consistent with the known mechanism of activating adhesion molecules to recruit neutrophils and platelets in response to endothelial damage.⁵¹ While it is known that SARS-CoV-2 infection activates p38MAPK and the downstream signaling, possibly leading to cell death⁵², the pathways leading to this activation in preeclampsia remain to be elucidated. Indeed, a preclinical study in a SARS-CoV-2 mouse model showed protective effects of p38MAPK inhibition pointing out its potential

therapeutic effect⁵³ These data suggest that, despite their different pathophysiology, both preeclampsia and COVID-19 finally activate common pathways of endothelial dysfunction explaining similarities in the clinical scenario.

Strengths and limitations

The main strength of this study is the prospective recruitment of well characterized COVID-19 cases in pregnant women that were matched for baseline characteristics with SARS-CoV-2 negative pregnancies both normotensive and preeclamptic. In addition, a large panel of endothelial damage markers has been investigated. The small sample size should be considered a limitation of the present study. Indeed, it hindered the detection of heterogeneity -if present- between early- and late-onset preeclampsia. On the other hand, we acknowledge that longitudinal changes in the studied biomarkers were not explored in the current study. Given the complexity and clinical heterogeneity of these conditions, future studies are warranted to confirm the similarities and differences in the endothelial and angiogenic profiles of these entities.

Conclusion, clinical and research implications

In conclusion, this study suggests a differential profile of circulating biomarkers with a similar end-stage *in vitro* induced endothelial dysfunction. Soluble biomarkers of coagulopathy [VWF], endothelial inflammation [sTNFRI], barrier damage [HS] and angiogenesis [sFlt1] seem to be highly specific to differentiate preeclampsia from severe COVID-19 in pregnancy. These findings hold the potential to improve our understanding of the pathophysiological

435 pathways in preeclampsia and COVID-19 in pregnancy. We also identify circulating
436 biomarkers that may be useful in the differential diagnosis of preeclampsia and SARS-CoV-2
437 infection in pregnancy. Given the difficulty of clinically differentiate some cases of
438 preeclampsia and COVID-19, a panel of circulating biomarkers for the differential diagnosis
439 could be of most help to optimize patient's management. Finally, this study also opens
440 opportunities for new therapeutic targets that could improve the underlying endothelial damage
441 observed in these entities.

442 **ACKNOWLEDGMENTS**

443 We thank all the patients that agreed to participate in the present study. We would like to thank
444 Biobanks of Fundació Sant Joan de Déu and Clínic-IDIBAPS for valuable management of
445 samples. We are indebted to Fundació Glòria Soler for its support to the COVIDBANK
446 initiative and to the HCB-IDIBAPS Biobank for the biological human samples and data
447 procurement.

REFERENCES

1. Duley L. The Global Impact of Pre-eclampsia and Eclampsia. *Semin Perinatol.* 2009;33(3):130-137. doi:10.1053/j.semperi.2009.02.010
2. Ghulmiyyah L, Sibai B. Maternal Mortality From Preeclampsia/Eclampsia. *Semin Perinatol.* 2012;36(1):56-59. doi:10.1053/j.semperi.2011.09.011
3. Steegers EAP., Von Dadelszen P., Duvekot JJ., Pijnenborg R. Pre-eclampsia. *Lancet.* 2010;376(9741):631-644.
4. Erez O, Romero R, Jung E, et al. Preeclampsia and eclampsia: the conceptual evolution of a syndrome. *Am J Obstet Gynecol.* 2022;226(2):S786-S803. doi:10.1016/j.ajog.2021.12.001
5. Jung E, Romero R, Yeo L, et al. The etiology of preeclampsia. *Am J Obstet Gynecol.* 2022;226(2):S844-S866. doi:10.1016/j.ajog.2021.11.1356
6. Melchiorre K, Giorgione V, Thilaganathan B. The placenta and preeclampsia: villain or victim? *Am J Obstet Gynecol.* Published online 2021:1-9. doi:10.1016/j.ajog.2020.10.024
7. Rana S, Burke SD, Karumanchi SA. Imbalances in circulating angiogenic factors in the pathophysiology of preeclampsia and related disorders. *Am J Obstet Gynecol.* 2021;1:1-16. doi:10.1016/j.ajog.2020.10.022
8. Burwick RM, Feinberg BB. Complement activation and regulation in preeclampsia and hemolysis, elevated liver enzymes, and low platelet count syndrome. *Am J Obstet Gynecol.* 2020;6. doi:10.1016/j.ajog.2020.09.038
9. Dupont A, Rauch A, Staessens S, et al. Vascular Endothelial Damage in the Pathogenesis of Organ Injury in Severe COVID-19. *Arterioscler Thromb Vasc Biol.* 2021;(May):1760-1773. doi:10.1161/ATVBAHA.120.315595
10. Rauch A, Dupont A, Goutay J, et al. Endotheliopathy Is Induced by Plasma From Critically Ill Patients and Associated With Organ Failure in Severe COVID-19. *Circulation.* 2020;142(19):1881-1884. doi:10.1161/CIRCULATIONAHA.120.050907

- 475 11. Castro P, Palomo M, Moreno-Castaño AB, et al. Is the Endothelium the Missing Link
476 in the Pathophysiology and Treatment of COVID-19 Complications? *Cardiovasc drugs*
477 *Ther.* Published online 2021. doi:10.1007/s10557-021-07207-w
- 478 12. Fernández S, Moreno-Castaño AB, Palomo M, et al. Distinctive Biomarker Features in
479 The Endotheliopathy of COVID-19 and Septic Syndromes. *Shock*. Published online
480 June 24, 2021. doi:10.1097/shk.0000000000001823
- 481 13. Conde-Agudelo A, Romero R. *SARS-COV-2 Infection during Pregnancy and Risk of*
482 *Preeclampsia: A Systematic Review and Meta-Analysis*. Elsevier Inc.; 2021.
483 doi:10.1016/j.ajog.2021.07.009
- 484 14. Lai J, Romero R, Tarca AL, et al. SARS-CoV-2 and the subsequent development of
485 preeclampsia and preterm birth: evidence of a dose-response relationship supporting
486 causality. *Am J Obstet Gynecol.* 2021;225(6):689-693.e1.
487 doi:10.1016/j.ajog.2021.08.020
- 488 15. Giorgione V, Thilaganathan B. SARS-CoV-2 related myocardial injury might explain
489 the predisposition to preeclampsia with maternal SARS-CoV-2 infection. *Am J Obstet*
490 *Gynecol.* 2022;226(2):279-280. doi:10.1016/j.ajog.2021.09.043
- 491 16. Tong M, Jiang Y, Xia D, et al. Elevated expression of serum endothelial cell adhesion
492 molecules in COVID-19 patients. *J Infect Dis.* 2020;222(6):894-898.
493 doi:10.1093/infdis/jiaa349
- 494 17. Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in
495 COVID-19. *Lancet.* 2020;395(10234):1417-1418. doi:10.1016/S0140-6736(20)30937-
496 5
- 497 18. Torres-Torres J, Espino-y-Sosa S, Poon LC, et al. Increased levels of soluble fms-like
498 tyrosine kinase-1 are associated with adverse outcomes in pregnant women with
499 COVID -19 . *Ultrasound Obstet Gynecol.* Published online 2021.
500 doi:10.1002/uog.24798
- 501 19. Karumanchi SA. Angiogenic factors in preeclampsia: From diagnosis to therapy.
502 *Hypertension.* 2016;67(6):1072-1079. doi:10.1161/HYPERTENSIONAHA.116.06421

20. Zeisler H, Llurba E, Chantraine F, et al. Predictive Value of the sFlt-1:PlGF Ratio in Women with Suspected Preeclampsia. *N Engl J Med*. 2016;374(1):13-22. doi:10.1056/nejmoa1414838
21. Serradell M, Diaz-ricart M, Cases A, et al. Uremic medium causes expression, redistribution and shedding of adhesion molecules in cultured endothelial cells. 2002;87(10).
22. Palomo M, Diaz-Ricart M, Carbo C, et al. The Release of Soluble Factors Contributing to Endothelial Activation and Damage after Hematopoietic Stem Cell Transplantation Is Not Limited to the Allogeneic Setting and Involves Several Pathogenic Mechanisms. *Biol Blood Marrow Transplant*. 2009;15(5):537-546. doi:10.1016/j.bbmt.2009.01.013
23. Hanzu FA, Palomo M, Kalko SG, et al. Translational evidence of endothelial damage in obese individuals: Inflammatory and prothrombotic responses. *J Thromb Haemost*. 2011;9(6):1236-1245. doi:10.1111/j.1538-7836.2011.04285.x
24. Crovetto F, Crispi F, Llurba E, et al. Impact of SARS-CoV-2 Infection on Pregnancy Outcomes: A Population-Based Study. *Clin Infect Dis*. 2021;ciab104. doi:10.1093/cid/ciab104
25. Gestational Hypertension and Preeclampsia: ACOG Practice Bulletin, Number 222. *Obstet Gynecol*. 2020;135(6):e237-e260. doi:10.1097/AOG.0000000000003891
26. Serra B, Mendoza M, Scazzocchio E, et al. A new model for screening for early-onset preeclampsia. *Am J Obstet Gynecol*. 2020;222(6):608.e1-608.e18. doi:10.1016/j.ajog.2020.01.020
27. Robinson HP, Fleming JEE. A CRITICAL EVALUATION OF SONAR "CROWN-RUMP LENGTH" MEASUREMENTS. *BJOG An Int J Obstet Gynaecol*. 1975;82(9):702-710. doi:10.1111/j.1471-0528.1975.tb00710.x
28. Figueras F, Meler E, Iraola A, et al. Customized birthweight standards for a Spanish population. *Eur J Obstet Gynecol Reprod Biol*. 2008;136(1):20-24. doi:10.1016/j.ejogrb.2006.12.015
29. Moreno-Castaño AB, Ramos A, Pino M, et al. Diagnostic challenges in von Willebrand

disease. Report of two cases with emphasis on multimeric and molecular analysis.

Platelets. 2020;32(5):1-4. doi:10.1080/09537104.2020.1784403

30. Verlohren S, Dröge LA. The diagnostic value of angiogenic and antiangiogenic factors in differential diagnosis of preeclampsia. *Am J Obstet Gynecol*. Published online 2020. doi:10.1016/j.ajog.2020.09.046

31. Crispi F, Domínguez C, Llurba E, Martín-Gallán P, Cabero L, Gratacós E. Placental angiogenic growth factors and uterine artery Doppler findings for characterization of different subsets in preeclampsia and in isolated intrauterine growth restriction. *Am J Obstet Gynecol*. 2006;195(1):201-207. doi:10.1016/j.ajog.2006.01.014

32. Heyl W, Heintz B, Reister F, et al. Increased soluble VCAM-1 serum levels in preeclampsia are not correlated to urinary excretion or circadian blood pressure rhythm. *J Perinat Med*. 2005;33(2):144-148. doi:10.1515/JPM.2005.027

33. Kim SY, Ryu HM, Jae HY, et al. Maternal serum levels of VCAM-1, ICAM-1 and E-selectin in preeclampsia. *J Korean Med Sci*. 2004;19(5):688-692. doi:10.3346/jkms.2004.19.5.688

34. Kong DH, Kim YK, Kim MR, Jang JH, Lee S. Emerging roles of vascular cell adhesion molecule-1 (VCAM-1) in immunological disorders and cancer. *Int J Mol Sci*. 2018;19(4):13-17. doi:10.3390/ijms19041057

35. David S. Injury to the endothelial glycocalyx in critically ill patients with COVID-19. *Am J Respir Crit Care Med*. 2020;208(8):1178-1181. doi:10.1164/rccm.202007-2676LE

36. Molvarec A, Rigó J, Böze T, et al. Increased plasma von Willebrand factor antigen levels but normal von Willebrand factor cleaving protease (ADAHTS13) activity in preeclampsia. *Thromb Haemost*. 2009;101(2):305-311. doi:10.1160/TH08-05-0330

37. Bergmann F, Rotmensch S, Rosenzweig B, How H, Chediak J. The role of von Willebrand factor in pre-eclampsia. *Thromb Haemost*. 1991;66(5):525-528. doi:10.1055/s-0038-1646453

38. Dicke C, Schneppenheim S, Holstein K, et al. Distinct mechanisms account for

acquired von Willebrand syndrome in plasma cell dyscrasias. *Ann Hematol.* 2016;95(6):945-957. doi:10.1007/s00277-016-2650-x

39. Sucak GT, Acar K, Sucak A, Kirazli S, Haznedar R. Increased global fibrinolytic capacity as a clue for activated fibrinolysis in pre-eclampsia. *Blood Coagul Fibrinolysis.* 2006;17(5):347-352. doi:10.1097/01.mbc.0000233364.72863.a0
40. Palomo M, Blasco M, Molina P, et al. Complement activation and thrombotic microangiopathies. *Clin J Am Soc Nephrol.* 2019;14(12):1719-1732. doi:10.2215/CJN.05830519
41. Youssef L, Miranda J, Blasco M, et al. Complement and coagulation cascades activation is the main pathophysiological pathway in early-onset severe preeclampsia revealed by maternal proteomics. *Sci Rep.* 2021;11(3048). doi:10.1038/s41598-021-82733-z
42. Arcanjo A, Logullo J, Menezes CCB, et al. The emerging role of neutrophil extracellular traps in severe acute respiratory syndrome coronavirus 2 (COVID-19). *Sci Rep.* 2020;10(1):1-11. doi:10.1038/s41598-020-76781-0
43. Gupta AK, Hasler P, Holzgreve W, Hahn S. Neutrophil NETs: A novel contributor to preeclampsia-associated placental hypoxia? *Semin Immunopathol.* 2007;29(2):163-167. doi:10.1007/s00281-007-0073-4
44. Ackermann M, Anders H-J, Bilyy R, et al. Patients with COVID-19: in the dark-NETs of neutrophils. *Cell Death Differ.* Published online 2021. doi:10.1038/s41418-021-00805-z
45. Foley JH. Examining coagulation-complement crosstalk: complement activation and thrombosis. *Thromb Res.* 2016;141:S50-S54. doi:10.1016/S0049-3848(16)30365-6
46. Servante J, Swallow G, Thornton JG, et al. Haemostatic and thrombo-embolic complications in pregnant women with COVID-19: a systematic review and critical analysis. *BMC Pregnancy Childbirth.* 2021;21(1):1-14. doi:10.1186/s12884-021-03568-0
47. Scheres LJJ, Lijfering WM, Groenewegen NFM, et al. Hypertensive Complications of

Pregnancy and Risk of Venous Thromboembolism. *Hypertension*. Published online
2020;781-787. doi:10.1161/HYPERTENSIONAHA.119.14280

48. Levine RJ, Maynard SE, Qian C, et al. Circulating Angiogenic Factors and the Risk of
Preeclampsia. *N Engl J Med*. 2004;350:672-683. doi:10.1056/NEJMoa031884

49. Schaarschmidt W, Rana S, Stepan H. The course of angiogenic factors in early-vs. late-
onset preeclampsia and HELLP syndrome. *J Perinat Med*. 2013;41(5):511-516.
doi:10.1515/jpm-2012-0248

50. Mendoza M, Garcia-Ruiz I, Maiz N, et al. Pre-eclampsia-like syndrome induced by
severe COVID-19: a prospective observational study. *BJOG An Int J Obstet Gynaecol*.
2020;127(11):1374-1380. doi:10.1111/1471-0528.16339

51. Rao RM, Yang L, Garcia-Cardena G, Luscinskas FW. Endothelial-dependent
mechanisms of leukocyte recruitment to the vascular wall. *Circ Res*. 2007;101(3):234-
247. doi:10.1161/CIRCRESAHA.107.151860b

52. Kopecky-Bromberg SA, Martinez-Sobrido L, Palese P. 7a Protein of Severe Acute
Respiratory Syndrome Coronavirus Inhibits Cellular Protein Synthesis and Activates
p38 Mitogen-Activated Protein Kinase. *J Virol*. 2006;80(2):785-793.
doi:10.1128/jvi.80.2.785-793.2006

53. Grimes JM, Grimes K V. p38 MAPK inhibition: A promising therapeutic approach for
COVID-19. *J Mol Cell Cardiol*. 2020;144(April):63-65.
doi:10.1016/j.yjmcc.2020.05.007

608 **Tables**609 **Table 1. Baseline and perinatal characteristics of the study populations.**

610

	Controls n=10	Mild COVID-19 n=9	Severe COVID-19 n=8	Preeclampsia n=13
<i>Maternal characteristics</i>				
Age (years)	36.9 (31.6 – 38.7)	36 (30.6 – 37.7)	35.2 (24.7 – 39.1)	29 (26 – 35.9)
Ethnicity				
White	8 (80)	5 (55.6)	4 (50)	4 (30.8)
African	0 (0)	0 (0)	1 (12.5)	2 (15.4)
Latin	2 (20)	3 (33.3)	2 (25)	2 (15.4)
Asian	0 (0)	1 (11.1)	1 (12.5)	5 (38.5) *
Pre-gestational body mass index (Kg/m ²)	22.4 (21.1 – 25.6)	22.7 (20.3 – 28.7)	21.8 (21 – 23.9)	25.9 (21.9 – 28.4)
Nulliparity	7 (70)	5 (55.6)	2 (25)	7 (53.8)
Use of assisted reproductive technologies	2 (20)	0 (0)	0 (0)	0 (0)
Smoking during pregnancy	0 (0)	0 (0)	1 (12.5)	0 (0)
<i>Perinatal outcomes</i>				
Gestational age at delivery (weeks)	40.2 (38.9 – 41)	39.1 (38.7 – 39.6)	39.2 (38.3 – 41.1)	39.1 (35.1 – 39.6) *
Preterm delivery#	1 (10)	1 (11.1)	2 (25)	4 (30.8)
Cesarean section	1 (10)	3 (33.3)	2 (25)	5 (38.5)
Female gender	4 (40)	4 (44.4)	5 (62.5)	6 (46.1)
Birthweight (g)	2975 (2780 – 3220)	3280 (2940 – 3335)	3290 (2780 – 3670)	2558 (2010 – 3268)
Small-for-gestational	3 (30)	0 (0)	0 (0)	7 (53.8)

age Ψ

APGAR score 5 min <7	0 (0)	0 (0)	1 (12.5)	1 (7.7)
Umbilical artery pH	7.21 (7.15 – 7.23)	7.18 (7.12 – 7.21)	7.17 (7.12 – 7.2)	7.22 (7.17 – 7.24)
Admission to neonatal intensive care unit	1 (10)	0 (0)	1 (12.5)	5 (38.5)

611

612 Data are median (interquartile range) or n (%) as appropriate.

613 # Preterm delivery defined as delivery occurring before 37 weeks of gestation.

614 Ψ Small for gestational age defined as birthweight below the 10th centile according to local
 615 standards.

616 * $p < 0.05$ by Mann Whitney U test, Pearson χ^2 or Fisher exact tests as appropriate, as
 617 compared to controls.

FIGURE LEGENDS

Figure 1. Scattered boxplots showing the levels of soluble endothelial damage and immune response markers in the study populations. The line in the boxes depicts the sample median and the boxes are the 1st and 3rd quartiles. The whiskers point to the maximum and the minimum values of the sample. For a better visualization of data points distribution and to show possible outliers, a second layer of information is included in the figure with all data points scattered along the y axis. Significant differences of adjusted p values (Student's t-test, Benjamini-Hochberg procedure for multiple pairwise comparisons) are noted as * $p < 0.05$ and ** $p < 0.01$ vs. Control, \$ $p < 0.05$ and \$\$ $p < 0.01$ vs. mild COVID-19, and # $p < 0.05$ and ## $p < 0.01$ vs. severe COVID-19. Controls (C, $n=10$), mild COVID-19 (mcovid-19, $n=9$), severe COVID-19 (scovid-19, $n=8$), preeclampsia (PE, $n=13$).

Figure 2. Analysis of the differential profile of soluble biomarkers among the study groups. Through statistical methods previously described the variability of all the soluble markers analyzed was transformed into the following: 2A) 2-dimensional principal component analysis (to visualize the distribution in 2 dimensions of the variability in the different study groups) 2B) Unsupervised hierarchical clustering based on univariate analysis comparing severe COVID-19 vs. Preeclampsia. In this analysis, a z-score transformation was performed on the intensity of each biomarker across all samples and each sample z-score is displayed in the heatmap. Biomarkers (in rows) and samples (in columns) are clustered by Euclidean distance and Ward linkage. Controls (C, $n=10$), mild COVID-19 (mcovid-19, $n=9$), severe COVID-19 (scovid-19, $n=8$), preeclampsia (PE, $n=13$). VWFAg, VWF antigen; VWFGPIbM, VWF activity; $\alpha 2AP$, $\alpha 2$ -antiplasmin; sFlt-1, soluble fms-like tyrosine kinase-1; PlGF, placental growth factor; Ang2, Angiopoietin 2; VCAM-1, vascular cell adhesion molecule-1, TNFRI, soluble TNF- α receptor I, HS, heparan sulfate; NETS, neutrophil extracellular traps; PAI, plasminogen activator inhibitor; TM, thrombomodulin.

Figure 3. VWF multimeric analysis in preeclampsia and severe COVID-19 patients. This analysis was performed to confirm the qualitative defects of this protein suggested by the low VWF:Ag/VWF:GPIb detected in PE patients. A normalized 1.2% multimer gel (A) and densitometry (B) of plasma VWF multimers from normal control (C), patient with known von Willebrand disease Type 2A as a positive control (VWD2A) (characterized by a loss of high molecular weight multimers and an increase in the low molecular weight multimers), 4 preeclampsia patients (PE), and 2 pregnant women with severe COVID-19 (sCOVID-19). Of note, each sample dilution was performed following the antigenic concentration (VWF:Ag) previously obtained (additionally it was an indirect confirmation of the results). Sample identification is followed by the dilution used to resolve VWF multimeric pattern.

Figure 4. Expression of ICAM-1 and VWF in cultured endothelial cells: effect of COVID-19 and preeclampsia sera. Changes in inflammation and thrombogenic phenotypes induced by the study conditions were explored through an *in vitro* approach consisting of the exposure of endothelial cells in culture to patients' sera. Representative fluorescence micrographs showing ICAM-1 expression (in green, on the left panel) on cell surface and VWF release (in red, on the right panel) on endothelial cells in culture supplemented with serum from controls (C) or mild and severe COVID-19 (m and sCOVID-19) and preeclampsia (PE). The boxplots represent the quantitative assessment of ICAM-1 expression and VWF release. The line in the boxes depicts the sample median and the boxes are the 1st and 3rd quartiles. The whiskers point to the maximum and the minimum values of the sample ($n = 6$, $*P < 0.05$ vs. Controls and $\#P < 0.05$ comparison between sCOVID-19 and PE).

Figure 5. Inflammatory signaling pathways in endothelial cells exposed to mild and severe COVID-19 and preeclampsia milieu. The analysis of signaling pathways activation was

performed by exposing resting endothelial cells to the sera under study. In the present study, activation of p38 MAPK (a kinase with a key role in inflammatory cellular responses to injurious stress) in endothelial cells in culture exposed to sera from controls (C), mild and severe COVID-19 (m and sCOVID-19), and preeclampsia (PE) patients for 5 minutes. Immunoblot image shows phosphorylated p38 MAPK and B-actin, and the boxplots represent the relative quantification of p38 MAPK / B-actin compared to controls (n=3, *P < 0.05 vs. controls).

Glossary of terms

- A disintegrin and metalloproteinase with thrombospondin type 1 motif, 13 (ADAMTS-13): is primarily synthesized in the liver, and its main function is to cleave von Willebrand factor (VWF) anchored on the endothelial surface, in circulation, and at the sites of vascular injury.
- Angiopoietin 2 (Ang2): is produced by endothelial cells and acts as an autocrine regulator mediating vascular destabilization and regulating vascular homeostasis.
- α 2-antiplasmin (α 2AP): is a serine protease inhibitor (serpin) responsible for inactivating plasmin.
- Endothelium: composed by endothelial cells plays an important role in inflammation by regulating vascular permeability for macromolecules and leukocytes, vascular tone and hemostasis, and by binding and producing and inflammatory mediators such as cytokines.
- *Ex vivo* approach: to quantify the degree of endothelial activation is of interest when evaluating inflammation. Due to the localization of this type of cells, this evaluation cannot be carried out directly and a number of indirect measures such as the measurement of soluble molecules released by the endothelium has been employed instead.


- Soluble fms-like tyrosine kinase-1 (sFlt-1): is a circulating antiangiogenic protein synthesized by the placenta, which acts as an antagonist of vascular endothelial growth factor (VEGF) and placental growth factor (PlGF) and is upregulated in preeclampsia.
- sFlt-1/PlGF ratio: an imbalance in these two biomarkers levels has been reported to be involved in preeclampsia pathogenesis. An elevated sFlt-1/PlGF seems to be highly predictive of preeclampsia.
- Heparan sulfate (HS): is glycosaminoglycan from endothelial glycocalyx used by viral pathogens such as SARS-CoV-2 for the initial interaction with host cells.
- *In vitro* approach: consists in a well characterized *in vitro* model of endothelial dysfunction, in which endothelial cells in culture are exposed to patient's sera in order to assess its capacity to modulate the endothelial phenotype. This analysis is performed through the quantification of changes in inflammatory and thrombogenicity markers together with the activation of certain intracellular signaling pathways.
- Intercellular adhesion molecule-1 (ICAM-1): adhesion molecule that is upregulated during endothelial activation and mediates lymphocyte binding. This molecule is not only released from endothelium, but also from lymphocytes, monocytes and eosinophils. Elevated levels of soluble ICAM-1 have been reported in preeclampsia.
- Neutrophil extracellular traps (NETS): are extracellular webs of chromatin, microbicidal proteins, and oxidant enzymes that are released by neutrophils to fight

against infections and that, in elevated concentrations, have the potential to propagate inflammation and microvascular thrombosis.


- Placental growth factor (PlGF): is a member of the vascular endothelial growth factor (VEGF) family and is predominantly expressed in the placenta. The circulating levels of this molecule have been postulated as a useful screening tool in the prediction preeclampsia.
- Plasminogen activator inhibitor (PAI): is a member of the serine protease inhibitor (serpin) superfamily and constitutes a central molecule linking pathogenesis and progression of thrombotic vascular events.
- Principal component analysis: is a statistical method that aims to reduce the dimensionality of large data sets by transforming them into a smaller ones. This method preserves as much information as possible and the resulting data set become easier to explore and visualize than the original one.
- p38 mitogen-activated protein kinase (P38MAPK): plays a pivotal role mediating cellular responses to injurious stress and immune signaling partly through the activation of gene expression.
- soluble Complement 5b-9 (C5b9): is also known as soluble membrane attack complex and constitutes a marker of complement activation. This molecule creates a transmembrane channel on the surface of targeted cell that leads to cell lysis and death.

- 748 • Soluble TNF receptor type I (sTNFRI): is one of the two soluble receptors of TNF-alpha
749 (TNF α), a proinflammatory cytokine that plays a central role in inflammation, which
750 act as physiological attenuator of TNF α activity.
751
- 752 • Thrombomodulin (TM): is a thrombin receptor on endothelial cells that is involved in
753 promoting activation of the anticoagulant protein C pathway during blood coagulation.
754
- 755 • Vascular Cell Adhesion Molecule -1 (VCAM-1): adhesion molecule that is upregulated
756 during endothelial activation and mediates lymphocyte binding. Elevated levels of
757 soluble VCAM-1 have been reported in preeclampsia.
758
- 759 • von Willebrand Factor (VWF): a multimeric blood protein primarily synthesized, stored
760 and secreted by endothelial cells. It constitutes a marker of acute and chronic
761 inflammation. The analysis of this protein implies both antigen concentration
762 (VWF:Ag) and functionality (VWF:GPIbM).
763
- 764 • von Willebrand Factor multimeric analysis: is a method carried out by electrophoresis
765 of plasma samples using non-reducing agarose gels in the presence of different
766 concentrations of sodium dodecyl sulphate. This analysis aims to identify qualitative
767 defects of this protein and is usually performed after functional and immunological
768 VWF assays indicate a potential abnormality.


Biomarkers of endothelial damage




Biomarkers of coagulopathy/fibrinolysis




Biomarkers of innate immune response



Biomarkers of angiogenesis



A**B**