

Attenuation of *in vitro* host–pathogen interactions in quinolone-resistant *Salmonella* Typhi mutants

Clara Ballesté-Delpierre¹, Anna Fàbrega¹, Mario Ferrer-Navarro¹, Ramkumar Mathur², Sankar Ghosh² and Jordi Vila^{1*}

¹ISGlobal, Barcelona Ctr. Int. Health Res. (CRESIB), Hospital Clínic - Universitat de Barcelona, Rosselló 149-153 Barcelona, 08036, Spain; ²Department of Microbiology and Immunology, College of Physicians and Surgeons, Columbia University, 701 West 168th Street, New York City, NY 10032, USA

*Corresponding author. Tel: +34-932-27-55-22; Fax: +34-932-27-93-72; E-mail: jvila@ub.edu

Received 21 May 2015; returned 23 June 2015; revised 7 July 2015; accepted 22 August 2015

Objectives: The relationship between quinolone resistance acquisition and invasion impairment has been studied in some *Salmonella enterica* serovars. However, little information has been reported regarding the invasive human-restricted pathogen *Salmonella* Typhi. The aim of this study was to investigate the molecular mechanisms of quinolone resistance acquisition and its impact on virulence in this serovar.

Methods: Two antibiotic-resistant mutants (Ty_c1 and Ty_c2) were generated from a *Salmonella* Typhi clinical isolate (Ty_wt). The three strains were compared in terms of antimicrobial susceptibility, molecular mechanisms of resistance, gene expression of virulence-related factors, ability to invade eukaryotic cells (human epithelial cells and macrophages) and cytokine production.

Results: Multidrug resistance in Ty_c2 was attributed to AcrAB/TolC overproduction, decreased OmpF (both mediated by the *mar* regulon) and decreased OmpC. The two mutants showed a gradually reduced expression of virulence-related genes (*invA*, *hilA*, *hilD*, *fliC* and *fimA*), correlating with decreased motility, reduced infection of HeLa cells and impaired uptake by and intracellular survival in human macrophages. Moreover, Ty_c2 also showed reduced *tviA* expression. Additionally, we revealed a significant reduction in TNF- α and IL-1 β production and decreased NF- κ B activation.

Conclusions: In this study, we provide an in-depth characterization of the molecular mechanisms of antibiotic resistance in the *Salmonella* Typhi serovar and evidence that acquisition of antimicrobial resistance is concomitantly detected with a loss of virulence (epithelial cell invasion, macrophage phagocytosis and cytokine production). We suggest that the low prevalence of clinical isolates of *Salmonella* Typhi highly resistant to ciprofloxacin is due to poor immunogenicity and impaired dissemination ability of these isolates.

Introduction

Typhoid fever is a human-restricted systemic infection caused by *Salmonella enterica* serovar Typhi. This pathogen causes ~21 million infections causing typhoid fever each year, leading to ~200 000 deaths.¹ The highest incidence rates are seen in Asia, where this pathogen remains a major public health problem. The transmission route is through exposure to food and water contaminated with human faeces. Although it belongs to the same species as *S. enterica* serovar Typhimurium, their pathogenesis substantially differs. While salmonellosis caused by *Salmonella* Typhimurium infection is usually self-limiting, typhoid fever is a systemic infectious disease with much more severe consequences. Once bacteria reach the small intestine, they adhere to the mucosa and mostly invade M cells, which mediate internalization of the pathogen and allow transportation to Peyer's

patches. This process is followed by phagocytosis by dendritic cells and macrophages, thus favouring dissemination through the mesenteric lymph nodes and eventual spread to the liver, spleen and bone marrow where *Salmonella* Typhi is able to survive and replicate.^{2–4}

The treatment of typhoid fever has been changing and adapting together with the spread of drug-resistant strains. Chloramphenicol, trimethoprim/sulfamethoxazole and ampicillin were initially used until the emergence and spread of plasmid-mediated resistance to these antimicrobial agents. As a consequence, these drugs were replaced by ceftriaxone and ciprofloxacin as well as other fluoroquinolones such as ofloxacin.⁵ However, dissemination in the past decades, mostly in Asia, of strains showing both resistance to nalidixic acid (MIC >256 mg/L) and decreased ciprofloxacin susceptibility (DCS) (MIC range, 0.125–1 mg/L) has occurred.^{5–7} In addition, some reports have already shown the

emergence of highly fluoroquinolone-resistant isolates.^{8–10} This scenario is forcing the introduction of new-generation fluoroquinolones (e.g. gatifloxacin) and alternative drugs such as azithromycin.^{5,11}

Quinolones act through inhibition of DNA gyrase and topoisomerase IV, hampering DNA replication and transcription as well as interfering with cell division. In *S. enterica*, resistance to quinolones is mainly due to point mutations in the quinolone resistance-determining regions (QRDRs) of the target genes, with amino acid substitutions at positions 83 and 87 of GyrA being the most frequently reported.^{4,12,13} Mutations affecting the internal accumulation of the drug, by decreasing the expression of porins (e.g. OmpF and OmpC) and/or increasing the expression of efflux pumps such as AcrAB/TolC,^{13–15} are also of great concern. Expression of the efflux pump AcrAB is controlled by its local repressor AcrR and by three homologous transcriptional activators, MarA, SoxS and RamA. The contribution of these regulators to the MDR phenotype has been reported in laboratory mutants and clinical isolates in *S. enterica*.^{16–20} There is only a single exception, since results concerning the role of MarA have been demonstrated in *Escherichia coli*,^{21–23} but not yet in clinical isolates of *Salmonella*. Similarly, activation of the transcription of *micF*, an antisense RNA that inhibits synthesis of the outer membrane porin OmpF,²⁴ has yet to be supported with clinical data for *Salmonella*. In addition, plasmid-encoded genes [e.g. *aac(6′)-Ib-cr* and the *qnr* genes] are also responsible for quinolone resistance in *Salmonella* spp.¹³

Pathogenicity is primarily mediated by a number of virulence factors, the genes of which are organized within particular regions of the genome, named *Salmonella* pathogenicity islands (SPIs). To date, 15 SPIs have been identified in *Salmonella* Typhi, among which SPI-1, homologous to that described in *Salmonella* Typhimurium, contains genes encoding the type III secretion system-1 (T3SS-1) and secreted effector proteins needed for the invasion process.^{25–27} Furthermore, *Salmonella* Typhi-specific SPI-7 contains genes involved in the biosynthesis, regulation and export of the Vi capsular antigen. This virulence factor plays a key role in the attempt of *Salmonella* Typhi to avoid host defences and is therefore important in enhancing infectivity and virulence.²⁸ Moreover, innate immune responses such as inflammation are also activated through the interaction of pathogen-associated molecular patterns, e.g. LPS and flagellin, with the Toll-like receptors (TLRs) TLR-4 and TLR-5, respectively.^{4,27,29} Stimulation of TLRs induces activation of the transcriptional regulator NF-κB, thus triggering the production of proinflammatory cytokines such as TNF-α, IL-6 and IL-1β.³⁰

Previous reports conducted in non-Typhi *S. enterica* have shown that acquisition of quinolone resistance is related to decreased expression of virulence factors and impaired adherence to and invasion into the host cell.^{31,32} Even though there is an epidemiological relatedness between Typhi and non-Typhi serovars, many differences have been reported in terms of virulence and colonization behaviour, suggesting that adequate experiments should be conducted to address this relationship in *Salmonella* Typhi. Thus, the main objective of this study was to investigate if this loss of virulence, in terms of invasion and induction of host immune responses, also occurred in *Salmonella* Typhi upon acquisition of quinolone resistance. Moreover, despite the above-mentioned mechanisms of quinolone resistance being well studied in non-typhoidal *Salmonella*, an in-depth

characterization of the molecular mechanisms involved in antibiotic resistance of two *Salmonella* Typhi mutants was also carried out.

Materials and methods

Strains

A clinical isolate of *Salmonella* Typhi was recovered from a patient diagnosed with spondylodiscitis at the Hospital Clinic of Barcelona (Spain). This isolate (Ty_wt) was used to generate *in vitro* antibiotic-resistant mutants in a multistep selection process, as previously described,³¹ by exposure of the bacteria to increasing concentrations of ciprofloxacin, starting at 0.25 mg/L. The mutants studied in this work were selected at 1 and 2 mg/L ciprofloxacin, named Ty_c1 and Ty_c2, respectively. When indicated overnight cultures were grown for 16–18 h.

Antimicrobial susceptibility

Antimicrobial susceptibility to nalidixic acid, ciprofloxacin, norfloxacin, ampicillin, ceftriaxone, ceftiofloxacin, erythromycin, chloramphenicol, tetracycline, trimethoprim and kanamycin was assessed using Etests (bioMérieux) on Mueller–Hinton II plates (Becton Dickinson) following the manufacturer's recommendations. At least three replicates of each susceptibility test were performed.

Sequencing of quinolone resistance genes

DNA amplification of the target genes *gyrA*, *gyrB*, *parC* and *parE* as well as the global regulators *acrR*, *marAB*, *soxRS* and *ramR* was performed using the primers listed in Table S1 (available as Supplementary data at JAC Online). Sequencing was performed by Beckman Coulter Genomics (Essex, UK) and the sequences analysed by alignment with the template sequence of *Salmonella* Typhi (RefSeq NC 016832.1).

Bacterial growth

Fresh cultures were grown at 37°C with shaking, as previously described,³¹ and OD readings at 600 nm (OD₆₀₀) were done every 15 min for 24 h by means of an iEMS Multiskan Reader MF (Thermo Fisher Scientific). Each plate included four replicates of each sample and the assay was repeated three times.

Motility

Bacterial cultures were grown overnight in LB medium at 37°C with shaking and 10 μL was 'stab inoculated' into soft agar plates containing 0.5% agar. Plates were incubated for 7 days at 37°C in a humid chamber. Motility was assessed by measuring and comparing the growth diameter of the three tested strains.

Relative expression of resistance and virulence genes

RNA extraction from exponential cultures of Ty_wt, Ty_c1 and Ty_c2 was performed as reported by Fábrega *et al.*³¹ A two-step RT-PCR was performed as previously described by our group³³ following the 2^{-ΔΔCT} method. Briefly, the obtained RNA was reverse transcribed to cDNA using the PrimeScript RT Reagent Kit (Takara) followed by PCR reaction (SYBR® Premix Ex Taq Tli RNase H Plus Kit, Takara) under universal thermal cycling conditions. Expression of the efflux-related genes *acrB*, *tolC*, *emrB* and *acrF*, the *marA* regulator, the *ompF* porin and the virulence-related genes *hilA*, *hilD*, *invA*, *fliC*, *firmA* and *tvfA* was studied. The primers used (Table S2) were designed using Primer Express® software (Applied Biosystems). The 2^{-ΔΔCT} method was applied to measure the gene

Q4
235 expression, defined as relative quantification (RQ) of the target genes in the two mutant strains, normalized with the 16S rRNA as reference gene and the Ty_wt as calibrator strain. Five independent extractions were performed and differences of >2-fold were considered relevant. Standard deviation was calculated and only reported when this value was >0.1.

240 **Epithelial cell invasion assay**

The invasion assay was performed according to Fàbrega *et al.*³¹ with some modifications. Briefly, HeLa cells (ECACC 84211901) were seeded into 6-well tissue culture-treated plates (Corning) in order to obtain a monolayer corresponding to $\sim 5 \times 10^5$ cells/well at 37°C in a 5% CO₂ atmosphere. Infection was then performed with bacteria grown overnight at 37°C without shaking at an moi of 100. Plates were incubated at 37°C/5% CO₂ for 2 h followed by an additional 2 h of incubation in the presence of 100 mg/L gentamicin (Life Technologies) to kill extracellular bacteria. Wells were then washed and 1 mL of chilled sterile water was added and kept at 4°C for 30 min to lyse the cells. This volume was recovered for intracellular bacterial counting by plating several dilutions in LB agar. At least three independent experiments were performed with intraexperiment duplicates. The invasion ability of each strain was determined by calculating the ratio between the number of intracellular bacteria and the inoculum.

255 **Human macrophage infection assay**

Human-derived monocytes (U-937 cell line) were used upon differentiation into adherent macrophages with 50 µg/L phorbol 12-myristate 13-acetate. A total of 5×10^5 cells/well were seeded into 24-well plates (Becton Dickinson) and incubated at 37°C/5% CO₂ with RPMI-1640 (Life Technologies) supplemented with 10% FBS (Life Technologies). After 24 h, wells were washed three times with PBS (Life Technologies) and new medium was added in order to eliminate the non-adhered cells. Infection was carried out with *Salmonella* Typhi strains grown overnight at 37°C without shaking at an moi of 50. Plates were then centrifuged for 5 min at 2500 g at room temperature and incubated for 20 min at 37°C/5% CO₂. Afterwards, cells were washed again and new medium containing 100 mg/L gentamicin was added before incubating for 2 h under the same conditions as described above. After this time, bacterial entry into host cells was determined by eliminating the supernatant, incubating the cells for 15 min at 4°C with 1 mL of chilled sterile water and then plating the appropriate dilutions on LB agar plates in order to allow bacterial counting. Alternatively, to determine the uptake rate, after 2 h of incubation with 100 mg/L gentamicin, the medium was replaced with new RPMI containing a lower dose of gentamicin (12 mg/L) and maintained for 24 h. Then, wells were washed and treated with 1 mL of chilled sterile water for cell lysis and bacterial counting. The intracellular survival ability was determined by calculating the ratio between the number of bacteria recovered at 24 and 2 h post-infection.

280 **Cytokine assay**

Production levels of TNF-α and IL-1β were measured in the supernatants during the human macrophage infection assay. After the two incubation periods (2 and 24 h), supernatants were collected and levels of TNF-α (from the 2 h post-infection plates) and IL-1β (from the 24 h post-infection plates) were measured by ELISA according to the manufacturer's recommendations (BD OptEIA).

290 **NF-κB luciferase assay**

HEK293 cells stably transfected with an NF-κB luciferase reporter construct, previously obtained by Koblansky *et al.*,³⁴ were plated onto 48-well plates. After 24 h of incubation at 37°C/5% CO₂, cells were

stimulated with heat-killed *Salmonella* Typhi as well as with 0.2 µg/L TNF-α as a positive control for 6 h. Luciferase activity was measured with the Luciferase Reporter Assay System (Promega). Heat-killed bacteria were prepared from an overnight culture adjusted to OD₆₀₀=0.7. An 800 µL aliquot of the bacterial culture was centrifuged at 13 500 g for 10 min, washed twice with PBS and resuspended in 1 mL of the same solution. Then, 1 h incubation at 65°C was followed and the bacterial solution was vortexed for 3 min and centrifuged for 5 min at 2500 g to discard protein debris. For stimulation, 25 µL of the supernatant was added to each well. Each sample was done in triplicate and three independent assays were performed.

Outer membrane protein analysis

Extraction of outer membrane proteins was performed with *N*-lauroyl sarcosinate. Briefly, bacteria were harvested from 200 mL of exponential culture (OD₆₀₀=0.6) by centrifugation at 4°C and 3500 g. Cells were washed twice in PBS in order to remove any residual medium. Then, cells were resuspended in 6 mL of 10 mM Tris, pH 8.0/1% NaCl. At this point, cells were sonicated for 15 min (cycles of 59 s 'on' and 59 s 'off'). After cell disruption, samples were centrifuged at 4°C and 3500 g in order to remove any cell debris. Supernatants were collected and transferred into ultracentrifugation tubes and samples were centrifuged at 100 000 g for 1 h at 4°C in a Sorvall MS-150 microultracentrifuge (Thermo Scientific) using an S50-ST rotor. After this centrifugation step, the supernatant was discarded and the pellet resuspended in 1% sarcosyl solution and incubated for 60 min at room temperature with gentle agitation. After incubation, samples were again centrifuged at 100 000 g for 1 h at 4°C and the resulting pellet was cleaned twice with 10 mM Tris, pH 8.0/1% NaCl. Finally, the pellet was carefully resuspended in 500 mL of milliQ water and the protein concentration estimated with the 2D-Quant Kit (GE Healthcare). Protein extracts were separated by SDS-PAGE on 12.5% isocratic Laemmli gels using a mid-size gel casting system (Hoefer SE600 Chroma). Gels were run at 25 mA following the manufacturer's recommendations. Bands showing differential expression between strains were recovered and further identified through MALDI-TOF MS performed using an Ultraflex instrument (Bruker Daltonics).

Statistics

Data were analysed using IBM SPSS Statistics 20 software. As data were normally distributed, multiple comparisons were performed using the one-way ANOVA test. *P* values <0.05 were considered to be significant.

Results

Selection of the strains

A *Salmonella* Typhi clinical isolate (Ty_wt) was selected to generate mutants able to grow at inhibitory concentrations of ciprofloxacin. First, Ty_wt was plated onto MacConkey agar plates containing 0.25 mg/L ciprofloxacin, corresponding to 0.5× its MIC (0.5 mg/L). From this point onwards, the concentration of ciprofloxacin was doubled at each step, in which a single colony was selected and plated again in the following selection stage, reaching a ciprofloxacin concentration of 16 mg/L. The mutants selected for the study were Ty_c1 and Ty_c2, which corresponded to colonies grown on plates containing 1 and 2 mg/L ciprofloxacin with MICs of ciprofloxacin of 1 and 8 mg/L, respectively. The mutant recovered at 4 mg/L ciprofloxacin, with an MIC of 8 mg/L, was not considered for the study as it showed the same antimicrobial susceptibility profile as Ty_c2, suggesting that no additional resistance mechanisms were selected. Mutants able to grow

at greater concentrations (8 and 16 mg/L) were also recovered (MICs 12–24 mg/L), although they could not be further studied as they showed important growth impairment. Resistant colonies appeared at frequencies of 2×10^{-6} and 7×10^{-6} mutants per cfu for Ty_c1 and Ty_c2, respectively. When bacteria were exposed to a concentration of 8 mg/L ciprofloxacin, the frequency of selection decreased to 5.5×10^{-7} and in the presence of 16 mg/L it was further reduced to 6×10^{-8} mutants per cfu.

To assess reversion of the resistance phenotype, the selected resistant mutants Ty_c1 and Ty_c2 were plated onto MacConkey agar plates without antibiotic. However, no revertant colonies were obtained after 35 consecutive passages.

Bacterial growth

A link between fluoroquinolone resistance and fitness has been stated previously. Several studies carried out in *Salmonella* Typhimurium and *S. enterica* serovar Enteritidis have shown that acquisition of high levels of fluoroquinolone resistance have a fitness cost.^{31,35,36} In order to evaluate this feature in our strains, bacterial growth was examined. No differences between the three strains were observed, indicating that acquisition of antimicrobial resistance in these mutants (ciprofloxacin MICs of 1 and 8 mg/L) did not impose an energy cost on bacterial growth. However, the mutants with ciprofloxacin MICs of 12 and 32 mg/L (named Ty_c8 and Ty_c32) showed a much longer lag phase than the previous strains and Ty_c32 did not reach the same OD in the stationary phase as the previous mutants (Figure 1).




Figure 1. Mutants Ty_c8 and Ty_c32 show impaired growth. Bacterial growth of all the mutants obtained was assessed. The most resistant mutants (Ty_c8 and Ty_c32) were rejected as they presented growth deficiencies. Results correspond to the mean of three independent experiments.

Table 1. Antimicrobial susceptibility testing of the three strains studied

Strain	MIC (mg/L)										
	NAL	CIP	NOR	ERY	CHL	TET	TMP	AMP	FOX	CRO	KAN
Ty_wt	250	0.5	1	96	4	0.5	0.094	0.094	1	0.064	1
Ty_c1	1000	1	8	128	4	1.5	0.25	0.75	4	0.125	1
Ty_c2	2000	8	24	>256	128	4	0.75	4	128	0.5	1

NAL, nalidixic acid; CIP, ciprofloxacin; NOR, norfloxacin; ERY, erythromycin; CHL, chloramphenicol; TET, tetracycline; TMP, trimethoprim; AMP, ampicillin; FOX, cefoxitin; CRO, ceftriaxone; KAN, kanamycin.

465 previous study and were suggested to contribute to the quinolone
resistance phenotype,³⁷ sequencing of the entire target genes
(*gyrA*, *gyrB*, *parC* and *parE*) was also performed, but no additional
change was found.

470 *Multidrug resistance profile*

The absence of new mutations acquired in the target genes of the
two mutants suggested the acquisition of broad-spectrum resist-
ance mechanisms such as increased efflux. In this context, the
antimicrobial susceptibility profile of Ty_wt, Ty_c1 and Ty_c2
was assessed against a broader collection of antibiotics including
erythromycin, ampicillin, ceftriaxone, ceftiofloxacin, chloramphenicol,
tetracycline, trimethoprim and kanamycin. The results obtained,
reported in Table 1, showed a progressive, but different, increase
in the MIC of most of the antibiotics tested. In comparison with
Ty_wt, Ty_c1 showed the highest increase in the MIC of ampicillin
(8-fold), modest increments (2- to 4-fold) in the MICs of tetracy-
cline, trimethoprim, ceftiofloxacin and ceftriaxone, whereas no major
change (<1.5-fold) was seen for chloramphenicol and erythromy-
cin. Contrarily, a different response was observed for strain Ty_c2.
This mutant could be classified as MDR according to the ECDC.³⁸ It
acquired resistance to ceftiofloxacin and chloramphenicol with a
32-fold increase in the MIC compared with Ty_c1. Moreover, and
even though the MICs of the remaining drugs tested did not reach
the resistance breakpoints, increases of 3- to almost 5-fold were
recorded in all cases. The only antibiotic tested for which no
changes in susceptibility were seen in any of the mutants was
kanamycin.

495 *Efflux and permeability are altered in Ty_c2*

In *Salmonella* Typhimurium, the predominant transport system
involved in multidrug resistance is AcrAB/TolC.¹⁵ Other less
frequently detected drug transporters such as AcrF, highly homolo-
gous to AcrB, and EmrB have also been reported to extrude
quinolones^{13,39} and have been shown to be overexpressed in
quinolone-resistant *Salmonella* Typhimurium mutants.⁴⁰ Expression
of the *acrB*, *tolC*, *acrF*, *emrB* and *ompF* genes was tested in the
three strains by real-time PCR (RT-PCR). Results revealed overex-
pression of *acrB* and *tolC* only in Ty_c2 (RQ=3.91 for *acrB* and
RQ=3.42 for *tolC*). Surprisingly, the other efflux-related genes,
acrF and *emrB*, were repressed in this strain compared with
Ty_wt (3- and 3.9-fold, respectively). Additionally, the porin
ompF was >6-fold repressed in Ty_c2. In the case of Ty_c1, no
major change was detected and considered to be relevant for
any of these genes despite statistically significant differences
were seen for *acrF* and *emrB* (fold change <2). Thus, these results
suggest the involvement of AcrAB/TolC and OmpF in the multidrug
resistance profile of Ty_c2 (Table 2).

Analysis of the bacterial outer membrane proteins further con-
firmed part of these changes in gene expression: two bands iden-
tified as AcrA and TolC were overproduced only in Ty_c2, together
with a decreased production of the outer membrane protein
OmpC (Figure 2).

520 *Role of MarA in the antimicrobial resistance profile of Ty_c2*

Mutations in AcrR have been shown to impair its repressive effect,
hence leading to derepression of the *acrAB* genes.¹⁶⁻¹⁸ The

Table 2. Transcriptional levels of genes involved in antimicrobial resistance and virulence

Strain	Efflux-related genes					Regulatory genes					Virulence genes				
	<i>acrB</i>	<i>tolC</i>	<i>acrF</i>	<i>emrB</i>	<i>ompF</i>	<i>marA</i>	<i>soxS</i>	<i>ramA</i>	<i>acrR</i>	<i>hilD</i>	<i>hilA</i>	<i>invA</i>	<i>tvfA</i>	<i>fimA</i>	<i>fljC</i>
Ty_wt	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Ty_c1	-1.08	-1.64	-1.60*	-1.90**	-1.20*	-1.28	1.25	-1.32*	-1.89**	-2.50**	-2.59**	-3.75**	-1.26*	-3.22**	-9.89**
Ty_c2	3.91**	3.42**	-3.06*	-3.96**	-6.37**	108.19**	-2.38**	-2.70**	-5.88**	-9.19**	-5.04**	-8.55**	-27.85**	-6.09**	-121.09**

*P<0.5 and **P<0.01.

repressors MarR and RamR and the activator SoxR are regulators of transcriptional factors MarA, RamA and SoxS, respectively. Mutations in RamR and SoxR have been reported in different *Salmonella* serovars to increase expression of the homologue activators and, therefore, increase protein levels of AcrAB and down-regulate OmpF.¹⁶⁻²⁰ However, studies describing mutations in MarR in strains selected in the clinical setting have been reported in *E. coli*, but not in *Salmonella* spp.^{21,22}

In this study, sequencing of the known regulatory regions of AcrAB expression (*acrR*, *marRAB*, *soxRS* and *ramR*) was carried out. Indeed, an important alteration in the *marR* locus was detected in the Ty_c2 mutant corresponding to a deletion of 445 nucleotides. The fragment deleted included the first 411 nucleotides of the *marR* gene and 34 nucleotides upstream of the gene (Figure 3). In *E. coli*, the existence of two binding sites (Site I and Site II) for MarR in the *marO* region has been reported.⁴¹ In Ty_c2, the entire Site II was deleted although Site I was completely preserved as well as the -35 and -10 elements of the promoter. Interpretation of this finding suggests that MarA expression is not only possible since the promoter is still intact, but also overexpression is likely to be detected due to a lack of the MarR protein. Consistently, RT-PCR transcription analysis of *marA* revealed high overexpression (RQ 108.19 ± 0.16) in the MDR strain Ty_c2 compared with Ty_wt and Ty_c1. Nonetheless,

despite no further changes being seen in the sequences of the other regulators, repression of *acrR*, *soxS* and *ramA* ranging from 2- to almost 6-fold was observed in Ty_c2 (Table 2). The results obtained in this section suggest that overexpression of MarA is responsible, at least in part, for the resistance phenotype observed in Ty_c2.

Evaluation of virulence in the quinolone-resistant *Salmonella* Typhi mutants

Gradual reduction of motility seen in Ty_c1 and Ty_c2

Intact motility of *Salmonella* Typhi has been determined to participate in the invasion process and is therefore related to the virulence potential of this pathogen.^{42,43} For this reason, the motility of the three strains was measured by inoculation into soft agar plates. Although the clinical strain (Ty_wt) was already poorly motile, with a colony diameter of 13 mm, a decrease in motility was seen for the mutants, showing diameters of 9 mm for Ty_c1 and 7 mm for Ty_c2 (data not shown).

As flagellin is the major component of the bacterial flagellar filament, the expression profile of the flagellin-encoding gene *fliC* was analysed by RT-PCR and revealed statistically significant differences between the parental strain and the mutants (Table 2). Ty_c1 showed a 10-fold reduction in *fliC* expression levels in comparison with Ty_wt and an additional 10-fold decreased expression was seen for Ty_c2 (RQ=0.01). These results are in accordance with the phenotype observed in the soft agar assay.

In order to determine whether the differences in *fliC* expression could be attributed to the acquisition of mutations within the gene, *fliC* as well as the regulators *fliA* and *flhDC*, involved in flagella biosynthesis,^{35,44} were sequenced. Amino acid substitutions at particular positions of the conserved site of flagellin have been reported to be essential for protofilament assembly, bacterial motility and TLR-5 recognition.⁴⁵ A single mutation in *fliC* was found in both Ty_c1 and Ty_c2 leading to the amino acid substitution T187I whereas no mutations in any of the regulators were detected. This change detected in the mutants is located outside of the conserved region, likely suggesting it to be trivial for the function and structure of flagellin.⁴⁵




Figure 2. Ty_wt, Ty_c1 and Ty_c2 have different outer membrane protein profiles. Proteins showing differential expression are indicated by arrows.




Figure 3. Schematic representation of the deleted region located in the *marRAB* regulatory region of Ty_c2. The predicted -35 and -10 promoter regions are shown in underlined bold letters. The two MarR-binding sites (Site I and Site II) are also indicated and extrapolated from *E. coli*.⁴¹ Grey letters in italics correspond to the deletion detected and numbers in grey indicate the first and last position of the deleted nucleotides.

Salmonella Typhi mutants are less able to invade epithelial cells

As described previously in several serovars of *S. enterica*,^{31,32,46} acquisition of quinolone resistance has been related to decreased invasion ability in an *in vitro* eukaryotic cell model. For this reason, *in vitro* invasion of HeLa cells was examined with the three studied strains (Ty_wt, Ty_c1 and Ty_c2) and revealed a gradual reduction in the invasion ability of the mutants compared with the parental isolate. The invasion rate of Ty_wt was 7.31%, that of Ty_c1 was 1.69% (~4 times lower) and the lowest rate of invasion (0.27%) was reported for Ty_c2, which was 6 times lower than the Ty_c1 mutant and 27 times lower than Ty_wt (Figure 4a).

Salmonella Typhi mutants show impaired internalization and survival in macrophages

Internalization of *Salmonella* Typhi by phagocytes once *Salmonella* Typhi has reached the submucosa as well as survival of the pathogen inside these host cells are crucial processes for systemic infection.² Thus, differentiated U-937 human macrophages were infected with the three strains. In order to study their ability to be phagocytosed by macrophages, the number of intracellular bacteria was determined at 2 h post-infection. Results were expressed as percentage values of recovered bacteria with respect to the total number of infecting bacteria. A reduction of 2.75-fold in Ty_c1 compared with Ty_wt and an additional 1.6-fold for Ty_c2, which corresponded to 4.37 times less than Ty_wt, was seen (Figure 4b). Moreover, the intracellular survival rate was also determined and expressed as the percentage bacterial count at 2 h versus bacterial load at 24 h post-infection. Similar results were observed for the two mutants and corresponded to a 4-fold reduction compared with the parental strain (Figure 4c). These results showed a gradual inability of the mutants to be recognized by macrophages and therefore be internalized. In addition, the internalized bacteria were less able to survive inside macrophages, showing no significant differences between the two mutants.

Transcription levels of virulence factors are low in *Salmonella* Typhi mutants

Among the virulence factors reported to play a key role in *Salmonella* Typhi–host interactions, several loci have been described. The invasion process of *Salmonella* is mainly driven by virulence determinants contained in SPI-1,^{25,26,36} such as the effector *invA* and the key regulators *hilD* and *hilA*.⁴⁷ Type 1 fimbriae are important for adhesion to eukaryotic cells and are encoded by *fim* genes, with *fimA* being responsible for the production of the major fimbrial subunit.³⁶ The Vi capsular antigen, a *Salmonella* Typhi-specific determinant encoded by the *tviA* gene located in SPI-7, is involved in systemic dissemination.²⁵ In this study, the gene expression profiles of *invA*, *hilD*, *hilA*, *fimA* and *tviA* were evaluated through RT-PCR analysis (Table 2). The results reflect sequential gene repression in the two mutants compared with the original strain for almost all the genes tested. In Ty_c1, *invA*, *hilD*, *hilA* and *fimA* were 2.5- to 3.75-fold less expressed than in Ty_wt. On comparing Ty_c2 versus Ty_c1, expression of these genes was reduced by 1.89- to 3.67-fold. Contrary to the results observed for these genes, *tviA* expression in Ty_c1 was




Figure 4. Ability to invade HeLa cells and uptake by and survival in macrophages is compromised upon infection with Ty_c1 and Ty_c2. Invasion was performed by infecting HeLa cells with the different bacteria (a). Uptake at 2 h (b) and survival at 24 h (c) was assessed by infecting human differentiated macrophages with the three strains studied. Results correspond to the mean of three independent experiments \pm standard error. Differences between groups were assessed using the one-way ANOVA test. * $P < 0.05$.

comparable to that seen for the parental strain (RQ=0.79). However, almost 30-fold less expression was observed in Ty_c2 (Table 2). These results suggest involvement of *invA*, *hilD*, *hilA* and *fimA* in the progressive loss of virulence in the two mutants; however, *tviA*, which showed decreased expression, would only be related to the phenotype of Ty_c2.

Induction of the immune response is compromised in *Salmonella* Typhi mutants

The levels of the proinflammatory cytokines TNF- α and IL-1 β produced by infected U-937 macrophages were measured. Levels of TNF- α were checked in the supernatants collected after 2 h of infection and revealed a gradual and significant decrease in the two mutants compared with Ty_wt. The levels of this cytokine consisted of 5181 pg/mL in Ty_wt, almost 4000 pg/mL in Ty_c1 and 3256 pg/mL in Ty_c2 (Figure 5a). A greater reduction was seen for IL-1 β , checked in the supernatants collected at 24 h post-infection: Ty_wt reached 3082 pg/mL whereas the levels in Ty_c1 and Ty_c2 were reduced by >50%, showing values of 1167 and 1067 pg/mL, respectively (Figure 5b).

Since transcription of these two cytokines is controlled by NF- κ B,^{2,27} activation of this nuclear transcription factor was checked upon stimulation of HEK293 cells with heat-killed bacteria by means of a luciferase reporter assay. Activation of NF- κ B significantly diminished in Ty_c1 compared with the parental strain and corresponded to a decrease of 16.67%. This ratio was much more significant for Ty_c2, with a 71% reduction of the NF- κ B activity in comparison with Ty_wt (Figure 5c). The results obtained in this section indicate that *Salmonella* Typhi




Figure 5. *Salmonella* Typhi mutants are less immunogenic than Ty_wt. Levels of TNF- α (a) and IL-1 β (b) were measured in infected macrophages and activation of NF- κ B (c) was assessed in stimulated HEK293 cells. Results correspond to the mean of three independent experiments \pm standard error. Differences between groups were assessed using the one-way ANOVA test. * $P < 0.05$ and ** $P < 0.01$.

mutants elicit a more discrete immunogenic reaction *in vitro* than the original clinical isolate, with Ty_c2 being significantly less reactive.

Discussion

In this study, we evaluated the process of quinolone resistance acquisition and the impact on virulence properties in the pathogen *Salmonella* Typhi. In the first part of the study, we characterized the molecular mechanisms of resistance since they have not been extensively studied in this *Salmonella* serovar. We obtained two mutants, Ty_c1 and Ty_c2, with increased MICs of ciprofloxacin (1 and 8 mg/L, respectively) selected from a clinical isolate (ciprofloxacin MIC of 0.5 mg/L). All studies reporting resistance or decreased susceptibility to ciprofloxacin (MIC > 0.25 mg/L) in *Salmonella* Typhi associate this phenotype with mutations in the QRDR of the target genes, particularly in *gyrA*.^{37,48} In this study, we found that Ty_wt already harboured a GyrA substitution (S83F), but no additional change in any of the entire quinolone target genes was detected in the two mutants. The profile of increased MICs was different for the two mutants. A moderate increase to several unrelated compounds was seen in Ty_c1 whereas more suggestive increases were reported for Ty_c2 affecting all compounds tested and leading to an MDR phenotype.

According to the results obtained from gene expression and protein analyses, we found that alteration of membrane permeability was seen in Ty_c2 due to a reduction in the production of OmpF and OmpC. Although this alteration in porin expression has been previously shown in *E. coli* and non-Typhi *Salmonella*,^{31,49,50} it has not been reported in *Salmonella* Typhi. More importantly, this MDR phenotype was primarily associated with hyperproduction of the major efflux pump AcrAB/TolC. Only few reports, such as that of Chiou *et al.*,¹⁰ have previously reported increased efflux (based on the use of an efflux pump inhibitor) among *Salmonella*

Typhi clinical isolates with resistance or decreased susceptibility to ciprofloxacin (MICs of 0.12–16 mg/L). Nonetheless, this is the first time that the efflux pump AcrAB/TolC has been characterized in this serovar. Our findings together with the data of Chiou *et al.*¹⁰ contrast with the results shown by Baucheron *et al.*,⁵¹ in which no involvement of efflux mechanisms was seen in a collection of *Salmonella* Typhi clinical isolates. It is worth mentioning that the majority of these strains showed MICs of ciprofloxacin ranging from 0.125 to 1 mg/L, although a single strain had a higher MIC of 8 mg/L, attributed to the acquisition of two further target gene mutations. In the study by Chiou *et al.*,¹⁰ clinical isolates for which increased efflux activity was reported had ciprofloxacin MICs of > 4 mg/L. In our study, Ty_c2 had a ciprofloxacin MIC of 8 mg/L. Therefore, we suggest that selection of increased efflux is acquired at ciprofloxacin MICs of 4–8 mg/L as a general trend. To better confirm this hypothesis, characterization of larger collections of ciprofloxacin-resistant clinical isolates is needed to accurately evaluate the relevance and prevalence of this mechanism. In an attempt to understand the increased levels of AcrAB/TolC detected in Ty_c2, we found that this was due to alterations in the *marRAB* regulatory region. A deletion of almost the entire *marR* gene detected in this mutant likely resulted in the absence of transcription of the repressor, hence leading to hyperproduction of MarA. This phenotype, in addition to causing increased amounts of AcrAB/TolC, would also explain the down-regulation of the OmpF porin detected in Ty_c2. This is the first demonstration in *Salmonella* Typhi of the molecular regulation of this efflux system, as it has been previously described only in non-typhoidal *Salmonella*.^{13,23,52} In addition, and to the best of our knowledge, it is also the first description of a *marR* mutation reported in *S. enterica* leading to increased AcrAB/TolC, since most of the reported mutations are located in *ramR*^{17,53} and, less frequently, in *acrR*⁵⁴ and *soxRS*.¹⁹

Surprisingly, the transcription levels of the regulators *acrR*, *soxS* and *ramA* were repressed in Ty_c2. Additionally, the alternative efflux pump candidates *emrB* and *acrF*, reported to be overexpressed in quinolone-resistant *Salmonella* Typhimurium isolates,⁴⁰ were also studied and revealed down-regulation only in this mutant. A similar observation has also been reported by Kang and Woo,⁵⁵ where expression levels of efflux pump-related genes were tested in ciprofloxacin-resistant *S. enterica* serovar Istanbul mutants. In all of the mutants, not only *soxS*, but also *acrF*, were down-regulated compared with the parental strain whereas *acrB*, *ramA* and/or *marA* were overexpressed. Despite another study conducted in *E. coli*⁵⁶ reporting an increase in the expression of *soxS* in first-step mutants selected with tetracycline, a decrease in the expression levels of this gene was observed in the most resistant strain. Therefore, the authors suggested that the SoxS response could be activated at the early stages of antibiotic resistance adaptation. Considering this proposal, the decreased expression levels of *soxS* observed in Ty_c2 could be partly explained by this phenomenon. According to all these results, including the findings reported in the present study, it seems that overexpression of a particular efflux system such as AcrAB may also lead to reduced expression of some of these regulators and alternative transport systems to avoid redundancy in efflux transportation, resulting in a reduction of the energy cost of the bacteria.

Contrary to the scenario reported for Ty_c2, the increased MICs detected for the intermediate resistant mutant Ty_c1 could not be attributed to the enhanced expression of AcrAB/TolC. Moreover,

gene expression of *acrF* and *emrB* was not affected neither. These results suggest that unknown TolC-independent mechanisms of resistance with reduced impact are likely to be involved in antibiotic resistance.

In the second part of this work, the association between virulence and quinolone resistance was investigated. This relation has been studied based on data reporting a rise in the number of *Salmonella* Typhi clinical isolates with DCS, considered endemic in countries such as India,⁵⁷ but a lower incidence of strains showing high levels of ciprofloxacin resistance. A study conducted in the UK from 2001 to 2004 reported low incidence (1%–2%) of ciprofloxacin-resistant *Salmonella* Typhi isolates (MIC \geq 1 mg/L) in contrast to an increase in the number of strains with DCS (from 35% in 2001 to 47% in 2004).⁵⁸ Nonetheless, some cases reporting high levels of ciprofloxacin in *Salmonella* Typhi have already emerged.^{9,59}

Moreover, previous studies conducted in different serovars of *S. enterica* have already associated quinolone resistance with a decreased invasion profile. Wang *et al.*³² reported a reduction in the expression of SPI-1 genes, decreased ability to invade epithelial cells and low replication inside macrophages in *Salmonella* Typhimurium and *Salmonella* Choleraesuis resistant mutants. Similarly, studies conducted with *Salmonella* Typhimurium and *Salmonella* Enteritidis also showed an association between resistance, reduced virulence and impaired expression of SPI-1 genes in quinolone-resistant mutants.^{16,31} Consistently with these findings, our results showed that upon acquisition of resistance to quinolones, the two mutants showed a progressively reduced ability to invade epithelial cells and macrophages. Moreover, the survival rate inside macrophages was also diminished. Regulation of these virulence determinants has been well established in *Salmonella* Typhimurium³⁶ and demonstrated to be mostly homologous in *Salmonella* Typhi by Faucher *et al.*²⁶ They identified that SPI-1 genes were involved in the invasion process when infecting macrophages. Moreover, Bishop *et al.*⁶⁰ showed that *Salmonella* Typhi strains lacking *invA* were 1000-fold less able to invade epithelial cells. Thus, the results of our gene expression analysis agreed with the phenotype observed, since several SPI-1-related genes (*invA*, *hilA* and *hilD*) showed a gradual reduction in their expression levels.

In an attempt to find an explanation for this effect on virulence, several hypotheses should be considered. First, DNA supercoiling has been proposed to influence the transcription process.⁶¹ Thus, acquisition of mutations in DNA gyrase has been considered to impair regular supercoiling activity and hence modify gene expression patterns.³² Nonetheless, our results do not support this as our parental strain already had a mutation in *GyrA*, but no other QRDR mutation was acquired in the mutants. Second, impaired bacterial growth has been described for quinolone-resistant mutants of *Salmonella* Typhimurium and *Salmonella* Enteritidis.^{31,46,62} This reduced ability to grow may shut off dispensable bacterial functions such as virulence properties. Nonetheless, the three strains reported in this study did not show any difference in terms of bacterial growth. On the contrary, a recent study performed in *Salmonella* Typhi associated mutations in the QRDR of the target genes with fitness benefits, with the S83F mutation in *GyrA*, which leads to DCS, being the most advantageous.⁶³ The presence of this mutation in our strains may explain the unchanged growth observed. Third, high production levels of AcrAB/TolC have been suggested to be the reason for

impaired virulence.⁴⁶ As quorum-sensing signal molecules have been shown to be extruded by efflux pumps⁶⁴ and such molecules can activate virulence genes,⁶⁵ an impaired quorum-sensing homeostasis triggered by increased efflux activity may lead to impaired gene transcription. To strengthen this hypothesis, a study conducted by Bailey *et al.*⁶⁶ showed that high overexpression of *ramA* was accompanied by overexpression of efflux pumps as well as a decrease in the expression of virulence genes and impaired host–pathogen interactions. Taking into account these findings, we suggest that a similar situation could be happening in the present work triggered by the high overexpression of *marA* (>100-fold) in the Ty_c2 mutant.

Additional virulence genes were also studied in the present work. On the one hand, *Salmonella* Typhimurium and *Salmonella* Typhi have been demonstrated to share a number of pathogenesis determinants, such as the bacterial flagellum. Mutations in the flagellar regulatory genes *flhDC* and *fliA* have been associated with severely decreased entry into eukaryotic cells and reduced cytotoxicity in macrophages.⁶⁷ Moreover, uptake and survival defects in macrophages have also been seen in *fliC* and *flhCD* mutants as reported by Sabbagh *et al.*⁶⁸ A correlation between reduced motility, diminished *fliC* levels and impaired invasion of epithelial cells was seen in our mutants. On the other hand, *Salmonella* Typhi-specific virulence genes have also been identified. Among these factors, the most relevant are the genes located in the *Salmonella* Typhi-specific pathogenicity island SPI-7.^{25,69} The Vi capsular polysaccharide is encoded by the *viaB* locus, from SPI-7, and has been reported to be necessary for *Salmonella* Typhi survival in macrophages, serum resistance and systemic dissemination.^{25,70} An important repression of the *tviA* gene was reported in the MDR mutant Ty_c2. Thus, it seems reasonable that repression of these virulence factors (*FliC* and *TviA*) has also contributed to the impaired virulence phenotype observed in the two mutants, concerning invasion of epithelial cells as well as uptake by and survival inside human macrophages.

It has been previously shown that *TviA* represses important virulence factors including genes encoding flagella and T3SS-1 through the *flhDC*–*fliZ*–*hilD*–*hilA* axis.^{60,71} In our study, transcription of *tviA* was only repressed in Ty_c2 whereas the expression levels of *fliC* and the SPI-1 encoded genes, were already decreased in the Ty_c1 mutant. Thus, these results support the existence of a *tviA*-independent regulation of flagellin expression, in accordance with the representative number of regulators governing flagella expression reported in other *Salmonella* serovars. However, we should not underestimate the role of *TviA* in this pattern of down-regulation of *fliC* and the SPI-1 encoded genes in the Ty_c2 mutant to link the changes observed in these sets of genes.

In the last part of this work, we evaluated the ability of the mutants to trigger an immunogenic response when infecting human macrophages. The flagellin protein *FliC* has been reported to activate TLR-5 inside macrophages, thus decreasing the signalling pathway of NF- κ B, which elicits cytokine production (TNF- α and IL-1 β). Our results showed that Ty_c1 and Ty_c2 triggered a poor innate immune response as reflected by a decrease in the production of these two proinflammatory cytokines as well as in the nuclear transcriptional regulator NF- κ B. This attenuated immunogenic profile is most likely linked to reduced levels of *FliC*, as described previously.^{72,73} Moreover, we attribute the more noticeable reduction observed for IL-1 β (>2-fold reduction

1045 compared with TNF- α) to the additive effects of the previously reported effects and alternative TLR-independent secretion pathway of IL-1 β through caspase-1 activation, which is also stimulated by flagellin.³⁰

1050 Conclusions

The results obtained in the present study provide evidence, for the first time in *Salmonella* Typhi, of a link between the MDR phenotype and increased levels of AcrAB/TolC caused by the hyperproduction of its regulator MarA. Moreover, we determined that when *Salmonella* Typhi acquires resistance to quinolones and other unrelated antibiotics it becomes less virulent. Concerning the hypothesis to explain this impaired virulence and host-pathogen interactions, our results rule out previously reported explanations such as acquisition of QRDR mutations or reduced growth. Our findings, particularly those obtained from Ty_c2, reinforce the idea that high efflux pump activity may affect quorum-sensing homeostasis, eventually leading to changes in virulence gene expression. Although more research is needed to clarify this phenomenon and consider also the situation reported for Ty_c1, our results contribute to the complex understanding of quinolone resistance and virulence. On the other hand, with the aim to explain the lower prevalence of ciprofloxacin-resistant *Salmonella* Typhi causing illness in the clinical setting, we hypothesize that, in comparison with susceptible bacteria, when quinolone-resistant *Salmonella* Typhi reaches the gut only a small proportion of the bacterial load is able to cross the mucosa and survive inside the host due to: (i) poor invasion of enterocytes due to a mild activation of the invasion machinery; (ii) reduced phagocytosis by dendritic cells and macrophages; and (iii) reduced survival rate inside macrophages. This situation, together with decreased activation of the innate immune response, may result in impaired dissemination of the resistant pathogen. However, the emergence of isolates able to overcome antibiotic pressure needs to be taken into account and efforts should be made to fight against them. In order to do so, the use of this attenuated strain able to trigger a poor, but still immunogenic, response as a potential vaccine needs to be considered and further explored.

1085 Acknowledgements

We acknowledge Thomas Most for his advice on the macrophage infection procedure.

1090 Funding

This study was supported by grant 2014SGR0653 from the Departament de Universitats, Recerca i Societat de la Informació de la Generalitat de Catalunya, by the Ministerio de Economía y Competitividad, Instituto de Salud Carlos III, co-financed by European Regional Development Fund (ERDF) 'A Way to Achieve Europe,' the Spanish Network for Research in Infectious Diseases (REIPI RD12/0015), EUROSALUD (EUS2008-03616) and FIS 11/02024. We also thank the Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica for the travel grant awarded to C. B.-D. that allowed her to perform the work at Columbia University of the City of New York. A. F. is sponsored by the Barcelona Institute for Global Health (ISGlobal).

Transparency declarations

None to declare.

Supplementary data

Tables S1 and S2 are available as Supplementary data at JAC Online (<http://jac.oxfordjournals.org/>).

References

- 1 WHO. Typhoid vaccines. *Wkly Epidemiol Rec* 2008; **83**: 49–60. 1115
- 2 De Jong HK, Parry CM, van der Poll T *et al.* Host-pathogen interaction in invasive salmonellosis. *PLoS Pathog* 2012; **8**: e1002933.
- 3 Monack DM, Mueller A, Falkow S. Persistent bacterial infections: the interface of the pathogen and the host immune system. *Nat Rev Microbiol* 2004; **2**: 747–65. 1120
- 4 Bhan MK, Bahl R, Bhatnagar S. Typhoid and paratyphoid fever. *Lancet* 2005; **366**: 749–62. 1125
- 5 Butler T. Treatment of typhoid fever in the 21st century: promises and shortcomings. *Clin Microbiol Infect* 2011; **17**: 959–63.
- 6 Chau TT, Campbell JI, Galindo CM *et al.* Antimicrobial drug resistance of *Salmonella enterica* serovar Typhi in Asia and molecular mechanism of reduced susceptibility to the fluoroquinolones. *Antimicrob Agents Chemother* 2007; **51**: 4315–23. 1130
- 7 Threlfall EJ, Ward LR. Decreased susceptibility to ciprofloxacin in *Salmonella enterica* serotype Typhi, United Kingdom. *Emerg Infect Dis* 2001; **7**: 448–50. 1135
- 8 Dutta S, Sur D, Manna B *et al.* Emergence of highly fluoroquinolone-resistant *Salmonella enterica* serovar Typhi in a community-based fever surveillance from Kolkata, India. *Int J Antimicrob Agents* 2008; **31**: 385–7.
- 9 Lee C-J, Su L-H, Huang Y-C *et al.* First isolation of ciprofloxacin-resistant *Salmonella enterica* serovar Typhi in Taiwan. *J Microbiol Immunol Infect* 2013; **46**: 469–73. 1140
- 10 Chiou C-S, Lauderdale T-L, Phung DC *et al.* Antimicrobial resistance in *Salmonella enterica* serovar Typhi isolates from Bangladesh, Indonesia, Taiwan, and Vietnam. *Antimicrob Agents Chemother* 2014; **58**: 6501–7. 1145
- 11 Parry C, Tran TH, Dougan G *et al.* Typhoid fever. *N Engl J Med* 2002; **347**: 1770–82.
- 12 Renuka K, Kapil A, Kabra SK *et al.* Reduced susceptibility to ciprofloxacin and *gyrA* gene mutation in North Indian strains of *Salmonella enterica* serotype Typhi and serotype Paratyphi A. *Microb Drug Resist* 2004; **10**: 146–53. 1150
- 13 Fàbrega A, Madurga S, Giralte E *et al.* Mechanism of action of and resistance to quinolones. *Microb Biotechnol* 2009; **2**: 40–61.
- 14 Giraud E, Cloeckaert A, Kerboeuf D *et al.* Evidence for active efflux as the primary mechanism of resistance to ciprofloxacin in *Salmonella enterica* serovar Typhimurium. *Antimicrob Agents Chemother* 2000; **44**: 1223–8. 1155
- 15 Cloeckaert AC, Ancla EC. Mechanisms of quinolone resistance in *Salmonella*. *Vet Res* 2001; **32**: 291–300.
- 16 O'Regan E, Quinn T, Pagès J-M *et al.* Multiple regulatory pathways associated with high-level ciprofloxacin and multidrug resistance in *Salmonella enterica* serovar Enteritidis: involvement of RamA and other global regulators. *Antimicrob Agents Chemother* 2009; **53**: 1080–7.
- 17 Abouzeed YM, Baucheron S, Cloeckaert A. *ramR* mutations involved in efflux-mediated multidrug resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob Agents Chemother* 2008; **52**: 2428–34. 1160
- 18 Kehrenberg C, Cloeckaert A, Klein G *et al.* Decreased fluoroquinolone susceptibility in mutants of *Salmonella* serovars other than Typhimurium:

detection of novel mutations involved in modulated expression of *ramA* and *soxS*. *J Antimicrob Chemother* 2009; **64**: 1175–80.

19 Koutsolioutsou A, Martins EA, White DG *et al*. A *soxRS*-constitutive mutation contributing to antibiotic resistance in a clinical isolate of *Salmonella enterica* (serovar Typhimurium). *Antimicrob Agents Chemother* 2001; **45**: 38–43.

20 Zheng J, Tian F, Cui S *et al*. Differential gene expression by *RamA* in ciprofloxacin-resistant *Salmonella* Typhimurium. *PLoS One* 2011; **6**: e22161.

21 Oethinger M, Podglajen I, Kern WV *et al*. Overexpression of the *marA* or *soxS* regulatory gene in clinical topoisomerase mutants of *Escherichia coli*. *Antimicrob Agents Chemother* 1998; **42**: 2089–94.

22 Alekshun MN, Levy SB. Regulation of chromosomally mediated multiple antibiotic resistance: the *mar* regulon. *Antimicrob Agents Chemother* 1997; **41**: 2067–75.

23 Grkovic S, Brown MH, Skurray RA. Regulation of bacterial drug export systems. *Microbiol Mol Biol Rev* 2006; **66**: 671–701.

24 Delilhas N, Forst S. *micF*: an antisense RNA gene involved in response of *Escherichia coli* to global stress factors. *J Mol Biol* 2001; **313**: 1–12.

25 Daigle F. Typhi genes expressed during infection or involved in pathogenesis. *J Infect Dev Ctries* 2008; **2**: 431–7.

26 Faucher SP, Porwollik S, Dozois CM *et al*. Transcriptome of *Salmonella enterica* serovar Typhi within macrophages revealed through the selective capture of transcribed sequences. *Proc Natl Acad Sci USA* 2006; **103**: 1906–11.

27 Kaur J, Jain SK. Role of antigens and virulence factors of *Salmonella enterica* serovar Typhi in its pathogenesis. *Microbiol Res* 2012; **167**: 199–210.

28 Winter SE, Winter MG, Godinez I *et al*. A rapid change in virulence gene expression during the transition from the intestinal lumen into tissue promotes systemic dissemination of *Salmonella*. *PLoS Pathog* 2010; **6**: e1001060.

29 Hayashi F, Smith KD, Ozinsky A *et al*. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* 2001; **410**: 1099–103.

30 Franchi L, Amer A, Body-Malapel M *et al*. Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1 β in *Salmonella*-infected macrophages. *Nat Immunol* 2006; **7**: 576–82.

31 Fàbrega A, du Merle L, Le Bouguéne C *et al*. Repression of invasion genes and decreased invasion in a high-level fluoroquinolone-resistant *Salmonella* Typhimurium mutant. *PLoS One* 2009; **4**: e8029.

32 Wang Y-P, Li L, Shen J-Z *et al*. Quinolone-resistance in *Salmonella* is associated with decreased mRNA expression of virulence genes *invA* and *avrA*, growth and intracellular invasion and survival. *Vet Microbiol* 2009; **133**: 328–34.

33 Ballesté-Delpierre C, Solé M, Domènech Ò *et al*. Molecular study of quinolone resistance mechanisms and clonal relationship of *Salmonella enterica* clinical isolates. *Int J Antimicrob Agents* 2013; **43**: 121–5.

34 Koblansky AA, Jankovic D, Oh H *et al*. Recognition of profilin by Toll-like receptor 12 is critical for host resistance to *Toxoplasma gondii*. *Immunity* 2013; **38**: 119–30.

35 Cummings L, Wilkerson WD, Bergsbaken T *et al*. In vivo, *flhC* expression by *Salmonella enterica* serovar Typhimurium is heterogeneous, regulated by ClpX, and anatomically restricted. *Mol Microbiol* 2006; **61**: 795–809.

36 Fàbrega A, Vila J. *Salmonella enterica* serovar Typhimurium skills to succeed in the host: virulence and regulation. *Clin Microbiol Rev* 2013; **26**: 308–41.

37 Capoor MR, Nair D, Walia NS *et al*. Molecular analysis of high-level ciprofloxacin resistance in *Salmonella enterica* serovar Typhi and *S. Paratyphi A*: need to expand the QRDR region? *Epidemiol Infect* 2009; **137**: 871–8.

38 Magiorakos A, Srinivasan A, Carey RB *et al*. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2011; **18**: 268–81.

39 Eaves DJ, Ricci V, Piddock LJV. Expression of *acrB*, *acrF*, *acrD*, *marA*, and *soxS* in *Salmonella enterica* serovar Typhimurium: role in multiple antibiotic resistance. *Antimicrob Agents Chemother* 2004; **48**: 1145–50.

40 Chen S, Cui S, McDermott PF *et al*. Contribution of target gene mutations and efflux to decreased susceptibility of *Salmonella enterica* serovar Typhimurium to fluoroquinolones and other antimicrobials. *Antimicrob Agents Chemother* 2007; **51**: 535–42.

41 Martin RG, Rosner JL. Binding of purified multiple antibiotic-resistance repressor protein (MarR) to *mar* operator sequences. *Proc Natl Acad Sci USA* 1995; **92**: 5456–60.

42 Liu SL, Ezaki T, Miura H *et al*. Intact motility as a *Salmonella* Typhi invasion-related factor. *Infect Immun* 1988; **56**: 1967–73.

43 Kalai K, Chai LC, Thong KL. Variations in motility and biofilm formation of *Salmonella enterica* serovar Typhi. *Gut Pathog* 2014; **6**: 1–10.

44 Kutsukake K, Ohya Y, Iino T. Transcriptional analysis of the flagellar regulon of *Salmonella* Typhimurium. *J Bacteriol* 1990; **172**: 741–7.

45 Smith KD, Andersen-Nissen E, Hayashi F *et al*. Toll-like receptor 5 recognizes a conserved site on flagellin required for protofilament formation and bacterial motility. *Nat Immunol* 2003; **4**: 1247–53.

46 O'Regan E, Quinn T, Frye JG *et al*. Fitness costs and stability of a high-level ciprofloxacin resistance phenotype in *Salmonella enterica* serotype Enteritidis: reduced infectivity associated with decreased expression of *Salmonella* pathogenicity island 1 genes. *Antimicrob Agents Chemother* 2010; **54**: 367–74.

47 Boddicker JD, Knosp BM, Jones BD. Transcription of the *Salmonella* invasion gene activator, *hilA*, requires HilD activation in the absence of negative regulators. *J Bacteriol* 2003; **185**: 525–33.

48 Wu W, Wang H, Lu J *et al*. Genetic diversity of *Salmonella enterica* serovar Typhi and Paratyphi in Shenzhen, China from 2002 through 2007. *BMC Microbiol* 2010; **10**: 1–7.

49 Ma L, Siu LK, Lin J *et al*. Updated molecular epidemiology of carbapenem-non-susceptible *Escherichia coli* in Taiwan: first identification of KPC-2 or NDM-1-producing *E. coli* in Taiwan. *BMC Infect Dis* 2013; **13**: 1–8.

50 Chenia HY, Pillay B, Pillay D. Analysis of the mechanisms of fluoroquinolone resistance in urinary tract pathogens. *J Antimicrob Chemother* 2006; **58**: 1274–8.

51 Baucheron S, Monchaux I, Le Hello S *et al*. Lack of efflux mediated quinolone resistance in *Salmonella enterica* serovars Typhi and Paratyphi A. *Front Microbiol* 2014; **5**: 1–6.

52 Sulavik MC, Dazer M, Miller PF. The *Salmonella* Typhimurium *mar* locus: molecular and genetic analyses and assessment of its role in virulence. *J Bacteriol* 1997; **179**: 1857–66.

53 Fàbrega A, Soto SM, Ballesté-Delpierre C *et al*. Impact of quinolone-resistance acquisition on biofilm production and fitness in *Salmonella enterica*. *J Antimicrob Chemother* 2014; **69**: 1815–24.

54 Olliver A, Valle M, Chaslus-Dancla E *et al*. Role of an *acrR* mutation in multidrug resistance of in vitro-selected fluoroquinolone-resistant mutants of *Salmonella enterica* serovar Typhimurium. *FEMS Microbiol Lett* 2004; **238**: 267–72.

55 Kang HW, Woo GJ. Increase of multidrug efflux pump expression in fluoroquinolone-resistant *Salmonella* mutants induced by ciprofloxacin selective pressure. *Res Vet Sci* 2014; **97**: 182–6.

56 Viveiros M, Dupont M, Rodrigues L *et al*. Antibiotic stress, genetic response and altered permeability of *E. coli*. *PLoS One* 2007; **2**: e365.

- 1280 **57** Gaind R, Paglietti B, Murgia M *et al.* Molecular characterization of ciprofloxacin-resistant *Salmonella enterica* serovar Typhi and Paratyphi A causing enteric fever in India. *J Antimicrob Chemother* 2006; **58**: 1139–44.
- 58** Threlfall EJ, Day M, de Pinna E *et al.* Drug-resistant enteric fever in the UK. *Lancet* 2006; **367**: 1576.
- 59** Capoor MR, Nair D, Deb M *et al.* Enteric fever perspective in India: emergence of high-level ciprofloxacin resistance and rising MIC to cephalosporins. *J Med Microbiol* 2007; **56**: 1131–2.
- 1285 **60** Bishop A, House D, Perkins T *et al.* Interaction of *Salmonella enterica* serovar Typhi with cultured epithelial cells: roles of surface structures in adhesion and invasion. *Microbiology* 2008; **154**: 1914–26.
- 61** Dorman CJ, Corcoran CP. Bacterial DNA topology and infectious disease. *Nucleic Acids Res* 2009; **37**: 672–8.
- 1290 **62** Giraud E. Fitness cost of fluoroquinolone resistance in *Salmonella enterica* serovar Typhimurium. *J Med Microbiol* 2003; **52**: 697–703.
- 63** Baker S, Duy PT, Nga TVT *et al.* Fitness benefits in fluoroquinolone-resistant *Salmonella* Typhi in the absence of antimicrobial pressure. *Elife* 2013; **2**: e01229.
- 1295 **64** Pearson JP, Van Delden C, Iglewski BH. Active efflux and diffusion are involved in transport of *Pseudomonas aeruginosa* cell-to-cell signals. *J Bacteriol* 1999; **181**: 1203–10.
- 65** Sperandio V, Torres AG, Kaper JB. Quorum sensing *Escherichia coli* regulators B and C (QseBC): a novel two-component regulatory system involved in the regulation of flagella and motility by quorum sensing in *E. coli*. *Mol Microbiol* 2002; **43**: 809–21.
- 66** Bailey AM, Ivens A, Kingsley R *et al.* RamA, a member of the AraC/XylS family, influences both virulence and efflux in *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2010; **192**: 1607–16. 1335
- 67** Eichelberg K, Galán JE. The flagellar sigma factor FliA (σ^{28}) regulates the expression of *Salmonella* genes associated with the centisome 63 type III secretion system. *Infect Immun* 2000; **68**: 2735–43. 1340
- 68** Sabbagh SC, Lepage C, McClelland M *et al.* Selection of *Salmonella enterica* serovar Typhi genes involved during interaction with human macrophages by screening of a transposon mutant library. *PLoS One* 2012; **7**: e36643.
- 69** Seth-Smith HMB. SPI-7: *Salmonella*'s Vi-encoding pathogenicity island. *J Infect Dev Ctries* 2008; **2**: 267–71. 1345
- 70** Virlogeux I, Waxin H, Ecobichon C *et al.* Role of the *viaB* locus in synthesis, transport and expression of *Salmonella* Typhi Vi antigen. *Microbiology* 1995; **141**: 3039–47.
- 71** Eaves DJ, Randall L, Gray DT *et al.* Prevalence of mutations within the quinolone resistance-determining region of *gyrA*, *gyrB*, *parC*, and *parE* and association with antibiotic resistance in quinolone-resistant *Salmonella enterica*. *Antimicrob Agents Chemother* 2004; **48**: 4012–5. 1350
- 72** Wyant TL, Tanner MK, Sztein MB. *Salmonella* Typhi flagella are potent inducers of proinflammatory cytokine secretion by human monocytes. *Infect Immun* 1999; **67**: 3619. 1355
- 73** Finamore E, Vitiello M, D'Isanto M *et al.* Evidence for IL-6 promoter nuclear activation in U937 cells stimulated with *Salmonella enterica* serovar Typhimurium porins. *Eur Cytokine Netw* 2009; **20**: 140–7. 1360

1360

1365

1370

1375

1380

1385

1390