

# Quinoa (*Chenopodium quinoa* Willd), from Nutritional Value to Potential Health Benefits: An Integrative Review

Bastidas EG<sup>1</sup>, Roura R<sup>2</sup>, Rizzolo DAD<sup>3</sup>, Massanés T<sup>2</sup> and Gomis R<sup>1,3,5,\*</sup>

<sup>1</sup>Department of Endocrinology and Nutrition, Hospital Clínic, Barcelona, Spain

<sup>2</sup>Fundació Alicia, Barcelona, Spain

<sup>3</sup>Diabetes and Obesity Research Laboratory, IDIBAPS, Barcelona, Spain

<sup>4</sup>CIBER in Diabetes and Associated Metabolic Disorders (CIBERDEM), Barcelona, Spain

<sup>5</sup>University of Barcelona, Barcelona, Spain

\*Corresponding author: Gomis R, IDIBAPS, C/Rosselló, 149-153, 08036 - Barcelona, Spain, Tel: 34933129411; E-mail: [ramon.gomis@idibaps.org](mailto:ramon.gomis@idibaps.org)

Received date: Mar 02, 2016; Accepted date: Apr 21, 2016; Published date: Apr 26, 2016

Copyright: © 2016 Bastidas EG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

*Chenopodium quinoa* Willd, known as quinoa, has been cultivated and consumed by humans for the last 5,000-7,000 years. Quinoa was important to pre-Columbian Andean civilizations, as the Incas considered it a gift from their gods. Quinoa has potential health benefits and exceptional nutritional value: a high concentration of protein (all essential amino acids highly bioavailable), unsaturated fatty acids, a low glycemic index; vitamins, minerals and other beneficial compounds; it is also gluten-free; furthermore, quinoa is a sustainable food, as plants exhibit a carbon and water food print that is between 30 and 60 times lower than that of beef. Quinoa is easy to cook, has versatility in preparation, and could be cultivated in different environments. For these reasons, quinoa, previously considered a food of low social prestige, is now the focus of attention of many countries worldwide. However, few studies exist on quinoa or quinoa compounds, *in vitro*, *in vivo* and clinical trials, for assessing its potential clinical applications supported by strong scientific evidence; thus, there is a need for well-designed clinical trials and increased scientific research in this field.

**Keywords:** Quinoa; Glycemic index; Gluten free; Essential amino acids; Healthy food

## Abbreviations

20E: 20-Hydroxyecdysone; ACE: Angiotensin I Converting Enzyme; Ca: Calcium; Cx43: Connexin 43; DPPH: Ferric Thiocyanate and 1,1-Diphenyl-2-Picrylhydrazyl; Fe: Iron; GI: Glycemic Index; HbA1c: Hemoglobin A1c; HF: High Fat Diet; HF20E: High Fat Diet+20-Hydroxyecdysone; HFQ: High Fat Diet+Quinoa; IgE: Immunoglobulin E; K: Potassium; kcals: Kilocalories; Mg: Magnesium; MTT: 3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide; NASA: National Aeronautics and Space Administration; P: Phosphorus; PAI-1: Plasminogen Activator Inhibitor-1; PEPCK: Phosphoenolpyruvate Carboxykinase

## Introduction

Quinoa is a grain with exceptional nutritional value; it has been cultivated for the last 5,000-7,000 years in the Andean region of Bolivia and Peru. 2013 was declared by the United Nations as the International Year of Quinoa as recognition of its significant potential [1-4].

Quinoa has high concentrations of protein, all essential amino acids, unsaturated fatty acids, and a low glycemic index (GI); it also contains vitamins, minerals and other beneficial compounds, and is gluten-free by nature. Quinoa is easy to cook and has versatility in preparation [1-4].

In this article, we present an extensive review of quinoa, starting with its history, analyzing its botanical characteristics, and covering its

nutritional profile. In addition, we offer a critical comparison with other grains and describe quinoa culinary applications and uses. We also provide an updated review of the scientific information available on quinoa or quinoa compound properties through *in vivo*, *in vitro* and clinical studies, in order to evaluate its potential translational applications.

## History

Quinoa has been cultivated for thousands of years in the Andean region of Bolivia and Peru [5,6]. It is known by different local names, or simply quinoa or quinoa (quinoa is in Quechua) [4]. This plant was called “the mother grain” by the Incas and was considered a gift from their gods, used even for treating medical issues. Traditionally, quinoa seeds were roasted and cooked, added to soups, used as a cereal, and even fermented into beer or chichi (traditional drink of the Andes) [4,7,8].

After the Spanish conquest of South America, the colonists looked down on quinoa as a peasant food or food of the Indians; consequently, quinoa has been considered a food of low social prestige. In addition, the Catholic Church actively suppressed its cultivation after discovering that quinoa was used as a sacred drink (Mudai) during indigenous religious ceremonies. Thus, quinoa remained only where Europeans could not reach and replace it with other grains [4].

## Botanical description and cultivation

The botanical name of quinoa is *Chenopodium quinoa* Willd and belongs to the Goosefoot family “Chenopodiaceae”, which includes: Swiss chard (*Beta sp.*), spinach (*Spinacia oleracea*) and Lamb’s quarters

(*Chenopodium album*). Quinoa is a dicot plant that can grow from 1 to 3m high; it is considered a pseudo-cereal, not a true grain but rather a fruit. The seeds are round and flat, about 1.5-4.0 mm in diameter and their color varies from white to grey and black, with tones of yellow,

rose, red, purple and violet (Figure 1). Quinoa has demonstrated a strong tolerance to salty, acid or alkaline soils, in both cold (-5°C) or hot climates (up to 35°C) [4,5].



**Figure 1:** Quinoa plant, Quinoa is a pseudo-cereal. Its seeds can be different colors; its leaves are shaped like a goose's foot.

Quinoa is distributed worldwide and includes 250 varieties. Its classification is based on the color of the plant and fruits, or on plant morphology [4,5]. This grain has been authorized to be sown in Europe, North America, Asia and Africa. In Europe the project “Quinoa: a multipurpose crop for the European Community” was approved in 1993 [4].

### Nutritional profile

The effects of globalization and urbanization have influenced dietary patterns and lifestyle behaviors among population groups throughout the world. Traditional food patterns rich in complex carbohydrates, micronutrients, fiber and phytochemicals are being replaced with diets high in animal fats and refined carbohydrates and oils, a situation that has made a direct impact on the prevalence of certain chronic diseases [9]. For this reason, many researchers devote their efforts to analyzing food or food components that may prove to be healthy for human consumption [5]; one example is the work of the

HEALTHGRAIN Consortium, which, of note, has included quinoa in its list of healthy grains [10].

Quinoa is considered one of the best vegetal protein sources, as its protein levels are similar to those found in milk and higher than those present in cereals such as wheat, rice and maize. Quinoa also has been used by the National Aeronautics and Space Administration (NASA) due to its versatility in meeting the needs of humans during space missions [3,4,7,11,12].

### Protein

Proteins and amino acids are major biological macromolecules that serve as structural constituents and as catalysts for enzymatic reactions, energy sources and protein synthesis in the body [13,14].

The protein biological value measures the proportion of protein absorbed from a food which then becomes incorporated into the proteins of the body. Quinoa has a high biological value (73%), similar



to that of beef (74%), and higher than those of white rice (56%), wheat (49%) and corn (36%). Quinoa also contains all ten essential amino acids, and its protein content ranges from 12.9 to 16.5% [4,15-17]. Of primary interest is the high lys value, an essential amino acid that is deficient in many grains [16,18]. Quinoa is also high in the essential amino acid met, which is deficient in many legumes [19,20]. According to the daily recommended amounts of amino acids indicated by the Food and Agriculture Organization (FAO) of the United Nations and by the World Health Organization (WHO), quinoa fulfills the amino acid requirements for adults: 180% of histidine (his), 274% of isoleucine (ile), 338% of lysine (lys), 212% of methionine+cysteine (met+cys), 320% of phenylalanine+tyrosine (phe+tyr), 331% of threonine (thr), 228% of tryptophan (trp) and 323% of valine (val) (20). For these reasons, quinoa could represent a valuable source of nutrition, especially for infants and children, and may be used in nutritive foods and beverages [21].

Another feature of note is the high sustainability of plant food consumption due to their low carbon, water and ecological food prints. In this sense, quinoa is an excellent protein source such as beef, but it has a carbon and water food print that is between 30 and 60 times lower in value [22].

## Carbohydrates

Carbohydrates are organic compounds comprised of carbon, hydrogen, and oxygen. Carbohydrates act as signaling molecules, energy sources and structural components [14].

Starch, as a carbohydrate, provides the major source of physiological energy in the human diet. The content of starch in quinoa ranges from 58.1% to 64.2% of dry matter, of which 11% is amylose. Moreover, quinoa has a high content of D-xylose and maltose and a low content of glucose and fructose.

**100 g of quinoa contains:** Glucose 1.70 mg, fructose 0.20 mg, saccharose 2.90 mg and maltose 1.40 mg [4,5,17]. In addition, there are studies that suggest that quinoa polysaccharides have antioxidant properties [23].

## Lipids

Lipids are concentrated sources of energy as well as structural components of cell membranes that the body uses for performing a variety of normal functions [2,4]. A higher intake of vegetables is associated with a reduced risk of type 2 diabetes, due to their high unsaturated fat content, which is associated with lower inflammation [24,25].

The quality of lipids is very important. For example, it is known that omega-6 stimulates inflammatory activity in the body, while omega-3 performs anti-inflammatory functions. A lower ratio of omega-6: omega-3 fatty acids are more desirable for reducing the risk of cardiovascular disease, cancer, and inflammatory and autoimmune diseases [2,4,26].

The total lipid content of quinoa is 14.5%, with approximately 70%-89.4% being unsaturated (38.9%-57% of linoleic acid, 24.0%-27.7% of oleic acid and 4% of  $\alpha$ -linolenic acid). The unsaturated fatty acid content is protected by vitamin E in this plant. The ratio between omega-6 and omega-3 in quinoa is about 6:1 [2,4].

## Fiber

Dietary fiber is the indigestible portion of food derived from plants and has two main components: soluble and insoluble. Soluble fiber dissolves in water, is readily fermented in the colon into gases and physiologically active products, and has prebiotic properties. Insoluble fiber, which does not dissolve in water, is either metabolically inert and provides bulking mass, or it can be prebiotic and metabolically ferment in the large intestine. Bulking fibers absorb water, easing defecation [3,27].

Greater consumption of fiber-rich whole grains is associated with a lower risk of type 2 diabetes [24] and cardiovascular disease [28]. Quinoa is an excellent source of dietary fiber, comprising about 2.6%-10% of the total weight of the grain; about 78% of its fiber content is insoluble and 22% soluble [3,5,17,27,29].

## Glycemic index (GI)

The GI is a ranking of carbohydrates on a scale from 0 to 100 according to their impact on blood sugar levels during the 2 h following consumption. Low GI (<55) foods produce gradual rises in blood sugar and insulin levels. Low GI diets have been shown to improve glucose and lipid levels and weight control because they help control appetite [30,31]. Low GI diets also reduce insulin resistance and the risk of cardiovascular diseases, diabetes and some cancers [9,24,32]. It is reported that fasting insulin is lower in individuals with higher dietary fiber intakes [33] and that the ingestion of complex carbohydrates promotes longevity [14]; and also that a high GI diet increases the levels of inflammation biomarkers. In patients with celiac disease, a reduced inflammatory state could provide certain protective mechanisms, thus, following a low GI diet could meliorate this disease [34].

The GI for quinoa ranges 35-53 depending on the cooking time; 150 g of quinoa, cooked, refrigerated and reheated in the microwave for 1.5 min has a GI of 53. Therefore, even when more or less "overcooked", quinoa maintains a low GI [32].

In Table 1 we present a summary of the nutritional profile of cooked quinoa, in comparison with common cereals (also cooked).

Food	Energy (kcal)	Proteins (g)	Essential amino acids (number)	Carbohydrates (g)	Lipids (g)	Unsaturated fatty acids (g)	Fiber (g)	GI
Quinoa	120	4.4	42653	21.3	1.92	1.61	2.8	35-53
Rice (white, medium grain)	130	2.38	42623	28.59	0.21	0.12	0.3	75-89
Wheat (soft, white)	113	3.6	42653	25.12	0.66	0.35	4.23	48
Corn (sweet, yellow)	96	3.41	42623	20.98	1.5	0.98	2.4	60

**Table 1:** General nutritional profile of quinoa, we observe the overall content of quinoa in comparison with that of other foods [35,36].

\*Portion size is 100 g of each cooked food; as a norm, cooked food increases its size 3 times. kcal: Kilocalories, GI: Glycemic index.

## Vitamins

Vitamins are compounds essential for the health of humans. Quinoa has many vitamins, with 100 g of this grain containing: 0.4 mg of thiamine, 78.1 mg of folic acid, 1.4 mg of vitamin C, 0.20 mg of vitamin B6, and 0.61 mg of pantothenic acid [4]. Its vitamin E content ranges 37.49-59.82 µg/g. Tocopherol isoforms have also been detected in this seed: γ-tocopherol (47-53 µg/g), α-tocopherol (17-26 µg/g), and β- and δ-tocopherol (<5 µm/g) [2,5].

## Minerals

Dietary minerals are essential chemical elements that play a role in regulating electrolyte balance, glucose homeostasis, the transmission of nerve impulses and enzyme cofactors in the body [37].

Calcium, magnesium and potassium in quinoa are found in sufficient quantities and bioavailable forms necessary for maintaining a balanced human diet. Quinoa contains: 874 mg/kg of calcium (Ca) [4,5,38], iron (Fe) 948.5 mg/kg, phosphorus (P) 2735.0 mg/kg-4543.3 mg/kg, potassium (K) 9562.2 mg/kg and magnesium (Mg) 1901.5 mg/kg [3,5,39].

## Betaine

Betaine is an essential osmolyte source of methyl groups and finds its source either in diet or in the oxidation of choline. Its metabolism methylates homocysteine into methionine, also producing N,N-dimethylglycine. Mammals use betaine to assist in cell volume regulation, and as a methyl donor for the remethylation of homocysteine into methionine [27,40-42].

Hyperhomocysteinemia makes a person more prone to endothelial cell injury by activating inflammation pathways and is thus a possible risk factor for coronary artery disease. For this reason, betaine consumption could prove beneficial, as it converts homocysteine into methionine. Nevertheless, urinary excretion of betaine is minimal, even following a large betaine dose [43,44]. It is generally agreed that most people consume between 100 mg/day and 300 mg/day [5,41,45].

Betaine content of foods: cooked white rice 0 µg/g; cooked oatmeal 139 µg/g; cooked wheat spaghetti 287 µg/g; cooked quinoa approximately 1310 µg/g-2000 µg/g [45].

## Polyphenols

Polyphenols are natural organic chemicals with large multiples of phenol structural units. They are commonly found in plant foods and represent the most abundant antioxidants found in the human diet [46].

Quinoa presents at least 23 phenolic compounds. The total phenolic content (mg/kg quinoa) is 466.99, 634.66 and 682.05 for white, red and black quinoa, respectively. The most abundant phenols are ferulic acid and quercetin [1,6,47]. Quinoa contains more phenols than whole cereals, including wheat, barley, millet, rice and buckwheat [11].

## Isoflavones

Isoflavones are organic compounds that have a role in influencing sex hormone metabolism and biological activity through intracellular enzymes, protein synthesis, growth factor actions, malignant cell proliferations, differentiation and angiogenesis [48]. In particular, quinoa contains the isoflavones daidzein and genistein; these hormones could be recognized by human estrogen receptors and act as antagonists of vessel contraction and reduce arterial resistance [4].

## Carotenoids

Carotenoids are organic pigments found in the chloroplasts and chromoplasts of plants. The function of carotenoids in plants is to protect chlorophyll from photodamage. In humans, they exhibit antioxidant activity [27,49].

The total carotenoid content of white, red and black quinoa seeds are 11.87, 14.97 and 17.61 µg/g, respectively [50,51]. The β-carotene content of quinoa leaves ranges from 4.3 µg/1 g-19.5 µg/1 g [38].

## Gluten

Gluten is one of the most abundant components of food (including grains) [52]. Gluten is a composite of the proteins gliadin and glutenin [51].

Celiac disease is an immune-mediated reaction to gluten, characterized by an insufficient T- cell-mediated immune response that causes inflammatory injury to the small intestine [17,52]. Celiac disease patients must consume food with gluten substitutes or gluten-free grains [17,34,51,52].

Several gluten-free foodstuffs contain more salt and fat (saturated fat) but fewer minerals and vitamins than their gluten-containing counterparts. For this reason, quinoa is an excellent gluten-free grain, with a high content of vitamins and minerals that make it a potentially essential part of any healthy, gluten-free diet [7,17,34,53]; also quinoa complies with the Codex Alimentarius nomenclature of gluten-free products (gluten content <20 mg/kg) [51].

In Table 2, we present a summary of the principal quinoa compounds.

Vitamins		Minerals		Polyphenols	
Thiamine mg/100 g	0.4	Fe mg/kg	948.5	Total Phenolic Content mg/kg	467-682
Folic acid mg/100 g	78.1	Ca mg/kg	874	Ferulic acid mg/kg	37.5-58.4
Vitamin C mg/100 g	1.4	P mg/kg	2735	Quercetin mg/kg	5.3-13
B6 mg/100 g	0.2	K mg/kg	9562	Isoflavones	

Pantothenic acid mg/100 g	0.61	Mg mg/kg	1903	Daidzein mg/100 g	0.78-2.05
Vitamin E µg/1 g	37.49-59.82			Carotenoids	
Total carotenoids µg/g					11.87-17.61

**Table 2:** Summary of quinoa compounds. Quinoa contains several vitamins, minerals and antioxidant molecules. Fe: Iron, Ca: Calcium, P: Phosphorus, K: Potassium, Mg: Magnesium.

## Anti-nutritional factors

Saponins are natural detergents made of glycosylated secondary metabolites that are utilized by the plant as a predator repellent [54].

Quinoa contains saponins with a bitter taste, which could cause gastric irritation; its content varies between 0.1% and 5%; quinoa could even be classified according to its free saponin content, into either sweet (<0.11%) or bitter quinoa (>0.11%). In addition, there are studies of methods for saponin removal without any significant modification of nutritional value, such as thoroughly washing it in cold water [55].

On the other hand, isolated saponins could also have certain health benefits, such as anti-inflammatory activity [4,5,56-58].

Phytic acid is a saturated cyclic acid and is the principal storage form of phosphorus in many plant tissues. If the content of phytic acid is high in the food, this acid binds minerals, thereby rendering them unavailable for metabolism [27,59]. But, if the phytic acid content is low, it could act as an antioxidant: chelates various metals (for example, suppresses damaging Fe-catalysed redox reactions), inhibits xanthine oxidase, suppresses oxidant damage to the intestinal epithelium, and interferes with the formation of ADP-Fe-oxygen complexes that initiate lipid peroxidation; prevents the formation of carcinogens and blocks the interaction of carcinogens with cells; controls cell division and reduces cell proliferation rate; increases the immune response by enhancing the activity of natural killer cells; may be involved in cellular and nuclear signalling pathways; important source of P; inhibitor for renal stone development; hypoglycaemic (for example, by chelating Ca, an  $\alpha$ -amylase cofactor) and cholesterol-lowering (by binding Zn and decreasing Zn:Cu ratio) effects; affects the metabolic and detoxification capacity of the liver; reduces blood glucose and lipid, and hepatic lipid levels; inhibits calcification of cardiovascular system (lower level of Ca in aorta); prevention of platelet aggregation; high affinity for hydroxyapatite; adsorption onto Ca-based crystals; *in vitro* effect on gene expression through chromatin remodelling [27].

The general content of phytic acid in quinoa is low and ranges from 10.5 mg to 13.5 mg, in comparison with corn that contains 720 mg, wheat 390 mg and rice 60 mg [4].

Protease inhibitors are proteins that form very stable complexes with proteolytic enzymes in the body. The content of protease inhibitors in quinoa seeds is <50 ppm, which is very low and does not pose any serious concerns [4].

## Allergenicity

Asao and others observed that quinoa consumption did not show a positive reaction against wheat protein antibodies and suggested that quinoa could be ingested by people who have allergies against wheat proteins [11]. However, Astier and others reported a case of anaphylaxis to quinoa in France: a 52 year old man developed a systemic reaction consisting of dysphagia, dysphonia, urticaria and

angioedema after the ingestion of quinoa with fish and bread. They tested samples of the ingested food, and only quinoa showed Immunoglobulin E (IgE) reactivity in the patient's serum [60].

## Current status in experimental and clinical research

As we have seen above, quinoa has outstanding potential for human nutrition. Therefore, in this section we will provide an updated review of scientific information available on quinoa or quinoa compounds in different models.

### *In vitro*

**Betaine:** Lee and others tested a betaine pretreatment (2 mM) in H2.35 mouse hepatocyte cells. They observed an increase in cytochrome C oxidase activity and improved mitochondrial respiration (mitochondrial dysfunction is a commonly found in many pathological conditions and ageing) [40].

**Polyphenols:** Gawlik-Dziki and others cultured AT-2 and MAT-LyLu prostate cancer cells with quinoa leaf extract (rich in ferulic, sinapinic and gallic acid). They observed that cell proliferation was blocked, the expression of Connexin 43 (Cx43) was decreased, and lipid oxidation was prevented, inhibiting lipoxygenase activity. They also concluded that 80% of phenolics are available *in vitro* [61].

**Quinoa flour:** Extracts from quinoa flour and from that of whole cereals (wheat, barley, millet, rice, buckwheat) were used to evaluate antioxidant potential. The ferric thiocyanate and 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity methods were used; also, the Angiotensin I Converting Enzyme (ACE) inhibitory activity was measured. Quinoa presented higher antioxidant potential and higher ACE inhibition activity (23.3%) in comparison with the same whole cereals. These results suggested that quinoa possesses potent free radical-scavenging compounds [11].

### *In vivo*

**Saponins:** In mice, Verza and others reported that no lethality or local toxicity (local swelling, loss of hair, and piloerection) was observed after administering 300 µg of quinoa saponins [62].

**20-hydroxyecdysone:** Ecdysteroids are steroid hormones which have been implicated in plant repellence against insects, and in the control of the molting and reproduction processes of arthropods. The main phytoecdysteroid found in plants is 20-hydroxyecdysone (20E). Quinoa contains about 1.9% of 20E. These ecdysteroids have anabolic properties and help in controlling blood glucose levels. Graf and others tested an extract from quinoa seeds in obese mice and observed that fasting blood glucose levels were significantly lower [63].

Moreover, Foucault and others conducted a study on six-week-old male C57BL/6J mice. Mice were fed with a high-fat diet (HF) *ad libitum*, for 3 weeks. The high-fat diet group was supplemented or not

with either quinoa powder (HFQ) or pure 20E (HF20E). The epididymal adipose tissue following treatment was significantly lower in the HFQ and in the HF20E vs. HF group. The HFQ showed a trend of reduced lipid storage capacity in the adipose tissue, increased LPL and PPAR- $\gamma$  mRNA levels, and reduced expression of Phosphoenolpyruvate carboxykinase (PEPCK) and Plasminogen activator inhibitor-1 (PAI-1) in the adipose tissue, as well as decreased adipocyte hypertrophy [64,65].

Extract from seeds of quinoa: Meneguetti and others analyzed the effects of an extract from quinoa seeds in male Wistar rats assigned randomly to different groups: (1) a sedentary supplemented group, which received 2,000 mg/kg of the extract; (2) a sedentary control group, non-supplemented; (3) an exercise supplemented group with 2,000 mg/kg of the extract; and (4) an exercise control group non-supplemented. Follow-up was performed for 30 days. They observed beneficial effects in both supplemented groups (sedentary and exercise), as these groups showed a tendency of decreased food intake, body weight, fat deposition, and blood triglycerides (TG) levels [15].

**Quinoa seeds in lipid profile and glucose levels:** Pasko and others conducted a five week study with 24 male Wistar rats divided into different groups: a control group (corn starch); a control group with 31% of its dieting consisting of fructose; a group fed with a diet of quinoa seeds (310 g/kg); and a group fed with a diet of quinoa seeds (310 g/kg) and 31% of fructose. The group fed with quinoa showed significantly reduced serum total cholesterol (26%), LDL (57%), TG (11%) and glucose (10%). Fructose significantly decreased HDL (15%) levels in the control group, but when quinoa seeds were added to the diet, the decrease in HDL levels was inhibited [66]. The same research group analyzed fructose administration effects in the same animal groups, and they reported an increase in oxidative stress (high malondialdehyde) in the control group with 31% of its dieting consisting of fructose. In addition, the co-administration of quinoa seeds (310 g/kg fodder) decreased malondialdehyde in plasma [67].

## Clinical studies

**Quinoa and serum lipids:** A prospective and double-blind study was conducted in overweight, postmenopausal women testing the effects of quinoa flakes vs corn flakes (25 g for 4 weeks). The group that consumed quinoa flakes showed a significant reduction in serum TG, a tendency of reduced total cholesterol and LDL and increased glutathione [68].

**Quinoa and celiac disease:** Zevallos and others conducted a study in nineteen celiac patients who consumed 50 g/day of quinoa for 6 weeks as part of their usual gluten-free diet. Duodenal biopsies and blood

samples were obtained. Quinoa consumption was well-tolerated and did not exacerbate the clinical presentation of celiac disease. The study showed a positive trend toward improved histological parameters (the ratio of villus height to crypt depth improved from slightly below normal values 2.8:1 to normal levels 3:1, surface-enterocyte cell height improved from 28.76  $\mu$ m to 29.77  $\mu$ m and the number of intra-epithelial lymphocytes per 100 enterocytes decreased from 30.3 to 29.7) and serum total cholesterol [51].

**Betaine and homocysteine in plasma:** McRae conducted a meta-analysis of randomized placebo-controlled trials that used betaine supplementation. This analysis included five randomized controlled trials, published between 2002 and 2010. The studies used healthy adult participants who were supplemented with at least 4 g/d of betaine for a period between 6 and 24 weeks. The meta-analysis concluded that betaine supplementation decreased plasma homocysteine 1.23  $\mu$ mol/L [69].

**Betaine and metabolic syndrome:** There is a significant negative association between markers of obesity (BMI, percent body fat and waist circumference) and plasma betaine concentrations in human cross-sectional data; in addition, more than 20% of patients with diabetes mellitus excrete abnormal amounts of betaine in their urine (41). Higher concentrations of plasma betaine were associated also with lower non-HDL cholesterol, TG, homocysteine and markers of inflammation [70].

**Quinoa and diabetes:** Jenkins and others tested a low GI diet (quinoa included). For 6 months, they performed follow-up on 210 patients divided into 2 groups (high cereal fiber diet or low GI diet). They observed a greater decrease of Hemoglobin A1c (HbA1c) in the low GI diet (-0.50%) and an increase in HDL (1.7 mg/dL) in a statistically significant manner [71].

To summarize the results presented in this section, quinoa emerges as a food of particular interest to celiac patients, as the potential cornerstone of a healthy, gluten-free diet. Furthermore, we hypothesize that including quinoa in the diet could decrease oxidative stress, improve serum lipid profile, help to control body weight and serum glucose, and decrease cardiovascular disease and type 2 diabetes risk factors; quinoa may even prove beneficial in reversing the effects of these diseases. However, until now, very few studies using quinoa or quinoa compounds *in vitro*, *in vivo* or clinical trials have taken place for determining translational applications based on strong scientific evidence.

In Table 3, we provide a complete summary of the potential health benefits of quinoa.

Compound	Dose	Model	Effect	Reference
Betaine	2 mM	H2.35 mouse hepatocyte cells	Improves mitochondrial respiration	[40]
Quinoa leaves extract	WI	AT-2 and MAT-LyLu prostate cancer cells	Cell proliferation was blocked, expression of Cx43 decreased, and lipid oxidation was prevented.	[61]
Quinoa flour	WI	Analysis of quinoa flour	ACE inhibition activity (23.3%), potent free radical-scavenging.	[11]
Saponins	300 $\mu$ g	Mice	No lethality or local toxicity.	[62]
20-hydroxyecdysone	WI	C57BL/6J mice	Decreased epididymal adipose tissue. Trend of reduced lipid storage capacity in the adipose tissue, increased LPL and P-PARG, reduced	[64,65]



			expression of PEPCK and PAI-1 in adipose tissue, and also decreased adipocyte hypertrophy.	
Extract from seeds of quinoa	2,000 mg/kg	Male Wistar rats	Tendency of decreased food intake, body weight, fat deposition, and blood TG levels.	[15]
Quinoa seeds	310 g/kg	Male Wistar rats	Reduction in total cholesterol (26%), LDL (57%), TG (11%), glucose (10%), and malondialdehyde in plasma.	[66,67]
Quinoa flakes	25 g	Postmenopausal women who were overweight	Reduction in serum TG, and tendency of decreased total cholesterol and LDL and increased glutathione.	[68]
Betaine	4 g	Healthy adults	Decreased plasma homocysteine 1.23 µmol/L.	[69]
Quinoa	50 g	Celiac patients	Good tolerance without any exacerbation of celiac disease. Trend toward improved histological parameters and serum total cholesterol.	[51]
Low GI diet that included quinoa	WI	Patients with type 2 diabetes	HbA1c decreased 50% and HDL increased 1.7 mg/dL.	[71]

**Table 3:** Potential health benefits of quinoa, we review the effects of quinoa or its components in *in vitro* and *in vivo* studies and in clinical trials. WI: Without information, ACE: Angiotensin I Converting Enzyme, LPL: Lipopolysaccharides, PAI-1: Plasminogen Activator Inhibitor-1, Hb1Ac: Hemoglobin A1c.

## Culinary Issues

Quinoa is traditionally cooked as a whole grain similar to rice or milled into flour and made into pasta and breads [8]. This grain can also be processed by extrusion, drum-drying and autoclaving. Quinoa is very versatile in terms of culinary preparation: whole grains, uncooked or roasted flour, small leaves, meal, and instant powder can be prepared in a number of ways. There are numerous recipes available featuring approximately 100 preparations (Figure 2). Commercially available quinoa products exist such as pasta, bread, cookies, muffins, cereal, snacks, drinks, flakes, baby food, and diet supplements; in general, these have approximately 20% quinoa content [5,11,72-80]. The fresh leaves and tender shoots of the plant are eaten raw in salads, or cooked and eaten as a vegetable [1,5,20].

Quinoa cooking times varied from 11.9 to 19.2 min depending on the kind of seed (longer cooking time also correlated with higher protein content). Water uptake ratio varied from 2.5 to 4.0 fold [73].

## Organoleptic properties

Organoleptic properties are the aspects of food as experienced by the senses, including taste, sight, smell, and touch. Food texture refers to those qualities of a food that can be felt with the fingers, tongue, palate, or teeth. In addition, the replacement of refined flour by whole grains such as quinoa could produce changes in the organoleptic properties of foods, like darker color due to the presence of bran, and may influence the consumer decisions.

Cooked quinoa normally has a texture described as creamy and smooth but, at the same time, slightly crunchy [79].

## Conclusion

The use of pseudocereals such as quinoa represents a promising area of research, as its use could improve the intake of certain macromolecules and phytochemicals that are known to be beneficial to human health. Quinoa is an attractive, gluten-free alternative available

to celiac patients, and including quinoa in the diet may prove to be a good strategy for consuming high biological value proteins as well as all bioavailable essential amino acids, something that other grains rarely offer. Quinoa also contains unsaturated lipids, fiber, complex carbohydrates and other beneficial compounds such as betaine, and may be used to improve the metabolic risk factor profile and help control type 2 diabetes. Quinoa may be useful for some other medical complications such as cancer, obesity and dyslipidemia. Furthermore, quinoa, as a plant food, is highly sustainable, with a carbon and water food print that is 30 to 60 times lower than that of beef.

Moreover, the anti-nutritional factors in the quinoa grain could easily be inactivated or reduced to safe health levels when appropriate techniques for industrial processing or household preparation of this food are used. As for culinary applications, the replacement of refined flour by whole grains such as quinoa raises several challenges that need to be addressed, including changes in organoleptic properties.

Finally, quinoa could represent a strategic crop used to complement the diet in rural/marginal regions where energy-protein malnutrition affects a greater part of the population in certain developing countries.

In summary, there are few *in vitro* or *in vivo* studies or clinical trials using quinoa or quinoa compounds; there is a justified need for well-designed clinical trials and increased scientific research in this field, in order to determine the most effective translational applications of quinoa based on solid scientific evidence. Quinoa, known as the “mother grain”, represents an exotic and healthy rediscovered food in the developed world.

## Acknowledgement

All authors have read and approved the final manuscript.

The authors thank Kimberly Katte for providing linguistic assistance and also Laura Brugnara and Serafin Murillo for manuscript proofreading.



**Figure 2:** Quinoa salad, quinoa is very versatile in culinary preparation; in this picture quinoa is boiled in water and then mixed with vegetables.

### Sources of Financial Support

This work was supported by Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS) and Fundació Catalunya-La Pedrera, both in Barcelona, Spain.

EGB is recipient of a postdoctoral fellowship from CONACYT (México). DADR has a doctoral fellowship from Resercaixa (Barcelona, Spain).

### Potential Conflicts of Interest

The authors (EGB, ER, DADR, TM and RG) have no potential conflicts of interest relevant to this work.

### References

1. Tang Y, Li X, Chen PX, Zhang B, Hernandez M, et al. (2015b) Characterisation of fatty acid, carotenoid, tocopherol/tocotrienol compositions and antioxidant activities in seeds of three *Chenopodium quinoa* Willd. genotypes. Food chemistry 174: 502-508.
2. Tang Y, Li X, Zhang B, Chen PX, Liu R, et al. (2015) Characterisation of phenolics, betanins and antioxidant activities in seeds of three *Chenopodium quinoa* Willd. genotypes. Food Chem 166: 380-388.
3. González Martín MI, Wells Moncada G, Fischer S, Escuredo O (2014) Chemical characteristics and mineral composition of quinoa by near-infrared spectroscopy. J Sci Food Agric 94: 876-881.
4. Vega-Gálvez A, Miranda M, Vergara J, Uribe E, Puente L, et al. (2010) Nutrition facts and functional potential of quinoa (*Chenopodium quinoa* Willd.), an ancient Andean grain: a review. J Sci Food Agric 90: 2541-2547.
5. Jancurová M, Minarovicova L, Dandar A (2009) Quinoa - A Review. Czech J Food Sci 27: 71-79.
6. Galvez Ranilla L, Apostolidis E, Genoveses MI, Lajolo FM, Shetty K (2009) Evaluation of Indigenous Grains from the Peruvian Andean Region for Antidiabetes and Antihypertension Potential Using *In Vitro* Methods. J Med Food 12: 704-13.
7. Cooper R (2015) Re-discovering ancient wheat varieties as functional foods. J Tradit Complement Med 5: 138-143.
8. Bazile D, Bertero D, Nieto C (2014) Estado del arte de la quinua en el mundo en 2013: FAO (Santiago de Chile) y CIRAD (Montpellier, Francia). Perspectivas Nutracéuticas de la Quinua: Propiedades Biológicas y aplicaciones funcionales. FAO pp: 341-57.
9. Schaffer-Lequart C, Lehmann U, Ross AB, Roger O, Eldridge AL, et al. (2015) Whole Grain in Manufactured Foods: Current use, Challenges and the way forward. Crit Rev Food Sci Nutr (ahead of print).
10. van der Kamp JW, Poutanen K, Seal CJ, Richardson DP (2014) The HEALTHGRAIN definition of 'whole grain'. Food Nutr Res 58.
11. Asao M, Watanabe K (2010) Functional and Bioactive Properties of Quinoa and Amaranth. Food Sci Technol Res 16: 163-8.
12. Kozioł MJ (1992) Chemical Composition and Nutritional Evaluation of Quinoa (*Chenopodium quinoa* Willd.). J Food Comp Anal 5: 35-68.
13. Morrison CD, Laeger T (2015) Protein-dependent regulation of feeding and metabolism. Trends Endocrinol Metab 26: 256-262.
14. Lee D, Hwang W, Artan M, Jeong DE, Lee SJ (2015) Effects of nutritional components on aging. Aging Cell 14: 8-16.
15. Meneguetti QA, Brenzan MA, Batista MR, Bazotte RB, Silva DR, et al. (2011) Biological effects of hydrolyzed quinoa extract from seeds of *Chenopodium quinoa* Willd. J Med Food 14: 653-657.



16. Gesinski K, Nowak K (2011) Comparative analysis of the biological value of protein of *Chenopodium quinoa* Willd. and *Chenopodium album* L. Part I Amino acid composition of the seed protein. Acta Sci Pol Agricultura 10: 47-56.
17. Saturni L, Ferretti G, Bacchetti T (2010) The gluten-free diet: safety and nutritional quality. Nutrients 2: 16-34.
18. Yang H, Ludewig U (2014) Lysine catabolism, amino acid transport, and systemic acquired resistance: what is the link? Plant Signal Behav 9: e28933.
19. Sánchez-Chino X, Jiménez-Martínez C, Dávila-Ortiz G, Álvarez-González I, Madrigal-Bujaidar E (2015) Nutrient and Non-nutrient Components of legumes, and Its Chemopreventive Activity: A Review. Nutr Cancer 24: 1-10.
20. Schlick G, Bubenheim DL (1993) (Ames Research Center, Moffett Field, California). Quinoa: An Emerging "New" Crop with Potential for CELSS. NASA (US). Report No: 3422, pp: 1-6.
21. Abugoch LE, Romero N, Tapia CA, Silva J, Rivera M (2008) Study of some physicochemical and functional properties of quinoa (*Chenopodium quinoa* Willd) protein isolates. J Agric Food Chem 56: 4745-4750.
22. Ruini LE, Ciati R, Pratesi CA, Marino M, Principato L, et al. (2015) Working toward healthy and sustainable diets: the "Double Pyramid Model" developed by the Barilla center for Food and Nutrition to raise awareness about the environmental and nutritional impact of foods. Frontiers in Nutrition 2: 1-6.
23. Yao Y, Shi Z, Ren G (2014a) Antioxidant and Immunoregulatory Activity of Polysaccharides from Quinoa (*Chenopodium quinoa* Willd.). International Journal of Molecular Sciences 15: 19307-19318.
24. Maki KC, Phillips AK (2015) Dietary substitutions for refined carbohydrate that show promise for reducing risk of type 2 diabetes in men and women. J Nutr 145: 159S-163S.
25. Da Silva MS, Rudkowska I (2015) Dairy nutrients and their effect on inflammatory profile in molecular studies. Mol Nutr Food Res 59: 1249-1263.
26. Guadarrama-López AL, Valdés-Ramos R, Martínez-Carrillo BE (2014) Type 2 Diabetes, PUFAs, and Vitamin D: Their Relation to Inflammation. J Immuno Res 1-13.
27. Fardet A (2010) New hypotheses for the health-protective mechanisms of whole-grain cereals: what is beyond fibre? Nutr Res Rev 23: 65-134.
28. Wu Y, Qian Y, Pan Y, Li P, Yang J, et al. (2015) Association between dietary fiber intake and risk of coronary heart disease: A meta-analysis. Clin Nutr 34: 603-611.
29. Lamothe LM, Srichuwong S, Reuhs BL, Hamaker BR (2015) Quinoa (*Chenopodium quinoa* W.) and amaranth (*Amaranthus caudatus* L.) provide dietary fibres high in pectic substances and xyloglucans. Food Chem 167: 490-496.
30. Thomas D, Elliott EJ (2009) Low glycaemic index, or low glycaemic load, diets for diabetes mellitus. Cochrane Database Syst Rev: CD006296.
31. Thomas DE, Elliott EJ, Baur L (2007) Low glycaemic index or low glycaemic load diets for overweight and obesity. Cochrane Database Syst Rev: CD005105.
32. Atkinson FS, Foster-Powell K, Brand-Miller JC (2008) International tables of glycemic index and glycemic load values: 2008. Diabetes Care 31: 2281-2283.
33. Pereira MA, Jacobs DR Jr, Pins JJ, Raatz SK, Gross MD, et al. (2002) Effect of whole grains on insulin sensitivity in overweight hyperinsulinemic adults. Am J Clin Nutr 75: 848-855.
34. Pellegrini N, Agostoni C (2014) Nutritional aspects of gluten-free products. J Sci Food Agric 95: 1-6.
35. USDA (2015) National Nutrient database for Standard Reference Release 27. National Agricultural Library. Basic Report 20137 (consulted on March 2015).
36. The University of Sydney (2014) GI Foods advanced research. Sydney University's Glycemic Index Research Service (consulted on July 2015).
37. Granados-Silvestre Mde L, Ortiz-López MG, Montúfar-Robles I, Menjivar-Iraheta M (2014) Micronutrients and diabetes, the case of minerals. Cir Cir 82: 119-125.
38. Sharma KD, Bindal G, Rathour R, Rana JC (2012).  $\beta$ -carotene and mineral content of different *Chenopodium* species and the effect of cooking on micronutrient retention. Int J F Sci Nutr 63: 290-295.
39. Nascimento AC, Mota C, Coelho I, Gueifao S, Santos M, et al. (2014) Characterisation of nutrient profile of quinoa (*Chenopodium quinoa*), amaranth (*Amaranthus caudatus*), and purple corn (*Zea mays* L.) consumed in the North of Argentina: Proximates, minerals and trace elements. Food Chem 148: 420-426.
40. Lee I (2015) Betaine is a positive regulator of mitochondrial respiration. Biochem Biophys Res Commun 456: 621-625.
41. Lever M, Slow S (2010) The clinical significance of betaine, an osmolyte with a key role in methyl group metabolism. Clin Biochem 43: 732-744.
42. Shabala L, Mackay A, Tian Y, Jacobsen SE, Zhou D, et al. (2012) Oxidative stress protection and stomatal patterning as components of salinity tolerance mechanism in quinoa (*Chenopodium quinoa*). Physiologia Plantarum 146: 26-38.
43. Schartum-Hansen H, Ueland PM, Pedersen ER, Meyer K, Ebbing M, et al. (2013) Assessment of urinary betaine as a marker of diabetes mellitus in cardiovascular patients. PLoS One 8: e69454.
44. Wijekoon EP, Brosnan ME, Brosnan JT (2007) Homocysteine metabolism in diabetes. Biochem Soc Trans 35: 1175-1179.
45. Ross AB, Zangger A, Guiraud SP (2014) Cereal foods are the major source of betaine in the Western diet--analysis of betaine and free choline in cereal foods and updated assessments of betaine intake. Food Chem 145: 859-865.
46. Jakobek L (2015) Interactions of polyphenols with carbohydrates, lipids and proteins. Food Chem 175: 556-567.
47. Repo-Carrasco-Valencia R, Hellstrom JK, Pihlava JM, Mattila PH (2010) Flavonoids and other phenolic compounds in Andean indigenous grains: Quinoa (*Chenopodium quinoa*), kañiwa (*Chenopodium pallidicaule*) and kiwicha (*Amaranthus caudatus*). Food Chemistry 120:128-133.
48. Messina M (2014) Soy foods, isoflavones, and health of postmenopausal women. Am J Clin Nutr 1: S423-S430.
49. Müller L, Caris-Veyrat C, Lowe G, Böhm V (2015) Lycopene and Its Antioxidant Role in the Prevention of Cardiovascular Diseases - A Critical Review. Crit Rev Food Sci Nutr .
50. Tang Y, Li X, Chen PX, Zhang B, Hernandez M, et al. (2014) Lipids, Tocopherols, and Carotenoids in Leaves of Amaranth and Quinoa Cultivars, and a New Approach to Overall Evaluation of Nutritional Quality Traits. J Agric Food Chem 62: 12610-12619.
51. Zevallos VF, Herencia LI, Chang F, Donnelly S, Ellis HJ, et al. (2014) Gastrointestinal effects of eating quinoa (*Chenopodium quinoa* Willd.) in celiac patients. Am J Gastroenterol 109: 270-278.
52. Tovoli F, Masi C, Guidetti E, Negrini G, Paterini P, et al. (2015) Clinical and diagnostic aspects of gluten related disorders. World J Clin Cases 3: 275-284.
53. Peñas E, Uberti F, di Lorenzo C, Ballabio C, Brandolini A, et al. (2014) Biochemical and immunochemical evidences supporting the inclusion of quinoa (*Chenopodium quinoa* Willd.) as a gluten-free ingredient. Plant Foods Hum Nutr 69: 297-303.
54. Elekofehinti OO (2015) Saponins: Anti-diabetic principles from medicinal plants - A review. Pathophysiology 22: 95-103.
55. Maradini Filho AM, Ribeiro Pirozi M, Da Silva Borges JT, Pinheiro Sant 'Ana HM, Paes Chaves JB, et al. (2015) Quinoa: Nutritional, functional and anti-nutritional aspects. Critical Reviews in Food Science and Nutrition (ahead of print).
56. Gómez-Caravaca AM, Iafelice G, Verardo V, Marconi E, Caboni MF (2014) Influence of pearling process on phenolic and saponin content in quinoa (*Chenopodium quinoa* Willd). Food Chem 157: 174-178.
57. Gómez-Caravaca AM, Iafelice G, Lavini A, Pulvento C, Caboni ME, et al. (2012) Phenolic compounds and saponins in quinoa samples

- (*Chenopodium quinoa* Willd.) grown under different saline and nonsaline irrigation regimens. J Agric Food Chem 60: 4620-4627.
58. Yao Y, Yang X, Shi Z, Ren G (2014b) Anti-Inflammatory Activity of Saponins from Quinoa (*Chenopodium quinoa* Willd.) Seeds in Lipopolysaccharide-Stimulated RAW 264.7 Macrophages Cells. J Food Sci 79: H1018-H1023.
  59. Gupta RK, Gangoliya SS, Singh NK (2015) Reduction of phytic acid and enhancement of bioavailable micronutrients in food grains. J Food Sci Technol 52: 676-684.
  60. Astier C, Moneret-Vautrin DA, Puillandre E, Bihain BE (2009) First case report of anaphylaxis to quinoa, a novel food in France. Allergy 64: 819-820.
  61. Gawlik-Dziki U, Ąswieca M, SuĄkowski M, Dziki D, Baraniak B, et al. (2013) Antioxidant and anticancer activities of *Chenopodium quinoa* leaves extracts - in vitro study. Food Chem Toxicol 57: 154-160.
  62. Verza SG, Silveira F, Cibulski S, Kaiser S, Ferreira F, et al. (2012) Immunoadjuvant activity, toxicity assays, and determination by UPLC/Q-TOF-MS of triterpenic saponins from *Chenopodium quinoa* seeds. J Agric Food Chem 60: 3113-3118.
  63. Graf BL, Poulev A, Kuhn P, Grace MH, Lila MA, et al. (2014) Quinoa seeds leach phytoecdysteroids and other compounds with anti-diabetic properties. Food Chem 163: 178-185.
  64. Foucault AS, Even P, Lafont R, Dioh W, Veillet S, et al. (2014) Quinoa extract Enriched in 20-Hydroxyecdysone affects energy homeostasis and intestinal fat absorption in mice fed a high-fat diet. Physiology and Behavior 128: 226-231.
  65. Foucault AS, Mathé V, Lafont R, Even P, Dioh W, et al. (2011) Quinoa extract enriched in 20-hydroxyecdysone protects mice from diet-induced obesity and modulates adipokines expression. Obesity 20: 270-277.
  66. Pasko P, Zagrodzki P, Barton H, Chlopicka J, Gorinstein S (2010a) Effect of quinoa seeds (*Chenopodium quinoa*) in diet on some biochemical parameters and essential elements in blood of high fructose-fed rats. Plant Foods Hum Nutr 65: 333-338.
  67. Pasko P, Barton H, Zagrodzki P, Izewska A, Krosniak M, et al. (2010) Effect of diet supplemented with quinoa seeds on oxidative status in plasma and selected tissues of high fructose-fed rats. Plant Foods Hum Nutr 65: 146-151.
  68. De Carvalho FG, Ovidio PP, Padovan GJ, Jordao Junior AA, Marchini JS, et al. (2014) Metabolic parameters of postmenopausal women after quinoa or corn flakes intake- a prospective and double-blind study. Int J Food Sci Nutr 65: 380-385.
  69. McRae MP (2013) Betaine supplementation decreases plasma homocysteine in healthy adult participants: a meta-analysis. J Chiropr Med 12: 20-25.
  70. Ross AB, Bruce SJ, Blondel-Lubrano A, Oguey-Araymon S, Beaumont M, et al. (2011) A whole-grain cereal-rich diet increases plasma betaine, and tends to decrease total and LDL-cholesterol compared with refined-grain diet in healthy subjects. British Journal of Nutrition 105: 1-12.
  71. Jenkins DJ, Kendall CW, McKeown-Eyssen G, Josse RG, Silverberg J, et al. (2008) Effect of a low-glycemic index or a high-cereal fiber diet on type 2 diabetes: a randomized trial. JAMA 300: 2742-2753.
  72. Carciochi RA, Manrique GD, Dimitrov K (2015) Optimization of antioxidant phenolic compounds extraction from quinoa (*Chenopodium quinoa*) seeds. J Food Sci Technol 52: 4396-4404.
  73. Wu G, Morris CF, Murphy KM (2014) Evaluation of texture differences among varieties of cooked quinoa. J Food Sci 79: S2337-2345.
  74. Bianchi F, Rossi EA, Gomes RG, Sivieri K (2015) Potentially synbiotic fermented beverage with aqueous extracts of quinoa (*Chenopodium quinoa* Willd) and soy. Food Sci Technol Int 21: 403-415.
  75. Medina WT, de la Llera AA, Condori JL, Aguilera JM (2011) Physical properties and microstructural changes during soaking of individual corn and quinoa breakfast flakes. J Food Sci 76: E254-265.
  76. Schumacher AB, Brandelli A, Macedo FC, Pieta L, Klug TV, et al. (2010) Chemical and sensory evaluation of dark chocolate with addition of quinoa (*Chenopodium quinoa* Willd.). J Food Sci Technol 47: 202-206.
  77. Schoenlechner R, Drausinger J, Ottenschlaeger V, Jurackova K, Berghofer E (2010) Functional properties of gluten-free pasta produced from amaranth, quinoa and buckwheat. Plant Foods Hum Nutr 65: 339-349.
  78. Del Castillo V, Lescano G, Armada M (2009) Foods formulation for people with celiac disease based on quinoa (*Chenopodium quinoa*), cereal flours and starches mixtures. Arch Latinoam Nutr 59: 332-336.
  79. Abugoch L, Castro E, Tapia C, Añón MC, Gajardo P, et al. (2009) Stability of quinoa flour proteins (*Chenopodium quinoa* Willd.) during storage. Int J Food Sci Technol 44: 2010-2013.
  80. Ruales J, de Grijalva Y, Lopez-Jaramillo P, Nair BM (2002) The nutritional quality of an infant food from quinoa and its effect on the plasma level of insulin-like growth factor-1 (IGF-1) in undernourished children. Int J Food Sci Nutr 53: 143-154.