


Whole-Genome Sequence of *Aeromonas hydrophila* Strain AH-1 (Serotype O11)

Gabriel Forn-Cuní, Juan M. Tomás,  Susana Merino

Departamento de Genética, Microbiología y Estadística, Sección Microbiología, Virología y Biotecnología, Facultad de Biología, Universidad de Barcelona, Barcelona, Spain

***Aeromonas hydrophila* is an emerging pathogen of aquatic and terrestrial animals, including humans. Here, we report the whole-genome sequence of the septicemic *A. hydrophila* AH-1 strain, belonging to the serotype O11, and the first mesophilic *Aeromonas* with surface layer (S-layer) to be sequenced.**

Received 7 July 2016 Accepted 18 July 2016 Published 1 September 2016

Citation Forn-Cuní G, Tomás JM, Merino S. 2016. Whole-genome sequence of *Aeromonas hydrophila* strain AH-1 (serotype O11). *Genome Announc* 4(5):e00920-16. doi:10.1128/genomeA.00920-16.

Copyright © 2016 Forn-Cuní et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Susana Merino, smerino@ub.edu.

Aeromonas hydrophila is a water-borne opportunistic pathogen of poikilothermic animals, including fish, reptiles, and mammals (1). Although it is an integral part of the intestinal flora of healthy fish (2), some *A. hydrophila* strains can cause severe outbreaks of motile aeromonad septicemia (MAS), causing huge economic losses in the aquaculture industry (3). In humans, *A. hydrophila* pathogenesis involves gastrointestinal and wound infections and even septicemia in immunocompromised patients (4). *Aeromonas* strains are considered a contaminant agent by the U.S. Environmental Protection Agency and are routinely monitored in drinking water. The virulence of *A. hydrophila* is multifactorial and has been associated with several pathogenic factors, including, but not limited to, surface polysaccharides, surface layers (S-layers), secretion systems, and flagella (5).

S-layers are bacterial cell surface proteins associated with different pathogenic functions, such as cell adhesion, antigenic properties, and protection against host complement system lysis and phagocytes (6). *A. hydrophila* AH-1 is a septicemic strain and the first mesophilic *A. hydrophila* strain with an S-layer to be sequenced to date. The analysis of the genome from this strain will allow in-depth understanding of the evolution and lateral transfer of genes involved in the S-layer production in bacteria and its importance in virulence and pathogenicity.

The genome of *A. hydrophila* AH-1 strain was fully sequenced using Illumina MiSeq II, generating a total of 6,856,223 paired reads with 82× coverage. Read quality analysis and trimming were done with Prinseq 0.20.4 (7). *De novo* assembly with SPAdes 3.6.0 (8) resulted in 218 scaffolds larger than 500 kb.

Genome annotation was performed both via the NCBI Prokaryotic Genome Annotation Pipeline (PGAAP) and Rapid Annotations using Subsystems Technology (RAST). The complete genome of *A. hydrophila* AH-1 is 5,123,179 bp, with 60.9% G+C content, and codes for 4,773 predicted genes, eight rRNAs, and 95 tRNA sequences.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession

no. [LYXN00000000](https://ncbi.nlm.nih.gov/bioproject/PRJNA323709) (BioProject PRJNA323709). The version described in this paper is the first version, LYXN01000000.

ACKNOWLEDGMENTS

This work was supported by the Plan Nacional de I+D+i (Ministerio de Economía y Competitividad, Spain) and the Generalitat de Catalunya (Centre de Referència de Biotecnologia).

We thank Maite Polo for her technical assistance.

FUNDING INFORMATION

This work, including the efforts of Susana Merino, was funded by Ministerio de Economía y Competitividad (MINECO) (BIO2013-47198-P).

REFERENCES

- Janda JM, Abbott SL. 2010. The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clin Microbiol Rev* 23:35–73. <http://dx.doi.org/10.1128/CMR.00039-09>.
- Li J, Zhang XL, Liu YJ, Lu CP. 2011. Development of an *Aeromonas hydrophila* infection model using the protozoan *tetrahymena thermophila*. *FEMS Microbiol Lett* 316:160–168. <http://dx.doi.org/10.1111/j.1574-6968.2010.02208.x>.
- Pang M, Jiang J, Xie X, Wu Y, Dong Y, Kwok AH, Zhang W, Yao H, Lu C, Leung FC, Liu Y. 2015. Novel insights into the pathogenicity of epidemic *Aeromonas hydrophila* ST251 clones from comparative genomics. *Sci Rep* 5:9833. <http://dx.doi.org/10.1038/srep09833>.
- Parker JL, Shaw JG. 2011. *Aeromonas* spp. clinical microbiology and disease. *J Infect* 62:109–118. <http://dx.doi.org/10.1016/j.jinf.2010.12.003>.
- Tomás JM. 2012. The main *Aeromonas* pathogenic factors. *ISRN Microbiol* 2012:256261. <http://dx.doi.org/10.5402/2012/256261>.
- Beveridge TJ, Pouwels PH, Sára M, Kotiranta A, Lounatmaa K, Kari K, Kerosuo E, Haapasalo M, Egelseer EM, Schocher I, Sleytr UB, Morelli L, Callegari ML, Nomellini JF, Bingle WH, Smit J, Leibovitz E, Lemaire M, Miras I, Salamitou S, Béguin P, Ohayon H, Gounon P, Matuschek M, Koval SF. 1997. Functions of S-layers. *FEMS Microbiol Rev* 20:99–149.
- Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863–864. <http://dx.doi.org/10.1093/bioinformatics/btr026>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.