

Schrödinger equations for higher order nonrelativistic particles and N -Galilean conformal symmetry

Joaquim Gomis*

*Departament d'Estructura i Constituents de la Matèria and Departament de Física, Universitat de Barcelona, Diagonal 647, 08028 Barcelona, Spain
and C. E. R. for Astrophysics, Particle Physics and Cosmology, Barcelona Spain*

Kiyoshi Kamimura†

*Department of Physics, Toho University, Miyama, Funabashi, 274-8510, Japan
(Received 26 September 2011; published 15 February 2012)*

We consider Schrödinger equations for a nonrelativistic particle obeying an $N + 1$ -th order higher derivative classical equation of motion. These equations are invariant under N (odd)-extended Galilean conformal algebras in general $d + 1$ dimensions. In $2 + 1$ dimensions, the exotic Schrödinger equations are invariant under N (even)-Galilean conformal algebras.

DOI: 10.1103/PhysRevD.85.045023

PACS numbers: 03.65.-w, 02.20.Sv, 45.20.D-

I. INTRODUCTION

The Schrödinger equation for a nonrelativistic free particle

$$\left(i\partial_t + \frac{1}{2M}\nabla^2\right)\psi(t, \mathbf{x}) = 0 \quad (1)$$

is invariant under the scalar projective representation of the Galilei group [1,2]. In addition, the maximal symmetry of (1) is the Schrödinger algebra [3,4] that includes extra generators associated with dilatation, expansion (special conformal transformations in one dimension) and a central charge. The Schrödinger equation is obtained by the canonical quantization of the nonrelativistic free particle whose action $S = \int dt \frac{M}{2} \left(\frac{d\mathbf{X}}{dt}\right)^2$ is also invariant under the Schrödinger group.

In this paper we will generalize above result ($N = 1$ case) and show that the higher order nonrelativistic particle model given by the Lagrangian¹

$$\mathcal{L}_X = \frac{M}{2} \left(\frac{d^{(N+1)/2}\mathbf{X}}{dt^{(N+1)/2}}\right)^2, \quad N = 1, 3, 5, \dots \quad (2)$$

has the N -Galilean conformal (NGC) symmetry [7,8].² M is the constant with the dimension $[M] = (\text{mass})^{2-N}$. Furthermore the corresponding Schrödinger equations are projective representation of the NGC symmetries. For even N case we consider only $2 + 1$ dimensions where the central extension of the NGC algebra exists.

*gomis@ecm.ub.es

†kamimura@ph.sci.toho-u.ac.jp

¹The $N = 3$ case was considered in [5] as an example of a realization of the Galilei algebra with zero mass. The model was also considered in [6].

²In [9] it was conjectured that $N + 1$ -th order free equations of motion are NGC invariant for any N , odd and even.

The NGC algebra [7,8] is the finite dimensional extension of the Galilean algebra for positive integer N . Their commutators are

$$\begin{aligned} [D, H] &= iH, & [C, H] &= 2iD, & [D, C] &= -iC, \\ [H, \mathcal{P}_j] &= -ij\mathcal{P}_{j-1}, & [D, \mathcal{P}_j] &= -i\left(j - \frac{N}{2}\right)\mathcal{P}_j, \\ [C, \mathcal{P}_j] &= -i(j - N)\mathcal{P}_{j+1}, & [\mathbf{J}_{ab}, \mathcal{P}_{j,c}] &= i\delta_{c[b}\mathcal{P}_{j,a]}, \\ [\mathbf{J}_{ab}, \mathbf{J}_{cd}] &= i(\delta_{c[b}\mathbf{J}_{a]d} - \delta_{d[b}\mathbf{J}_{a]c}). \end{aligned} \quad (3)$$

It has subalgebras, (H, D, C) , one dimensional conformal algebra and $(H, \mathcal{P}_0, \mathcal{P}_1, \mathbf{J})$, the unextended Galilean algebra and the acceleration extended Galilean algebra [6] with higher order accelerations \mathcal{P}_j , ($j = 2, \dots, N$). N is interpreted in terms of the dynamical exponent z under the dilation D of the coordinates as

$$t \rightarrow \lambda^z t, \quad \mathbf{X} \rightarrow \lambda \mathbf{X}, \quad z = \frac{2}{N}. \quad (4)$$

Recently it gets much attention on the Galilean conformal symmetry and its extensions especially in condensed matter physics and gravity [9–12]. It is interesting to see how such symmetries are realized in a simple particle models.

In Sec. II we introduce a particle action invariant under the central extension of the NGC algebra (3). In Sec. III we discuss how the symmetry is realized in the Schrödinger equation in quantum theory. The even- N cases in $2 + 1$ dimensions are briefly commented in Sec. IV and the summary and discussion are in Sec. V. In the Appendix we explain how the invariant actions are derived using the method of nonlinear realization.

II. NGC INVARIANT PARTICLE MODEL

In order to quantize the particle model described by the Lagrangian (2) we construct the Hamiltonian associated with it. Since the theory is higher order in time derivative,

we introduce auxiliary coordinates $\mathbf{X}^j = \dot{\mathbf{X}}^{j-1}$, ($1 \leq j \leq \frac{N-1}{2}$), $\mathbf{X}^0 \equiv \mathbf{X}$ and the Lagrange multipliers \mathbf{Y}_j to get the Lagrangian as

$$\mathcal{L}_X = \frac{M}{2} (\dot{\mathbf{X}}^{(N-1)/2})^2 + \sum_{j=0}^{(N-3)/2} \mathbf{Y}_j (\dot{\mathbf{X}}^j - \mathbf{X}^{j+1}). \quad (5)$$

The Ostrogradsky momenta [13] are

$$\mathbf{P}_j = \mathbf{Y}_j, \quad \left(j = 0, \dots, \frac{N-3}{2} \right), \quad (6)$$

$$\mathbf{P}_{(N-1)/2} = M \dot{\mathbf{X}}^{(N-1)/2}.$$

If we use the second class constraint $(\mathbf{Y}_j, \mathbf{P}_j) = (\mathbf{P}_j, 0)$, the independent canonical pairs are $(\mathbf{X}^j, \mathbf{P}_j)$, ($0 \leq j \leq \frac{N-1}{2}$), and the Hamiltonian becomes

$$\mathcal{H} = \sum_{j=0}^{(N-3)/2} \mathbf{P}_j \mathbf{X}^{j+1} + \frac{1}{2M} (\mathbf{P}_{(N-1)/2})^2. \quad (7)$$

$$H = -\mathcal{H}, \quad D = -t\mathcal{H} + \sum_{j=0}^{(N-1)/2} \left(\frac{N}{2} - j \right) \frac{\{\mathbf{X}^j, \mathbf{P}_j\}_+}{2},$$

$$C = -t^2 \mathcal{H} + \sum_{j=0}^{(N-1)/2} \left((N-2j)t \frac{\{\mathbf{X}^j, \mathbf{P}_j\}_+}{2} + (N-j+1)j \mathbf{X}^{j-1} \mathbf{P}_j \right) - \frac{M}{2} \left(\frac{N+1}{2} \right)^2 (\mathbf{X}^{(N-1)/2})^2,$$

$$\mathcal{P}_j = j! \left(\sum_{\ell=0}^j \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{P}_\ell \right), \quad \left(j = 0, \dots, \frac{N-1}{2} \right),$$

$$\mathcal{P}_j = j! \left(\sum_{\ell=0}^{(N-1)/2} \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{P}_\ell - M \sum_{\ell=(N+1)/2}^j (-)^{((N+1)/2)+\ell} \times \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{X}^{N-\ell} \right), \quad \left(j = \frac{N+1}{2}, \dots, N \right),$$

$$\mathbf{J}_{ab} = \sum_{j=0}^{(N-1)/2} (\mathbf{X}_b^j \mathbf{P}_{ja} - \mathbf{X}_a^j \mathbf{P}_{jb}), \quad (10)$$

where $\{\mathbf{X}^j, \mathbf{P}_k\}_+ = \mathbf{X}^j \mathbf{P}_k + \mathbf{P}_k \mathbf{X}^j$. These generators, G , which have explicit t dependence, are constant of motion, i.e., $\frac{d}{dt} G = -i[G, \mathcal{H}] + \frac{\partial G}{\partial t} = 0$. It proves the transformations generated by G 's are symmetry of the Lagrangian (5). Transformations of \mathbf{Y}_j are given by those of \mathbf{P}_j . Their commutators are those of the NGC algebra (3). In addition, \mathcal{P}_j 's are no longer commuting and it appears a central charge $Z = M$ [14],

$$[\mathcal{P}_j^a, \mathcal{P}_k^b] = -i \delta^{ab} \delta^{N,j+k} (-1)^{(k-j+1)/2} k! j! Z. \quad (11)$$

In the Appendix we show that in a nonlinear realization method [15] the Lagrangian (5) appears as the left invariant one-form associated with the central charge Z in (11).

III. SCHRÖDINGER EQUATION

Now we consider the Schrödinger equation associated with the Hamiltonian (7). The canonical pairs $(\mathbf{X}^j, \mathbf{P}_j)$

Using the canonical commutators

$$[\mathbf{X}^j, \mathbf{P}_k] = i \delta_{j,k}, \quad [\mathbf{X}^j, \mathbf{X}^k] = [\mathbf{P}_j, \mathbf{P}_k] = 0, \quad (8)$$

the Heisenberg equations

$$\dot{\mathbf{X}}^j = \mathbf{X}^{j+1}, \quad \dot{\mathbf{P}}_{j+1} = -\mathbf{P}_j, \quad \left(0 \leq j \leq \frac{N-3}{2} \right), \quad (9)$$

$$\dot{\mathbf{X}}^{(N-1)/2} = \frac{1}{M} \mathbf{P}_{(N-1)/2}, \quad \dot{\mathbf{P}}_0 = 0$$

reproduce the Euler-Lagrangian equation of (2), $\frac{d^{N+1}}{dt^{N+1}} \mathbf{X}^0 = 0$. The Lagrangian (5) is invariant under the NGC symmetry whose Hermitian canonical generators are

satisfying (8) are realized as the Hermitian linear operators $(\mathbf{x}^j, -i\nabla_j)$ on complex wave functions $\psi(t, \mathbf{x}^0, \dots, \mathbf{x}^{(N-1)/2})$ with the inner product

$$\langle \psi_1 | \psi_2 \rangle = \int d^d \mathbf{x}^0 \dots d^d \mathbf{x}^{(N-1)/2} \overline{\psi_1(t, \mathbf{x}^j)} \psi_2(t, \mathbf{x}^j), \quad (12)$$

where $\bar{\psi}$ is the complex conjugate function of ψ . The Schrödinger equation $i \partial_t \psi = \mathcal{H} \psi$ for the wave function $\psi(t, \mathbf{x}^0, \dots, \mathbf{x}^{(N-1)/2})$ is

$$\Phi_S \psi(t, \mathbf{x}^0, \dots, \mathbf{x}^{(N-1)/2}) = 0, \quad (13)$$

where Φ_S is the Schrödinger differential operator defined by

$$\Phi_S = i \frac{\partial}{\partial t} - \mathcal{H} = i \frac{\partial}{\partial t} + \left(i \sum_{j=0}^{(N-3)/2} \mathbf{x}^{j+1} \nabla_j + \frac{1}{2M} (\nabla_{(N-1)/2})^2 \right). \quad (14)$$

The Schrödinger equation can be deduced from the action

$$I = \int dt d^d \mathbf{x}^j \overline{\psi(t, \mathbf{x}^j)} \Phi_S \psi(t, \mathbf{x}^j), \quad (15)$$

which is invariant under NGC transformations in the scalar projective representation of the wave function. The NGC algebra (3) is realized by the generators (10) in the form of Hermitian vector fields on the wave functions $\psi(t, \mathbf{x}^1, \dots, \mathbf{x}^{(N-1)/2})$,

$$\begin{aligned} H &= -\mathcal{H}, & D &= -t\mathcal{H} + \sum_{j=0}^{(N-1)/2} \left(\frac{N}{2} - j\right) \mathbf{x}^j (-i\nabla_j) - id \frac{(N+1)^2}{16}, \\ C &= -t^2 \mathcal{H} + \sum_{j=0}^{(N-1)/2} ((N-2j)t\mathbf{x}^j + (N-j+1)j\mathbf{x}^{j-1}) (-i\nabla_j) - \frac{M}{2} \left(\frac{N+1}{2}\right)^2 (\mathbf{x}^{(N-1)/2})^2 - id \frac{(N+1)^2}{8} t, \\ \mathcal{P}_j &= j! \left(\sum_{\ell=0}^j \frac{t^{j-\ell}}{(j-\ell)!} (-i\nabla_\ell) \right), & (j &= 0, \dots, \frac{N-1}{2}), \\ \mathcal{P}_j &= j! \left(\sum_{\ell=0}^{(N-1)/2} \frac{t^{j-\ell}}{(j-\ell)!} (-i\nabla_\ell) - M \sum_{\ell=(N+1)/2}^j (-)^{((N+1)/2)+\ell} \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{x}^{N-\ell} \right), & (j &= \frac{N+1}{2}, \dots, N), \quad Z = M, \end{aligned} \quad (16)$$

where d is the spatial dimensions. They are satisfying the NGC algebra (3) and (11). These generators commute with the Φ_S , showing that they are constant of motion. Therefore, the Schrödinger Eq. (13) remains invariant under the NGC transformations. When $\psi(t, \mathbf{x}^j)$ is a solution of the Schrödinger equation, then $\psi'(t, \mathbf{x}^j) = e^{i\alpha G} \psi(t, \mathbf{x}^j)$ is also a solution for $G = (H, D, C, \mathcal{P}_j, \mathbf{J}_{ab})$. Since the generators are Hermitian, the transformations $U = e^{i\alpha G}$ are unitary. The transformed wave function $\psi'(t, \mathbf{x}^j)$ is also written in a form

$$\psi'(t, \mathbf{x}^j) = e^{i\alpha G} \psi(t, \mathbf{x}^j) = e^{\mathcal{A}+i\mathcal{B}} \psi(t', \mathbf{x}^{j'}), \quad (17)$$

using the Schrödinger equation. Here \mathcal{A} , \mathcal{B} are real functions of the coordinates and the transformation parameters

α 's. $(t', \mathbf{x}^{j'})$ are the coordinates transformed by the NGC transformation.

The H transformation is time translation,

$$\psi'(t, \mathbf{x}^j) = e^{iaH} \psi(t, \mathbf{x}^j) = e^{a(\partial_t)} \psi(t, \mathbf{x}^j) = \psi(t+a, \mathbf{x}^j). \quad (18)$$

For the *finite* scale transformation,

$$\psi'(t, \mathbf{x}^j) = e^{i\lambda D} \psi(t, \mathbf{x}^j) = e^{\lambda d(N+1)^2/16} \psi(e^\lambda t, e^{\lambda((N/2)-j)} \mathbf{x}^j), \quad (19)$$

under which the action (15) is invariant. For the *finite* conformal transformation, $\alpha G = \kappa C$,

$$\psi'(t, \mathbf{x}^j) = e^{i\kappa C} \psi(t, \mathbf{x}^j) = e^{\mathcal{A}+i\mathcal{B}} \psi(t', \mathbf{x}^{j'}), \quad (20)$$

with

$$\begin{aligned} t' &= \frac{t}{1 - \kappa t}, & \mathbf{x}^{j'} &= \sum_{r=0}^j \frac{j C_r r! \kappa^r C_{N-j+r} C_r}{(1 - \kappa t)^{N-2j+r}} \mathbf{x}^{j-r}, & e^{\mathcal{A}} &= (1 - \kappa t)^{-(d/2)/(N+1/2)^2}, & \mathcal{B} &= -\frac{\kappa M}{2} \left(\frac{N+1}{2}\right)^2 \frac{g_N}{(1 - \kappa t)}, \\ g_N &\equiv \sum_{r,s=0}^{(N-1)/2} \frac{\gamma_r \gamma_s}{r! s! (r+s+1)} \left(\frac{\kappa}{1 - \kappa t}\right)^{r+s} \mathbf{x}^{((N-1)/2)-r} \mathbf{x}^{((N-1)/2)-s}, & \gamma_r &\equiv_{(N-1)/2} C_{((N-1)/2)-r} (r!)_{((N+1)/2)+r} C_{(N+1)/2}. \end{aligned} \quad (21)$$

For the *finite* \mathcal{P}_j transformations,

$$\psi'(t, \mathbf{x}^j) = e^{i \sum_{j=0}^N \beta^j \mathcal{P}_j} \psi(t, \mathbf{x}^j) = e^{-i2\pi\omega_1(t, \mathbf{x}; \boldsymbol{\beta})} \psi(t, \mathbf{x}^j). \quad (22)$$

The nontrivial projective phase [1] ω_1 associated with the \mathcal{P}_j transformations is given by

$$2\pi\omega_1(t, \mathbf{x}; \boldsymbol{\beta}) = M \sum_{k=0}^{(N-1)/2} (-)^{((N-1)/2)+k} \left(\mathbf{x}^k + \frac{\tilde{\boldsymbol{\beta}}^k}{2} \right) \tilde{\boldsymbol{\beta}}^{N-k}, \quad (23)$$

where the transformed coordinates are given by

$$\mathbf{x}^{0\prime} = \mathbf{x}^0 + \tilde{\boldsymbol{\beta}}^0, \quad \mathbf{x}^{\ell\prime} = \mathbf{x}^\ell + \tilde{\boldsymbol{\beta}}^\ell, \quad \tilde{\boldsymbol{\beta}}^\ell(t) \equiv \frac{d^\ell}{dt^\ell} \sum_{j=0}^N t^j \boldsymbol{\beta}^j. \quad (24)$$

We have a projective representation of the NGC group.

Under successive \mathcal{P}_j transformations we get the non-trivial 2-cocycle $\omega_2(\boldsymbol{\beta}, \boldsymbol{\beta}')$,

$$U(\boldsymbol{\beta})U(\boldsymbol{\beta}') = e^{-i2\pi\omega_2(\boldsymbol{\beta}, \boldsymbol{\beta}')}U(\boldsymbol{\beta} + \boldsymbol{\beta}'), \quad (25)$$

$$2\pi\omega_2(\boldsymbol{\beta}, \boldsymbol{\beta}') = M \sum_{k=0}^{(N-1)/2} \frac{(-)^{((N-1)/2)+k}}{2} \times (\tilde{\boldsymbol{\beta}}^k \tilde{\boldsymbol{\beta}}^{N-k} - \tilde{\boldsymbol{\beta}}'^k \tilde{\boldsymbol{\beta}}'^{N-k}). \quad (26)$$

The projective invariance of the Schrödinger equation is a one to one correspondence with the fact that the higher order Lagrangian (2), or the corresponding Lagrangian (5) is invariant up to a total derivative under the *finite* \mathcal{P}_j transformations [16],

$$\mathcal{L}'_X = \mathcal{L}_X + \frac{d}{dt}(2\pi\omega_1), \quad (27)$$

where transformations of \mathbf{X}^j and \mathbf{Y}_j are

$$\mathbf{X}^{\ell\prime} = \mathbf{X}^\ell + \tilde{\boldsymbol{\beta}}^\ell, \quad \mathbf{Y}'_j = \mathbf{Y}_j + M(-)^{((N-1)/2)+j} \tilde{\boldsymbol{\beta}}^{N-j}. \quad (28)$$

These two properties are related to the fact that NGC algebra has a central extension (11) [17].

IV. EVEN N MODEL IN 2 + 1 DIMENSIONS

In 2 + 1 dimensions we can construct a local higher order Lagrangian given by

$$\mathcal{L}_X = \frac{M}{2} \epsilon^{ab} \frac{d^{(N/2)} \mathbf{X}_a}{dt^{(N/2)}} \frac{d^{(N/2)+1} \mathbf{X}_b}{dt^{(N/2)+1}}, \quad (29)$$

where N is any positive even integer.³ The Lagrangian equivalent to (29) is

$$\mathcal{L}_X = \frac{M}{2} \epsilon^{ab} \mathbf{X}_a^{(N/2)} \dot{\mathbf{X}}_b^{(N/2)} + \sum_{j=0}^{(N/2)-1} \mathbf{Y}_j (\dot{\mathbf{X}}^j - \mathbf{X}^{j+1}). \quad (30)$$

The Ostrogradsky momenta [13] are

$$\mathbf{P}_j = \mathbf{Y}_j, \quad \left(j=0, \dots, \frac{N}{2}-1\right), \quad \mathbf{P}_{(N/2)}^b = \frac{M}{2} \mathbf{X}_a^{(N/2)} \epsilon^{ab}. \quad (31)$$

Using the second class constraints

$$\chi^a = \mathbf{P}_{(N/2)}^a + \frac{M}{2} \epsilon^{ab} \mathbf{X}^{(N/2)b} = 0, \quad a = 1, 2, \quad (32)$$

³The case $N = 2$ was analyzed in [18].

the variables $\mathbf{X}_a^{(N/2)}$, ($a = 1, 2$) can be expressed in terms of the independent $\mathbf{P}_{(N/2)}^a$, ($a = 1, 2$) satisfying

$$[\mathbf{P}_{(N/2)}^a, \mathbf{P}_{(N/2)}^b] = -i \frac{M}{4} \epsilon^{ab}. \quad (33)$$

Note that the model has built in a noncommutative structure in phase space. The Hamiltonian is

$$\mathcal{H} = \sum_{j=0}^{(N/2)-1} \mathbf{P}_j \mathbf{X}^{j+1} \quad (34)$$

and the equations of motion for $\mathbf{X} \equiv \mathbf{X}^0$ gives $\frac{d^{N+1}}{dt^{N+1}} \mathbf{X}^0 = 0$. Note that in this case the order of derivatives, $(N + 1)$, is odd. The canonical generators of NGC algebra are

$$\begin{aligned} H &= -\mathcal{H}, \quad D = -t\mathcal{H} + \sum_{j=0}^{(N/2)-1} \left(\frac{N}{2}-j\right) \frac{\{\mathbf{X}^j, \mathbf{P}_j\}_+}{2}, \\ C &= -t^2 \mathcal{H} + \sum_{j=0}^{(N/2)-1} \left((N-2j)t \frac{\{\mathbf{X}^j, \mathbf{P}_j\}_+}{2} \right. \\ &\quad \left. + (N-j+1)j \mathbf{X}^{j-1} \mathbf{P}_j \right) + N \left(\frac{N}{2}+1\right) \mathbf{P}_{(N/2)} \mathbf{X}^{(N/2)-1}, \\ \mathcal{P}_j &= j! \left(\sum_{\ell=0}^j \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{P}_\ell \right), \quad \left(j=0, \dots, \frac{N}{2}-1\right), \\ \mathcal{P}_j^b &= j! \left(\sum_{\ell=0}^{(N/2)} \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{P}_\ell^b - M \right. \\ &\quad \left. \times \sum_{\ell=(N/2)+1}^j (-)^\ell \times \frac{t^{j-\ell}}{(j-\ell)!} \mathbf{X}_a^{N-\ell} \epsilon^{ab} \right), \\ &\quad \left(j = \frac{N}{2}, \dots, N\right), \\ \mathbf{J}_{ab} &= \sum_{j=0}^{(N/2)} (\mathbf{X}_b^j \mathbf{P}_{ja} - \mathbf{X}_a^j \mathbf{P}_{jb}), \quad Z = M, \end{aligned} \quad (35)$$

where the central charge Z appears in

$$[\mathcal{P}_j^a, \mathcal{P}_k^b] = -i \epsilon^{ab} \delta^{N, j+k} (-1)^{(j-k)/2} k! j! Z. \quad (36)$$

As in the odd N case we can prove that these generators are constant of motion verifying the central extended NGC algebra. We can also prove that the associated Schrödinger equation is invariant under a scalar projective representation of the NGC group.

V. SUMMARY AND DISCUSSION

In this paper we have shown that the Schrödinger equation associated to the higher order nonrelativistic particle is

invariant under a projective representation of $N(\text{odd})$ GCA for any dimension. This results generalize the well-known result [3,4] that the ordinary Schrödinger equation is invariant under the Schrödinger (conformal) group. In the case of $2 + 1$ dimensions we have seen that the exotic Schrödinger equation is invariant under $N(\text{even})$ conformal algebras. We expect these results could be useful among other areas in the holographic description of condensed matter physics.

The Hamiltonians (7) and (34) of present models are not positive semidefinite as is known in general for higher time derivative Lagrangian systems.⁴ In quantum theory generically the system will contain ghost degrees of freedom. One possible procedure is to change the scalar product that has been applied to higher order harmonic oscillator [20] in Ref. [21]. Another possibility is to eliminate the ghost spectrum by imposing a Becchi-Rouet-Stora-Tyutin like operator on the physical states [22].

Possible extensions of the work is to consider the case of the NGC algebra for even N in any dimensions. We could also study the symmetry properties of the fourth order derivative harmonic oscillator [20], and its generalizations. We expect in this case that we will have a realization of the Newton-Hooke NGC algebra [14]. There are also possible higher order extensions of the Levy-Leblond equation [23] and the associated superconformal algebra [24].

ACKNOWLEDGMENTS

We thank J. de Azcárraga, R. Casalbuoni, J. Gomis, J. Lukierski, and P. Townsend for comments. We acknowledge partial financial support from Ministry of Education and Science FP2010-20807-C02-01, CPAN Consolider CSD 2007-00042 and from Generalitat Catalunya 2009SGR502.

APPENDIX: PARTICLE MODEL ACTION FROM NONLINEAR REALIZATION

Here we show how the NGC invariant action (5), for odd N , is derived using the nonlinear realization of the group G/H [15] (see also [25] and references therein), where G is the centrally extended algebra (3) with (11). The left invariant Maurer Cartan (MC) form $\Omega = -ig^{-1}dg$ is expanded as

$$\Omega = HL_H + DL_D + CL_C + \mathbf{P}_j L_{\mathbf{P}_j} + \frac{1}{2} \mathbf{J}_{ab} L_{\mathbf{J}^{ab}} + ZL_Z \tag{A1}$$

and is satisfying the MC equation $d\Omega + i\Omega \wedge \Omega = 0$. Using the NGC algebra (3) and (11) their components satisfy the MC equations

⁴A fourth order gravity in $(2 + 1)$ dimensions without ghost has been studied in [19].

$$\begin{aligned} dL_H &= -L_D L_H, & dL_D &= -2L_C L_H, & dL_C &= L_D L_C, \\ dL_{\mathbf{P}_j}^a &= (j+1)L_H L_{\mathbf{P}_{j+1}}^a + \left(j - \frac{N}{2}\right)L_D L_{\mathbf{P}_j}^a \\ &\quad + (j-1-N)L_C L_{\mathbf{P}_{j-1}}^a - L_{\mathbf{J}_b}^a L_{\mathbf{P}_j}^b, \\ dL_{\mathbf{J}_b}^a &= -L_{\mathbf{J}_c}^a L_{\mathbf{J}_b}^c, & dL_Z &= \frac{1}{2} \sum_{j=0}^N \frac{(-1)^j}{N C_j} L_{\mathbf{P}_j} \wedge L_{\mathbf{P}_{N-j}}. \end{aligned} \tag{A2}$$

They are closed under "d" that is equivalent with that the Jacobi identities of the algebra are satisfied. The right-hand side of dL_Z is the Wess-Zumino two-form closed and invariant in the nonextended algebra.

We parametrize the coset element as

$$g = e^{iHt} e^{i\mathbf{P}_j \tilde{\mathbf{X}}^j} e^{iC\sigma} e^{iD\rho} e^{iZc}. \tag{A3}$$

$(t, \tilde{\mathbf{X}}^0)$ for the generators (H, \mathbf{P}_0) are identified to the $D + 1$ dimensional space-time coordinates. The left invariant MC form components are

$$L_H = e^{-\rho} dt, \quad L_D = (d\rho - 2\sigma dt), \quad L_C = e^\rho (d\sigma + \sigma^2 dt), \tag{A4}$$

$$\begin{aligned} L_{\mathbf{P}_j} &= e^{(j-(N/2))\rho} \sum_{k=0}^j {}_{N-k}C_{j-k} (-\sigma)^{j-k} (d\tilde{\mathbf{X}}^k - (k+1)\tilde{\mathbf{X}}^{k+1} dt), \\ L_Z &= dc - \frac{\delta^{ab}}{2} \sum_{j=0}^N \left(\frac{(-1)^j}{N C_j} \tilde{\mathbf{X}}_a^{N-j} d\tilde{\mathbf{X}}_b^j \right. \\ &\quad \left. + \frac{(-1)^j}{N C_{j-1}} \tilde{\mathbf{X}}_a^{N-j+1} \tilde{\mathbf{X}}_b^j dt \right). \end{aligned} \tag{A5}$$

Here and hereafter when $\tilde{\mathbf{X}}^{-1}$ and $\tilde{\mathbf{X}}^{N+1}$ appear, they are understood to be zero by definition. Note in the present parametrization of the coset L_Z does not depend on ρ and σ .

In the method of nonlinear realization the particle action is constructed from \mathbf{J} invariant one-forms. They are L_H, L_D, L_C, L_Z and we can use their linear combination as the invariant action

$$\begin{aligned} I &= \int (\mathcal{L}_c + \mathcal{L}_X), & \mathcal{L}_c &= (b_H L_H + b_D L_D + b_C L_C)^*, \\ & & \mathcal{L}_X &= (a L_Z)^*, \end{aligned} \tag{A6}$$

where * means pullback to the particle world line parametrized by τ . The first term \mathcal{L}_c depends only on the $\text{su}(1, 1)$ variables, (t, σ, ρ) , and giving one-dimensional conformal mechanics Lagrangian [26].

We take the second term \mathcal{L}_X as the particle Lagrangian now depending on t and $\tilde{\mathbf{X}}^j$ in the present parametrization of the coset (A3). Using (A5)

$$\begin{aligned} \mathcal{L}_X &= a \left[\dot{c} - \frac{\delta^{ab}}{2} \left(\sum_{j=0}^N \frac{(-1)^j}{N C_j} \tilde{\mathbf{X}}_a^{N-j} \dot{\tilde{\mathbf{X}}}_b^j \right) \right. \\ &\quad \left. + \sum_{j=1}^N \frac{(-1)^j}{N C_{j-1}} \tilde{\mathbf{X}}_a^{N-j+1} \tilde{\mathbf{X}}_b^j \right] d\tau \end{aligned} \tag{A7}$$

and subtracting a surface term, it becomes

$$\mathcal{L}_X = a \left[\sum_{j=0}^{(N-3)/2} \frac{(-1)^j (j+1)}{N C_j} \tilde{\mathbf{X}}^{N-j} \cdot \left(\tilde{\mathbf{X}}^{j+1} i - \frac{1}{j+1} (\dot{\tilde{\mathbf{X}}}^j) \right) + \frac{(-1)^{(N-1)/2} (N+1)!^2}{2N! i} \left(\tilde{\mathbf{X}}^{(N+1)/2} i - \frac{2}{N+1} \dot{\tilde{\mathbf{X}}}^{(N-1)/2} \right)^2 - \frac{(-1)^{(N-1)/2} (N+1)!^2}{2N! i} (\dot{\tilde{\mathbf{X}}}^{(N-1)/2})^2 \right] d\tau. \quad (\text{A8})$$

Here $\tilde{\mathbf{X}}^{N-j}$ in the first term runs over $\tilde{\mathbf{X}}^{(N+3)/2}, \dots, \tilde{\mathbf{X}}^N$ and they play the roles of Lagrange multipliers giving their equations of motion,

$$\tilde{\mathbf{X}}^{j+1} = \frac{1}{j+1} \left(\frac{\dot{\tilde{\mathbf{X}}}^j}{i} \right), \quad j = 0, \dots, \frac{N-3}{2}. \quad (\text{A9})$$

$\tilde{\mathbf{X}}^{(N+1)/2}$ equation of motion determines $\tilde{\mathbf{X}}^{(N+1)/2}$ as

$$\tilde{\mathbf{X}}^{(N+1)/2} = \frac{2}{N+1} \left(\frac{\dot{\tilde{\mathbf{X}}}^{(N-1)/2}}{i} \right). \quad (\text{A10})$$

Using it in the Lagrangian (A8) the second term can be dropped. $\tilde{\mathbf{X}}^{(N+1)/2}, \dots, \tilde{\mathbf{X}}^1$ are solved iteratively in terms of $\tilde{\mathbf{X}}^0$ and its derivatives,

$$\tilde{\mathbf{X}}^j = \frac{1}{j!} D^j \tilde{\mathbf{X}}^0, \quad D \equiv \frac{1}{i} \frac{d}{d\tau}, \quad j = 1, \dots, \frac{N+1}{2}. \quad (\text{A11})$$

If we use them in the Lagrangian (A8) we obtain

$$\mathcal{L}_X = a \frac{(-1)^{(N+1)/2}}{2N!} (D^{(N+1)/2} \tilde{\mathbf{X}}^0)^2 i d\tau. \quad (\text{A12})$$

In a static gauge $i = 1$ the Lagrangians (A8) and (A12) become (5) and (2), respectively, by identifications $a = (-1)^{(N+1)/2} N! M$ and

$$\begin{aligned} \tilde{\mathbf{X}}^j &= \frac{\mathbf{X}^j}{j!}, & \left(j = 0, \dots, \frac{N-1}{2} \right), \\ \tilde{\mathbf{X}}^j &= \frac{(-1)^j N!}{a j!} \mathbf{Y}^{N-j}, & \left(j = \frac{N+3}{2}, \dots, N \right). \end{aligned} \quad (\text{A13})$$

For the even N case the central extension is possible only in $D = 1 + 2$ dimensions (36). Applying similar discussions, we arrive at the actions (29) and (30).

-
- [1] V. Bargmann, *Ann. Math.* **59**, 1 (1954).
[2] J.-M. Lévy-Leblond, *J. Math. Phys.* (N.Y.) **4**, 776 (1963).
[3] U. Niederer, *Helv. Phys. Acta* **45**, 802 (1972).
[4] C. R. Hagen, *Phys. Rev. D* **5**, 377 (1972).
[5] J. P. Gauntlett, J. Gomis, and P. K. Townsend, *Phys. Lett. B* **249**, 255 (1990).
[6] J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, *Phys. Lett. B* **650**, 203 (2007).
[7] J. Negro and M. A. Olmo, *J. Math. Phys.* (N.Y.) **38**, 3786 (1997); J. Negro, M. A. Olmo, and A. Rodoriguez-Marco, *J. Math. Phys.* (N.Y.) **38**, 3810 (1997).
[8] M. Henkel, *Phys. Rev. Lett.* **78**, 1940 (1997).
[9] C. Duval and P. Horvathy, *arXiv:1104.1502*.
[10] M. Henkel, R. Schott, S. Stoimenov, and J. Unterberger, *arXiv:math-ph/0601028*.
[11] S. Bhattacharyya, S. Minwalla, and S. R. Wadia, *J. High Energy Phys.* **08** (2009) 059.
[12] A. Bagchi and R. Gopakumar, *J. High Energy Phys.* **07** (2009) 037.
[13] M. Ostrogradski, *Mem. Ac. St. Petersburg* **6**, 385 (1850); As quoted in E. T. Whittaker, *Analytic Dynamics* (Cambridge University Press, Cambridge, England, 1937), p. 265.
[14] A. Galajinsky and I. Masterov, *Phys. Lett. B* **702**, 265 (2011).
[15] S. R. Coleman, J. Wess, and B. Zumino, *Phys. Rev.* **177**, 2239 (1969); C. G. Callan, S. R. Coleman, J. Wess, and B. Zumino, *Phys. Rev.* **177**, 2247 (1969).
[16] J.-M. Lévy-Leblond, *Commun. Math. Phys.* **12**, 64 (1969).
[17] G. Marmo, G. Morandi, A. Simoni, and E. C. G. Sudarshan, *Phys. Rev. D* **37**, 2196 (1988).
[18] P. C. Stichel and W. J. Zakrzewski, *Ann. Phys.* (N.Y.) **310**, 158 (2004).
[19] S. Deser, *Phys. Rev. Lett.* **103**, 101302 (2009).
[20] A. Pais and G. E. Uhlenbeck, *Phys. Rev.* **79**, 145 (1950).
[21] C. M. Bender and P. D. Mannheim, *Phys. Rev. Lett.* **100**, 110402 (2008).
[22] A. A. Slavnov, *Phys. Lett. B* **258**, 391 (1991).
[23] J.-M. Lévy-Leblond, *Commun. Math. Phys.* **6**, 286 (1967).
[24] J. P. Gauntlett, J. Gomis, and P. K. Townsend, *Phys. Lett. B* **248**, 288 (1990).
[25] J. Gomis, K. Kamimura, and P. C. West, *Classical Quantum Gravity* **23**, 7369 (2006).
[26] V. de Alfaro, S. Fubini, and G. Furlan, Report No. IC-77-7, 1977.