

STEREODYNAMICAL STUDIES OF VELOCITY ALIGNED PHOTOFRAGMENTS

M. BROUARD, S. P. DUXON, P. A. ENRIQUEZ, R. SAYOS†
and J. P. SIMONS

Chemistry Department, The University, Nottingham NG7 2RD, UK

The state resolved stereodynamics of bimolecular reactions can be probed using velocity aligned photofragments as reagents, and polarised, Doppler resolved laser detection techniques for the products. The new strategy and its application to the reaction $O(^1D) + N_2O \rightarrow NO + NO$ are outlined.

KEY WORDS: Stereodynamics, vector correlations, photodissociation.

1. INTRODUCTION

Polarised laser probing allows unprecedented levels of reagent state specification/selection and product state resolution and, with very narrow line lasers, of product velocity resolution *via* Doppler resolved spectroscopic techniques.^{1–4} Such methods, first initiated for the stereodynamics of photodissociation, have now been extended to create a new experimental strategy for probing the stereodynamics of *bi*-molecular reactions.^{5–7} Polarised photodissociation is used to generate superthermal “beams” of velocity aligned photofragments and the state-to-state stereodynamics of their subsequent collisions are probed at high resolution and sensitivity in a “simple” bulb experiment. The new strategy both complements and amplifies the more conventional molecular beam scattering technique. Doppler resolution of the polarised laser induced fluorescence (or ionisation) spectrum of the nascent *bi*-molecular collision products can allow determination of their quantum state distributions, scalar pair correlations *and* angular vector correlations, referenced initially to the laboratory frame, but after a simple laboratory to centre of mass transformation, to the *bi*-molecular collision frame.

The highly exothermic reaction



which follows the photodissociation of N_2O at 193 nm leads to the generation of rotationally excited and aligned NO molecules. The excitation into all vibrational levels up to and beyond the thermo-chemical limit of $\nu_{NO} = 16$ establishes the

† Present address: Departamento di Quimica Fisica, Universidad de Barcelona, 08028, Barcelona, Spain.

primary generation of superthermal, velocity aligned O(¹D) atoms.¹⁷ The current status of our understanding of the reaction follows a brief outline of the basic theory for velocity aligned photofragment dynamics.

2. THEORY

Translation Alignment. Figure 1 shows the Newton diagram for the collision of a superthermal velocity aligned atom A with a stationary target molecule BC. The LAB frame is defined by the polarisation vector of the incident photolysis beam $\epsilon_p \parallel Z$. The CM collision frame is defined by the collision velocity vector $\mathbf{k} \parallel z \parallel \mathbf{v}$, the photofragment recoil velocity; for a stationary target, $\mathbf{v} \parallel \mathbf{k} \parallel \mathbf{v}_{\text{CM}}$, the centre of mass velocity.

The LAB angular distribution of the bimolecular reaction products, AB, C, separating with a relative velocity \mathbf{k}' is⁵

$$I(\hat{\epsilon}_p \cdot \hat{\mathbf{k}}') \sim 1 + BP_2(\hat{\epsilon}_p \cdot \hat{\mathbf{k}}') \quad (2)$$

The coefficient B, reflecting their LAB translational anisotropy is related to β , the reagent translational anisotropy, *via* the equation

$$B = \beta \langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{k}}') \rangle \equiv 5 \langle P_2(\hat{\epsilon}_p \cdot \hat{\mathbf{k}}') \rangle \quad (3)$$

Determination of the CM translational alignment $\langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{k}}') \rangle$ simply involves laboratory frame measurement of the product and reagent translational anisotropies. This can be done by using Doppler resolved laser induced fluorescence (or ionisation) spectroscopy, to probe the reaction products and the recasting of Eq. (2)




Figure 1 Schematic Newton diagram for the collision of superthermal velocity aligned atoms, with a stationary target molecule.

in terms of the spectral lineshape function²

$$g(\bar{\nu}) \sim 1 + \beta P_2(\hat{\epsilon}_p \cdot \hat{\mathbf{k}}_a) P_2(\chi_D) \quad (4)$$

\mathbf{k}_a represents the probe laser wave vector and χ_D , the relative displacement from the line centre, is defined as

$$\chi_D = \left(\frac{\bar{\nu} - \bar{\nu}_O}{\bar{\nu}_O} \right) \frac{c}{v} \equiv \frac{\bar{\nu} - \bar{\nu}_O}{\Delta \bar{\nu}_D} \quad (5)$$

Rotational alignment. The classical angular distribution of reaction product rotational vectors, \mathbf{j}' is†

$$I(\hat{\epsilon}_p \cdot \hat{\mathbf{j}}') \sim 1 + A P_2(\hat{\epsilon}_p \cdot \hat{\mathbf{j}}') \quad (6)$$

where

$$A = \beta \langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{j}}') \rangle = 5 \langle P_2(\hat{\epsilon}_p \cdot \hat{\mathbf{k}}') \rangle \quad (7)$$

In terms of the rotational alignment parameter $A_O^{(2)}$, conventionally defined as^{2,8}

$$A_O^{(2)}(\text{LAB}) \equiv 2 \langle P_2(\hat{\epsilon}_p \cdot \hat{\mathbf{j}}') \rangle \quad \text{or} \quad A_O^{(2)}(\text{CM}) \equiv 2 \langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{j}}') \rangle \quad (8)$$

the alignment coefficient

$$A \equiv \frac{5}{2} A_O^{(2)}(\text{LAB}) \equiv \frac{1}{2} \beta A_O^{(2)}(\text{CM}) \quad (9)$$

in the high j limit.

In practice, estimation of the translational anisotropy and rotational alignment parameters B and A *via* simulation of the experimental Doppler profiles, requires several levels of convolution, to accommodate the probe laser line profile, the thermal motion of the precursor and target molecules and any spread in the superthermal reagent speeds introduced by the dynamics of the primary photodissociation process.⁶

Other vector correlations. Doppler resolved probing of photoninitiated *bi*-molecular collision/reaction products can reveal several additional correlations. As with direct photodissociation a total of nine bipolar moments are required to characterise fully the correlated angular distributions of \mathbf{j}' and \mathbf{k}' about ϵ_p , in a linearly polarised pump-probe experiment.² Provided the reagent translational anisotropy is known, the distributions of \mathbf{j}' and \mathbf{k}' can be referenced to the \mathbf{k} vector rather than ϵ_p or the transition dipole μ and the Doppler profiles can be analysed to obtain the full set of bipolar moments.² Note however, the important correlation \mathbf{k}' , \mathbf{j}' , reflecting angle bending torques at the transition state, is invariant to the LAB \rightarrow CM transformation.

3. STEREODYNAMICS OF THE REACTION $O(^1D) + N_2O \rightarrow NO + NO$

Quantum State Distributions and Scalar Pair Correlations

Photodissociation of N_2O at 193 nm initiates the highly exothermic reaction



† In Eqs. (3) and (7) the factor $\beta/5$ may be equated with $\langle P_2(\hat{\epsilon}_p \cdot \hat{\mathbf{k}}) \rangle$, arising from the LAB \rightarrow CM transformation.





Figure 2 LIF spectra of NO($v=17$) recorded under (a) relaxed, and (b) near single collision conditions.

which proceeds with near unit collision efficiency to generate NO molecules in a very broad spectrum of rovibrational states up to and including $\nu_{\text{NO}} = 18$.⁷ Fully dispersed LIF spectra of NO have been recorded under both single collision and partially relaxed conditions, *via* the γ ($\nu \leq 12$) and β ($10 \leq \nu \leq 18$) band systems. Full assignment of the populated levels is a daunting task, because of the very broad and dense rovibrational product state distributions (see Figure 2) but the task is now largely complete. Apart from those in the level $\nu = 0$ (*vide infra*) they are all endowed with high levels of rotational excitation (see Figure 2). Population of the most highly excited molecules, i.e. those generated in $\nu = 18$ two quanta above the thermochemical limit, require a collision energy $\langle E_{\text{CM}} \rangle \leq 70 \text{ kJ mol}^{-1}$. This can only be introduced *via* the translational excitation of the reagent O(¹D) atoms generated in the primary photo-dissociation step.



If the N₂ were formed predominantly in $\nu_{\text{N}_2} = 0$, for $\langle E_{\text{CM}} \rangle \leq 70 \text{ kJ mol}^{-1}$ energy conservation would require the primary excitation of the N₂ into rotational levels $J \leq 90$, accounting for $\leq 55\%$ of the available energy.† The analogous photodissociation of the isoelectronic molecule HN₃ generates N₂ in $\nu_{\text{N}_2} = 0$ with $\sim 50\%$ of the available energy in rotation.^{9,10} Molecules generated in $\nu_{\text{NO}} = 0$ are rotationally cold (mean rotational temperatures $T_{\text{R}} \approx 320 \text{ K}$ —see Figure 3) *and* translationally unexcited (Doppler with $\Delta\tilde{\nu}_{\text{D}} = 0.044 \text{ cm}^{-1}$ —equivalent to a LAB kinetic energy $\sim 110 \text{ cm}^{-1}$ —see Figure 4). NO molecules in levels $\nu \geq 1$ are rotationally highly excited; e.g. those in $\nu = 1$ have a rotational temperature, $T_{\text{R}} \sim 5500 \text{ K}$, those in higher vibrational levels appear to be even hotter. They also carry higher translational excitation (see Figure 4). These data allow estimates of the mean energy disposals and by energy conservation, of the scalar pair correlations summarised in Table 1.

The exceptionally cold distribution in NO($\nu = 0$) requires that it be partnered by NO molecules carrying very high levels of internal excitation, predominantly in the “super-excited” levels $16 \leq \nu_{\text{NO}} \leq 18$. The narrow spread in the Doppler profiles of NO($\nu = 0$), reflects a narrow spread in the internal energy of the excited partner; the broad distribution over internal rovibrational levels in NO($16 \leq \nu \leq 18$) must reflect the spread of collision energies in the hot atom reaction. A “near-stripping” mechanism for the channel leading to NO($\nu = 0$) would accord with the behaviour observed.

The NO molecules produced in levels $\nu = 1-3$ must also be accompanied by highly excited partners but the dramatic jump in their rotational and translational energies suggests the operation of an alternative dynamical pathway. High internal excitation in *both* NO partners suggests the operation of a short-lived complex mechanism: this view is reinforced by the absence of any measureable translational vector correlation, $\langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{k}}') \rangle \approx 0$ (see below).

† A direct measurement of the discussion dynamics by Huber’s group gives $\langle (E_{\text{int}}(\text{N}_2)) \rangle = 58\%$, and $\beta[\text{O}(\text{¹D})] = +0.48$ (P. Felder, B.-M. Haas and R. Huber, private communication).




Figure 3 Rotational population distributions in NO; (a) $\nu = 0$; (b) $\nu = 1$.

Vector correlations. The LAB rotational alignments, $A_{\text{O}}^{(2)}(\text{LAB})$ for $\text{NO}(\nu = 0)$ are shown in Figure 5; despite the (inevitable) scatter, they are all negative, with $A_{\text{O}}^{(2)}(\text{LAB}) \sim -0.1$. Assuming the product alignment $\langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{j}}') \rangle > 0$, the photofragment alignment must be positive, with $2 \geq \langle \beta \rangle \geq 1/2$.[†] If the cold (“old”) $\text{NO}(\nu = 0)$ were a spectator in a pure stripping reaction, no alignment would be expected; there must be some residual interaction between the two NO species at the transition state. The negative alignment could be introduced by reagent orbital angular momentum transfer.

A near stripping mechanism requires $\hat{\mathbf{k}} \cdot \hat{\mathbf{k}}' \sim 1$, and a LAB translational alignment $B[\text{NO}] \approx \langle \beta[\text{O}(^1\text{D})] \rangle$ (cf. Eq. (3)). The $\hat{\mathbf{k}}, \hat{\mathbf{k}}'$ correlation can be estimated from the Doppler profiles; the rotational lines in the slow moving $\text{NO}(\nu = 0)$ are too narrow for the experimental resolution to reveal any changes in line shape when the excitation-detection geometry is changed from coaxial to perpendicular but their

[†] A direct measurement of the discussion dynamics by Huber’s group gives $\langle \langle E_{\text{int}}(\text{N}_2) \rangle \rangle = 58\%$, and $\beta[\text{O}(^1\text{D})] = +0.48$ (P. Felder, B.-M. Haas and R. Huber, private communication).




Figure 4 Doppler profiles for NO (a) $\nu = 0$, $N = 20$; (b) $\nu = 1$, $N = 35$; (c) $\nu = 3$, $N = 31$; all Case A; (d) $\nu = 10$, $N = 26$; Case A—circles, Case D—crosses.

Table 1 Energy disposal and pair correlations

$NO(\nu_i)$	E_ν, cm^{-1}	$\langle E_R \rangle / \text{cm}^{-1}$	$\langle E_T \rangle^{L,AB} / \text{cm}^{-1}$	$NO(\langle \nu_2 \rangle)$
0	0	200	110	16–18
1	1,876	4,000	690	15
3	5,540	>4,000	1,350	11
10	17,600	not determined	1,280	≤ 4
15	26,145	4,000	700 ^a	1
16	27,600	3–4,000	$\geq 1,600$	0




Figure 5 Rotational alignments for $\text{NO}(v=0)$ generated in the reaction of velocity aligned $\text{O}(^1\text{D})$ with N_2O .

forward scattered partners, $\text{NO}(v \sim 16\text{--}18)$ are endowed with much higher LAB velocities, $V_{\text{NO}} \geq 1.52 \text{ km s}^{-1}$ (for stripping dynamics $V_{\text{NO}} \geq 2 V_{\text{CM}} \approx 1.8 \text{ km s}^{-1}$). The NO molecules excited into intermediate vibrational levels also have larger Doppler widths but their resolved parallel and perpendicular detection profiles are identical within experimental precision (see Figure 4(d)) implying a near isotropic angular distribution, with $\langle P_2(\hat{\mathbf{k}} \cdot \hat{\mathbf{k}}') \rangle \approx 0$, and the intermediacy of a short-lived collision complex in the channels leading to vibrationally excited NO pairs.

Acknowledgements

R. Sayos thanks the British Council and the Spanish Ministry of Education and Science for the award of a Fleming Fellowship, P. A. Enriquez also thanks the Spanish Ministry of Education and Science for postgraduate research support.

References

1. "Orientation and Polarisation Effects in Reactive Collisions," Faraday Symposium, *J. Chem. Soc. Faraday Trans. 2*, 1989, **85**, 925–1376.
2. R. N. Dixon, *J. Chem. Phys.* **95**, 1866 (1986).
3. J. P. Simons, *J. Phys. Chem.* **91**, 5378 (1987).
4. G. E. Hall and P. L. Houston, *Ann. Revs. Phys. Chem.* **40**, 375 (1989).
5. G. W. Johnston, S. Satyapal, R. Bersohn and B. Katz, *J. Chem. Phys.* **92**, 206 (1990); B. Katz *et al. Faraday Discussion Chem. Soc.* **91** (1991) in press.
6. F. Green, G. Hancock and A. J. Orr-Ewing, *Faraday Discussion Chem. Soc.* **91** (1991) in press; F. Green *et al. Chem. Phys. Lett.* (1991) in press.
7. M. Brouard, S. P. Duxon, P. A. Enriquez, R. Sayos and J. P. Simons, *J. Phys. Chem.* (1991) in press.
8. R. N. Zare, "Angular Momentum: Understanding Spatial Aspects in Chemistry and Physics," Wiley-Interscience: New York 1988.
9. K.-H. Gericke, R. Theinl and F. J. Comes, *J. Chem. Phys.* **92**, 6548 (1990).
10. J. J. Chu, P. Marcus and P. Dagdigian, *J. Chem. Phys.* **93**, 257 (1990).

Special Issue on Lasers in the Preservation of Cultural Heritage

Call for Papers

During the Laser Optics Conference held in St. Petersburg, Russia, a workshop on Lasers in the Preservation of Cultural Heritage was organized.

A special issue based on this workshop aims to focus on the following topics:

- Needs and approaches in conservation: what are the needs, approaches, and limitations?
- Documentation and diagnostics:
 - Analysis of composition
 - Structural diagnostics by NDT
- Laser interventions: cleaning and restorations
- Case studies

Authors should follow the LC manuscript format described at <http://www.hindawi.com/GetJournal.aspx?journal=LC>. Prospective authors should submit an electronic copy of their complete manuscript through the LC manuscript tracking system at <http://www.hindawi.com/mts/>, according to the following timetable:

Manuscript Due	September 15, 2006
Acceptance Notification	October 31, 2006
Final Manuscript Due	November 30, 2006
Publication Date	4th Quarter, 2006

GUEST EDITORS:

Costas Fotakis, Institute of Electronic Structure and Laser (IESL), Foundation for Research and Technology - Hellas (FORTH), P.O. Box 1385, Vassilika Vouton, 71110 Heraklion, Crete, Greece; fotakis@iesl.forth.gr

Wolfgang Kautek, Institut of Physical Chemistry, University of Vienna, 1010 Vienna, Austria; wolfgang.kautek@univie.ac.at

Marta Castillejo, Spanish National Research Council (CSIC), Serrano 117, 28006 Madrid, Spain; marta.castillejo@iqfr.csic.es

Prof. Costas Fotakis
Institute of Electronic
Structure and Laser (IESL)
Foundation for Research and
Technology – Hellas (FORTH)
P.O. Box 1385, Vassilika Vouton
GR-711 10 Heraklion, Crete
Greece

Heraklion, August 23, 2006

Dear Colleague,

Laser Chemistry (LC) is resuming publication as an open access electronic journal. Reflecting the expanding research activities in this field, LC aims to provide a forum for the activities in laser interactions and laser induced chemical processes in fields of chemistry, physics, biology and materials science. The journal publishes original theoretical and applied papers as well as overviews covering a large number of topics, with particular focus on the following: laser selective chemistry; reactive scattering and state-to-state dynamics; photo dissociation and photoexcitation phenomena; laser-assisted collisions and reactions in bulk and on surfaces; processes in photochemistry and chemiluminescence; molecular relaxation processes; dephasing, energy redistribution and ultrafast dynamics; new laser techniques; applications of lasers in biology and materials science.


LC is now an open access journal; hence the full text of all articles is freely available on the journal's website immediately after publication. The main advantage of open access journals is that their readership is not limited to the subscribing institutes, leading to higher rates of downloads and citations than comparable subscription-based journals. In addition, authors retain the copyright of their papers by signing a Creative Commons Attribution License, which gives the readers the right to download, print, and redistribute any article as long as it is properly cited.

Another important issue is the review and publication speed of the journal. The publisher's electronic Manuscript Tracking System (MTS) helps reduce the review time significantly, since it eliminates many of the delays that occur during peer-review. We are then aiming for an average publication time of a few weeks following acceptance.

As the Editor-in-Chief of Laser Chemistry, I would like to invite you to submit your papers for publication to the journal and maximize the readership and impact of your research articles. You can submit your contributions at <http://www.hindawi.com/mts/>.

Please do not hesitate to contact me at lc@hindawi.com if you need further information.

Sincerely yours,



Prof. Costas Fotakis
Editor-in-Chief of LC