

1 **Photocatalytic degradation of sulfamethoxazole using TiO₂ in simulated**
2 **seawater: evidence for direct formation of reactive halogen species and**
3 **halogenated by-products**

4
5 Oriol Porcar-Santos*, Alberto Cruz-Alcalde, Núria López-Vinent, Dimitrios Zanganas,
6 Carme Sans

7 Department of Chemical Engineering and Analytical Chemistry, Faculty of Chemistry, Universitat de
8 Barcelona, C/Martí i Franqués 1, 08028 Barcelona, Spain

9
10 *Corresponding author: oriol.porcar@ub.edu
11
12
13

14 **Keywords:** Titanium dioxide nanoparticles, saline water, photocatalysis, RHS, ROS, halogenation

15
16 **Abstract**

17 Nowadays photoactivation mechanism of titanium dioxide nanoparticles (TiO₂ NPs) and
18 reactive species involved in saline waters is not sufficiently established. In this study, TiO₂
19 photocatalytic process under simulated solar irradiation was evaluated in synthetic seawater
20 and compared with deionized water, using sulfamethoxazole (SMX) as model organic
21 compound. For a TiO₂ concentration of 100 mg L⁻¹, SMX degradation resulted two times slower
22 in seawater than in deionized water by the determination of their pseudo-first order rate
23 constants of 0.020 min⁻¹ and 0.041 min⁻¹, respectively. Selected scavenging experiments
24 revealed no significant contribution of hydroxyl radicals ([•]OH) on the degradation process in
25 seawater, while these radicals contributed to circa 60% on the SMX depletion in deionized
26 water. Instead, the involvement of reactive halogen species (RHS) as main contributors for the
27 SMX degradation in seawater could be established. A mechanism for the RHS generation was
28 proposed, whose initiation reactions involve halides with the TiO₂ photogenerated holes,
29 yielding chlorine and bromine radicals (Cl[•] and Br[•]) that may later generate other RHS.
30 Production of RHS was further confirmed by the identification of SMX transformation products

31 (TPs) and their evolution over time, carried out by liquid chromatography-mass spectrometry
32 (LC-MS). SMX transformation was conducted through halogenation, dimerization and
33 oxidation pathways, involving mainly RHS. Most of the detected transformation products
34 accumulated over time (up to 360 min of irradiation). These findings bring concerns about the
35 viability of photocatalytic water treatments using TiO₂ NPs in saline waters, as RHS could be
36 yielded resulting in the generation and accumulation of halogenated organic byproducts.

37

38 **1. Introduction**



39 In recent years, titanium dioxide as nanoparticles (TiO₂ NPs), has been one of the most
40 employed photocatalysts in heterogeneous Advanced Oxidation Processes (AOPs) for water
41 treatment, due to its low cost and toxicity, and also to its high reactivity and chemical stability
42 (Riaz and Park, 2020). One of the advantages of heterogeneous photocatalysis compared with
43 other AOPs is the possibility to use natural light, like sunlight irradiation, as energy source
44 (Martins et al., 2019). Heterogeneous photocatalysis has been proved to efficiently remove a
45 wide range of micropollutants, such as pesticides and pharmaceuticals (Byrne et al., 2018;
46 Mahmoud et al., 2017). This degradation of trace organic compounds is known to occur by the
47 oxidizing activity of reactive oxygen species (ROS), which is characteristic of the
48 photoactivation process of TiO₂ in water.

49

50 In pure water, the generation mechanism of ROS during TiO₂ photoactivation such as hydroxyl
51 ([•]OH) and superoxide (O₂^{•-}) radicals, as well as hydrogen peroxide (H₂O₂) and singlet oxygen
52 (¹O₂) is a well-known process, summarized in reactions 1 to 6 (Fujishima et al., 2008; Nosaka
53 and Nosaka, 2017; Rodríguez et al., 2014). TiO₂ is known to have semiconductor properties.
54 Thus, when this material absorbs radiation with enough energy to overcome its band gap,
55 electrons (e⁻) are excited from the valence band to the conduction band, generating holes (h⁺)

56 in the valence band (reaction 1). Then, on one hand, the promoted e^- of the conduction band
 57 can be captured by dissolved O_2 in its triplet state, promoting the generation of $O_2^{\bullet-}$ (reaction
 58 2) and avoiding the recombination process between e^- and h^+ . Additionally, at acidic pH
 59 conditions, $O_2^{\bullet-}$ may be also present in its protonated form (HO_2^{\bullet} , $pK_a = 4.8$) and lead to
 60 further production of H_2O_2 and $\bullet OH$ (reactions 3 and 4). On the other hand, in the
 61 photogenerated hole, $\bullet OH$ is produced by the oxidation of H_2O (reaction 5), whereas 1O_2 can
 62 be produced by the oxidation of the previously generated $O_2^{\bullet-}$ (reaction 6).

63



64

65

66 On the other hand, photocatalytic treatments with TiO_2 are considered as treatment option for
 67 micropollutants abatement in saline waters and seawater (D. Camacho-Muñoz, 2020) and also
 68 for disinfection (Romero-Martínez et al., 2014; Rubio et al., 2013). However, the
 69 photoactivation mechanism of TiO_2 in seawater has not been completely established.
 70 According to the previous knowledge regarding the TiO_2 -UV mechanisms in pure water, it
 71 could be hypothesized that similar processes involving ROS generation could be also observed
 72 in seawater. However, recent studies suggest that the high content in halides ions of seawater
 73 may modify the chemistry of the process, this resulting in the generation and subsequent
 74 participation of reactive halogen species (RHS). This was indirectly inferred by Hao and co-
 75 workers, who observed that TiO_2 NPs promoted halogenation of dissolved organic matter
 76 (DOM) under sunlight irradiation in seawater (Hao et al., 2018). As known, many halogenated

77 organic compounds may display toxic properties and thus become a risk to the environment
78 and human health (Jiang et al., 2017; Liu and Zhang, 2014). For this reason, understanding the
79 underlying photoactivation mechanisms taking place in photocatalytic treatments of saline
80 waters is essential for a better assessment of the potential risks derived from the release of
81 halogenated transformation products into the environment.

82

83 Sulfamethoxazole (SMX) has been one of the most studied micropollutants in water treatment
84 in a wide variety of advanced oxidation processes. Among others, SMX was used in
85 photocatalytic degradation experiments with TiO₂, under solar irradiation and artificial light
86 sources (Borowska et al., 2019; Martins et al., 2019). Consequently, SMX degradation
87 mechanism by photocatalysis in pure water is well known.

88

89 The aim of this work was to investigate the degradation mechanisms taking place when
90 photocatalytic treatment with TiO₂ is implemented in saline waters such as seawater. To do so,
91 photocatalytic degradation experiments with SMX were conducted in the presence of TiO₂ NPs
92 under controlled irradiation conditions (*i.e.*, simulated sunlight) in a standard seawater matrix.
93 The relative contribution of the different generated reactive species was completed by the
94 application of selective radical scavengers in degradation tests. These experiments, together
95 with a comprehensive identification and monitoring of reaction intermediates formed during
96 SMX degradation, allowed the elucidation of the corresponding reactive halide species
97 generation mechanism and their main reaction pathways.

98

99 **2. Materials and methods**

100 *2.1 Chemicals and reagents*

101 Sulfamethoxazole and *tert*-butanol were purchased from Sigma-Aldrich (Germany). Titanium
102 dioxide (Degussa P25), was acquired from Evonik (Germany). Sodium chloride, magnesium

103 chloride hexahydrate, calcium chloride dihydrate, potassium chloride, sodium hydrogen
104 carbonate, potassium bromide, boric acid, sodium hydroxide, sodium azide and acetonitrile
105 were supplied by Panreac (Spain). Sodium sulfate and *p*-benzoquinone were purchased from
106 Probus (Spain) and Merck Schuchardt OHG (Germany), respectively. Pure nitrogen gas
107 (99.995%) was acquired from Abelló Linde (Spain).

108

109 *2.2 Photocatalytic experiments*

110 Degradation experiments were performed in simulated seawater (SW) and in deionized water
111 (DW), for comparison purposes. Simulated seawater was prepared according to the ASTM
112 D1141-98 standard (ASTM international, 2013). The chemical composition of SW is detailed in
113 Table S1 of the Supplementary Information. Its pH was adjusted to 8.2 by the addition of
114 NaOH. For DW, unless otherwise specified, the pH of the solution (pH = 6) was not modified.

115 TiO₂ NPs used in this work (Degussa P25) have a content of 80% of anatase and 20% of rutile,
116 and a BET surface between 35 and 65 m² g⁻¹. The band gap is 3.2 eV and particles have a mean
117 diameter of 21 nm.

118 The concentration of SMX and TiO₂ NPs used for degradation experiments was 1 mg L⁻¹ and
119 100 mg L⁻¹, respectively. For the transformation products characterization study, the
120 concentrations increased to 10 mg L⁻¹ of SMX and 1000 mg L⁻¹ of TiO₂ NPs, for a better
121 identification of the formed byproducts.

122 Irradiation experiments were conducted as follows: aqueous solutions were placed into a 1L
123 jacketed reservoir tank with continuous magnetic stirring. After that, these were continuously
124 pumped with a peristaltic pump, into a Duran glass tubular photoreactor (diameter and length
125 of 2 and 24.5 cm, respectively), placed inside the solar simulation chamber Xenoterm-1500RF,
126 CCI (Spain), and then recirculated back to the reservoir tank. The solar simulation chamber was
127 equipped with a Xenon lamp (1.5 kW) emulating the solar spectrum. Radiation with
128 wavelengths below 290 nm was cut-off by the Duran glass of the photoreactor. An *o*-

129 nitrobenzaldehyde actinometry was employed to establish the photonic flow in the
130 wavelength range of 290-400 nm, being this $0.66 \mu\text{Einstein s}^{-1}$ (Bustos et al., 2019). A
131 thermostatic bath was used to keep the tank temperature at $20 \text{ }^{\circ}\text{C}$ during the irradiation
132 process. The irradiation time was 120 min for all the experiments, except for the intermediates
133 study where it was extended up to 360 min. In the first case, aliquots of 10 mL were extracted
134 from the reservoir tank at different irradiation times and then filtered with $0.45 \mu\text{m}$ PVDF
135 filters. For degradation experiments, data were obtained, at least, in duplicates. The values
136 plotted showed deviations below 5%. In the intermediates study, due the observed
137 interference of the salts in the mass spectrometry analysis and for a suitable detection of the
138 intermediates generated, a solid phase extraction (SPE) treatment was employed. Aliquots of
139 40 mL were extracted from the reservoir tank at the irradiation times of 0, 30, 120, 240 and
140 360 min, and filtered with $0.45 \mu\text{m}$ PVDF filters. After that, SPE cartridges Oasis HLB 20 cc g^{-1}
141 acquired from Waters Corporation (USA), were loaded with the corresponding aliquots, then
142 washed with 40 mL of deionized water. Extraction of organic components (*i.e.*, SMX and
143 reaction intermediates) was finally achieved with 8 mL of acetonitrile. This way, salts were
144 removed and the organic components in original samples were concentrated 3 times.

145

146 *2.3 Analytical methods*

147 The SMX degradation in irradiation experiments was evaluated by an Infinity 1260 HPLC
148 provided by Agilent Technologies (USA). The employed column was a Mediterranea Sea 18
149 ($250 \text{ mm} \times 4.6 \text{ mm}$ and $5 \mu\text{m}$ particle size) supplied by Teknokroma (Spain). The mobile phases
150 used consisted of volumetric mixtures containing 60% of acetonitrile and 40% of ultrapure
151 water adjusted with orthophosphoric acid at pH 3. The flux employed was 1.1 mL min^{-1} and the
152 UV detector was set at 270 nm. The injection volume was set to $50 \mu\text{L}$ and the column
153 temperature was set to $30 \text{ }^{\circ}\text{C}$.

154

155 The intermediates identification was performed by liquid chromatography-mass spectrometry
156 (LC-MS) with an Agilent 1100 HPLC coupled to G1969A LC/MSD-TOF mass spectrometer. All
157 intermediates were detected in negative electrospray ionization mode (ESI (-)) with a
158 fragmentor voltage of 150 V and collected in a scan range of 50-1100 m/z. Data in extracted
159 ion chromatograms (EIC) were employed for the monitoring of the intermediates in the
160 different samples. The chromatographic separation was performed with the same column
161 used in the Infinity 1260 HPLC analyses and a volumetric mixture containing 40% of milli Q
162 water and 60% of acetonitrile as mobile phase. As for the SMX degradation experiments, in the
163 byproducts study the column temperature was also set to 30 °C but in this case the injection
164 volume was set to 2 µL.

165

166 **3. Results and discussion**


167 *3.1 Evaluation of photoactivation of TiO₂ NPs in seawater under simulated sunlight irradiation*

168 Photoactivation of TiO₂ NPs under simulated sunlight was studied assessing the degradation of
169 SMX as a model organic compound. Initially, a control experiment was conducted in order to
170 test SMX adsorption with TiO₂ NPs under dark conditions in the seawater matrix. Photolysis of
171 SMX by simulated sunlight irradiation was also assessed. After 120 min, SMX degradation was
172 1% and 3%, respectively, suggesting the lack of SMX adsorption and the negligible photolysis
173 contribution to the SMX degradation.

174

175 Photocatalytic degradation of 1 mg L⁻¹ SMX with TiO₂ in the SW matrix was evaluated by the
176 addition of different concentrations of TiO₂ (10, 100 and 1000 mg L⁻¹). Although concentrations
177 of TiO₂ NPs and micropollutants in the aquatic environment are typically lower (*i.e.*, in the
178 range from ng to µg L⁻¹), these experimental conditions were selected to magnify the process
179 so that this could be studied in a reasonable lapse of time. The final degradation after 120 min
180 of irradiation and the corresponding pseudo-first order rate constants obtained were 14%

181 (0.001 min⁻¹), 91% (0.020 min⁻¹) and 100% (0.079 min⁻¹), respectively, which demonstrate the
 182 photodegradation activity of the process in seawater. According to the results, a concentration
 183 of 100 mg L⁻¹ of TiO₂ was selected for performing degradation experiments. Results in SW were
 184 compared with DW using 100 mg L⁻¹ of TiO₂. After 120 min of irradiation, a complete SMX
 185 degradation was obtained, with a pseudo-first order rate constant of 0.041 min⁻¹, (Fig. 1), that
 186 is two times faster than that in SW.
 187



188
 189 **Figure 1.** Degradation of SMX using different concentration of TiO₂ in SW compared with 100
 190 mg L⁻¹ in DW.

191
 192 *3.2 Identification of reactive species generated from photoactivation of TiO₂ NPs in seawater*

193 The study of the contribution of the different reactive oxygen species (ROS), such as [•]OH, O₂^{•-},
 194 and other possible radical and oxidizing agents formed in the photocatalytic process was
 195 carried out by the addition of selected radical scavengers. For the determination of the [•]OH
 196 contribution in the bulk solution, photocatalytic experiments were performed with the
 197 presence of *tert*-butanol (TBA), which has a reaction rate constant with [•]OH of 6.0 x 10⁸ M⁻¹ s⁻¹
 198 (Buxton et al., 1988). Dissolved oxygen promotes the photocatalytic reaction by trapping the
 199 excited electron of the TiO₂ conduction band, generating superoxide radical (O₂^{•-}) and
 200 avoiding the recombination process. 1,4-benzoquinone (BQ) has been used by different
 201 researchers as scavenger of O₂^{•-} (Jedsukontorn et al., 2018; Xiao et al., 2016; Zhang et al.,


202 2017) because of its high reactivity with this species (rate constant value of $9.8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$
203 (Bielski et al., 1985)). However, BQ has also a high reactivity with $\bullet\text{OH}$ ($6,6 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ (Buxton
204 et al., 1988)). Thus, if $\bullet\text{OH}$ is expected to be involved in the degradation process, the whole
205 interaction with BQ should be considered (Rodríguez et al., 2014). Moreover, by removing
206 dissolved oxygen by bubbling the solution with N_2 , the overall contribution of dissolved oxygen
207 on the promotion of radical species can be assessed (García-Fernández et al., 2015; Rodríguez
208 et al., 2014). Another possible ROS generated in the photocatalytic process is singlet oxygen
209 ($^1\text{O}_2$), which mainly comes from the oxidation of $\text{O}_2^{\bullet-}$ by the hole (h^+). To determine the $^1\text{O}_2$
210 participation on the SMX degradation, many works in literature report the use of sodium azide
211 (NaN_3) as scavenger of this oxidizing agent (Fotiou et al., 2016; Tiwari et al., 2019; Xiao et al.,
212 2016) because of the high reactivity between both species (rate constant of $2.1 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$)
213 (Catalán et al., 2004). However, N_3^- has also a high rate constant with $\bullet\text{OH}$ ($1.2 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$),
214 generating $\bullet\text{N}_3$, also a reactive radical (Buxton et al., 1988). Thus, a double scavenging effect of
215 N_3^- should be considered if $\bullet\text{OH}$ is expected to be generated in the photocatalytic process
216 (Rodríguez et al., 2014). The h^+ participation in the SMX degradation can be evaluated by the
217 addition of formic acid (FA), which is able to suppress all the h^+ processes (Doudrick et al.,
218 2013). These include: 1) the direct oxidation of SMX by h^+ , 2) the possible generation of $^1\text{O}_2$
219 from $\text{O}_2^{\bullet-}$, 3) the generation of $\bullet\text{OH}$ through the oxidation of H_2O and 4) the production of
220 other oxidant species (Cavalcante et al., 2016; Ribao et al., 2019; Zheng et al., 2010).
221 Additionally, if $\bullet\text{OH}$ is generated in the system through other pathways, FA is also able to
222 scavenge this oxidant because of the high rate constant value of their corresponding reaction
223 ($3.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ as formate ion at neutral pH (Buxton et al., 1988)).

224


225 Results of the addition of the different scavengers (3 mM of TBA, 3 mM of NaN_3 , 11 and 560
226 mM of FA as well as 0.075 and 0.25 mM of BQ), with 100 mg L^{-1} of TiO_2 and 1 mg L^{-1} of SMX, in
227 SW and also in DW, are shown in Fig. 2A and 2B, respectively. Unless otherwise specified, the

228 concentration of scavenger employed in each test was selected to represent at least 95% of
229 the total scavenging effect exerted over the oxidant under evaluation.

230



231



232

233 **Figure 2.** Effect in the SMX degradation by the addition of different scavengers in A) SW and B)
234 DW.

235

236 The most significant difference between activation of TiO₂ NPs in seawater and deionized
237 water was for the tests performed with TBA. In SW, no inhibition of SMX degradation was
238 observed, indicating that the degradation process is not mediated by [•]OH in the bulk solution.
239 This result confirms that the mechanism of the photocatalytic degradation process in SW is
240 different from DW, where circa 60% of the SMX degradation occurred by the [•]OH in the bulk
241 solution, as it can be seen in Fig. 2B.

242

243 Experiments with FA in SW also generated significant findings. Tests were performed at two
244 different concentrations, 11 mM and 560 mM. The employment of 11 mM is based on other
245 studies that used similar concentration to block the h^+ pathway in DW (Zheng et al., 2010). On
246 the other hand, the experiment with 560 mM of FA was conducted to reach the same
247 concentration of chlorine anions (Cl^-) in SW, with the aim of overcoming potential competition
248 of this species in the FA adsorption on the TiO_2 surface. Results for 11 mM and 560 mM of FA
249 showed that, after 120 min, the SMX degradation was inhibited from 91% to 52% and from
250 91% to 5%, respectively. Thus, 560 mM of FA were able to avoid the h^+ pathway, suggesting
251 that, in SW, halide species may directly participate in the redox mechanism that takes place in
252 the hole. In the case of experiments in DW, only the 11 mM FA test was carried out. In this
253 assay in particular, the pH of the solution was adjusted to 6 with the addition of NaOH in order
254 to keep the same pH of the other experiments in DW. Results showed an almost complete
255 inhibition of the SMX degradation, from 100% to 14%, which was also attributed to the
256 impediment of the h^+ pathway avoiding the generation of the $\bullet OH$.

257

258 Other tests with selected scavengers also provided complementary information about TiO_2
259 NPs photoactivation in seawater. The removal of dissolved oxygen from the solution, by
260 bubbling it with N_2 , led to almost total inhibition of the SMX degradation, resulting just 10%
261 removal after 120 min of irradiation. This result remarks the importance of the presence of
262 dissolved oxygen to inhibit the recombination of e^-/h^+ and thus allow the performance of the
263 photocatalytic reactions. In the case of DW, the inhibition was weaker compared with SW,
264 allowing 40% removal in 120 min.

265

266 The experiment performed with a BQ concentration of 0.075 mM resulted in a SMX
267 degradation inhibited from 91% to 52%. By increasing the BQ concentration to 0.25 mM, SMX
268 degradation inhibition was even more significant, down to 18%. These differences in the BQ

269 inhibition capacity suggest that, in addition to the reaction of BQ with $O_2^{\bullet-}$, this reagent may
270 undergo reactions with other oxidant species, which in fact would be the main responsible for
271 the observed SMX degradation. In similar experiments conducted with DW and 0.075 mM of
272 BQ, SMX degradation was inhibited from 100% to 27%. This strong inhibition can be explained
273 by the presence of $\bullet OH$ in the bulk solution, as BQ may react with both $O_2^{\bullet-}$ and $\bullet OH$, resulting
274 this in a double inhibition effect over the SMX degradation.


275

276 Finally, NaN_3 test in SW revealed no participation of 1O_2 in the SMX degradation. In DW, the
277 SMX degradation was inhibited from 100% to 82% in 120 min. However, the reactivity of $^-N_3$
278 with $\bullet OH$ generate $\bullet N_3$, which may as well behave as an oxidant (An et al., 2010; Bancirova,
279 2011) and thus participate on the SMX degradation in DW.

280

281 According to the obtained results, when TiO_2 NPs are present in seawater, the oxidizing species
282 produced in the redox process at the hole may involve the participation of halide ions that are
283 found at high concentration in this water matrix (560 mM and 0.8 mM for Cl^- and Br^- ,
284 respectively). Therefore, the oxidative role of $\bullet OH$, which is the main responsible of the SMX
285 degradation in deionized water, may be replaced by reactive halogen species in saline waters.
286 Different authors previously studied the reactivity of halide ions with the TiO_2 photogenerated
287 hole to promote the formation of free halogen radicals (reactions 7 and 8), which may later
288 generate other reactive halogen species (RHS) (Boutiti et al., 2017; Daimon et al., 2008). RHS
289 comprise radical and non-radical species, such as X^\bullet , $X_2^{\bullet-}$, $HOX^{\bullet-}$, HOX , ^-OX and X_2 , where X
290 corresponds to Cl or Br, as presented in reactions 7 to 41 (Grebel et al., 2009). These species
291 are less reactive but more selective oxidants than $\bullet OH$, being mainly involved in reactions with
292 compounds containing electron-rich functional groups (Grebel et al., 2009).

293



294


295 RHS could also result from the reaction of $\bullet\text{OH}$, in case of its primary generation in the h^+ , with
 296 halides (reaction 9 and 10). However, because of the lack of inhibition of SMX degradation in
 297 the TBA experiment, this pathway is unlikely to occur. This observation, together with the
 298 complete inhibition of the h^+ pathway by FA at high concentration (similar to Cl^-), points out
 299 that the direct oxidation of halides in the photoactivated holes in saline matrices are the main
 300 promoters of RHS. Between halides, Br^- is expected to be more reactive with the hole than Cl^-

301 because of its lower reduction potential ($(E(\text{Cl}^\bullet/\text{Cl}^-) = 2.5 V_{\text{NHE}} > (E(\text{Br}^\bullet/\text{Br}^-) = 2.0 V_{\text{NHE}})$) (Zhang
302 and Parker, 2018).

303 At the seawater pH (8.2), the equilibrium between Cl^\bullet and $\bullet\text{OH}$ (reaction 9) is displaced to the
304 $\bullet\text{OH}$ formation (von Sonntag and von Gunten, 2012). Consequently, TBA experiments should
305 also have shown some inhibition of SMX degradation by $\bullet\text{OH}$ scavenging, which was not the
306 observed situation. However, it must be taken into account that the equilibrium between Br^\bullet
307 and $\bullet\text{OH}$ is displaced to the production of $\text{BrOH}^{\bullet-}$ under the same conditions (reaction 10),
308 subsequently yielding $\text{BrCl}^{\bullet-}$ (reaction 30) (Zhang and Parker, 2018). Hence, the lack of
309 inhibition of SMX degradation in the TBA experiment may be justified by the scavenging effect
310 of Br^- towards $\bullet\text{OH}$ generated through reaction 9 that would compete with TBA, minimizing its
311 scavenging effect. This is also in accordance with the higher kinetic constant of Br^- with $\bullet\text{OH}$
312 ($1.1 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$) compared with the TBA with $\bullet\text{OH}$ ($6.0 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$).

313 Additionally, Daimon and co-workers, observed a competition between halogen radicals (Cl^\bullet
314 and Br^\bullet) and dissolved O_2 for the photoexcited electron of the conduction band (Daimon et al.,
315 2008). This fact was observed with the inhibition of SMX degradation when dissolved O_2 was
316 removed from the solution (Fig. 2A), suggesting that halogen radicals recovered the electron
317 previously donated to h^+ , inactivating RHS production. This fact supports the suggested
318 primary pathway for the RHS generation by the direct reactivity of halides with the
319 photogenerated hole. Thus, once RHS are produced in the hole, more RHS could be formed
320 through the reactivity of $\text{O}_2^{\bullet-}$ with other RHS (reactions 21, 22 and 39).

321 In view of all these considerations, TiO_2 photoactivation mechanism is proposed and presented
322 in Fig. 3.



323
324

325 **Figure 3.** Proposed TiO_2 photoactivation mechanism in SW for the generation of RHS.

326

327 In short, results obtained confirm the different reactive species whose generation is
 328 photoinduced by solar radiation and nanosized TiO_2 , when this pollutant is present in different
 329 environmental water matrices. The mechanisms taking place in the aqueous environment are
 330 expected to lie between a process governed by ROS generation, in the case of low salinity
 331 waters, to another characterized and controlled by RHS production in the case of seawater, as
 332 proposed in Fig. 3. Between these two mechanisms, increasing contribution of RHS to derived
 333 reactions such as those concerning transformation of organic compounds is expected to occur
 334 as salinity of the aqueous medium increases.

335

336 *3.3 Generation and evolution of sulfamethoxazole transformation products over seawater*
 337 *photocatalytic treatment*

338 In order to corroborate the observed contribution of RHS in the photocatalytic degradation of
 339 SMX in SW, HPLC-MS analyses were performed in samples withdrawn at different irradiation
 340 times to identify the main intermediate products generated. To do so, SMX and TiO₂
 341 concentrations employed were increased to 10 mg L⁻¹ and 1000 mg L⁻¹, respectively, for a
 342 better detection of transformation products.




343











344 A total of 17 TPs were distinguished during the SMX degradation in SW, as shown in Table 1.


345 The relative evolution of the formed intermediates during the photocatalytic process in SW
 346 was followed through min-max normalization of the areas obtained by the extracted ion
 347 chromatogram. These data are represented in Fig. 4. Based on identified intermediate
 348 products and their corresponding evolutions, a mechanism for SMX transformation in the
 349 sunlight-TiO₂ process in SW was proposed and presented in Fig. 5.

350

351 **Table 1.** SMX and TPs identified in HPLC-MS analysis with ESI (-).

Molecular structure, molar mass (g/mol) and designation	Molecular Formula	LC-MS [M-H] ⁻ (m/z)
 SMX M = 253.28	C ₁₀ H ₁₁ N ₃ O ₃ S	252.04
 SMX-Cl M = 287.72	C ₁₀ H ₁₀ ClN ₃ O ₃ S	286.01-288.01
 SMX-Br M = 332.17	C ₁₀ H ₁₀ BrN ₃ O ₃ S	329.96-331.96

 <p>SMX-2Br M = 411.07</p>	$C_{10}H_9Br_2N_3O_3S$	407.87-409.86-411.86
 <p>SMX-Br-OH M = 348.17</p>	$C_{10}H_{10}BrN_3O_4S$	345.95-347.95
 <p>SMX-Br-NCN M = 276.11</p>	$C_7H_6BrN_3O_2S$	273.93-275.93
 <p>SMX-NO M = 267.26</p>	$C_{10}H_9N_3O_4S$	266.03
 <p>SMX-NO2 M = 283.26</p>	$C_{10}H_9N_3O_5S$	282.02
 <p>SMX-OH M = 269.28</p>	$C_{10}H_{11}N_3O_4S$	268.04
 <p>SMX-NCN M = 197.21</p>	$C_7H_7N_3O_2S$	196.02
 <p>SMX-Br-NO M = 346.16</p>	$C_{10}H_8BrN_3O_4S$	343.95-345.95
 <p>SMX-NO-NCN M = 211.20</p>	$C_7H_5N_3O_3S$	210.00
 <p>SMX-NO2-NCN M = 227.20</p>	$C_7H_5N_3O_4S$	225.99



352

353



354 According to the MS data, SMX transformation occurred through three primary pathways,
 355 halogenation, oxidation and dimerization, mainly driven by RHS.


356

357 The halogenation and dimerization pathways (Fig. 5, red arrows and blue arrows, respectively)
 358 may initiate through H-abstraction of the phenylamine nitrogen by halogen radical species,
 359 such as Cl^\bullet , Br^\bullet , $HOCl^{\bullet-}$, $HOBr^{\bullet-}$, $BrCl^{\bullet-}$, $Cl_2^{\bullet-}$ and $Br_2^{\bullet-}$, leading to the formation of the SMX^\bullet
 360 molecule. Further halogen radical species addition to the generated SMX^\bullet molecule would end
 361 in the formation of primary halogenated TPs. At the same time, two SMX^\bullet molecules could
 362 interact yielding SMX dimerization. The individuality of the proposed halogenation and
 363 dimerization mechanisms is justified in view of the simultaneous formation of their primary
 364 TPs (SMX-Cl and SMX-Br) and (2SMX(-)), that reached their maximum concentrations at 120
 365 min of irradiation (see Fig. 4A and 4C).

366 Five halogenated TPs were detected (Fig. 5, red arrows). The ion fragment at 233.92-235.92
 367 m/z for $(M-H)^-$ (Fig. S1 of Supplementary Information) containing the Br atom in its structure
 368 suggests that this addition took place in the phenylamine moiety, most probably at the
 369 nitrogen position (Dodd and Huang, 2004; Gao et al., 2014). From the SMX-Br intermediate,
 370 three secondary TPs were generated (SMX-2Br, SMX-Br-OH and SMX-Br-NCN). The second
 371 bromination to form SMX-2Br possibly took place again in the nitrogen atom of the
 372 phenylamine moiety. The observed hydroxylation to generate the SMX-Br-OH intermediate
 373 could probably occur in the phenyl ring. Finally, the detected SMX-Br-NCN intermediate may
 374 result from the fragmentation of the isoxazole ring. This opening ring mechanism is also
 375 reported in literature (Cai and Hu, 2017; Yuan et al., 2019), through a first hydroxylation of the
 376 ring followed by its fragmentation.


377





380

381 **Figure 4.** Normalized area of the TPs and SMX over the time by the EIC of the mass
 382 spectrometer for A) halogenation, B) oxidation and C) dimerization pathways.



383

384 **Figure 5.** SMX intermediates formation mechanism. Red, green and blue arrows refer to the
 385 degradation process initiated by halogenation, oxidation and dimerization pathways,
 386 respectively.

387 The dimerization pathway (Fig. 5, blue arrows) generated 4 identified TPs. From the primary
388 dimer 2SMX(-), three secondary TPs were identified (2SMX(=), SMX(-)-Br and SMX(=)-OH). The
389 2SMX(-)-Br generation might occur by a bromination of one of the nitrogen atoms of the
390 2SMX(-) hydrazine group. The 2SMX(=) dimeric intermediate is expected to come from a
391 second H-abstraction in the nitrogen atoms of the 2SMX(-) hydrazine group, giving rise to the
392 formation of the azo functional group. Subsequently, the hydroxylation of one of the phenyl
393 groups in the 2SMX(=) TP might lead to the formation of 2SMX(=)-OH.

394

395 Oxidation pathway exhibits different mechanisms and TPs. Despite the TBA scavenger test
396 showed no significant participation of the $\bullet\text{OH}$ in the SMX degradation, hydroxylation of the
397 SMX molecule occurred in its degradation process. Among other possibilities, the equilibrium
398 of the reaction 9 and 10 involving $\text{HOCl}^{\bullet-}$ and $\text{HOBr}^{\bullet-}$ could balance to some $\bullet\text{OH}$ production.
399 Other possibility would be the direct hydroxylation performed by some of the RHS as $\text{HOCl}^{\bullet-}$,
400 $\text{HOBr}^{\bullet-}$, HOCl/OCl^- and HOBr/OBr^- or even by some secondary reaction involving the $\text{O}_2^{\bullet-}$. In
401 any case, 10 oxidized TPs were detected (Fig. 5, green arrows). The primary intermediate SMX-
402 NO may result from the oxidation of the phenylamine group of the SMX molecule to nitroso
403 benzene moiety, which may be conducted by HOCl/OCl^- and HOBr/OBr^- as suggested in
404 previous studies (Gao et al., 2014). The oxidation of the phenylamine amine group is
405 supported by the fragment ion detected of 169.99 m/z for $(\text{M}-\text{H})^-$ (Fig. S2 of the
406 Supplementary Information).

407 Formation of the primary intermediate SMX-OH may be apparently performed through
408 hydroxylation of the phenylamine moiety, was corroborated through the identification of the
409 ion fragment of 172.01 m/z for $(\text{M}-\text{H})^-$ (Fig. S3 of the Supplementary Information). The OH
410 addition probably occurred in the phenyl group, as other authors concluded in previous studies
411 (Dodd and Huang, 2004; Gao et al., 2014; Yuan et al., 2019). Additionally, and even though it
412 was not a primary TP, SMX-NCN was rapidly generated and reached the maximum

413 concentration after 120 min of irradiation (see Fig. 4B). The hydroxylation and fragmentation
414 of the isoxazole ring could thus have been part of the SMX-NCN generation.

415 According to Figure 5, four secondary TPs generated from intermediate SMX-NO were
416 identified (SMX-NO₂, SMX-Br-NO, SMX-NO-NCN and SMX-NO₂-NCN). Formation of the species
417 SMX-NO₂ might be attributed to the oxidation of the nitrosobenzene to nitrobenzene moiety,
418 which was in fact corroborated by the detected ion fragment of 185.99 m/z for (M-H)⁻ (Fig. S4
419 of the Supplementary Information). Bromination of the nitrobenzene ring could generate SMX-
420 Br-NO TP, and the subsequent hydroxylation and fragmentation of the isoxazole ring might
421 have generated the SMX-NO-NCN TP. Both SMX-NO₂ and SMX-NO-NCN are precursors of the
422 intermediate SMX-NO₂-NCN, respectively by their hydroxylation/fragmentation and oxidation.
423 From the primary intermediate SMX-OH, two main secondary TPs were identified (SMX-Br-OH
424 and SMX-SO₃H). The intermediate SMX-Br-OH was formed from the bromination of the
425 nitrogen atom of the phenyl amine moiety, a common product within the oxidation pathway.
426 On the other hand, SMX-SO₃H may be generated after hydroxylation followed by
427 fragmentation of the phenyl moiety. In the case of the SMX-NCN intermediate, three
428 secondary TPs were detected (SMX-Br-NCN, SMX-NO-NCN and SMX-NO₂-NCN). SMX-Br-NCN is
429 a common intermediate in the halogenation pathway, formed by bromination of the
430 phenylamine nitrogen moiety. SMX-NO-NCN generation might be attributed to the oxidation
431 of the SMX-NCN phenylamine nitrogen to the nitrosobenzene group. Then, the generated
432 SMX-NO-NCN can be further oxidized from the nitrosobenzene group to nitrobenzene group,
433 forming the SMX-NO₂-NCN intermediate.

434

435 As can be observed in Fig. 4A, Fig. 4B and Fig. 4C, primary halogenated TPs (SMX-Cl and SMX-
436 Br), dimeric 2SMX(-) TP, and oxidized TPs (SMX-OH, SMX-NCN and SMX-NO) presented their
437 maximum concentration at 120 min of sunlight irradiation, while secondary TPs accumulated
438 along the 6 hours of SMX degradation process. Detection of sulfamethoxazole intermediates

439 through different transformation pathways contributed to demonstrate the generation of RHS
440 induced by solar irradiation of TiO₂ nanosized particles dispersed in seawater. Moreover, the
441 proposed transformation mechanisms support the possible halogenation and dimerization of
442 dissolved organic matter and micropollutants by photocatalytic treatments.

443

444 **Conclusions**

445 Results presented in this work demonstrate the photogeneration of reactive halogen species
446 through the irradiation of TiO₂ NPs with simulated sunlight in synthetic seawater. Selected
447 scavenging tests and the identification of generated intermediates allowed the proposal of the
448 TiO₂ photoactivation mechanism in this water matrix, this suggesting that RHS may result from
449 the direct reactivity of halides with photogenerated holes. The efficiency of these active
450 species in the degradation of organic pollutants, represented in this work by the model
451 compound SMX, may be markedly slower compared with the reactive oxygen species
452 generated in deionized water, which mainly involves hydroxyl radical oxidation. The
453 photocatalytic SMX transformation in seawater occurred by halogenation, oxidation and
454 dimerization as primary pathways, through reactions between this compound and the RHS
455 generated in the TiO₂ photo-activation process. The evidence of RHS generation in the
456 photocatalytic process studied in this work brings concerns about the feasibility of
457 photocatalytic water treatments using TiO₂ NPs in saline waters, as RHS could later lead to the
458 generation and accumulation of potentially toxic compounds, such as halogenated and
459 dimerized transformation products.

460

461 **Acknowledgements**

462 This work was financially supported by MINECO/FEDER, UE (project CTQ2017-86466-R), Oriol
463 Porcar's FPI fellowship (PRE2018-084184) and the agency for Management of University and
464 Research Grants of the Government of Catalonia (project 2017SGR131).

465

466 **References**

- 467 An, T., Yang, H., Li, G., Song, W., Cooper, W.J., Nie, X., 2010. Kinetics and mechanism of
468 advanced oxidation processes (AOPs) in degradation of ciprofloxacin in water. *Appl.*
469 *Catal. B Environ.* 94, 288–294. <https://doi.org/10.1016/j.apcatb.2009.12.002>
- 470 ASTM D1141-98, 2013. Standard for the Preparation of Substitute Ocean Water, ASTM
471 International, West Conshohocken, PA, 2013. <https://doi.org/10.1520/D1141-98R13>
- 472 Bancirova, M., 2011. Sodium azide as a specific quencher of singlet oxygen during
473 chemiluminescent detection by luminol and Cypridina luciferin analogues. *Luminescence*
474 26, 685–688. <https://doi.org/10.1002/bio.1296>
- 475 Bielski, B.H.J., Cabelli, D.E., Arudi, R.L., Ross, A.B., 1985. Reactivity of HO₂/O₂⁻ Radicals in
476 Aqueous Solution. *J. Phys. Chem. Ref. Data* 14, 1041–1100.
477 <https://doi.org/10.1063/1.555739>
- 478 Borowska, E., Gomes, J.F., Martins, R.C., Quinta-Ferreira, R.M., Horn, H., Gmurek, M., 2019.
479 Solar Photocatalytic Degradation of Sulfamethoxazole by TiO₂ Modified with Noble
480 Metals. *Catalysts* 9, 500.
- 481 Boutiti, A., Zouaghi, R., Bendjabeur, S.E., Guittonneau, S., Sehili, T., 2017. Photodegradation of
482 1-hexyl-3-methylimidazolium by UV/H₂O₂ and UV/TiO₂: Influence of pH and chloride. *J.*
483 *Photochem. Photobiol. A Chem.* 336, 164–169.
484 <https://doi.org/10.1016/j.jphotochem.2016.12.030>
- 485 Bustos, N., Cruz-Alcalde, A., Iriel, A., Fernández Cirelli, A., Sans, C., 2019. Sunlight and UVC-254
486 irradiation induced photodegradation of organophosphorus pesticide dichlorvos in

487 aqueous matrices. *Sci. Total Environ.* 649, 592–600.
488 <https://doi.org/10.1016/j.scitotenv.2018.08.254>

489 Buxton, G. V., Greenstock, C.L., Helman, W.P., Ross, A.B., 1988. Critical Review of rate
490 constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals
491 ($\cdot\text{OH}/\cdot\text{O}^-$) in Aqueous Solution. *J. Phys. Chem. Ref. Data* 17, 513–886.
492 <https://doi.org/10.1063/1.555805>

493 Byrne, C., Subramanian, G., Pillai, S.C., 2018. Recent advances in photocatalysis for
494 environmental applications. *J. Environ. Chem. Eng.* 6, 3531–3555.
495 <https://doi.org/10.1016/j.jece.2017.07.080>

496 Cai, Q., Hu, J., 2017. Decomposition of sulfamethoxazole and trimethoprim by continuous
497 UVA/LED/TiO₂ photocatalysis: Decomposition pathways, residual antibacterial activity
498 and toxicity. *J. Hazard. Mater.* 323, 527–536.
499 <https://doi.org/10.1016/j.jhazmat.2016.06.006>

500 Catalán, J., Díaz, C., Barrio, L., 2004. Analysis of mixed solvent effects on the properties of
501 singlet oxygen ($^1\Delta_g$). *Chem. Phys.* 300, 33–39.
502 <https://doi.org/10.1016/j.chemphys.2004.01.010>

503 Cavalcante, R.P., Dantas, R.F., Bayarri, B., González, O., Giménez, J., Esplugas, S., Machulek, A.,
504 2016. Photocatalytic mechanism of metoprolol oxidation by photocatalysts TiO₂ and TiO₂
505 doped with 5% B: Primary active species and intermediates. *Appl. Catal. B Environ.* 194,
506 111–122. <https://doi.org/10.1016/j.apcatb.2016.04.054>

507 D. Camacho-Muñoz, L.L. and C.E., 2020. Degradation of okadaic acid in seawater by UV/TiO₂
508 photocatalysis – Proof of concept. *Sci. Total Environ.*
509 <https://doi.org/10.1016/j.colsurfa.2020.124658>

510 Daimon, T., Hirakawa, T., Kitazawa, M., Suetake, J., Nosaka, Y., 2008. Formation of singlet
511 molecular oxygen associated with the formation of superoxide radicals in aqueous
512 suspensions of TiO₂ photocatalysts. *Appl. Catal. A Gen.* 340, 169–175.

513 <https://doi.org/10.1016/j.apcata.2008.02.012>

514 Dodd, M.C., Huang, C.H., 2004. Transformation of the antibacterial agent sulfamethoxazole in
515 reactions with chlorine: Kinetics, mechanisms, and pathways. *Environ. Sci. Technol.* 38,
516 5607–5615. <https://doi.org/10.1021/es035225z>

517 Doudrick, K., Yang, T., Hristovski, K., Westerhoff, P., 2013. Photocatalytic nitrate reduction in
518 water: Managing the hole scavenger and reaction by-product selectivity. *Appl. Catal. B*
519 *Environ.* 136–137, 40–47. <https://doi.org/10.1016/j.apcatb.2013.01.042>

520 Fotiou, T., Triantis, T.M., Kaloudis, T., O’Shea, K.E., Dionysiou, D.D., Hiskia, A., 2016.
521 Assessment of the roles of reactive oxygen species in the UV and visible light
522 photocatalytic degradation of cyanotoxins and water taste and odor compounds using C-
523 TiO₂. *Water Res.* 90, 52–61. <https://doi.org/10.1016/j.watres.2015.12.006>

524 Fujishima, A., Zhang, X., Tryk, D.A., 2008. TiO₂ photocatalysis and related surface phenomena.
525 *Surf. Sci. Rep.* 63, 515–582. <https://doi.org/10.1016/j.surfrep.2008.10.001>

526 Gao, S., Zhao, Z., Xu, Y., Tian, J., Qi, H., Lin, W., Cui, F., 2014. Oxidation of sulfamethoxazole
527 (SMX) by chlorine, ozone and permanganate-A comparative study. *J. Hazard. Mater.* 274,
528 258–269. <https://doi.org/10.1016/j.jhazmat.2014.04.024>

529 García-Fernández, I., Fernández-Calderero, I., Inmaculada Polo-López, M., Fernández-Ibáñez,
530 P., 2015. Disinfection of urban effluents using solar TiO₂ photocatalysis: A study of
531 significance of dissolved oxygen, temperature, type of microorganism and water matrix.
532 *Catal. Today* 240, 30–38. <https://doi.org/10.1016/j.cattod.2014.03.026>

533 Grebel, J.E., Pignatello, J.J., Song, W., Cooper, W.J., Mitch, W.A., 2009. Impact of halides on the
534 photobleaching of dissolved organic matter. *Mar. Chem.* 115, 134–144.
535 <https://doi.org/10.1016/j.marchem.2009.07.009>

536 Hao, Z., Yin, Y., Wang, J., Cao, D., Liu, J., 2018. Formation of organobromine and organoiodine
537 compounds by engineered TiO₂ nanoparticle-induced photohalogenation of dissolved
538 organic matter in environmental waters. *Sci. Total Environ.* 631–632, 158–168.

539 <https://doi.org/10.1016/j.scitotenv.2018.03.027>

540 Jedsukontorn, T., Ueno, T., Saito, N., Hunsom, M., 2018. Mechanistic aspect based on the role
541 of reactive oxidizing species (ROS) in macroscopic level on the glycerol photooxidation
542 over defected and defected-free TiO₂. *J. Photochem. Photobiol. A Chem.* 367, 270–281.
543 <https://doi.org/10.1016/j.jphotochem.2018.08.030>

544 Jiang, J., Zhang, X., Zhu, X., Li, Y., 2017. Removal of Intermediate Aromatic Halogenated DBPs
545 by Activated Carbon Adsorption: A New Approach to Controlling Halogenated DBPs in
546 Chlorinated Drinking Water. *Environ. Sci. Technol.* 51, 3435–3444.
547 <https://doi.org/10.1021/acs.est.6b06161>

548 Liu, J., Zhang, X., 2014. Comparative toxicity of new halophenolic DBPs in chlorinated saline
549 wastewater effluents against a marine alga: Halophenolic DBPs are generally more toxic
550 than haloaliphatic ones. *Water Res.* 65, 64–72.
551 <https://doi.org/10.1016/j.watres.2014.07.024>

552 Mahmoud, W.M.M., Rastogi, T., Kümmerer, K., 2017. Application of titanium dioxide
553 nanoparticles as a photocatalyst for the removal of micropollutants such as
554 pharmaceuticals from water. *Curr. Opin. Green Sustain. Chem.* 6, 1–10.
555 <https://doi.org/10.1016/j.cogsc.2017.04.001>

556 Martins, R.C., Domingues, E., Bosio, M., Quina, M.J., Gmurek, M., Quinta-ferreira, R.M.,
557 Gomes, J., 2019. Effect of Different Radiation Sources and Noble Metal Doped onto TiO₂
558 for Contaminants of Emerging Concern Removal. *Water* 11, 894.
559 <https://doi.org/10.3390/w11050894>

560 Nosaka, Y., Nosaka, A.Y., 2017. Generation and Detection of Reactive Oxygen Species in
561 Photocatalysis. *Chem. Rev.* 117, 11302–11336.
562 <https://doi.org/10.1021/acs.chemrev.7b00161>

563 Riaz, S., Park, S.J., 2020. An overview of TiO₂-based photocatalytic membrane reactors for
564 water and wastewater treatments. *J. Ind. Eng. Chem.* 84, 23–41.

565 <https://doi.org/10.1016/j.jiec.2019.12.021>

566 Ribao, P., Corredor, J., Rivero, M.J., Ortiz, I., 2019. Role of reactive oxygen species on the
567 activity of noble metal-doped TiO₂ photocatalysts. *J. Hazard. Mater.* 372, 45–51.
568 <https://doi.org/10.1016/j.jhazmat.2018.05.026>

569 Rodríguez, E.M., Márquez, G., Tena, M., Álvarez, P.M., Beltrán, F.J., 2014. Determination of
570 main species involved in the first steps of TiO₂ photocatalytic degradation of organics
571 with the use of scavengers: The case of ofloxacin. *Appl. Catal. B Environ.* 178, 44–53.
572 <https://doi.org/10.1016/j.apcatb.2014.11.002>

573 Romero-Martínez, L., Moreno-Andrés, J., Acevedo-Merino, A., Nebot, E., 2014. Improvement
574 of ballast water disinfection using a photocatalytic (UV-C + TiO₂) flow-through reactor for
575 saltwater treatment. *J. Chem. Technol. Biotechnol.* 89, 1203–1210.
576 <https://doi.org/10.1002/jctb.4385>

577 Rubio, D., Casanueva, J.F., Nebot, E., 2013. Improving UV seawater disinfection with
578 immobilized TiO₂: Study of the viability of photocatalysis (UV₂₅₄/TiO₂) as seawater
579 disinfection technology. *J. Photochem. Photobiol. A Chem.* 271, 16–23.
580 <https://doi.org/10.1016/j.jphotochem.2013.08.002>

581 Tiwari, A., Shukla, A., Lalliansanga, Tiwari, D., Lee, S.M., 2019. Au-nanoparticle/nanopillars TiO₂
582 meso-porous thin films in the degradation of tetracycline using UV-A light. *J. Ind. Eng.*
583 *Chem.* 69, 141–152. <https://doi.org/10.1016/j.jiec.2018.09.027>

584 von Sonntag, C., von Gunten, U., 2012. *Chemistry of Ozone in Water and Wastewater*
585 *Treatment: From Basic Principles to Applications*, IWA Publishing.
586 <https://doi.org/10.2166/9781780400839>

587 Xiao, J., Xie, Y., Han, Q., Cao, H., Wang, Y., Nawaz, F., Duan, F., 2016. Superoxide radical-
588 mediated photocatalytic oxidation of phenolic compounds over Ag⁺/TiO₂: Influence of
589 electron donating and withdrawing substituents. *J. Hazard. Mater.* 304, 126–133.
590 <https://doi.org/10.1016/j.jhazmat.2015.10.052>

591 Yuan, R., Zhu, Y., Zhou, B., Hu, J., 2019. Photocatalytic oxidation of sulfamethoxazole in the
592 presence of TiO₂: Effect of matrix in aqueous solution on decomposition mechanisms.
593 Chem. Eng. J. 359, 1527–1536. <https://doi.org/10.1016/j.cej.2018.11.019>

594 Zhang, K., Parker, K.M., 2018. Halogen Radical Oxidants in Natural and Engineered Aquatic
595 Systems. Environ. Sci. Technol. 52, 9579–9594. <https://doi.org/10.1021/acs.est.8b02219>

596 Zhang, Y., Li, J., Bai, J., Li, L., Xia, L., Chen, S., Zhou, B., 2017. Dramatic enhancement of organics
597 degradation and electricity generation via strengthening superoxide radical by using a
598 novel 3D AQS/PPy-GF cathode. Water Res. 125, 259–269.
599 <https://doi.org/10.1016/j.watres.2017.08.054>

600 Zheng, S., Cai, Y., O’Shea, K.E., 2010. TiO₂ photocatalytic degradation of phenylarsonic acid. J.
601 Photochem. Photobiol. A Chem. 210, 61–68.
602 <https://doi.org/10.1016/j.jphotochem.2009.12.004>

603