

Activation of p21 limits acute lung injury and induces early senescence after acid aspiration and mechanical ventilation

Jorge Blázquez-Prieto , Covadonga Huidobro , Inés López-Alonso ,
Laura Amado-Rodríguez , Paula Martín-Vicente ,
Cecilia López-Martínez , Irene Crespo , Cristina Pantoja ,
Pablo J Fernandez-Marcos , Manuel Serrano , Jacob I Sznajder ,
Guillermo M Albaiceta



PII: S1931-5244(21)00022-0
DOI: <https://doi.org/10.1016/j.trsl.2021.01.008>
Reference: TRSL 1515

To appear in: *Translational Research*

Received date: 14 August 2020
Revised date: 13 December 2020
Accepted date: 21 January 2021

Please cite this article as: Jorge Blázquez-Prieto , Covadonga Huidobro , Inés López-Alonso ,
Laura Amado-Rodríguez , Paula Martín-Vicente , Cecilia López-Martínez , Irene Crespo ,
Cristina Pantoja , Pablo J Fernandez-Marcos , Manuel Serrano , Jacob I Sznajder ,
Guillermo M Albaiceta , Activation of p21 limits acute lung injury and induces early senescence
after acid aspiration and mechanical ventilation, *Translational Research* (2021), doi:
<https://doi.org/10.1016/j.trsl.2021.01.008>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Activation of p21 limits acute lung injury and induces early senescence after acid aspiration and mechanical ventilation

Jorge Blázquez-Prieto¹, Covadonga Huidobro^{1,2}, Inés López-Alonso^{1,2,3}, Laura Amado-Rodríguez^{1,2,3,4}, Paula Martín-Vicente^{1,2,3}, Cecilia López-Martínez^{1,2,3}, Irene Crespo⁵, Cristina Pantoja⁶, Pablo J Fernandez-Marcos⁶, Manuel Serrano⁷, Jacob I Sznajder⁸, Guillermo M Albaiceta^{1,2,3,4,5}

¹*Instituto de Investigación Sanitaria del Principado de Asturias. Oviedo, Spain.*

²*Centro de Investigación Biomédica en Red (CIBER)-Enfermedades Respiratorias. Madrid, Spain.* ³*Instituto Universitario de Oncología del Principado de Asturias. Oviedo, Spain.* ⁴*Unidad de Cuidados Intensivos Cardiológicos. Hospital Universitario Central de Asturias. Oviedo, Spain.* ⁵*Departamento de Biología Funcional. Universidad de Oviedo. Oviedo, Spain.* ⁶*Metabolic Syndrome Group - BIOPROMET, Madrid Institute for Advanced Studies - IMDEA Food, CEI UAM+CSIC. Madrid, Spain.* ⁷*Institute for Research in Biomedicine (IRB Barcelona); Barcelona Institute of Science and Technology (BIST); Catalan Institution for Research and Advanced Studies (ICREA); Barcelona, Spain.* ⁸*Division of Pulmonary and Critical Care Medicine, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA.*

Author for correspondence:

Guillermo M Albaiceta

Instituto de Investigación Biosanitaria del Principado de Asturias

Avenida de Roma s/n

33011 Oviedo. Spain.

E-mail: gma@crit-lab.org

Running head: Effects of p21 in acute lung injury

Abbreviations:

ANOVA: Analysis of the variance

ARDS: Acute respiratory distress syndrome

SAHF: Senescence associated heterochromatin foci

SASP: Senescence-associated secretory phenotype

TUNEL: Terminal deoxynucleotidyl transferase dUTP nick end labeling

VILI: Ventilator-induced lung injury

Journal Pre-proof

Abstract

The p53/p21 pathway is activated in response to cell stress. However, its role in acute lung injury has not been elucidated. Acute lung injury is associated with disruption of the alveolo-capillary barrier leading to acute respiratory distress syndrome (ARDS). Mechanical ventilation may be necessary to support gas exchange in patients with ARDS, however, high positive airway pressures can cause regional overdistension of alveolar units and aggravate lung injury. Here, we report that acute lung injury and alveolar overstretching activate the p53/p21 pathway to maintain homeostasis and avoid massive cell apoptosis. A systematic pooling of transcriptomic data from animal models of lung injury demonstrates the enrichment of specific p53- and p21-dependent gene signatures and a validated senescence profile. In a clinically relevant, murine model of acid aspiration and mechanical ventilation, we observed changes in the nuclear envelope and the underlying chromatin, DNA damage and activation of the *Tp53/p21* pathway. Absence of *Cdkn1a* decreased the senescent response, but worsened lung injury due to increased cell apoptosis. Conversely, treatment with lopinavir/ritonavir led to *Cdkn1a* overexpression and ameliorated cell apoptosis and lung injury. The activation of these mechanisms was associated with early markers of senescence, including expression of senescence-related genes and increases in senescence-associated heterochromatin foci in alveolar cells. Autopsy samples from lungs of patients with ARDS revealed increased senescence-associated heterochromatin foci. Collectively, these results suggest that acute lung injury activates p53/p21 as an anti-apoptotic mechanism to ameliorate damage, but with the side effect of induction of senescence.

Keywords: Acute lung injury; Mechanical ventilation; Senescence; Apoptosis

Introduction

The lungs have a stereotypic response to acute injury, which is preserved among species and many etiological agents. Once damage is inflicted, lung cells trigger a host response which can include inflammation, matrix remodeling and different forms of cell death, including apoptosis ¹. Although a limited host response may help to clear the injurious agent and promote lung tissue repair ², an overexuberant host response can lead to severe injury and gas exchange worsening. Therefore, therapeutic strategies aimed to limit lung damage and interference with lung repair are important.

Lungs are exposed to mechanical load during every breath. In pathologic conditions, generation of higher pressure gradients necessary for adequate ventilation may cause excessive cell stretch ³. This is especially relevant during mechanical ventilation with high pressures, which can lead to the so-called ventilator-induced lung injury (VILI) ^{4,5}. In mechanically ventilated patients, a strategy aimed to limit VILI decreased mortality in patients with the acute respiratory distress syndrome (ARDS)⁶.

Mechanotransduction is thought to regulate the molecular steps in VILI pathogenesis ⁷. The nuclear envelope has been reported as an important cell mechanosensor and signal transducer ⁸. Mechanical stretch appears to increase Lamin-A in the nuclear envelope, leading to nuclear stiffening. These changes in the nuclear envelope can also activate p53-dependent pathways. Wildtype p53 is a master regulator of cell homeostasis and fate, and its activation may lead to a variety of responses, ranging from apoptosis to cell cycle arrest. Inhibition of this response has been shown to increase p21 (*Cdkn1a*) expression and decrease VILI in an experimental model ⁹.

p53 and its downstream factor p21 are triggers of senescence ¹⁰, a cell response characterized by a stable arrest of the cell cycle and a switch towards a senescence-associated secretory phenotype (SASP). It has been proposed that senescence facilitates the clearance of damaged cells and is required for tissue repair ¹¹. Interestingly, some of the molecules have a significant overlap with the proinflammatory response associated with VILI.

We hypothesized that p53-dependent pathways play a role in the maintenance of lung homeostasis during acute injury, and that senescence could be a side effect of their activation. To test this hypothesis, we developed a clinically relevant model of lung injury caused by acid aspiration and VILI to assess the activation of p53 and its downstream factors.

Material and methods

Meta-analysis of transcriptomic data

To explore the main hypothesis, a pooled analysis of published transcriptomic data was performed, using a previously validated 55-gene expression signature of senescence ¹² as main endpoint. Datasets reporting lung gene expression in mouse models of acute lung injury and mechanical ventilation were obtained from public repositories (Gene Omnibus Expression -<https://www.ncbi.nlm.nih.gov/geo/>- and ArrayExpress - <https://www.ebi.ac.uk/arrayexpress/>-) using the following terms: “Stretch”, “Cyclic strain”, “Mechanical Ventilation”, “Lung”, and “Alveolar”. Fifty-one datasets were manually reviewed. Studies lacking a control group with intact, spontaneously breathing animals and those reporting less than 40 genes from the endpoint signature were excluded, so 9 datasets were finally used (Supplementary Table 1). When available, raw data was downloaded and normalized using the

Robust Multiarray Average method (for Affymetrix microarrays) or normal-exponential background correction followed by quantile normalization (all the other platforms).

Normalized datasets were pooled using the Combat-Co-normalization using controls (COCONUT) algorithm¹³. This method normalizes gene expression of the different datasets using an empirical Bayes fitting, but applied only to control samples (in this case, spontaneously breathing animals with intact lungs). Then the obtained normalization parameters are applied to the cases (i.e., those with lung injury)(Supplementary Figure 1). Three different signatures were studied, corresponding to 116 genes upregulated by p53, 14 genes downregulated by p21¹⁴ and a set of 50 genes consistently up- and down-regulated in senescence¹². A meta-score was computed for each sample as the geometric mean of the upregulated genes minus the geometric mean of the downregulated genes in the signature¹³. Meta-scores were finally compared among controls and animals with lung injury and/or mechanical ventilation.

Animal models

Male, 12 week old C57Bl/6 mice, kept under pathogen-free conditions with free access to food and water, were used in all experiments. The Animal Research Committee of the Universidad de Oviedo evaluated and approved the study.

A two-hit lung injury model, based on chlorhydric acid instillation and mechanical ventilation, was studied. Animals were anesthetized with intraperitoneal ketamine and xylazine and orotracheally intubated using a 20G catheter, through which 50 μ L of chlorhydric acid (0.1N, pH=1.5) were instilled. Two hours after instillation, mice were randomly assigned to receive mechanical ventilation or not. Mice were

ventilated with a pressure-controlled mode (peak inspiratory pressure 17 cmH₂O, PEEP 2 cmH₂O, respiratory rate 100 breaths/min) for 120 minutes.

Three additional series of experiments were performed. Mice lacking *Tp53* or *Cdkn1a* (p21, an endogenous inhibitor of cyclin-dependent kinases involved in the senescent response triggered by *Tp53*) and their wildtype littermates were subjected to the same model of injury, including acid instillation and mechanical ventilation. Genotypes were confirmed by PCR. In separate experiments, wildtype animals were treated with a single dose (200/50 mg/Kg) of lopinavir/ritonavir (a protease inhibitor that inhibits Zmpste24 and disrupts Lamin-A nuclear scaffolding, activating senescence pathways) or saline, administered intraperitoneally immediately after acid instillation, and then ventilated with the parameters described above.

Tissue harvest

Mice were studied in three different conditions: baseline, 4 hours after chlorhydric acid instillation without mechanical ventilation and 4 hours after acid instillation including 2 hours of mechanical ventilation. Lungs were removed after exsanguination of anesthetized animals. A laparotomy was performed, the renal artery sectioned, the thorax opened and the heart-lungs removed *in bloc*. The left lung was instilled with 250 microliters of 4% phosphate-buffered paraformaldehyde, immersed in the same fixative for 24 hours, and then stored in 50% ethanol. The right lung was immediately frozen at -80°C for biochemical analyses.

Patient samples

Paraffin-embedded lung tissue from autopsies of patients were obtained from the tissue bank at Hospital Universitario Central de Asturias, after signed consent from patients' next of kin. ARDS was defined using the Kigali modification of the Berlin definition ¹⁵, to include patients with lung injury but without mechanical ventilation and those without an arterial line. 13 samples were recovered (Supplementary Table 2).

Histological studies

After fixation, tissues were embedded in paraffin and three slices with at least 1mm of separation between them were cut and stained with hematoxylin and eosin. A pathologist blinded to the experimental settings evaluated the degree and extension of lung damage using a predefined histological score ¹⁶.

Additional lung sections were processed as previously described ¹⁷ for detection of myeloperoxidase- and Ki-67-positive cells, using specific antibodies (See Supplementary Table 3 for references). Images from three random fields (x200) were taken and then number of positive cells averaged.

For immunofluorescence studies, slides were deparaffinated and antigens retrieved in citrate buffer 0.1M (pH=9). The autofluorescence of the tissue was diminished using a Sudan black B solution and sections were permeabilized (0.1% Triton X-100 in PBS for 15 minutes), blocked (1% BSA in PBS) and incubated overnight at 4°C with the primary antibody (Supplementary Table 3). After 24 hours, the slices were incubated with the corresponding secondary fluorescent antibody at room temperature for 1 hour. Images were taken using a confocal microscopy (Leica SP8) at 400x and 630x. The number of positive and negative nuclei were automatically quantified using ImageJ software (NIH, USA).

Apoptotic cells in lung slices were detected by terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) as previously described ⁹. Images from three random fields were acquired in a Leica SP8 confocal microscope and the positive nuclei were counted and expressed as percentage of the total nuclei count.

Western blot

Nuclei were extracted from fresh lung tissues and subsequently homogenized as described before ⁹. The total amount of protein from nuclear extracts was quantified (BCA Protein Assay Kit, Pierce) and 15 μ g of each sample were loaded in SDS-polyacrylamide gels, electrophoresed at 120mV and electrotransferred onto PVDF membranes. After blockade with 5% non-fat dry milk, the membranes were incubated with primary antibodies against Caspase-9, Lamin-A/C, Lamin-B1, γ H2AX, HP1 α or H3 (Supplementary Table 3) in 3% non-fat dry milk overnight at 4°C. After 24 hours, the membranes were incubated with the corresponding peroxidase-conjugated secondary antibodies in 2.5% non-fat dry milk. Proteins were detected by chemiluminescence in a LAS-4000 Imaging system. The intensity of each protein band was quantified using ImageJ software (NIH, USA).

Quantitative PCR

Lung fragments (2 mm x 2 mm) were homogenized with TRIZOL (Sigma, Poole, UK) and RNA precipitated by overnight incubation in isopropanol at -20°C. After 24 hours, samples were washed with ethanol and the RNA resuspended in RNase-free water and quantified. One μ g of total RNA was retrotranscribed into complementary cDNA using an RT-PCR kit (High capacity cDNA rt Kit, Applied

Biosystems). Quantitative PCRs were carried out in triplicate of each sample using 40 ng of cDNA per well. Expression of *Plk3*, *Gdnf*, *Meis1*, *Il6*, *Tp53*, *Cdkn1a* (p21), *Cdkn2a* (p16), *Rb*, and *Gapdh* was quantified using Sybr-green Power up, (Fisher Scientific) and 10uM of the corresponding primers (Supplementary Table 4). The relative expression of each gene was calculated as $2^{-\Delta\text{CT}(\text{gene of interest})-\Delta\text{CT}(\text{GAPDH})}$.

Statistical analysis

Data are shown as mean \pm standard error of the mean. Differences between two groups were studied using a T test. Differences among more than two groups were assessed using an analysis of the variance (ANOVA). For SAHF counts, three slides per animal were counted (considered as technical replicates) and analyzed using a mixed-effects ANOVA. When significant, pairwise comparisons were done using the Tukey's Honest Significant Difference test. A p-value lower than 0.05 was considered significant.

Results

Transcriptomic signatures of p53/p21 activation and senescence in lung injury

To test the hypothesis that the p53/p21 pathway is activated during acute lung injury and to identify early markers of a switch towards a senescent phenotype, data from 9 datasets of mouse lung injury and mechanical ventilation (Supplementary Table 1) were pooled and gene expression analyzed (Figure 1A). Three different transcriptomic signatures related to p53-dependent upregulation (116 genes, 85 available in the pooled data), p21-dependent downregulation (14 genes, 12 available)¹⁴ and senescence¹² (55 genes, 44 available) were analyzed. A

meta-score of expression of these genes was computed for each sample and compared to assess the effect of lung injury and mechanical ventilation.

Animals subjected to acid aspiration lung injury and mechanical ventilation showed higher expression of p53-dependent genes (ANOVA p-value<0.001, Figure 1B), lower expression of p21-downregulated genes (ANOVA p-value<0.001, Figure 1C) and a higher metascore (Figure 1D) in the senescence signature than spontaneously breathing controls (ANOVA p-value<0.001, Figure 1D). The expression of each gene of these signatures is shown in Supplementary Figure 2. These results support the notion that lung injury and mechanical ventilation activate p53/p21 pathways and the molecular mechanisms of senescence in acutely injured lungs.

Activation of the p53/p21 pathway in a clinically relevant model

To explore the mechanisms involved in the activation of p53-dependent signals, an experimental model of acid aspiration- and mechanical ventilation induced lung injury was tested. Chlorhydric acid instillation and mechanical ventilation induced a significant increase in lung damage and inflammation, assessed by histological scores (Figure 2A), neutrophilic infiltrates (Figure 2B) and *Il6* expression (Figure 2C). Lung damage increased the proportion of both proliferating and apoptotic cells in lung parenchyma (Figures 2D-E and Supplementary Figure 3). Immunohistochemical studies revealed that TUNEL-positive staining was localized in all the explored cell types, including type I and II pneumocytes, fibroblasts and endothelial cells (Supplementary Figure 4). In line with these findings, the abundance of cleaved caspase-9 in lung tissue was increased with lung injury (Figure 2F)

We then explored the putative activators of this response to acute injury. Lamins in the nuclear envelope act as cell mechanosensors, regulating chromatin organization in response to mechanical stress. We observed that Lamin-A/Lamin-B ratio increased after mechanical stretch (Figure 2G). Immunofluorescence studies confirmed the increase in Lamin-A in the nuclear envelope after mechanical stretch, but not after hydrochloric acid instillation alone (Figure 2H). These changes in the nuclear envelope coexisted with an increase in γ H2AX (Figure 2I) and HP1 α (Figure 2J), markers of DNA damage and chromatin remodeling respectively, in nuclear extracts from ventilated animals. Panel 2K shows representative Western blots of these parameters.

The expression of the canonical responders to DNA damage *Cdkn2a* (p16) and *Tp53* (p53) and their corresponding downstream factors *Rb* and *Cdkn1a* (p21) was also assessed. There were no differences in the levels of *Cdkn2a* (Figure 2L), whereas expression of *Rb* was significantly decreased (Figure 2M). However, we observed significant increases in *Tp53* (Figure 2N) and *Cdkn1a* (Figure 2O) expression with lung injury. The increase in P21 protein was observed mainly in type-I and type-II alveolar epithelial cells (positive for Aquaporin-5 or Surfactant-protein C respectively), and not in fibroblast or endothelial cells (positive for vimentin or Von-Willebrand factor respectively) (Supplementary Figure 5).

Increased lung damage in mice lacking p21

To address the role of p53 and p21 in acute lung damage, *Tp53*^{-/-}, *Cdkn1a*^{-/-} mice and their wildtype counterparts were subjected to acid instillation followed by mechanical ventilation. In preliminary experiments, absence of *Tp53* did not modify lung injury (histological score 2.4 \pm 1.6 vs 2.3 \pm 1.2, n=4/group, p=0.91,

Supplementary figure 6), so we focused on the downstream factor p21. In absence of p21, the mice had worse lung injury (Figure 3A) and higher counts of apoptotic cells (Figure 3B) and cleaved caspase-9 abundance (Figure 3C) compared to their wildtype counterparts. There were no differences in *Il6* or *Tp53* expression nor in abundance of γ H2AX or HP1 α between genotypes (Figures 3D-G respectively).

Lopinavir increases p21 and decreases lung damage

We have previously shown that HIV-protease inhibitors modify the nuclear response to mechanical stretch and protect against VILI⁹, an effect that could be due to the inhibition of the Lamin-A protease ZMPSTE24¹⁸. In our double-hit model, treatment with lopinavir/ritonavir impaired the structure of the nuclear lamina, decreasing the abundance of Lamin-A (Figure 4A), and decreased lung injury (Figure 4B), apoptotic cell count (Figure 4C) and cleaved caspase-9 (Figure 4D). Although abundance of γ H2Ax was not modified by this treatment (Figure 4E), there was a marked decrease in HP1 α (Figure 4F). Panel 4G shows representative blots of these measurements. Finally, treatment with lopinavir/ritonavir caused an increase in *Il6* expression (Figure 4H), with no changes in *Tp53* expression (Figure 4I) but an increase in *Cdkn1a* (p21, Figure 4J).

Early markers of senescence in acute lung injury

Then, we tried to identify early markers of senescence in our acute model. Acid aspiration and mechanical ventilation-induced lung injury was associated with an increase in the number of nuclei positive for Macro-H2A, a marker of senescence-associated heterochromatin foci (SAHF, Figure 5A), and changes in *Plk3* (Figure

5B), *Gdnf* (Figure 5C), and *Meis1* (Figure 5D), the genes from the senescence signature with the highest differential expression in the previous pooled analysis.

These markers of senescence were modified by manipulation of the p21 pathway.

Mutant animals lacking *Cdkn1a* exhibited a decreased number of SAHF after acid instillation and mechanical ventilation (Figure 5E) and in expression of the senescence-related gene *Plk3* (Figure 5F). In opposite, treatment with lopinavir/ritonavir (that increased *Cdkn1a* expression, Figure 4J) was related to lower counts of SAHF (Figure 5G), but increased *Plk3* expression (Figure 5H)

Finally, to confirm the incidence of SAHF in patients, lung tissue from autopsies of critically-ill patients with and without lung injury and mechanical ventilation (Supplementary Table 2) were stained with antibodies against Macro-H2A. There were no differences in age between the three groups of patients (62 ± 6 , 61 ± 11 and 54 ± 10 years for patients without ARDS or mechanical ventilation, with ARDS but without mechanical ventilation and ARDS and ventilation respectively, $p=0.17$ in ANOVA). Similarly to the animal model, nuclear Macro-H2A increased in those with severe lung injury and mechanical ventilation (Figure 5I).

Collectively, these findings suggest that lung injury and mechanical stretch trigger the appearance of early markers of senescence. The severity of lung injury and the abundance of senescence markers showed an inverse correlation after manipulation of p21 levels.

Discussion

We provide evidence that acute lung injury and its treatment with mechanical ventilation alters the nuclear envelope and causes DNA damage, activating the p53/p21 pathway. Activation of p21 plays a homeostatic role, limiting the extent of

apoptosis in response to injury. Moreover, this effect can be pharmacologically activated to ameliorate lung injury in a clinical setting. In spite of this beneficial effect, this pathway also leads to the appearance of early markers of senescence in lung tissue. Figure 6 summarizes the findings of this work.

The p53/p21 axis in acute injury

P53 and its downstream transcription factor p21 are major regulators of cell homeostasis. It has been shown that p53 regulates permeability in lung endothelial cells after an inflammatory insult ¹⁹. Similarly, activation of this pathway in response to hypertonic saline decreased lung injury and inflammation in human airway epithelial cells ²⁰. However, our observations in *Tp53*^{-/-} mice showed no differences in lung injury. Given the pleiotropic effects of p53 in cell homeostasis, this could be due to the existence of both protective and pathogenetic mechanisms. One of the main effects of the cyclin kinase inhibitor p21 is the blockade of apoptosis ²¹. Several pro-apoptotic pathways are activated during acute lung injury ²², as shown in our model. In this setting, p21 may have a compensatory role by avoiding the loss of a large amount of epithelial cells. Our observation of massive cell death in *Cdkn1a*^{-/-} mice suggests this homeostatic role. It has been proposed that caspase-9 is the downstream target responsible for the anti-apoptotic effects of p21 ²³. Overexpression of p21 increases the resistance to apoptosis of alveolar epithelial cells ²⁴, and the beneficial effects of lopinavir/ritonavir in VILI could represent the effects of the overexpression of this gene and the observed decrease in caspase-9. In contrast, absence of p21 was associated with more severe lung injury and increased numbers of apoptotic cells, as previously suggested ²⁵.

The role of mechanical stretch

Mechanical overstretch could be an important pathogenic factor involved in p53/p21 activation. The experimental model using a ventilatory strategy within the limits of protective ventilation (driving pressure 15 cmH₂O, avoidance of zero end-expiratory pressure) was chosen to increase the translational significance of our work. Although ventilation with very high inspiratory pressures and allowing expiratory collapse induces severe lung damage in less than one hour, even in healthy lungs, the clinical translation of these findings is less straightforward.

Several mechanisms have linked the mechanical load to a lung biological response, including oxidative stress and MAPK activation ²⁶. We focused on the role of the nuclear envelope as a critical structure regulating both mechanosensing and senescence. The mechanical load is transmitted from the extracellular matrix to the cytoskeleton and then to the nuclear membrane ²⁷. This causes a change in the nuclear lamina, reorganization of the underlying chromatin and DNA damage ²⁸, either mediated by MAPK activation ²⁹ or by a direct mechanical effect ³⁰. DNA damage is one of the triggers of the p53 pathway. In smooth muscle cells, stretch leads to p53 activation and upregulation of senescence markers ³¹, resembling our findings. Similarly, HIV protease inhibitors, such as lopinavir/ritonavir, inhibit ZMPSTE-24, a protease responsible for Lamin-A maturation ¹⁸, preserving nuclear compliance, increasing p21 expression and decreasing stretch induced apoptosis and VILI ⁹.

Our immunohistochemical studies suggest that although apoptosis occurs in all the explored cell lines (alveolar epithelium, endothelium and fibroblasts), activation of p21 pathway take place in the alveolar epithelium. Preexisting epithelial damage caused by acid instillation could amplify the mechanical load over the epithelium

by increasing heterogeneity of the lung parenchyma. Moreover, previous results in subacute models of lung damage have shown the need for two synergic hits to induce lung senescence ³².

Senescence in lung diseases

One of the known consequences of p53 activation is the cell switch towards a senescent phenotype. Lipopolysaccharide or bleomycin-induced lung injury increases the number of SA- β -galactosidase-positive cells and leads to cell cycle arrest ³³. It has been shown that this activation has no detrimental effects in acute inflammation. However, blockade of the cell cycle has been associated with increased collagen deposition ³⁴ and SASP may perpetuate lung inflammation ³⁵. Therefore, main features of abnormal lung repair after acute injury (limited cell proliferation, chronic inflammation and fibrosis) could be explained by a persistent senescent response ³⁶. Inhibition of this response by selective deletion of Tp53 in Club cells ameliorated lung damage related to chronic inflammation ³³, suggesting a novel mechanism amenable to treatment of lung diseases.

The acute nature of our model and its short-term lethality does not allow the identification of canonical senescence markers such as Senescence-associated β -galactosidase, as these require from days to weeks to be positive ³⁷. However, we identified a set of early markers including changes in chromatin structure and gene expression. In a model of repair after VILI, lung *Cdkn1a* expression remained elevated up to two days of spontaneous breathing after injury ⁹. Although it is unclear if these mechanisms may precipitate a full-blown senescent response in the long term, our results highlight the involvement of this molecular machinery in the early phase, and could be a therapeutic target to avoid late consequences.

Clinical implications

Our findings have several implications regarding the pathogenesis of lung injury and its long-term consequences. First, the described p21 response may be beneficial in the acute phase, and could be pharmacologically manipulated using lopinavir. In a recent clinical trial in patients with lung disease caused by the SARS-CoV-2 coronavirus, lopinavir did not reduce mortality, but decreased the risk of ARDS development ³⁸. However, the associated senescent response could worsen lung repair and long-term outcomes. Survivors after a prolonged ICU stay may have deleterious and prolonged sequels, including respiratory impairment ³⁹, neuropsychological disturbances ⁴⁰ and muscle atrophy ⁴¹, particularly the elderly ⁴². The mechanisms responsible for these sequels are largely unknown and no effective therapies are currently available. Local activation of senescence and its paracrine/systemic spread could contribute to the pathogenesis of these sequels ⁴³. As previously discussed, senescence may contribute to disordered lung repair. The confirmation of this framework could lead to the use of senolytics ^{44(p)} in critically ill patients. However, due to the protective nature of senescence in the early phase ⁴⁵, these treatments should be time-coordinated and modulated to optimize their effectiveness.

Conclusions

We provide new evidence suggesting that acute lung damage activates p21 to limit apoptosis. This response appears to be triggered by the induction of DNA damage and linked to chromatin changes caused by mechanical overstretch. Interaction with the nuclear lamina may enhance this anti-apoptotic response. Although p21

activation may be beneficial in the acute phase of lung injury, the long-term effects must be taken into consideration as they could explain some of the long term sequels of critically ill patients.

Acknowledgements

Funding: Supported by grants from Centro de Investigación Biomédica en Red (CIBER) and Instituto de Salud Carlos III (PI20/01360, Co-funded by European Regional Development Fund/European Social Fund). L.A.R. is the recipient of a grant from Instituto de Salud Carlos III (CM16/00128). C.H.F. is the recipient of a grant from Instituto de Salud Carlos III (CD16/00033). Instituto Universitario de Oncología del Principado de Asturias is supported by a grant from Fundación Cajastur-Liberbank. Work in the laboratory of P.J.F.M. was funded by the IMDEA Food Institute, by the AECC and Ramón Areces Foundations and by grants from the Spanish Ministry of Science, Innovation and Universities, co-funded by the European Regional Development Fund (ERDF) (SAF2017-85766-R and RYC-2017-22335). Work in the laboratory of M.S. was funded by the IRB, by “La Caixa” Foundation, and by grants from the Spanish Ministry of Economy co-funded by the European Regional Development Fund (ERDF) (SAF2017-82613-R) and from the European Research Council (ERC-2014-AdG/669622). J.I.S. is supported by NIH grants HL-147070, HL-71643 and AG-49665.

Conflicts of interest: All authors have read the journal’s authorship agreement and policy on disclosure of potential conflicts of interest. The authors have no conflicts of interest to disclose.

Brief commentary:

Background: P53 and its downstream factor p21 are involved in the cell response to stress. In addition, they are drivers of senescence. Although these pathways have been studied in chronic lung diseases, their role in acute lung injury has not been systematically explored.

Translational significance: Our findings show that the anti-apoptotic effects of p21 counteract the pro-apoptotic response triggered by acute lung injury and mechanical ventilation. Drug-induced overexpression of p21 decreases lung injury in this setting. However, activation of p21 also leads to an early senescent response within the lung, that may favor long-term side effects.

Authors' contributions

JBP, CHF, ILA and GMA designed the study. JBP, CHF, ILA, LAR, PMV, CLM, IC, CP and PJFM performed the experiments. GMA did the bioinformatic analysis. JBP, ILA, CHF, LAR, PJFM and GMA analyzed the results. CHF, ILA, LAR, PJFM, MS, JIS and GMA discussed the significance of the results. JIS, MS and GMA wrote the paper. GMA is the responsible of the integrity of the whole work.

Data statement

Raw data and R code used in this work are available from the corresponding author (GMA) upon reasonable request.

References

1. Herold S, Gabrielli NM, Vadász I. Novel concepts of acute lung injury and alveolar-capillary barrier dysfunction. *Am J Physiol Lung Cell Mol Physiol*. 2013;305(10):L665-681. doi:10.1152/ajplung.00232.2013
2. Blazquez-Prieto J, Lopez-Alonso I, Amado-Rodriguez L, et al. Impaired lung

- repair during neutropenia can be reverted by matrix metalloproteinase-9. *Thorax*. 2018;73(4):321-330. doi:10.1136/thoraxjnl-2017-210105
3. Perlman CE, Bhattacharya J. Alveolar expansion imaged by optical sectioning microscopy. *J Appl Physiol*. 2007;103(3):1037-1044. doi:10.1152/jappphysiol.00160.2007
 4. Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med*. 2013;369(22):2126-2136. doi:10.1056/NEJMra1208707
 5. Corbridge TC, Wood LD, Crawford GP, Chudoba MJ, Yanos J, Sznajder JI. Adverse effects of large tidal volume and low PEEP in canine acid aspiration. *Am Rev Respir Dis*. 1990;142(2):311-315. doi:10.1164/ajrccm/142.2.311
 6. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. 2000;342:1301-1308.
 7. Spieth PM, Bluth T, Gama De Abreu M, Bacelis A, Goetz AE, Kieffmann R. Mechanotransduction in the lungs. *Minerva Anesthesiol*. 2014;80(8):933-941.
 8. Swift J, Ivanovska IL, Buxboim A, et al. Nuclear lamin-A scales with tissue stiffness and enhances matrix-directed differentiation. *Science*. 2013;341(6149):1240104. doi:10.1126/science.1240104
 9. López-Alonso I, Blázquez-Prieto J, Amado-Rodríguez L, et al. Preventing loss of mechanosensation by the nuclear membranes of alveolar cells reduces lung injury in mice during mechanical ventilation. *Sci Transl Med*. 2018;10(456):eaam7598. doi:10.1126/scitranslmed.aam7598
 10. Varela I, Cadinanos J, Pendas AM, et al. Accelerated ageing in mice deficient in Zmpste24 protease is linked to p53 signalling activation. *Nature*. 2005;437(7058):564-568. doi:10.1038/nature04019
 11. Muñoz-Espín D, Serrano M. Cellular senescence: from physiology to pathology. *Nat Rev Mol Cell Biol*. 2014;15(7):482-496. doi:10.1038/nrm3823
 12. Hernandez-Segura A, de Jong TV, Melov S, Guryev V, Campisi J, Demaria M. Unmasking Transcriptional Heterogeneity in Senescent Cells. *Curr Biol CB*. 2017;27(17):2652-2660.e4. doi:10.1016/j.cub.2017.07.033
 13. Sweeney TE, Wong HR, Khatri P. Robust classification of bacterial and viral infections via integrated host gene expression diagnostics. *Sci Transl Med*. 2016;8(346):346ra91. doi:10.1126/scitranslmed.aaf7165
 14. Fischer M. Census and evaluation of p53 target genes. *Oncogene*. 2017;36(28):3943-3956. doi:10.1038/onc.2016.502
 15. Riviello ED, Kiviri W, Twagirumugabe T, et al. Hospital Incidence and Outcomes of the Acute Respiratory Distress Syndrome Using the Kigali Modification of the Berlin Definition. *Am J Respir Crit Care Med*. 2016;193(1):52-59. doi:10.1164/rccm.201503-0584OC
 16. Blázquez-Prieto J, López-Alonso I, Amado-Rodríguez L, Batalla-Solis E, González-López A, Albaiceta GM. Exposure to mechanical ventilation promotes tolerance to ventilator-induced lung injury by Ccl3 downregulation. *Am J Physiol Lung Cell Mol Physiol*. 2015;309(8):L847-56. doi:10.1152/ajplung.00193.2015
 17. González-López A, Astudillo A, García-Prieto E, et al. Inflammation and matrix remodeling during repair of ventilator-induced lung injury. *Am J Physiol Lung Cell Mol Physiol*. 2011;301(4):L500-9. doi:10.1152/ajplung.00010.2011
 18. Coffinier C, Hudon SE, Farber EA, et al. HIV protease inhibitors block the zinc metalloproteinase ZMPSTE24 and lead to an accumulation of prelamin A in cells. *Proc Natl Acad Sci U S A*. 2007;104(33):13432-13437.

doi:10.1073/pnas.0704212104

19. Barabutis N, Dimitropoulou C, Gregory B, Catravas JD. Wild-type p53 enhances endothelial barrier function by mediating RAC1 signalling and RhoA inhibition. *J Cell Mol Med.* 2018;22(3):1792-1804. doi:10.1111/jcmm.13460
20. Gamboni F, Anderson C, Mitra S, et al. Hypertonic Saline Primes Activation of the p53-p21 Signaling Axis in Human Small Airway Epithelial Cells That Prevents Inflammation Induced by Pro-inflammatory Cytokines. *J Proteome Res.* 2016;15(10):3813-3826. doi:10.1021/acs.jproteome.6b00602
21. Fielder E, von Zglinicki T, Jurk D. The DNA Damage Response in Neurons: Die by Apoptosis or Survive in a Senescence-Like State? *J Alzheimers Dis.* 2017;60(s1):S107-S131. doi:10.3233/JAD-161221
22. Martin TR, Hagimoto N, Nakamura M, Matute-Bello G. Apoptosis and epithelial injury in the lungs. *Proc Am Thorac Soc.* 2005;2(3):214-220.
23. Sohn D, Essmann F, Schulze-Osthoff K, Jänicke RU. p21 blocks irradiation-induced apoptosis downstream of mitochondria by inhibition of cyclin-dependent kinase-mediated caspase-9 activation. *Cancer Res.* 2006;66(23):11254-11262. doi:10.1158/0008-5472.CAN-06-1569
24. Inoshima I, Kuwano K, Hamada N, et al. Induction of CDK inhibitor p21 gene as a new therapeutic strategy against pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol.* 2004;286(4):L727-733. doi:10.1152/ajplung.00209.2003
25. Yamasaki M, Kang H-R, Homer RJ, et al. P21 regulates TGF-beta1-induced pulmonary responses via a TNF-alpha-signaling pathway. *Am J Respir Cell Mol Biol.* 2008;38(3):346-353. doi:10.1165/rcmb.2007-0276OC
26. Correa-Meyer E, Pesce L, Guerrero C, Sznajder JI. Cyclic stretch activates ERK1/2 via G proteins and EGFR in alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol.* 2002;282(5):L883-91.
27. Maurer M, Lammerding J. The Driving Force: Nuclear Mechanotransduction in Cellular Function, Fate, and Disease. *Ann Rev Biomed Eng.* 2019;21:443-468. doi:10.1146/annurev-bioeng-060418-052139
28. Yang Z, Maciejowski J, de Lange T. Nuclear Envelope Rupture Is Enhanced by Loss of p53 or Rb. *Mol Cancer Res.* 2017;15(11):1579-1586. doi:10.1158/1541-7786.MCR-17-0084
29. Upadhyay D, Correa-Meyer E, Sznajder JI, Kamp DW. FGF-10 prevents mechanical stretch-induced alveolar epithelial cell DNA damage via MAPK activation. *Am J Physiol Lung Cell Mol Physiol.* 2003;284(2):L350-359. doi:10.1152/ajplung.00161.2002
30. Nava MM, Miroshnikova YA, Biggs LC, et al. Heterochromatin-Driven Nuclear Softening Protects the Genome against Mechanical Stress-Induced Damage. *Cell.* 2020;181(4):800-817.e22. doi:10.1016/j.cell.2020.03.052
31. Mayr M, Hu Y, Hainaut H, Xu Q. Mechanical stress-induced DNA damage and rac-p38MAPK signal pathways mediate p53-dependent apoptosis in vascular smooth muscle cells. *FASEB J.* 2002;16(11):1423-1425. doi:10.1096/fj.02-0042fje
32. Mosteiro L, Pantoja C, Alcazar N, et al. Tissue damage and senescence provide critical signals for cellular reprogramming in vivo. *Science.* 2016;354(6315). doi:10.1126/science.aaf4445
33. Sagiv A, Bar-Shai A, Levi N, et al. p53 in Bronchial Club Cells Facilitates Chronic Lung Inflammation by Promoting Senescence. *Cell Rep.* 2018;22(13):3468-3479. doi:10.1016/j.celrep.2018.03.009
34. Waters DW, Blokland KEC, Pathinayake PS, et al. Fibroblast senescence in

- the pathology of idiopathic pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol*. 2018;315(2):L162-L172. doi:10.1152/ajplung.00037.2018
35. Kumar M, Seeger W, Voswinckel R. Senescence-associated secretory phenotype and its possible role in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol*. 2014;51(3):323-333. doi:10.1165/rcmb.2013-0382PS
36. He Y, Thummuri D, Zheng G, et al. Cellular senescence and radiation-induced pulmonary fibrosis. *Transl Res*. 2019;209:14-21. doi:10.1016/j.trsl.2019.03.006
37. Tominaga T, Shimada R, Okada Y, Kawamata T, Kibayashi K. Senescence-associated- β -galactosidase staining following traumatic brain injury in the mouse cerebrum. *PLoS One*. 2019;14(3):e0213673. doi:10.1371/journal.pone.0213673
38. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med*. 2020;382:1787-1799. doi:10.1056/NEJMoa2001282
39. Heyland DK, Groll D, Caeser M. Survivors of acute respiratory distress syndrome: relationship between pulmonary dysfunction and long-term health-related quality of life. *Crit Care Med*. 2005;33(7):1549-1556.
40. Girard TD, Thompson JL, Pandharipande PP, et al. Clinical phenotypes of delirium during critical illness and severity of subsequent long-term cognitive impairment: a prospective cohort study. *Lancet Respir Med*. 2018;6(3):213-222. doi:10.1016/S2213-2600(18)30062-6
41. Dos Santos C, Hussain SNA, Mathur S, et al. Mechanisms of Chronic Muscle Wasting and Dysfunction after an Intensive Care Unit Stay. A Pilot Study. *Am J Respir Crit Care Med*. 2016;194(7):821-830. doi:10.1164/rccm.201512-2344OC
42. Heyland DK, Garland A, Bagshaw SM, et al. Recovery after critical illness in patients aged 80 years or older: a multi-center prospective observational cohort study. *Intensive Care Med*. 2015;41(11):1911-1920. doi:10.1007/s00134-015-4028-2
43. Koons B, Greenland JR, Diamond JM, Singer JP. Pathobiology of frailty in lung disease. *Transl Res*. 2020;221:1-22. doi:10.1016/j.trsl.2020.04.001
44. Paez-Ribes M, González-Gualda E, Doherty GJ, Muñoz-Espín D. Targeting senescent cells in translational medicine. *EMBO Mol Med*. 2019;11(12):e10234. doi:10.15252/emmm.201810234
45. Chu X, Wen J, Raju RP. Rapid senescence-like response after acute injury. *Aging Cell*. Published online August 2, 2020:e13201. doi:10.1111/acel.13201

Figure legends.

Figure 1. Expression of gene signatures. A: Overview of the analysis. Eleven datasets (128 samples) reporting gene expression in animal models of lung injury were pooled and analyzed to calculate different Meta-scores summarizing the expression of genes included in specific signatures. B: Meta-score of a p53-dependent signature for each experimental group (2nd hit refers to any model of lung injury other than mechanical ventilation). C: Meta-score of a transcriptomic signature including genes downregulated by p21. D: Meta-score of a senescence-specific signature. Gray lines mark significant differences among groups ($p < 0.05$ in Tukey's post-hoc tests).

Journal Pre-proof




Figure 2. Characterization of lung injury. A: Acid instillation and mechanical ventilation caused lung damage assessed using a histological score (scale bar:100 μ). B: Myeloperoxidase-positive cell counts in histological sections, showing an increase of neutrophils in the injured lung. C: Expression of *Il6* in lung tissue. D: Quantification of Ki-67 positive cells in histological sections, as a marker of proliferation. E: TUNEL-positive cells in histological sections. F: Abundance of cleaved Caspase-9 in lung homogenates. G-H: Changes in Lamin-A/Lamin-B1 ratio in nuclei from lung tissue (G) and representative immunohistochemical sections (H, scale bar: 25 μ). I-J: Abundance of γ H2AX (I) and HP1 α (J), markers of DNA damage and heterochromatin respectively, in nuclei from lung tissue. K: Representative western blots of the previous quantifications. L-O: Changes in expression of the canonical senescence inducers *Cdkn2a* (p16, L), *Rb* (M), *Tp53* (N) and *Cdkn1a* (p21, O). N=4-6 animals per group. Gray lines mark significant differences among groups ($p < 0.05$ in Tukey's post-hoc tests).




Figure 3. Lung injury in wildtype and *Cdkn1a*^{-/-} animals. **A:** Histological score of lung damage in both genotypes (scale bar: 100 μ). **B:** Percentage of apoptotic (TUNEL+) cells (scale bar: 50 μ). **C:** Abundance of cleaved caspase-9 in lung homogenates from both genotypes. **D-E:** Expression of *Il6* (**D**) and *Tp53* (**E**) in wildtype and mutant mice. **F-G:** Abundance of γ H2AX (**F**) and HP1 α (**G**), with representative western blots, in lung homogenates. N=4-6 animals per group. Gray lines mark significant differences among groups (p<0.05 in T tests).




Figure 4. Effects of Lopinavir/Ritonavir on lung injury. A: Lamin-A abundance and staining in vehicle- and lopinavir/ritonavir treated animals (scale bar: 25 μ). B: Histological score of lung damage (scale bar: 100 μ). C: Apoptotic (TUNEL+) cell counts in both groups (scale bar: 50 μ). D-F: Abundance of Caspase-9 in tissue homogenates (D), γ H2AX (E) and HP1 α (F), with representative western blots (G) in lung homogenates. H-J: Expression of Il6 (H), *Tp53* (I) and *Cdkn1a* (p21, J). N=7-10 animals per group. Gray lines mark significant differences among groups ($p < 0.05$ in T tests).

Journal Pre-proof




Figure 5. Identification of early senescence markers in experimental models and patients. A: Counts of Senescence-associated heterochromatin foci (SAHF) in the experimental model of lung injury of acid instillation and mechanical ventilation (scale bar: 50 μ). B-D: Expression of *Plk3*, *Gdnf*, and *Meis1* in lung tissue. These senescence-associated genes were identified in the genomic analysis as those with the largest differences between control and injured samples. E-F: SAHF counts (E) and *Plk3* expression (F) in wildtype and *Cdkn1a*^{-/-} mice after lung injury. G-H: SAHF counts (G, scale bar: 25 μ) and *Plk3* expression (H) in vehicle and lopinavir/ritonavir (LPV)-treated mice after lung. I: Appearance of SAHF in autopsy samples from critically ill patients who died in the Intensive Care Unit with or without mechanical ventilation and acute respiratory distress syndrome (ARDS) (scale bar: 50 μ). N=4-7 animals per group, with three slides per animal as technical replicates in SAHF counts. Gray lines mark significant differences among groups ($p < 0.05$ in Tukey's post-hoc or in T tests).




Figure 6. The role of p21 pathway on apoptosis and senescence after acute lung injury. A: In control mice, lung injury and mechanical stretch cause DNA damage and changes in the nuclear envelope, activating the cell senescence program. The amount of apoptotic cells depends on the equilibrium between the activation of pro-apoptotic responses triggered by injury itself and the anti-apoptotic effects of the senescence inducer *Cdkn1a* (p21). B: In mice lacking *Cdkn1a*, absence of this anti-apoptotic factor leads to an increase in apoptosis and a more severe lung injury. C: Treatment with Lopinavir/ritonavir blocks the Lamin-A mediated chromatin remodeling, triggering a senescence-like response that increases p21 expression, thus decreasing apoptosis and lung damage.

