



Contents lists available at ScienceDirect

Metabolism Clinical and Experimental

journal homepage: www.metabolismjournal.com

High-density lipoprotein characteristics and coronary artery disease: a Mendelian randomization study

Albert Prats-Urbe^{a,b,c}, Sergi Sayols-Baixeras^{a,d,e,f}, Alba Fernández-Sanlés^{a,d,g}, Isaac Subirana^{a,h}, Robert Carreras-Torresⁱ, Gemma Vilahur^{e,j}, Fernando Civeira^{e,k}, Jaime Marrugat^{e,l}, Montserrat Fitó^{m,n}, Álvaro Hernández^{m,n,o,p,*}, Roberto Elosua^{a,e,q,**}

^a Cardiovascular Epidemiology and Genetics Research Group, Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain

^b Preventive Medicine and Public Health Unit, Parc de Salut Mar-Universitat Pompeu Fabra-ISGLOBAL, Barcelona, Spain

^c Centre for Statistics in Medicine, Botnar Research Centre, NDORMS, University of Oxford, Oxford, United Kingdom

^d Campus del Mar, Universitat Pompeu Fabra, Barcelona, Spain

^e Consorcio CIBER, M.P. Enfermedades Cardiovasculares (CIBERCV), Instituto de Salud Carlos III, Madrid, Spain

^f Molecular Epidemiology and Science for Life Laboratory, Department of Medical Sciences, Uppsala University, Uppsala, Sweden

^g MRC Integrative Epidemiology Unit, Bristol Medical School, University of Bristol, Bristol, United Kingdom

^h Consorcio CIBER, M.P. Epidemiología y Salud Pública (CIBERESP), Instituto de Salud Carlos III, Madrid, Spain

ⁱ Colorectal Cancer Group, ONCOBELL Program, Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Spain

^j Cardiovascular Program-ICCC, Research Institute-Hospital de la Santa Creu i Sant Pau, IIB-Sant Pau, Barcelona, Spain

^k Lipid Unit, Hospital Universitario Miguel Servet, IIS Aragon, Zaragoza, Spain

^l Girona Heart Registre Research Group (REGICOR), IMIM, Barcelona, Spain

^m Cardiovascular Risk and Nutrition Research Group, IMIM, Barcelona, Spain

ⁿ Consorcio CIBER, M.P. Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Instituto de Salud Carlos III, Madrid, Spain

^o Cardiovascular Risk, Nutrition, and Aging Research Unit, August Pi i Sunyer Biomedical Research Institute (IDIBAPS), Barcelona, Spain

^p Blanquerna School of Life Sciences, Universitat Ramon Llull, Barcelona, Spain

^q Medicine Department, Faculty of Medicine, University of Vic-Central University of Catalonia (UVic-UCC), Vic, Spain

ARTICLE INFO

Article history:

Received 3 July 2020

Accepted 31 August 2020

Available online xxxx

Keywords:

High-density lipoprotein

HDL quality

Coronary artery disease

Mendelian randomization

ABSTRACT

Background: To assess whether genetically determined quantitative and qualitative HDL characteristics were independently associated with coronary artery disease (CAD).

Methods: We designed a two-sample multivariate Mendelian randomization study with available genome-wide association summary data. We identified genetic variants associated with HDL cholesterol and apolipoprotein A-I levels, HDL size, particle levels, and lipid content to define our genetic instrumental variables in one sample (Kettunen et al. study, $n = 24,925$) and analyzed their association with CAD risk in a different study (CARDIoGRAMplusC4D, $n = 184,305$). We validated these results by defining our genetic variables in another database (METSIM, $n = 8372$) and studied their relationship with CAD in the CARDIoGRAMplusC4D dataset. To estimate the effect size of the associations of interest adjusted for other lipoprotein traits and minimize potential pleiotropy, we used the Multi-trait-based Conditional & Joint analysis.

Results: Genetically determined HDL cholesterol and apolipoprotein A-I levels were not associated with CAD. HDL mean diameter ($\beta = 0.27$ [95%CI = 0.19; 0.35]), cholesterol levels in very large HDLs ($\beta = 0.29$ [95%CI = 0.17; 0.40]), and triglyceride content in very large HDLs ($\beta = 0.14$ [95%CI = 0.040; 0.25]) were directly associated with CAD risk, whereas the cholesterol content in medium-sized HDLs ($\beta = -0.076$ [95%CI = -0.10; -0.052]) was inversely related to this risk. These results were validated in the METSIM-CARDIoGRAMplusC4D data.

Conclusions: Some qualitative HDL characteristics (related to size, particle distribution, and cholesterol and triglyceride content) are related to CAD risk while HDL cholesterol levels are not.

* Correspondence to: Á. Hernández, August Pi i Sunyer Biomedical Research Institute (IDIBAPS), IDIBAPS-Mallorca Offices, Carrer Mallorca 183, 08036, Barcelona, Spain.

** Correspondence to: R. Elosua, Cardiovascular Epidemiology and Genetics Research Group, Hospital del Mar Medical Research Institute (IMIM), Carrer Doctor Aiguader 88, 08003, Barcelona, Spain.

E-mail addresses: albert.prats@gmail.com (A. Prats-Urbe), ssayols@gmail.com (S. Sayols-Baixeras), alba.fernandez-sanles@bristol.ac.uk (A. Fernández-Sanlés), isubirana@imim.es (I. Subirana), rcarreras@idibell.cat (R. Carreras-Torres), gvilahur@santpau.cat (G. Vilahur), civeira@unizar.es (F. Civeira), jmarrugat@imim.es (J. Marrugat), mfito@imim.es (M. Fitó), alvaro.hernaez1@gmail.com (Á. Hernández), relosua@imim.es (R. Elosua).

¹ These authors contributed equally to this manuscript.

1. Introduction

The inverse association between high-density lipoprotein cholesterol (HDL-C) levels and the risk of coronary artery disease (CAD) has been reported in observational studies [1]. However, experimental and genetic studies question the causality of this association. On the one hand, drugs such as fibrates, niacin, and cholesteryl ester transfer protein inhibitors increase HDL-C levels but do not decrease CAD risk [2]. On the other hand, genetic predisposition to high HDL-C levels has not been linked to any decrease in the risk of cardiovascular events [3,4]. Thus, researchers are looking beyond HDL-C levels to disentangle this apparent contradiction. Anti-atherogenic properties of HDL particles seem to be determined by the quality or function of the lipoprotein [5]. HDL particle size and number have been linked to cardiovascular risk [6], and this association could be mediated through HDL functionality, which is predictive of cardiovascular risk [7]. The interplay between HDL-C and triglyceride levels, as two of the faces of atherogenic dyslipidemia, may also play a relevant role in their relationship with CAD [8,9]. Thus, further evidence of causal association between HDL characteristics and CAD risk would provide relevant data on the validity of these particles as therapeutic targets.

Mendelian Randomization (MR) studies have arisen as a powerful tool to ascertain the potential causality of the association between a biomarker and a disease [10]. These studies assess the association between the genetically determined lifelong values of a biomarker and the development of a clinical outcome. MR studies have already raised serious doubts on the causal role of quantitative HDL characteristics, such as HDL-C and apolipoprotein A-I (ApoA-I) levels, in CAD [3,11]. However, to date, the association between qualitative HDL characteristics and CAD has not been tested using a MR approach. HDL mean diameter, the concentration of HDL particles of each size subtype, the distribution of cholesterol across the HDL size subtypes, and the presence of other lipids in HDL particles (such as triglycerides, highly present in large HDLs) are some of these qualitative traits. Additionally, this evaluation must take into account the complexity of lipid metabolism and its potential genetic pleiotropic effects. HDL-C, low-density lipoprotein cholesterol (LDL-C), and triglyceride levels are highly interdependent and, therefore, the method used to test the association between HDL properties and CAD risk should take into account this inter-correlation [12].

This study had two aims: 1) to assess the potential causal association of quantitative and qualitative HDL characteristics with CAD risk, using a two-sample MR approach; and 2) to explore potential mechanisms explaining the observed associations.

2. Methods

2.1. Study design and data sources

We designed a two-sample MR study using aggregated summary data [10] from three published meta-analyses of genome-wide association studies. The main analysis was based on data from Kettunen et al. [13] ($n = 24,925$) and the CARDIoGRAMplusC4D consortium [14] ($n = 184,305$), and the validation analysis used the METSIM [15] ($n = 8372$) and the CARDIoGRAMplusC4D datasets. Regarding the GWASs on lipoprotein traits, we used the Kettunen et al. and the METSIM studies because these are the only ones with publicly available summary data. Regarding CARDIoGRAMplusC4D, we selected it because it had several quality characteristics that made it stand out among all GWAS on CAD with publicly available summary results: a large number of total participants ($n = 184,305$), a large number of CAD cases ($n = 60,801$), a large number of genetic variants analyzed (8.6 million), and a quite consistent definition of CAD in the participant studies. A more detailed description of the studies is available in Supplementary materials.

We centered our analysis on the genetic variants associated with: 1) the main lipid profile traits in serum (HDL-C, LDL-C, and triglyceride levels); 2) other measurements of HDL quantity (ApoA-I levels); 3) HDL mean diameter; 4) the quantities of cholesterol transported in small,

medium-sized, large and very large HDLs; 5) the quantity of other lipid species in HDL particles (triglycerides transported in very large HDLs); and 6) the levels of HDL particles according to the previous HDL size subtypes. Both the Kettunen et al. and the METSIM studies measured HDL qualitative characteristics by the same nuclear magnetic resonance spectroscopy technique [16].

2.2. Assessment of genetic variants linked to lipoprotein characteristics and CAD risk adjusted for other lipoprotein traits: Multi-trait-based Conditional & Joint analyses

To identify the genetic variants associated with each lipoprotein characteristic that were also linked to CAD considering the potential pleiotropy among lipid profile traits, we used the Multi-trait-based Conditional & Joint analysis [17] in both the main and the validation stage. This method enables the estimation of the magnitude of the association of each genetic variant with each lipoprotein characteristic and with CAD, independently from the other genetically determined lipoprotein traits (adjusted effect sizes). For example, if we considered HDL-C as “main variable” and LDL-C and triglyceride levels as “covariates”, we would obtain the betas and standard errors of the associations of genetic variants with (1) HDL-C (using the Kettunen and METSIM raw summary data) and (2) CAD (using the CARDIoGRAMplusC4D raw summary results), adjusted for the association of these same genetic variants with LDL-C and triglyceride concentrations. For this purpose, we defined six multivariate models a priori. Model 1 included HDL-C, LDL-C and triglyceride levels. Further models included the elements in Model 1 as covariates and the following parameters as main variables: ApoA-I levels (Model 2); HDL mean diameter (Model 3); the cholesterol content in each HDL size subtype (small, medium-sized, large, and very large HDL particles; Model 4); the levels of HDL particles of each size subtype (Model 5); and the triglyceride content in very large HDLs (Model 6). In model 4 and 5, we required for the presence of at least two of the HDL subtypes (small, medium-sized, large, and very large) traits to build the model.

The genetic correlation between traits was estimated by linkage disequilibrium score regressions using all genetic variants.

2.3. Mendelian randomization analyses

Based on the adjusted gene variant effects and their standard errors computed as previously described, we performed the MR analysis using the Generalized Summary-data-based Mendelian Randomization method [18]. The genetic variants to be considered were selected with the following criteria: 1) strong association with the lipid traits of interest ($p\text{-value} < 5 \cdot 10^{-8}$); 2) not in linkage disequilibrium ($R^2 < 0.01$, using the 1000 Genome project data – <http://www.1000genomes.org/phase-3-structural-variant-dataset> – as reference [19]); and 3) a minor allele frequency ≥ 0.05 .

As an additional approach to exclude potential pleiotropy, we also removed the variants with a significant result in the HEIDI-outlier test ($p\text{-value} < 0.01$). Finally, we explored and confirmed the exclusion of potential pleiotropic effects using Egger regressions [20]. Statistical significance of our results was corrected for multiple comparisons ($p\text{-value} = 0.05/\text{number of traits}$). A description of complementary sensitivity analyses using other MR analyses methods such as the median-based and inverse variance weighted, using the Global Lipid Genetic Consortium dataset [21] and *post-hoc* statistical power estimations [22] is available in Supplementary materials.

3. Results

3.1. Mendelian randomization results

3.1.1. Selected genetic variants

We identified genetic variants significantly associated with 13 lipoprotein characteristics in the data published by Kettunen et al. and

with 8 lipoprotein traits in the METSIM data. The number of genetic variants for each lipoprotein trait ranged from 6 (level of medium-sized HDL particles) to 22 (ApoA-I). Genetic variants included in the analyses and their unadjusted and adjusted effects are listed in Supplementary Excel File 1.

We observed high inverse genetic correlations (correlation coefficient ≤ -0.50) between triglyceride and HDL-C levels. Conversely, we observed very high direct genetic correlations (correlation coefficient ≥ 0.70) between the cholesterol content in each HDL size subpopulation and HDL-C concentrations, between HDL mean diameter and the level of very large HDL particles, and between ApoA-I and the level of very large HDL particles and their cholesterol content (Supplementary Fig. 1).

3.1.2. Main analysis

We defined as statistically significant those associations with a p -value $< 3.85 \cdot 10^{-3}$ (0.05/13). We observed a direct association between CAD risk and genetically determined levels of LDL-C ($\beta = 0.26$ [95% Confidence Interval]: [0.17; 0.35], p -value = $1.32 \cdot 10^{-8}$) and triglycerides ($\beta = 0.18$ [0.073; 0.29], p -value = $1.05 \cdot 10^{-3}$). Conversely, the genetically determined concentrations of HDL-C ($\beta = 0.008$ [-0.084; 0.099], p -value = 0.871) or ApoA-I ($\beta = 0.060$ [-0.015; 0.13], p -value = 0.116) were not associated with CAD risk (Figs. 1, 2A and B).

In qualitative HDL measurements, the genetically determined HDL mean diameter was directly associated with CAD risk ($\beta = 0.27$ [0.19; 0.35], p -value = $2.23 \cdot 10^{-11}$) (Figs. 1 and 2C). Cholesterol levels in very large HDLs was also positively linked to CAD risk ($\beta = 0.29$ [0.17; 0.40], p -value = $8.90 \cdot 10^{-7}$), whereas cholesterol in medium-sized HDLs was inversely related to this risk ($\beta = -0.076$ [-0.10; -0.052], p -value = $4.55 \cdot 10^{-11}$) (Figs. 1, 2D, and E). The genetically

determined levels of all subtypes of HDL particles showed an inverse trend towards an association with CAD risk, but only that between very large HDLs and CAD was statistically significant ($\beta = -0.22$ [-0.32; -0.12], p -value = $7.12 \cdot 10^{-6}$) (Fig. 1). Finally, the genetically determined levels of triglycerides in very large HDLs were directly related to CAD risk ($\beta = 0.14$ [0.040; 0.25], p -value = $6.84 \cdot 10^{-3}$) (Figs. 1 and 2F). Effect sizes of all the associations are available in Supplementary Table 1.

Genetic variants linked to both HDL qualitative traits and CAD risk were located within *LIPC*, the *APOE/C1/C4/C2* cluster, *PCIF1*, *TTC39B*, and *APOB* (Supplementary Excel File 1).

3.1.3. Validation analysis

In the validation analysis (Fig. 1), we confirmed the direct association between the genetically determined concentrations of LDL-C and CAD risk ($\beta = 0.24$ [0.17; 0.31], p -value = $1.21 \cdot 10^{-10}$), triglycerides and CAD risk ($\beta = 0.084$ [0.013; 0.15], p -value = 0.020), and the null link between the genetically determined HDL-C levels and CAD ($\beta = -0.025$ [-0.087; 0.036], p -value = 0.419).

Results regarding qualitative HDL traits were also replicated. We confirmed the direct association of genetically determined HDL mean diameter with CAD risk ($\beta = 0.34$ [0.23; 0.46], p -value = $4.47 \cdot 10^{-9}$). There was a positive link between the cholesterol content in large HDLs and CAD ($\beta = 0.45$ [0.34; 0.55], p -value = $1.61 \cdot 10^{-16}$) and an inverse relationship between the genetically determined cholesterol levels in medium-sized HDLs and CAD risk ($\beta = -0.46$ [-0.55; -0.37], p -value = $5.9 \cdot 10^{-23}$). Finally, high genetically determined levels of triglycerides in very large HDLs were nominally associated with greater CAD risk ($\beta = 0.081$ [0.008; 0.15], p -value = 0.030) (Fig. 1 and Supplementary Fig. 2). We could not assess the associations




Fig. 1. Association of the genetically determined lipid profile and HDL characteristics with coronary artery disease risk. Effect of genetically determined HDL cholesterol (HDL-C), LDL cholesterol (LDL-C) and triglyceride levels (Panel 1), apolipoprotein A-I concentrations (Panel 2), mean HDL diameter (Panel 3), cholesterol content in each HDL size subtype (Panel 4), number of particles of each HDL size subtype (Panel 5), and triglyceride levels in very large HDLs (Panel 6) on CAD, independently from the effect of genetically determined levels of classic lipid profile parameters. In all cases, main Mendelian randomization analyses (Kettunen-CARDIoGRAMplusC4D) appear first and the validation ones (METSIM-CARDIoGRAMplusC4D) appear below. NA: non-available association.




Fig. 2. Association of individual SNPs affecting lipoprotein traits with coronary artery disease risk in the main analysis. Estimates of the associations of individual SNPs related to (A) HDL cholesterol (HDL-C) levels, (B) apolipoprotein A-I concentrations, (C) mean HDL diameter, (D) cholesterol content in medium-sized HDLs, (E) cholesterol content in very large HDLs, and (F) triglyceride levels in very large HDLs with coronary artery disease risk. Estimates were derived from the study by Kettunen et al. and the CARDIoGRAMplusC4D meta-analyses (multivariate adjusted estimates). Error bars represent 95% confidence intervals. The slopes of the lines show the genetic instrumental variable regression estimates of the effect of the lipid characteristics on coronary artery disease risk.

between the levels of HDL particles and CAD due to the lack of genetic variants associated with at least two of these HDL traits in the METSIM study. The effect sizes of all the associations of interest are available in Supplementary Table 2.

3.1.4. Sensitivity analysis

Egger regression intercept estimates supported the absence of pleiotropic effects. Results of the median-based and inverse variance weighted methods confirmed the direction and significance of the main analyses (Supplementary Table 3).

Associations between genetically determined HDL-C, LDL-C and triglyceride levels and CAD risk identified in our main analysis were similar to those obtained from the Global Lipid Genetic Consortium (Supplementary Table 4).

3.1.5. Post-hoc statistical power estimation

Power estimation for the main analyses ranged from 2.4% to 96.9% (Supplementary Table 5).

4. Discussion

Our findings suggest a potential causal relationship between qualitative HDL characteristics and CAD risk, even though HDL-C and ApoA-I levels were not associated with CAD. In particular, genetically determined mean HDL size, the distribution of cholesterol across HDL size subpopulations, and the triglyceride content in HDL particles were related to CAD risk.

The relationship between HDL and cardiovascular risk is controversial [4]. Recent studies suggest that HDL functions and quality characteristics, rather than HDL-C concentration, are the main determinants of HDL anti-atherogenic properties [5]. Our data are consistent with previous evidence, and reflect that HDL-C and ApoA-I levels in the bloodstream are not causally related to CAD [3,11]. However, we observed a decrease in CAD risk when HDL-C was mainly transported in smaller HDLs, but an increase in CAD risk when HDL-C was carried by larger HDL particles (in both main and validation analyses, there is a gradient towards greater CAD risk as more cholesterol is transported in larger HDLs). The protective effect of cholesterol content in medium-sized HDLs and the increase in CAD risk due to cholesterol levels in larger particles observed in our data may contribute to explaining why the overall HDL-C levels are not causally associated with cardiovascular risk [3,11]. Our results could also help explain the therapeutic failure of the pharmacological agents known to increase HDL-C levels. Niacin or cholesteryl ester transfer protein inhibitors are effective in increasing HDL-C concentrations but not in reducing CAD risk [2]. This paradox could be explained by the promotion of the accumulation of cholesterol content in large HDLs after the use of these drugs [23,24]. In gemfibrozil-treated patients, changes in HDL-C levels accounted for a small proportion of the CAD risk reduction (<10%), whereas the increase in small HDLs was much more predictive of this risk reduction [25]. Finally, our results also concur with genetic studies analyzing variants in the SR-B1 gene, showing that individuals with loss-of-function variants have higher HDL-C concentrations, mainly in very large particles, but also higher CAD risk [26].

However, there is still controversy in the relationship between HDL size subtypes and cardiovascular risk: some authors advocate for small HDLs as indicators of lower CAD risk [27] while others suggest they are associated with increased CAD risk [28]. There are several possible explanations for this heterogeneity. First, baseline health conditions of the subjects affect HDL quality and function. Lipid-poor, protein-rich, small HDLs could be dysfunctional in pro-oxidative and pro-inflammatory pathological states due to post-translational modifications of their proteins and their enrichment in pro-inflammatory mediators (such as serum amyloid A or complement 3) [29]. Second, laboratory procedures to measure HDL size (nuclear magnetic resonance spectroscopy, electrophoresis, etc.) differed between the published studies, and there is low

concordance between these techniques [30]. Third, the statistical models used did not consider all the same confounding factors and did not always include as covariates the levels of HDL-C or other lipid profile parameters related to these lipoproteins (e.g. triglyceride concentrations).

Triglyceride and HDL-C levels may be two sides of the same coin, and this relationship may contribute to explaining why HDL-C is not causally related to CAD while triglycerides are. Hypertriglyceridemic states (generally due to high levels of very-low density lipoprotein concentrations in plasma) are linked to an increased activity of the cholesteryl ester transfer protein, an enzyme that exchanges triglycerides in very-low density lipoproteins for cholesterol in HDLs, resulting in an enrichment of HDLs in triglycerides [8]. Aged HDL particles may also become increasingly richer in triglycerides because this exchange is an essential process by which HDLs get rid of the cholesterol they have collected from peripheral cells and transfer it back to the liver [31]. In any case, triglyceride-rich HDLs have been shown to present their ApoA-I in an unstable conformation [32], which may be related to lower HDL function (lower cholesterol efflux capacity) and a greater disintegration of the HDL structure (lower HDL-C levels) [8]. Our results confirm that triglyceride-rich HDLs are causally related to higher CAD risk independently from the circulating levels of triglycerides and HDL-C. In addition, this mechanism also verifies the hypothesis that high triglycerides (in circulation and in HDL particles) are essential mediators of high cardiovascular risk and suggests that low HDL-C levels in these states may be a secondary consequence of this lipid disruption.

Both observational and experimental studies have more consistently found an inverse relationship between the number of HDL particles and cardiovascular risk, compared to HDL-C levels [7]. Similarly, we observed that the concentrations of HDL particles of all sizes were inversely related to CAD risk, although only the genetically determined levels of very large HDLs were significantly associated with it in the main analysis. Unfortunately, we could not validate these results due to the lack of valid genetic variants in the METSIM study.

Our results are mechanistically plausible and highlight novel potential therapeutic targets in cardiovascular disease, since genetic variants individually associated with HDL qualitative traits and CAD in our data were located within several HDL-related genes or gene clusters. *LIPC* encodes for hepatic lipase C, an enzyme that hydrolyzes triglycerides in circulating lipoproteins, including HDL particles [31]. Hydrolysis of HDL triglycerides by this enzyme generates small/medium-sized, triglyceride-depleted particles, considered to be more stable and functional than very large, triglyceride-rich HDLs [33]. Since triglyceride-rich HDLs were also causally linked to CAD in our data, this potential mechanism would contribute to explaining a decrease in cardiovascular risk. The *APOE/C1/C4/C2* cluster encodes apolipoproteins E, C-I, C-II, and C-IV and has been classically associated with blood lipid levels [34]. Particularly, apolipoprotein E is a pivotal mediator in reverse cholesterol transport [35] and apolipoprotein C-I is involved in the activation of lecithin-cholesterol acyltransferase and the inhibition of cholesteryl ester transfer protein, two key enzymes in HDL metabolism [36,37]. The third most relevant HDL-related locus encodes the PDX1 C-Terminal Inhibiting Factor 1 (*PCIF1*) and is located next to the *PLTP* gene, which expresses the phospholipid transfer protein, an enzyme involved in HDL remodeling/stabilization [31]. *PCIF1*-related gene variants have been shown to modulate phospholipid transfer protein function in other studies [38]. Finally, *TTC39B* encodes the tetratricopeptide repeat domain protein 39B, whose genetic variants had already been associated with HDL-C levels and CAD in previous works [34,39].

Our study presents some limitations. First, in order to use a MR approach, we had to make some assumptions [10], among which stands out the absence of pleiotropy. In our case, most of the genetic variants used as instruments were associated with more than one lipid trait. To solve this problem, we used a novel approach (Multi-trait based

Conditional & Joint analysis—Generalized Summary data-based Mendelian randomization methodology) to control for the confounding effects related to the close relationship between lipoprotein characteristics and to minimize pleiotropy [17]. Second, the interpretation of multivariable MR is challenging, especially when the covariate-biomarker lies on the causal pathway from the main-biomarker to disease, or when the covariate-biomarker measures the same entity as the main-biomarker [40]. Third, the population tested may exhibit significant genetic heterogeneity because of different ethnic origin. The Kettunen et al. study included European populations, the CARDIoGRAMplusC4D study included European and Asian populations, but the METSIM study included only Finnish. However, this heterogeneity is assumed to be minimal since the original GWASs excluded those genetic variants whose allele frequencies departed from Hardy-Weinberg equilibrium and all the variants included in our analyses, except rs73168081 (associated with cholesterol content in large HDL particles in Kettunen et al), presented a similar allele frequency in the three GWASs considered (Supplementary Excel File 1). In addition, the METSIM study only considered male population, and this may have partially distorted the findings of the validation analyses. Nevertheless, sex- and age-related bias had been originally addressed in the original GWASs as described in the Methodology section of each of the studies. Fourth, we have not been able to study whether other HDL functional properties (such as cholesterol efflux capacity, HDL antioxidant properties, and HDL particle type according to ApoA-I and ApoA-II content) are causally linked to CAD due to unavailability of GWAS studies on these traits with publicly available summary data. Fifth, the statistical power of our analyses was limited for some of the traits of interest. Finally, in the validation analysis we could not generate genetic instrumental variables for some of the lipoprotein traits. However, our study has several methodological strengths. First, our results are based in MR, a useful approach to explore the causality of the association between biomarkers and specific diseases. Second, we included two independent MR analyses to validate the results initially observed. Finally, the validity of the genetic variants for HDL-C, LDL-C and triglyceride levels initially generated was confirmed, supporting the validity of these datasets for the analysis of other genetic variants.

5. Conclusions

Several genetically and life-long qualitative HDL characteristics were related to CAD risk. Although HDL-C and ApoA-I levels were not causally linked to CAD, our results support a potential causality between higher mean HDL diameter, greater cholesterol levels in very large HDLs, and triglyceride-rich HDL particles and higher CAD risk, and between cholesterol levels in medium-sized HDLs and lower CAD risk. This relationship could be mediated by several HDL-related proteins, which are suggested as potential therapeutic targets for further exploration in cardiovascular prevention.

CRedit authorship contribution statement

Albert Prats-Uribe: Conceptualization, Formal analysis, Investigation, Methodology, Writing - original draft. **Sergi Sayols-Baixeras:** Investigation, Methodology, Writing - review & editing. **Alba Fernández-Sanlés:** Investigation, Methodology, Writing - review & editing. **Isaac Subirana:** Methodology, Writing - review & editing. **Robert Carreras-Torres:** Writing - review & editing. **Gemma Vilahur:** Writing - review & editing. **Fernando Civeira:** Writing - review & editing. **Jaume Marrugat:** Writing - review & editing. **Montserrat Fitó:** Writing - review & editing. **Álvaro Hernández:** Writing - original draft. **Roberto Elosua:** Conceptualization, Investigation, Methodology, Writing - review & editing.

Declaration of competing interest

The authors declare they do not have conflict of interest.

Acknowledgements

We thank Elaine M. Lilly, PhD, for her critical reading and revision of the English text. Data were downloaded from: www.computationalmedicine.fi/data#NMR_GWAS (Kettunen study), <http://csg.sph.umich.edu/boehnke/public/metsim-2017-lipoproteins/> (METSIM study), www.cardiogramplusc4d.org (CARDIoGRAMplusC4D), <http://csg.sph.umich.edu/abecasis/public/lipids2013/> (Global Lipid Genetic Consortium), <http://www.1000genomes.org/phase-3-structural-variant-dataset> (1000 Genome).

Funding sources

This work was supported by the Instituto de Salud Carlos III—European Regional Development Fund [grant numbers CD17/00122, IF114/00007, PI18/00017], the Medical Research Council [grant numbers MR/K501256/1, MR/N013468/1], the Spanish Ministry of Economy and Competitiveness [grant numbers BES-2014-069718, SAF2015-71653-R], the European Union's Horizon 2020 Research and Innovation Program [grant number 796216], and the Government of Catalonia through the Agency for Management of University and Research Grants [2017 SGR 222]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. CIBER Enfermedades Cardiovasculares (CIBERCV), Epidemiología y Salud Pública (CIBERESP), and Fisiopatología de la Obesidad y Nutrición (CIBEROBN) are initiatives of the Instituto de Salud Carlos III, Madrid, Spain, and financed by the European Regional Development Fund.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.metabol.2020.154351>.

References

- [1] Barter P, Gotto AM, LaRosa JC, Maroni J, Szarek M, Grundy SM, et al. HDL cholesterol, very low levels of LDL cholesterol, and cardiovascular events. *N Engl J Med*. 2007; 357:1301–10. <https://doi.org/10.1056/NEJMoa064278>.
- [2] Keene D, Price C, Shun-Shin MJ, Francis DP. Effect on cardiovascular risk of high density lipoprotein targeted drug treatments niacin, fibrates, and CETP inhibitors: meta-analysis of randomised controlled trials including 117,411 patients. *BMJ*. 2014;349: g4379. <https://doi.org/10.1136/bmj.g4379>.
- [3] Holmes MV, Asselbergs FW, Palmer TM, Drenos F, Lanktree MB, Nelson CP, et al. Mendelian randomization of blood lipids for coronary heart disease. *Eur Heart J*. 2015;36:539–50. <https://doi.org/10.1093/eurheartj/ehv571>.
- [4] Landmesser U, Hazen S. HDL-cholesterol, genetics, and coronary artery disease: the myth of the 'good cholesterol?'. *Eur Heart J*. 2018;39:2179–82. <https://doi.org/10.1093/eurheartj/ehy299>.
- [5] Sacks FM, Jensen MK. From high-density lipoprotein cholesterol to measurements of function. *Arterioscler Thromb Vasc Biol*. 2018;38:487–99. <https://doi.org/10.1161/ATVBAHA.117.307025>.
- [6] Superko HR, Pendyala L, Williams PT, Momary KM, King SB, Garrett BC. High-density lipoprotein subclasses and their relationship to cardiovascular disease. *J Clin Lipidol*. 2012;6:496–523. <https://doi.org/10.1016/j.jacl.2012.03.001>.
- [7] Rohatgi A, Khera A, Berry JD, Givens EG, Ayers CR, Wedin KE, et al. HDL cholesterol efflux capacity and incident cardiovascular events. *N Engl J Med*. 2014;371:2383–93. <https://doi.org/10.1056/NEJMoa1409065>.
- [8] Lamarche B, Uffelman KD, Carpentier A, Cohn JS, Steiner G, Barret PH, et al. Triglyceride enrichment of HDL enhances in vivo metabolic clearance of HDL apo A-I in healthy men. *J Clin Invest*. 1999;103:1191–9. <https://doi.org/10.1172/JCI5286>.
- [9] Musunuru K. Atherogenic dyslipidemia: cardiovascular risk and dietary intervention. *Lipids*. 2010;45:907–14. <https://doi.org/10.1007/s11745-010-3408-1>.
- [10] Zheng J, Baird D, Borges MC, Bowden J, Hemani G, Haycock P, et al. Recent developments in Mendelian randomization studies. *Curr Epidemiol Reports*. 2017;4:330–45. <https://doi.org/10.1007/s40471-017-0128-6>.
- [11] Karjalainen MK, Holmes MV, Wang Q, Anufrieva O, Kähönen M, Lehtimäki T, et al. Apolipoprotein A-I concentrations and risk of coronary artery disease: a Mendelian randomization study. *Atherosclerosis*. 2020;299:56–63. <https://doi.org/10.1016/j.atherosclerosis.2020.02.002>.
- [12] Würtz P, Kangas AJ, Soinen P, Lehtimäki T, Kähönen M, Viikari JS, et al. Lipoprotein subclass profiling reveals pleiotropy in the genetic variants of lipid risk factors for coronary heart disease: a note on Mendelian randomization studies. *J Am Coll Cardiol*. 2013;62:1906–8. <https://doi.org/10.1016/j.jacc.2013.07.085>.
- [13] Kettunen J, Demirkan A, Würtz P, Draisma HHM, Haller T, Rawal R, et al. Genome-wide study for circulating metabolites identifies 62 loci and reveals novel systemic effects of LPA. *Nat Commun*. 2016;7:11122. <https://doi.org/10.1038/ncomms11122>.

- [14] CARDIoGRAMplusC4D Consortium P, Deloukas P, Kanoni S, Willenborg C, Farrall M, Assimes TL, et al. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nat Genet.* 2013;45:25–33. <https://doi.org/10.1038/ng.2480>.
- [15] Davis JP, Huyghe JR, Locke AE, Jackson AU, Sim X, Stringham HM, et al. Common, low-frequency, and rare genetic variants associated with lipoprotein subclasses and triglyceride measures in Finnish men from the METSIM study. *PLoS Genet.* 2017;13:e1007079. <https://doi.org/10.1371/journal.pgen.1007079>.
- [16] Soininen P, Kangas AJ, Würtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ Cardiovasc Genet.* 2015;8:192–206. <https://doi.org/10.1161/CIRCGENETICS.114.000216>.
- [17] Zhu Z, Zheng Z, Zhang F, Wu Y, Trzaskowski M, Maier R, et al. Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat Commun.* 2018;9:224. <https://doi.org/10.1038/s41467-017-02317-2>.
- [18] Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 2011;88:76–82. <https://doi.org/10.1016/j.ajhg.2010.11.011>.
- [19] Sudmant PH, Rausch T, Gardner EJ, Handsaker RE, Abyzov A, Huddleston J, et al. An integrated map of structural variation in 2,504 human genomes. *Nature.* 2015;526:75–81. <https://doi.org/10.1038/nature15394>.
- [20] Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol.* 2015;44:512–25. <https://doi.org/10.1093/ije/dyv080>.
- [21] Willer CJ, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, Kanoni S, et al. Discovery and refinement of loci associated with lipid levels. *Nat Genet.* 2013;45. <https://doi.org/10.1038/ng.2797>.
- [22] Brion MJA, Shakhbazov K, Visscher PM. Calculating statistical power in Mendelian randomization studies. *Int J Epidemiol.* 2013;42:1497–501. <https://doi.org/10.1093/ije/dyt179>.
- [23] Ronsein GE, Hutchins PM, Isquith D, Vaisar T, Zhao XQ, Heinecke JW. Niacin therapy increases high-density lipoprotein particles and total cholesterol efflux capacity but not ABCA1-specific cholesterol efflux in statin-treated subjects. *Arterioscler Thromb Vasc Biol.* 2016;36:404–11. <https://doi.org/10.1161/ATVBAHA.115.306268>.
- [24] Chen Y, Dong J, Zhang X, Chen X, Wang L, Chen H, et al. Evacetrapib reduces pre β -1 HDL in patients with atherosclerotic cardiovascular disease or diabetes. *Atherosclerosis.* 2019;285:147–52. <https://doi.org/10.1016/j.atherosclerosis.2019.04.211>.
- [25] Robins SJ, Collins D, Wittes JT, Papademetriou V, Deedwania PC, Schaefer EJ, et al. Relation of gemfibrozil treatment and lipid levels with major coronary events: VA-HIT: a randomized controlled trial. *JAMA.* 2001;285:1585–91.
- [26] Zanon P, Khetarpal SA, Larach DB, Hancock-Cerutti WF, Millar JS, Cuchel M, et al. Rare variant in scavenger receptor BI raises HDL cholesterol and increases risk of coronary heart disease. *Science (80-)* 2016;351:1166–71. <https://doi.org/10.1126/science.aad3517>.
- [27] Otvos JD, Collins D, Freedman DS, Shalurova I, Schaefer EJ, McNamara JR, et al. Low-density lipoprotein and high-density lipoprotein particle subclasses predict coronary events and are favorably changed by gemfibrozil therapy in the Veterans Affairs High-Density Lipoprotein Intervention Trial. *Circulation.* 2006;113:1556–63. <https://doi.org/10.1161/CIRCULATIONAHA.105.565135>.
- [28] Rosenson RS, Otvos JD, Freedman DS. Relations of lipoprotein subclass levels and low-density lipoprotein size to progression of coronary artery disease in the Pravastatin Limitation of Atherosclerosis in the Coronary Arteries (PLAC-I) trial. *Am J Cardiol.* 2002;90:89–94.
- [29] Rosenson RS, Brewer HB, Ansell BJ, Barter P, Chapman MJ, Heinecke JW, et al. Dysfunctional HDL and atherosclerotic cardiovascular disease. *Nat Rev Cardiol.* 2016;13:48–60. <https://doi.org/10.1038/nrcardio.2015.124>.
- [30] Arsenault BJ, Lemieux I, Després JP, Wareham NJ, Stroes ESG, Kastelein JJP, et al. Comparison between gradient gel electrophoresis and nuclear magnetic resonance spectroscopy in estimating coronary heart disease risk associated with LDL and HDL particle size. *Clin Chem.* 2010;56:789–98. <https://doi.org/10.1373/clinchem.2009.140939>.
- [31] Rye KA, Barter PJ. Regulation of high-density lipoprotein metabolism. *Circ Res.* 2014;114:143–56. <https://doi.org/10.1161/CIRCRESAHA.114.300632>.
- [32] Sparks DL, Davidson WS, Lund-Katz S, Phillips MC. Effects of the neutral lipid content of high density lipoprotein on apolipoprotein A-I structure and particle stability. *J Biol Chem.* 1995;270:26910–7.
- [33] Chatterjee C, Sparks DL. Hepatic lipase, high density lipoproteins, and hypertriglyceridemia. *Am J Pathol.* 2011;178:1429–33. <https://doi.org/10.1016/j.ajpath.2010.12.050>.
- [34] Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, Koseki M, et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature.* 2010;466:707–13. <https://doi.org/10.1038/nature09270>.
- [35] Miettinen TA, Gylling H, Vanhanen H, Ollus A. Cholesterol absorption, elimination, and synthesis related to LDL kinetics during varying fat intake in men with different apoprotein E phenotypes. *Arterioscler Thromb Vasc Biol.* 1992;12:1044–52. <https://doi.org/10.1161/01.atv.12.9.1044>.
- [36] Weisgraber KH, Mahley RW, Kowal RC, Herz J, Goldstein JL, Brown MS. Apolipoprotein C-I modulates the interaction of apolipoprotein E with beta-migrating very low density lipoproteins (beta-VLDL) and inhibits binding of beta-VLDL to low density lipoprotein receptor-related protein. *J Biol Chem.* 1990;265:22453–9.
- [37] Gautier T, Masson D, Pais De Barros JP, Athias A, Gamber P, Aunis D, et al. Human apolipoprotein C-I accounts for the ability of plasma high density lipoproteins to inhibit the cholesteryl ester transfer protein activity. *J Biol Chem.* 2000;275:37504–9. <https://doi.org/10.1074/jbc.M007210200>.
- [38] Kim DS, Burt AA, Ranchalis JE, Vuletic S, Vaisar T, Li WF, et al. PLTP activity inversely correlates with CAAD: effects of PON1 enzyme activity and genetic variants on PLTP activity. *J Lipid Res.* 2015;56:1351–62. <https://doi.org/10.1194/jlr.P058032>.
- [39] Huang JH, Yin RX, Li WJ, Huang F, Chen WX, Cao XL, et al. Association of the TTC39B rs581080 SNP and serum lipid levels and the risk of coronary artery disease and ischemic stroke. *Int J Clin Exp Pathol.* 2017;10:11195–205.
- [40] Holmes MV, Davey Smith G. Challenges in interpreting multivariable Mendelian randomization: might “good cholesterol” be good after all? *Am J Kidney Dis.* 2018;71:149–53. <https://doi.org/10.1053/j.ajkd.2017.10.006>.