

Original Article

Dissecting Common and Unique Effects of Anti- $\alpha 4\beta 7$ and Anti-Tumor Necrosis Factor Treatment in Ulcerative Colitis

Marisol Veny,^{a,*} Alba Garrido-Trigo,^{a,*} Ana M. Corraliza,^a
Maria C. Masamunt,^a Helena Bassolas-Molina,^a Miriam Esteller,^a
Montserrat Arroyes,^a Eva Tristán,^b Agnès Fernández-Clotet,^a Ingrid Ordás,^a
Elena Ricart,^a Maria Esteve,^b Julian Panés,^a Azucena Salas^{a,Ⓞ}

^aInflammatory Bowel Disease Unit, Gastroenterology Department, Hospital Clinic of Barcelona, Barcelona, Spain; Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBERehd), Spain ^bHospital Universitari Mutua Terrassa, CIBERehd, Department of Gastroenterology, Terrassa, Spain

*These authors contributed equally.

Corresponding author: Azucena Salas, PhD, Center Esther Koplowitz, Rossello 149–153, Barcelona 08036, Spain. Tel: 34 93 2275400 ext 2436; Email: asalas1@clinic.cat

Abstract

Background and Aims: Vedolizumab is an anti- $\alpha 4\beta 7$ antibody approved for the treatment of ulcerative colitis [UC]. Although it is assumed that vedolizumab blocks intestinal homing of lymphocytes, its effects on different intestinal cell populations are not fully established. In order to establish the unique mechanisms of action of vedolizumab in UC patients, we compared its effects to those induced by anti-tumour necrosis factor [TNF].

Methods: Patients with active UC [endoscopic Mayo score >1] starting vedolizumab [$n = 33$] or anti-TNF [$n = 45$] and controls [$n = 22$] were included. Colon biopsies [at weeks 0, 14 and 46] and blood samples [at weeks 0, 2, 6, 14, 30 and 46] were used for cell phenotyping, transcriptional analysis [qPCR], and to measure receptor occupancy.

Results: Vedolizumab, in contrast to anti-TNF, significantly reduced the proportion of $\alpha 4\beta 7^+$ cells within intestinal T subsets while preserving the percentage of $\alpha 4\beta 7^+$ plasma cells. The marked decrease in $\alpha 4\beta 7$ did not change the percentage of colonic $\alpha E\beta 7^+$ cells [at 46 weeks]. Both vedolizumab and anti-TNF significantly downregulated inflammation-related genes in the colon of responders [Mayo score < 2]. Moreover, both treatments significantly decreased the percentage of intestinal, but not blood, total lymphocytes [T and plasma cells], as well as the proportion of $\alpha 4\beta 1^+$ cells within intestinal T lymphocytes.

Conclusions: Our data show that while vedolizumab and anti-TNF block two unrelated targets, they induce remarkably similar effects. On the other hand, vedolizumab's unique mechanism of action relies on blocking intestinal trafficking of $\alpha 4\beta 7$ T cells, despite effectively binding to B and plasma cells that express $\alpha 4\beta 7$.

Key Words: Vedolizumab; inflammatory bowel disease; $\alpha E\beta 7$; $\alpha 4\beta 1$

1. Introduction

Ulcerative colitis [UC] is a chronic remitting and relapsing inflammatory bowel disease [IBD] that affects the large intestine and is characterized by superficial mucosal inflammation that can lead to diarrhoea and rectal bleeding. Available treatments include 5-aminosalicylates for mild flares and corticosteroids and immunosuppressants for moderate-to-severe disease. Patients with inadequate response to conventional therapies are eligible to receive biological therapy including anti-tumour necrosis factor [TNF] antibodies, anti- α 4 β 7 or, more recently, a pan-Janus kinase inhibitor [tofacitinib].^{1,2} While these options target seemingly independent pathways, they can all have potent anti-inflammatory effects and promote healing of the colonic mucosa in a proportion of patients.^{3–6}

Anti-TNF blocking monoclonal antibodies [mAbs] interfere with the effects of TNF, a potent and pleiotropic cytokine that promotes phagocytosis, bacterial killing, leukocyte recruitment, antigen presentation and tissue remodelling. In addition, anti-TNF mAbs have been shown to bind to membrane-bound TNF and promote apoptosis of cells potentially involved in the inflammatory cascade.⁷ In contrast to anti-TNF, vedolizumab, approved for the treatment of both UC and Crohn's disease, binds to the α 4 β 7 integrin and blocks its binding to endothelial mucosal addressin cell adhesion molecule-1 [MAdCAM-1].^{8,9} α 4 β 7 is primarily expressed by certain T-cell subsets^{10,11} and B cells,^{12,13} including plasma cells,^{14–16} although it can also be expressed in innate cells such as granulocytes, natural killer cells⁸ and monocytes.¹⁷ Studies in mice,¹⁸ and later in primates,¹⁹ have demonstrated the involvement of this receptor in the specific homing of memory T cells towards gut-associated lymphoid tissues. α 4 β 7 binds to MAdCAM-1, a receptor that is highly upregulated in the intestinal vasculature of both UC and Crohn's disease patients²⁰ and that mediates leukocyte recruitment towards the lamina propria. In addition, α 4 β 7 and MAdCAM-1-dependent interactions have been reported to support naïve T- and B-cell recruitment towards induction sites [i.e. Peyer patches and intestinal lymphoid aggregates].^{21,22}

Despite the wide use of anti-TNF and anti-integrin therapy, the differential mechanisms of action underlying these drugs have not been completely characterized.

Here, we provide novel information on the effects that vedolizumab exerts on lymphoid cell circulation and accumulation in both the systemic and the colonic compartments of a group of UC patients receiving vedolizumab for up to 1 year. Our goal is to understand vedolizumab's mechanisms of action. By comparing with an independent cohort of patients receiving anti-TNF therapy we aim to differentiate between the specific, as well as non-specific, downstream effects of vedolizumab therapy.

2. Materials and methods

Additional details of the materials and methods used are given in the [Supplementary Materials](#).

2.1 Study subjects and sample collection

Patients with an established diagnosis of UC of at least 3 months' duration, and non-IBD controls [Ctrl] were included in the study between May 31, 2017 and November 5, 2019. All patients signed a written informed consent. Controls were subjects undergoing a colonoscopy for mild gastrointestinal symptoms, or screening for colorectal cancer that presented no mucosal lesions. UC patients were those starting treatment with an anti-TNF antibody [infliximab, adalimumab or golimumab] or vedolizumab and followed-up for a

maximum of 46 weeks. Patients underwent clinical and endoscopic evaluations at weeks 0, 14 and 46. All biopsy samples from controls and UC patients were obtained from the sigmoid colon [or rectum in the case of proctitis]. Follow-up samples were obtained whenever possible from the same locations. Blood was collected from UC patients at weeks 0, 2, 6, 14, 30 and 46. A total of 33 controls and 67 UC patients were recruited for this study. Whenever possible, both blood and biopsies were obtained from the same individuals. Patients who were switched to a different treatment due to lack of response or disease worsening dropped out of the study. [Table S1](#) shows the clinical and demographic characteristics of the subjects included in the study stratified according to the type of sample and analysis. Patients were recruited at the Department of Gastroenterology, Hospital Clinic Barcelona, and the Hospital Universitari Mutua Terrassa. The study protocol was approved by the ethics committees at both centres.

2.2 Assessment of disease activity

Endoscopic UC activity at the time of the colonoscopy was assessed according to the Mayo endoscopic subscore.²³ Active UC was defined as a Mayo endoscopic subscore of >1. Endoscopic improvement was defined as a Mayo endoscopic subscore \leq 1 at week 14 or 46.

2.3 Intestinal cell isolation

Colonic biopsies [$n = 2–3$ per patient] were collected from controls [$n = 10$] and UC patients [$n = 47$] and submitted to enzymatic digestion to obtain a cell suspension for flow analysis [[Table S1](#), Group1].

2.4 Cell staining

Single-cell suspensions obtained from digested biopsies were stained with a Zombie Aqua Fixable Viability Kit [BioLegend] to exclude dead cells and with the following antibodies: CD20 FITC, CD8 FITC, α 4[CD49d] PE [clone 9F10], CD3 PerCP, CD38 PerCP, CD4 PE-Cy7, α 4[CD49d] PE-Cy7 [clone 9F10], β 7 APC [clone FIB504], β 1[CD29] APC-Cy7, α E[CD103] BV421, CD19 BV421 [all from Biolegend] and α E[CD103] PE (Becton Dickinson [BD]).

Fresh blood from controls [$n = 20$] and UC patients [$n = 51$] was collected into an EDTA K2 Vacutainer [BD] and immediately processed or kept in a rocking platform at 4°C [<30 min] until processing [[Table S1](#), Group 3]. Two hundred microliters of blood was incubated for 15 min with Red Blood Cell Lysis Buffer [BioLegend] and washed twice with PBS. Cells were stained with CD8 FITC, CD19 FITC, β 1[CD29] PE, CD38 PerCP, α 4[CD49d] PE-Cy7 [clone 9F10], β 7 APC [clone FIB504], CD45RA APC-Fire750, IgD APC-Cy7, α E[CD103] BV421 [BioLegend] and CD4 PerCP [BD].

Intestinal and blood samples were fixed using BD Stabilizing Fixative [BD], and Precision Count Beads [BioLegend] were then added. Samples were acquired using a BD FACSCanto II flow cytometer [BD] and analysed with FlowJO software [BD].

2.5 Occupancy assay

In brief, cells were co-stained with the anti- β 7 PE [clone FIB540; BioLegend] antibody and vedolizumab labelled with Alexa Fluor 647.

2.6 RNA isolation

Biopsies [$n = 8$ controls and $n = 58$ patients] were placed in RNAlater RNA Stabilization Reagent [Qiagen] and stored at -80°C . Total RNA was extracted using a RNeasy Kit [Qiagen] according to the

manufacturer's instructions [Table S1, Group 2]. Blood [$n = 14$ patients] was collected in a PAXgene Blood RNA System [PreAnalytiX GmbH] and stored at -20°C . RNA was isolated using the PAXgene Blood RNA Kit [Qiagen] in a QIAcube instrument [Qiagen] following the manufacturer's instructions [Table S1, Group 2]. The purity and integrity of the total RNA were assessed with the 2100 Bioanalyzer [Agilent] and then quantified via a NanoDrop spectrophotometer [Nanodrop technologies]; only samples with an RNA integrity number [RIN] >7.0 were used.

2.7 Quantitative polymerase chain reaction

Total RNA was transcribed to cDNA using a reverse transcriptase [High Capacity cDNA Archive RT kit; Applied Biosystems]. For biopsy RNA, quantitative polymerase chain reaction [qPCR] was performed on selected genes including three endogenous controls [GAPDH, PPIC and UBA3] using custom-designed TaqMan Low Density Arrays [TLDA platform; Applied Biosystems]. For blood RNA, qPCR analysis was performed using a TaqMan Universal PCR Master Mix [Applied Biosystems] containing the probe of interest and β -actin [TaqMan primers and probes; Applied Biosystems]. Data are expressed as arbitrary units [AU], applying the $2^{-\text{dCt}}$ formula and correcting by the value of the endogenous genes.

2.8 Immunohistochemistry

Colonic biopsies were placed in 10% formalin for 24 h and then embedded in a paraffin block. Immunohistochemical staining of tissue sections was performed using commercially available rabbit anti-human CD103 antibody [1:1000; clone EPR4166(2); Abcam]. Image acquisition was performed on an Olympus BX51 microscope using CellF Software.

2.9 ELISA to measure vedolizumab levels

Serum was collected [Table S1, Group 4, $n = 51$] to determine vedolizumab concentration using a serum separator tube, and the sample was then stored at -80°C . The concentration of vedolizumab in serum samples was determined using the Promonitor-vedolizumab ELISA kit [Progenika] following the manufacturer's instructions.

2.10 Soluble $\alpha 4\beta 7$ ELISA

A home-made ELISA was developed to measure levels of soluble $\alpha 4\beta 7$ in serum [Group 4].

2.11 Statistical analysis

Graphs were made and statistical analysis was performed using Graphpad Prism [Graphpad Software]. Only data from available samples at the studied time point are included and provided as observed. Differences between groups were tested using the non-parametric Kruskal–Wallis test and correcting for multiple comparisons by controlling the false discovery rate [two-stage linear step-up procedure of Benjamini, Krieger and Yekutieli]. Corrected p -values <0.05 were considered statistically significant.

3. Results

3.1 Treatment with vedolizumab achieves full occupancy of $\alpha 4\beta 7$ on circulating and intestinal T and B lymphocytes

To quantify $\alpha 4\beta 7$ occupancy in patients receiving vedolizumab, we set up an assay using in-house fluorochrome-conjugated

vedolizumab together with a non-competing anti- $\beta 7$ antibody and measured both vedolizumab-free $\alpha 4\beta 7$ and total $\beta 7$ expression independently of bound vedolizumab. Baseline $\alpha 4\beta 7$ expression patterns, shown in Supplementary Figure 1 for the different blood and biopsy lymphocyte subsets, were in agreement with earlier studies.^{10,11,14,24} We quantified receptor occupancy before and after treatment with vedolizumab in blood [Supplementary Figure 2], as well as in colonic biopsies [Figure 1]. In agreement with previous studies involving humans²⁵ and primates,^{9,26} we found full occupancy [absence of free $\alpha 4\beta 7$] in blood T and B cells at weeks 2, 6, 14, 30 and 46 of vedolizumab treatment [Supplementary Figure 2]. Vedolizumab serum concentrations were also measured and compared between responders and non-responders [Supplementary Figure 3]. In agreement with published data,²⁷ serum levels at an early time point [week 6] were significantly increased in responders compared to non-responders, while at the end of induction [weeks 14 and 46] no significant differences were observed. In colonic cells, occupancy was also 100% within CD4⁺ T cells and CD19⁺ lymphocytes [Figure 1]. In contrast, $\alpha 4\beta 7$ was incompletely occupied [$<99\%$ occupancy] on intestinal CD8⁺ T cells in 54% of patients at week 14 and in 71% of patients at week 46. Nonetheless, the presence of vedolizumab-free $\alpha 4\beta 7$ on intestinal CD8⁺ cells was not associated with lack of response to vedolizumab [shown in Figure 1B as open circles] nor was it significantly correlated with lower serum vedolizumab concentrations [$p = 0.1603$]. Indeed, the percentage of intestinal CD8⁺ with unoccupied $\alpha 4\beta 7$ was not significantly different between responders and non-responders at weeks 14 or 46 [data not shown].

3.2 Vedolizumab and anti-TNF treatment can reduce total lymphocytes in the colon of UC patients

At baseline, patients with active UC showed a significant increase in the percentage of CD3⁺CD8⁺ and CD19⁺CD38⁺ cells within the inflamed colonic lamina propria compared to healthy controls [Figure 2A and B]. Patients in the vedolizumab cohort [a majority of them refractory to anti-TNF] showed significantly higher percentages of CD3⁺CD4⁺ T cells at week 0 compared to controls [Figure 2A]. This percentage was significantly downregulated at weeks 14 [$p < 0.05$] and 46 [$p < 0.001$] of vedolizumab treatment compared to week 0, reaching at the final time point levels similar to those of healthy controls. In addition, treatment with either vedolizumab or anti-TNF induced a significant decrease, most pronounced at week 46, in the percentage of colonic CD3⁺CD8⁺ T cells and CD19⁺CD38⁺ plasma cells, taking into account all patients in the group regardless of treatment response [Figure 2A and B].

Despite their different mechanism of action, vedolizumab and anti-TNF induced similar changes in the expression of a number of genes upregulated in active UC. Indeed, we confirmed that endoscopic improvement [Mayo endoscopic score ≤ 1] was associated with a significant downregulation of a panel of genes related to innate [neutrophil, monocyte and macrophages] and adaptive [lymphocytes] inflammatory mediators [CXCL9, CXCL10, FCGR3B, IL23A, IL17, IFNG] in both anti-TNF- and vedolizumab-treated patients [Figure 2C].

In contrast to their effects within the intestinal mucosa, neither vedolizumab nor anti-TNF had any significant effect on the percentages of circulating CD4⁺, CD8⁺ or CD19⁺ lymphocytes in peripheral blood up to 46 weeks after treatment initiation [Figure 3].

Overall, we observed that blocking either $\alpha 4\beta 7$ or TNF led to reductions in the sizes of T- and plasma cell populations in the colon of UC patients without affecting circulating subsets. While




Figure 1. Treatment with vedolizumab results in high $\alpha 4\beta 7$ occupancy within intestinal lymphocytes. Integrin occupancy was assessed by co-staining with fluorescently labelled vedolizumab and the non-competing anti- $\beta 7$ mAb. [A] Flow cytometry plots of intestinal CD4⁺ and CD8⁺T cells, as well as of CD19⁺CD38⁺ plasma cells at weeks 0, 14 and 46 after initiation of vedolizumab [VDZ] treatment of a representative UC patient. Gates show cells with unoccupied $\alpha 4\beta 7$ [VDZ⁺ $\beta 7$ ⁺] and the percentage on each cell subset is depicted for a representative sample. [B] Percentage of unoccupied cells within the different lymphocyte populations at weeks 0, 14 and 46. Patients are represented as responders [closed circles] and non-responders [open circles] at weeks 14 or 46. Plot lines show median and range.

vedolizumab appears to have a more pronounced effect on the intestinal CD4⁺ T cell subset, this may also stem from the fact that the vedolizumab cohort, but not the anti-TNF cohort, included in our study presented an increased percentage of CD4⁺ T cells at baseline.

3.3 Vedolizumab specifically reduces intestinal $\alpha 4\beta 7$ ⁺T cells and overall surface expression of $\alpha 4\beta 7$ in blood

We next measured the effects of treatment on the proportions of $\alpha 4\beta 7$ ⁺ cells within the different lymphocyte compartments. Vedolizumab significantly decreased the percentage of intestinal $\alpha 4\beta 7$ ⁺ within the CD4⁺ and CD8⁺ T-cell compartments at weeks 14 and 46 of follow-up, taking into account all patients in the group regardless of response [Figure 4A]. In contrast, despite decreasing total CD8⁺ intestinal lymphocytes [Figure 2B], anti-TNF treatment did not change the percentage of $\alpha 4\beta 7$ ⁺ cells within the CD8 compartment [Figure 4B].

The majority of plasma cells [CD19⁺CD38⁺] in the colon expressed high levels of $\alpha 4\beta 7$ [Supplementary Figure 1 and Figure 4]. Nonetheless, and despite full receptor occupancy [Figure 1], the proportion of $\alpha 4\beta 7$ ⁺ cells within the plasma cell compartment was not changed by vedolizumab or anti-TNF treatment [Figure 4]. These data demonstrate that while both treatments can reduce the total number of plasma cells in the intestinal mucosa, the proportion of those cells that express $\alpha 4\beta 7$ remains the same for up to 46 weeks of follow-up.

In addition, vedolizumab, but not anti-TNF treatment, selectively increased the percentage of $\alpha 4\beta 7$ ⁺ cells within memory CD4⁺CD45RA⁻ and CD19⁺IgD⁻ cells in the blood [Figure 5A] regardless of the response to treatment. This mild, but significant, increase was only detected at 1 year after initiation of the treatment. Nonetheless, in agreement with previous reports,²⁸ vedolizumab significantly decreased the surface expression of $\beta 7$ [measured by a decrease in the mean fluorescence intensity of a non-competing anti-beta7 mAb] on all circulating memory T and B lymphocytes [Figure 5B]. The effect of vedolizumab on integrin surface expression was already detectable at week 2, the earliest time point studied. This was not true, however, in anti-TNF-treated patients, suggesting that vedolizumab exerts specific and fast-acting effects on $\alpha 4\beta 7$ protein expression.

Vedolizumab-induced $\alpha 4\beta 7$ internalization has been previously suggested,^{29,30} and could indeed explain the reduction in $\beta 7$ expression *in vivo*. To test this hypothesis, we incubated control peripheral blood mononuclear cells [PBMCs] with vedolizumab at different doses [0.5–50 $\mu\text{g}/\text{mL}$], or an anti- $\beta 7$ antibody [clone FIB504; 0.5–10 $\mu\text{g}/\text{mL}$], and measured $\alpha 4\beta 7$ expression 24 h later. Even concentrations of 50 $\mu\text{g}/\text{mL}$ of vedolizumab [over therapeutic serum concentrations] did not induce receptor internalization [Supplementary Figure 4A]. In contrast, the anti- $\beta 7$ antibody [clone FIB504] caused a robust dose-dependent internalization of $\alpha 4\beta 7$ [Supplementary Figure 4A].³¹

In addition, when PBMCs from vedolizumab-treated patients were cultured *in vitro* they recovered expression of membrane $\alpha 4\beta 7$




Figure 2. Vedolizumab and anti-TNF reduce total intestinal lymphocytes and decrease inflammatory gene expression. Percentages of intestinal CD3⁺CD4⁺, CD3⁺CD8⁺ and CD19⁺CD38⁺ in controls [Ctrl] and UC patients treated with [A] vedolizumab or [B] anti-TNF at weeks 0, 14 and 46. Patients are represented as responders [closed circles] and non-responders [open circles] at weeks 14 or 46. Scatter dot plot lines show median \pm range taking into account responders and non-responders in the colon: in controls [Ctrl] and UC patients treated with vedolizumab or anti-TNF at weeks 0, 14 and 46. Differential gene expression analysis was performed for responders and non-responders separately. Data are expressed as arbitrary units [AU]. Bar graph lines show median \pm range. False discovery rate-corrected *p*-values: **p* < 0.05; ***p* < 0.01; and ****p* < 0.001.




Figure 3. Vedolizumab and anti-TNF have no significant effect on peripheral blood total circulating lymphocytes. Percentages of circulating CD3⁺CD4⁺, CD3⁺CD8⁺ and CD19⁺CD38⁺ lymphocytes from total blood mononuclear cells in controls [Ctrl], UC patients treated with vedolizumab [A] or anti-TNF [B] at weeks 0, 2, 6, 14, 30 and 46. Patients are represented as responders [closed circles] and non-responders [open circles] at weeks 14 or 46. Scatter dot plot lines show median \pm range taking into account responders and non-responders. False discovery rate-corrected p -value: ** < 0.01 .




Figure 4. Vedolizumab decreases the frequency of $\alpha 4\beta 7$ T cells but not plasma cells. Percentage of $\alpha 4\beta 7$ cells in CD3⁺CD4⁺, CD3⁺CD8⁺ and CD19⁺CD38⁺ intestinal lymphocytes determined by flow cytometry using co-staining with anti- $\alpha 4$ and anti- $\beta 7$ antibodies. UC patients receiving vedolizumab [A] or anti-TNF [B] at weeks 0, 14 and 46. Patients are represented as responders [closed circles] and non-responders [open circles] at weeks 14 or 46. Plot lines show median \pm range taking into account responders and non-responders. False discovery rate-corrected p -value: * < 0.05 ; ** < 0.01 ; and *** < 0.001 .

in a time-dependent manner [Figure 5C]. This new surface expression was prevented by cycloheximide [an inhibitor of protein synthesis], further suggesting that the integrin was not being recycled but rather synthesized *de novo* [Figure 5C]. We also tested the possibility of vedolizumab driving receptor shedding *in vivo*. In contrast to a recent study,³² we could not detect soluble $\alpha 4\beta 7$ in any of our samples [data not shown]. The test we used, however, had a detection limit of 26.9 ng/mL, which is higher than the concentrations previously reported [< 15 ng/mL].³² Thus, we cannot rule out a higher cleavage rate of the integrin as an explanation for the reduced density observed in the cellular surface.

Finally, we tested the hypothesis that vedolizumab may be decreasing the transcription of *ITGB7* and/or *ITGA4*, the genes encoding for the $\alpha 4$ and $\beta 7$ chain. Vedolizumab had no significant effect on *ITGB7* and *ITGA4* transcription in whole blood RNA for up to 46 weeks of treatment [Supplementary Figure 4B].

3.4 Vedolizumab does not significantly change the percentage of mucosal $\alpha E\beta 7$ lymphocytes

Expression of the αE [CD103] chain, which forms a heterodimer with the $\beta 7$ chain, is acquired in mucosal surfaces in response to epithelial-derived transforming growth factor- β [TGF- β]³³. Indeed,




Figure 5. Vedolizumab induces changes in the expression of $\alpha 4\beta 7$ in blood lymphocytes. [A] Percentage of $\alpha 4\beta 7^+$ CD4⁺ and CD8⁺ memory [CD45RA⁻] blood T and B [CD19⁺IgD⁻] cells determined by flow cytometry using co-staining with anti- $\alpha 4$ and anti- $\beta 7$ antibodies. [B] Mean fluorescence intensity [MFI] for $\beta 7$ is shown for memory blood T and B cells. For [A] and [B] the top row shows UC patients receiving vedolizumab at weeks 0, 2, 6, 14, 30 and 46; bottom row shows UC patients receiving anti-TNF at weeks 0, 2, 6, 14, 30 and 46. Patients are represented as responders [closed circles] and non-responders [opened circles] at weeks 14 or 46. Plot lines show median \pm range taking into account responders and non-responders. False discovery rate-corrected *p*-value: **p* < 0.05; ***p* < 0.01; and ****p* < 0.001. [C] Expression of $\alpha 4\beta 7$ determined by flow cytometry in blood memory CD4⁺CD45RA⁻ cells is shown at days 0, 1 and 4 after *in vitro* culture in complete media [CM] in the absence of vedolizumab. A representative sample is shown from a UC patient at week 46 after starting vedolizumab treatment. Cycloheximide [CHX] at 5 μ g/mL was added under the indicated conditions. One representative experiment out of three is shown.

$\alpha E\beta 7$ -expressing cells are abundant in the healthy intestinal lamina propria within both the intra-epithelial and the lamina propria compartments [Figure 6A]. In agreement with the literature,^{24,33} the lymphocyte subset with the highest percentage of $\alpha E\beta 7^+$ in the colon was CD8⁺ T cells, followed by CD4⁺ T cells [Figure 6B], while plasma cells did not express $\alpha E\beta 7$ [data not shown]. It has been proposed that $\alpha 4\beta 7^+$ T cells give rise to $\alpha E\beta 7^+$ T cells.³⁴ To test this hypothesis, we measured the effect that blocking $\alpha 4\beta 7$ would have on $\alpha E\beta 7$ cells. Remarkably, treatment with vedolizumab, despite significantly reducing the number of $\alpha 4\beta 7^+$ T cells in the mucosa, had no significant effect on the percentage of $\alpha E\beta 7^+$ T lymphocytes [Figure 6C], suggesting that $\alpha E\beta 7$ precursors may not be entering the mucosa primarily through $\alpha 4\beta 7$.

Nonetheless, vedolizumab, but not anti-TNF, did significantly increase the number of circulating $\alpha E\beta 7$ memory T cells as soon as 14 weeks after treatment initiation [Figure 6D]. This observation supports a role for $\alpha 4\beta 7$ on $\alpha E\beta 7^+$ T-cell recirculation.

Indeed, we looked into $\alpha 4\beta 7$ expression on both CD8⁺ and CD4⁺ $\alpha E\beta 7^+$ cells and found that while only a small percentage of $\alpha E\beta 7^+$ cells in the colon expressed $\alpha 4\beta 7$, over 50% of circulating $\alpha E\beta 7$ cells co-expressed the homing receptor $\alpha 4\beta 7$ [data not shown].

3.5 Vedolizumab's effects on intestinal $\alpha 4\beta 1^+$ cellular migration

In addition to $\alpha 4\beta 7$, leukocytes can employ $\alpha 4\beta 1$ to migrate into the inflamed intestine by binding to vascular cell-adhesion molecule [VCAM]-1. In contrast to $\alpha 4\beta 7$ [Figure 4A], the percentage of $\alpha 4\beta 1^+$ CD4⁺ T cells was significantly increased and to a similar extent in the mucosa of UC patients with active disease naïve to anti-TNF therapy or in those with prior anti-TNF failure [treated with vedolizumab] [week 0] [Figure 7]. Both treatments significantly reduced the percentage of $\alpha 4\beta 1^+$ CD4⁺ T cells, suggesting a non-specific effect on $\alpha 4\beta 1$ -expressing T-cell populations.

4. Discussion

Despite its approval in 2014 for the treatment of UC and Crohn's disease, the mechanisms of action underlying vedolizumab are still not fully understood. Vedolizumab was initially developed under the assumption that it would specifically block recruitment of mucosal homing T cells to the intestine,¹⁰ thereby dampening inflammation in a gut-restricted manner. Data from primates revealed rapid amelioration of colitis in cotton-top tamarins treated with an anti- $\alpha 4\beta 7$ mAb¹⁹ [ACT-1, a murine IgG1 version of vedolizumab] and a significant reduction in $\beta 7^+$ cells, as well as neutrophils and macrophages in the intestine of cynomolgus monkeys receiving vedolizumab.^{9,26} Both studies also showed the saturation of $\alpha 4\beta 7$ on blood lymphocytes by ACT-1 or vedolizumab. Subsequently, nearly full occupancy [$>95\%$ of $\alpha 4\beta 7$ being occupied by vedolizumab] in peripheral^{3,28} and intestinal cells²⁵ was reported in IBD patients. Our data confirm that vedolizumab administered at established clinical doses results in full $\alpha 4\beta 7$ occupancy in blood T and B cells, as well as in intestinal lymphocytes, except for a proportion of patients in whom we detected unoccupied $\alpha 4\beta 7$ on intestinal CD8⁺ cells, despite the full saturation of CD4⁺ and plasma cells in the same patient and tissue. A previous study had also shown incomplete occupancy of $\alpha 4\beta 7$ in intestinal memory CD3⁺ T cells.²⁵ This observation may reflect differential tissue antibody penetration depending on the cellular compartment [e.g. intra-epithelial compartment]. Importantly, incomplete CD8⁺ T-cell receptor occupancy in

tissue was not associated in our study with lack of therapeutic response during follow-up.

The number of lymphocytes populating the intestinal mucosa can change significantly, as we have confirmed here, in active UC due to increased cell recruitment and resident cell proliferation.³⁵⁻³⁸ Consequently, any therapy that targets inflammation and/or immune responses [immunosuppressors, corticosteroids, anti-TNF, anti-leukocyte trafficking therapy, etc.] could potentially have an impact on the total numbers and proportions of cells populating the intestinal mucosa of patients. While total lymphocytes are increased in active UC, we show that the proportion of lymphocytes [20-30% of T cells and 80% of plasma cells] that express $\alpha 4\beta 7$ was not significantly altered by inflammation. In contrast, there was an enrichment of $\alpha 4\beta 1$ -expressing CD4⁺ T cells in UC patients. $\alpha 4\beta 7$ and $\alpha 4\beta 1$, which are not co-expressed in most lymphocytes^{10,11} [and data not shown], mediate cell recruitment to the inflamed colonic mucosa by binding to independent receptors on the endothelial lining [MAdCAM-1 and VCAM-1, respectively] and therefore are independently regulated, as we confirm here.

The observation that both vedolizumab and anti-TNF decrease the proportion of $\alpha 4\beta 1^+$ CD4⁺ T cells in the mucosa suggests that this effect is secondary to the control of inflammation exerted by both antibodies, as neither of them directly targets $\alpha 4\beta 1$. In contrast to the $\beta 1$ integrin, we demonstrate that only vedolizumab, but not anti-TNF, specifically reduces the proportion of $\alpha 4\beta 7$ -expressing CD4⁺ and CD8⁺ T cells in the mucosa, while eventually increasing circulating memory CD4⁺ $\alpha 4\beta 7^+$ T cells and CD19⁺IgD⁻ $\alpha 4\beta 7^+$ B cells. These findings support the initial hypothesis that vedolizumab acts by reducing $\alpha 4\beta 7$ -mediated T-cell homing to the gut^{9,30} and are in agreement with recent data on mucosal CD4⁺ cells.³⁹

There is surprisingly little consensus on the effects of vedolizumab on resident tissue and tissue-infiltrating T and B cells, with no study thus far providing information beyond the 14-week induction period. Indeed, a recent article examined total CD3⁺ cells [as well as B cells and plasma cells] in lamina propria and the intra-epithelial compartment, separately, in a mixed cohort of UC and Crohn's disease patients starting vedolizumab, and reported no change in the percentages at the end of induction⁴⁰. In contrast, in an independent group of UC and Crohn's disease patients, the authors describe significant changes in total T-cell lamina propria populations after anti-TNF treatment. Inconsistencies in the total T-cell numbers, and therefore in the ability of a treatment to modify them significantly, can be explained by disease heterogeneity between cohorts, especially when including Crohn's disease patients, who are a highly heterogeneous population. More surprising is their observation that anti-TNF treatment had no effect on innate-related signatures, which is a consistently reported result in the literature.^{37,41-43} In addition, their study focused on total T cells and did not analyse the effects of treatment specifically on $\alpha 4\beta 7^+$ cells, which our current results show is the cell population in which greater differences can be observed between vedolizumab and anti-TNF-treated patients. In contrast, our results show that both anti-TNF and vedolizumab significantly modulate the transcriptional signature related to innate and acquired immune responses in responding patients. In agreement with these effects, the overall number of intestinal lymphocytes is markedly reduced by treatment with either therapy.

Beyond promoting homing to the intestinal mucosa, $\alpha 4\beta 7$ [but not $\alpha 4\beta 1$] mediates the recruitment of naïve T and B cells to gut-associated secondary lymphoid tissues,^{21,22} suggesting that vedolizumab could also impact primary responses to new antigens present in the intestinal lumen. As per the limitations of our




Figure 6. Differential effect of vedolizumab on intestinal or circulating $\alpha E\beta 7^+$ T cells. [A] Representative immunohistochemistry of a control colon stained with anti- αE antibody. Positive cells are abundant along the intestinal epithelium [white arrow] and the lamina propria [black arrow]. [B] Representative plots showing expression of αE and $\beta 7$ on intestinal $CD3^+CD4^+$ and $CD3^+CD8^+$ T cells. [C] Graphs showing the percentage of $\alpha E\beta 7^+$ in $CD3^+CD4^+$ and $CD3^+CD8^+$ intestinal lymphocytes determined by flow cytometry at weeks 0, 14 and 46. [D] Percentage of $\alpha E\beta 7^+$ within memory $CD3^+CD4^+$ and $CD3^+CD8^+$ peripheral blood circulating cells shown at weeks 0, 2, 6, 14, 30 and 46. All frequencies were measured in controls [Ctrl] and UC patients treated with vedolizumab or anti-TNF. Patients are represented as responders [closed circles] and non-responders [open circles] at weeks 14 or 46. Plot lines show median \pm range taking into account responders and non-responders. False discovery rate-corrected p -value: * < 0.05 ; ** < 0.01 ; and *** < 0.001 .




Figure 7. Vedolizumab and anti-TNF decrease the frequency of $\alpha 4\beta 1$ T cells but not plasma cells. Percentage of $\alpha 4\beta 1$ ⁺ cells in CD3⁺CD4⁺, CD3⁺CD8⁺ and CD19⁺CD38⁺ intestinal lymphocytes determined by flow cytometry using co-staining with anti- $\alpha 4$ and anti- $\beta 1$ antibodies. [A] UC patients receiving vedolizumab at weeks 0, 14 and 46. [B] UC patients receiving anti-TNF at weeks 0, 14 and 46. Patients are represented as responders [closed circles] and non-responders [open circles] at weeks 14 or 46. Plot lines show median \pm range taking into account responders and non-responders. False discovery rate-corrected *p*-value: * < 0.05 ; ** < 0.01 ; and *** < 0.001 .

observational study, we did not have access to inductive sites [i.e. lymph nodes draining the intestinal mucosa or Peyer patches in the small intestine], and thus we could not directly assess the effects of vedolizumab on leukocyte circulation within these sites. Nonetheless, previous evidence supports the role of $\alpha 4\beta 7$ -mediated interactions in maintaining those lymphoid aggregates abundant in the ileal mucosa.⁴⁴ Uzzan and colleagues studied the ileum and left colon [both uninvolved mucosa] of UC patients with concomitant chronic HIV infection. In this unique cohort they showed that vedolizumab, even in the absence of inflammation, decreased the number of naïve T cells and B cells in lymphoid aggregates. In line with this, another study demonstrated the role that $\alpha 4\beta 7$ -dependent cell recruitment plays in mounting humoral immune responses to oral vaccines, which was markedly reduced in vedolizumab-treated patients, while still preserving the response to systemic vaccination.⁴⁵ Altogether, these observations would support a role for vedolizumab that goes beyond the canonical view of exclusively interfering with effector memory T-cell recruitment to the lamina propria. Nonetheless, whether it plays a role in controlling chronic intestinal inflammation remains to be proven.

Our results, in agreement with others,⁴⁴ also rule out a role for $\alpha 4\beta 7$ on plasma cell recruitment to the intestinal mucosa. Indeed, while both vedolizumab and anti-TNF significantly reduced the overall plasma cell population in the colonic mucosa of UC patients, the remaining plasma cells showed high percentages of $\alpha 4\beta 7$ expression.⁴⁴ This observation suggests that, in contrast to other lymphocytes, plasma cells may not use $\alpha 4\beta 7$ as a receptor for intestinal homing. Beyond binding to endothelial adhesion molecules [MAdCAM-1 and VCAM-1], $\alpha 4\beta 7$ has been described to use the extracellular matrix protein fibronectin as ligand.⁴⁶ The functional role of $\alpha 4\beta 7$ -fibronectin interactions, specifically on plasma cells, has not been yet elucidated but we hypothesize that it could drive intracellular signals required for cell survival, differentiation or activation similarly to the integrin-fibronectin interactions⁴⁷.

An intriguing observation that can be made from our study is vedolizumab's apparent lack of effect on intestinal $\alpha E\beta 7$ ⁺ T cells. The role of $\alpha E\beta 7$ ⁺ T cells in IBD remains ambiguous. While some studies suggest that $\alpha E\beta 7$ ⁺ lymphocytes may promote inflammation in UC

patients,⁴⁸ these cells abundantly populate the healthy intestinal mucosa and intra-epithelial compartment and recent studies have found no changes during inflammation of this subset.^{33,49}

It has been assumed that $\alpha E\beta 7$ may arise from $\alpha 4\beta 7$ cells entering the gut and responding to environmental cues.³⁴ Nonetheless, the majority of $\alpha E\beta 7$ cells found within,⁵⁰ or migrating towards,⁵¹ the epithelial layer express low levels of $\alpha 4\beta 7$. Here, we noted that the majority of $\alpha E\beta 7$ ⁺ T cells in the colon did not co-express $\alpha 4\beta 7$. Moreover, vedolizumab treatment, while significantly decreasing the number of $\alpha 4\beta 7$ cells, had no clear impact on the intestinal $\alpha E\beta 7$ compartment. This would suggest that $\alpha E\beta 7$ can arise independently of gut homing $\alpha 4\beta 7$ ⁺ lymphocytes, a hypothesis that would require further confirmation.

In striking contrast to the intestinal subset and as previously reported,³³ most circulating $\alpha E\beta 7$ co-expressed $\alpha 4\beta 7$ and their numbers increased in the blood of patients treated with vedolizumab. However, the significance of circulating double $\beta 7$ -expressing T cells remains an unanswered question. We propose that they may represent a population of gut-imprinted cells that circulate between inductive sites and blood with the ability to enter the mucosa in an $\alpha 4\beta 7$ -dependent manner. In addition, $\alpha E\beta 7$ ⁺ in the blood were CD69⁻ [data not shown], supporting the view that they do not constitute tissue-resident T cells escaping mucosal tissues.⁵² While vedolizumab blocks the homing of these cells, as supported by their accumulation in blood, our data also support the conclusion that $\alpha 4\beta 7$ -independent pathways are required to maintain the $\alpha E\beta 7$ population in the gut. Indeed, to our knowledge no data support the likelihood that $\alpha E\beta 7$ can only be acquired by $\alpha 4\beta 7$ ⁺ cells; rather, all activated T cells can quickly upregulate αE and $\beta 7$ in response to TGF- β , with a bias of αE transcription towards CD8⁺ cells.⁵³

In summary, while vedolizumab can reduce the overall T- and plasma cell populations in the intestine, and downregulate the transcription of genes related to innate immune responses, we propose that these indirect effects are secondary to the control of inflammation. We base this conclusion on the fact that anti-TNF, which does not selectively block $\alpha 4\beta 7$, results in similar decreases in T-cell and plasma cell infiltration, as well as transcriptional downregulation of immune-related genes. In contrast, vedolizumab [but not anti-TNF]

depletes $\alpha 4\beta 7^+$ CD4⁺ and CD8⁺ T cells in the colon, while preserving the proportion of $\alpha 4\beta 7^+$ plasma cells. Vedolizumab also appears to block the homing of circulating $\alpha E\beta 7^+$ T cells, although this does not result in a significant change in $\alpha E\beta 7^+$ within the mucosa, suggesting that other mechanisms beyond $\alpha 4\beta 7$ are required to maintain that key pool of cells.

Finally, our study represents a first step towards understanding the specific and unique effects of anti- $\alpha 4\beta 7$ mAbs compared to anti-TNF. We believe further studies will help us decipher key predictors of response within the identified direct mechanisms of action that, in a manner similar to predictors, should be unique to each specific drug type.⁵⁴

Funding

This work was supported by the Agència de Gestió d'Ajuts Universitaris I de Recerca [AGAUR], Generalitat de Catalunya, Grant 2017SGR992 to A.S. and the Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas [CIBER-EHD], Spain.

Conflicts of Interest

M.V., A.G.-T., A.M.C., M.C.M., H.B.-M., Mi.E., M.A., E.T. and A.F.-C. have no conflicts of interest to declare. Ma.E. has acted as a consultant, assessor or scientific advisory member for AbbVie, MSD, Tillotts Pharma, Pfizer and Takeda and received research grants for MSD and AbbVie. E.R. has served as a speaker, a consultant, or an advisory member for MSD, AbbVie, Takeda, Janssen, Pfizer, Ferring, Fresenius-Kabi and Adaclyte. I.O. has received consulting fees from AbbVie and lecture fees from AbbVie, Takeda and Janssen. J.P. has received unrestricted institutional grants from AbbVie, MSD and Pfizer, and consultancy fees from AbbVie, Boehringer Ingelheim, Celgene, Genentech, GoodGut, GSK, Janssen, MSD, Nestle, Novartis, Oppilan, Pfizer, Progenity, Shire, Robarts, Roche, Takeda, Theravance and TiGenix. A.S. has served as a speaker or scientific advisor for Pfizer, Genentech, GSK, Boehringer Ingelheim and Roche, and has received grants from Roche, Genentech, AbbVie, Scipher Medicine, Robarts, Pfizer and GSK.

Acknowledgments

We thank Joe Moore for editorial assistance and members of the Salas lab for reviewing the final manuscript. We are also indebted to all the patients who participated in the study and to the Flow Cytometry Facility and Biobank at IDIBAPS for their support.

Author Contributions

M.V. and A.S. designed the study, analysed and interpreted data and drafted and revised the manuscript. M.V., A.G.-T., Mi.E., H.B.-M. and M.A. performed experiments and analysed the data. J.P. and Ma.E. carefully reviewed the manuscript for intellectual content. I.O., A.F.-C., E.R., J.P. and Ma.E. recruited patients and performed endoscopic and clinical follow-up. E.T. and M.C.M. collected samples and provided technical support. A.M.C. analysed the data and performed statistical analysis. A.S. supervised the study and obtained financial support.

Supplementary Data

Supplementary data are available at ECCO-JCC online.

References

- Harbord M, Eliakim R, Bettenworth D, et al. Third European evidence-based consensus on diagnosis and management of ulcerative colitis. Part 2: current management. *J Crohns Colitis* 2017;11:769–84.
- Feuerstein JD, Isaacs KL, Schneider Y, Siddique SM, Falck-Ytter Y, Singh S; AGA Institute Clinical Guidelines Committee. AGA Clinical Practice Guidelines on the management of moderate to severe ulcerative colitis. *Gastroenterology* 2020;158:1450–61.
- Feagan BG, Rutgeerts P, Sands BE, et al.; GEMINI 1 Study Group. Vedolizumab as induction and maintenance therapy for ulcerative colitis. *N Engl J Med* 2013;369:699–710.
- Reinisch W, Sandborn WJ, Hommes DW, et al. Adalimumab for induction of clinical remission in moderately to severely active ulcerative colitis: results of a randomised controlled trial. *Gut* 2011;60:780–7.
- Rutgeerts P, Sandborn WJ, Feagan BG, et al. Infliximab for induction and maintenance therapy for ulcerative colitis. *N Engl J Med* 2005;353:2462–76.
- Sandborn WJ, Su C, Sands BE, et al.; OCTAVE Induction 1, OCTAVE Induction 2, and OCTAVE Sustain Investigators. Tofacitinib as induction and maintenance therapy for ulcerative colitis. *N Engl J Med* 2017;376:1723–36.
- Atreya R, Zimmer M, Bartsch B, et al. Antibodies against tumor necrosis factor (TNF) induce T-cell apoptosis in patients with inflammatory bowel diseases via TNF receptor 2 and intestinal CD14⁺ macrophages. *Gastroenterology* 2011;141:2026–38.
- Soler D, Chapman T, Yang LL, Wyant T, Egan R, Fedyk ER. The binding specificity and selective antagonism of vedolizumab, an anti- $\alpha 4\beta 7$ integrin therapeutic antibody in development for inflammatory bowel diseases. *J Pharmacol Exp Ther* 2009;330:864–75.
- Fedyk ER, Wyant T, Yang LL, et al. Exclusive antagonism of the $\alpha 4\beta 7$ integrin by vedolizumab confirms the gut-selectivity of this pathway in primates. *Inflamm Bowel Dis* 2012;18:2107–19.
- Schweighoffer T, Tanaka Y, Tidswell M, et al. Selective expression of integrin $\alpha 4\beta 7$ on a subset of human CD4⁺ memory T cells with hallmarks of gut-trophism. *J Immunol* 1993;151:717–29.
- Erle DJ, Briskin MJ, Butcher EC, Garcia-Pardo A, Lazarovits AI, Tidswell M. Expression and function of the MAdCAM-1 receptor, integrin $\alpha 4\beta 7$, on human leukocytes. *J Immunol* 1994;153:517–28.
- Postigo AA, Sánchez-Mateos P, Lazarovits AI, Sánchez-Madrid F, de Landázuri MO. $\alpha 4\beta 7$ integrin mediates B cell binding to fibronectin and vascular cell adhesion molecule-1. Expression and function of $\alpha 4$ integrins on human B lymphocytes. *J Immunol* 1993;151:2471–83.
- Rott LS, Briskin MJ, Butcher EC. Expression of $\alpha 4\beta 7$ and E-selectin ligand by circulating memory B cells: implications for targeted trafficking to mucosal and systemic sites. *J Leukoc Biol* 2000;68:807–14.
- Farstad IN, Halstensen TS, Lazarovits AI, Norstein J, Fausa O, Brandtzaeg P. Human intestinal B-cell blasts and plasma cells express the mucosal homing receptor integrin $\alpha 4\beta 7$. *Scand J Immunol* 1995;42:662–72.
- Mei HE, Yoshida T, Sime W, et al. Blood-borne human plasma cells in steady state are derived from mucosal immune responses. *Blood* 2009;113:2461–9.
- Quiding-Järbrink M, Nordström I, Granström G, et al. Differential expression of tissue-specific adhesion molecules on human circulating antibody-forming cells after systemic, enteric, and nasal immunizations. A molecular basis for the compartmentalization of effector B cell responses. *J Clin Invest* 1997;99:1281–6.
- Schleier L, Wiendl M, Heibredere K, et al. Non-classical monocyte homing to the gut via $\alpha 4\beta 7$ integrin mediates macrophage-dependent intestinal wound healing. *Gut* 2020;69:252–63.
- Wang C, Hanly EK, Wheeler LW, Kaur M, McDonald KG, Newberry RD. Effect of $\alpha 4\beta 7$ blockade on intestinal lymphocyte subsets and lymphoid tissue development. *Inflamm Bowel Dis* 2010;16:1751–62.
- Hesterberg PE, Winsor-Hines D, Briskin MJ, et al. Rapid resolution of chronic colitis in the cotton-top tamarin with an antibody to a gut-homing integrin $\alpha 4\beta 7$. *Gastroenterology* 1996;111:1373–80.
- Briskin M, Winsor-Hines D, Shyjan A, et al. Human mucosal addressin cell adhesion molecule-1 is preferentially expressed in intestinal tract and associated lymphoid tissue. *Am J Pathol* 1997;151:97–110.
- Berlin C, Berg EL, Briskin MJ, et al. $\alpha 4\beta 7$ integrin mediates lymphocyte binding to the mucosal vascular addressin MAdCAM-1. *Cell* 1993;74:185–95.
- Wagner N, Löhler J, Kunkel EJ, et al. Critical role for $\beta 7$ integrins in formation of the gut-associated lymphoid tissue. *Nature* 1996;382:366–70.

23. Schroeder KW, Tremaine WJ, Ilstrup DM. Coated oral 5-aminosalicylic acid therapy for mildly to moderately active ulcerative colitis. A randomized study. *N Engl J Med* 1987;317:1625–9.
24. Farstad IN, Halstensen TS, Lien B, et al. Distribution of beta 7 integrins in human intestinal mucosa and organized gut-associated lymphoid tissue. *Immunology* 1996;89:227–37.
25. Ungar B, Kopylov U, Yavzori M, et al. Association of vedolizumab level, anti-drug antibodies, and $\alpha 4\beta 7$ occupancy with response in patients with inflammatory bowel diseases. *Clin Gastroenterol Hepatol* 2018;16:697–705.e7.
26. Calenda G, Keawvichit R, Arrode-Brusés G, et al. Integrin $\alpha 4\beta 7$ blockade preferentially impacts CCR6+ lymphocyte subsets in blood and mucosal tissues of naive rhesus macaques. *J Immunol* 2018;200:810–20.
27. Osterman MT, Rosario M, Lasch K, et al. Vedolizumab exposure levels and clinical outcomes in ulcerative colitis: determining the potential for dose optimisation. *Aliment Pharmacol Ther* 2019;49:408–18.
28. Boden EK, Shows DM, Chiorean MV, Lord JD. Identification of candidate biomarkers associated with response to vedolizumab in inflammatory bowel disease. *Dig Dis Sci* 2018;63:2419–29.
29. Wyant T, Yang L, Fedyk E. In vitro assessment of the effects of vedolizumab binding on peripheral blood lymphocytes. *MABS* 2013;5:842–50.
30. Rath T, Billmeier U, Ferrazzi F, et al. Effects of anti-integrin treatment with vedolizumab on immune pathways and cytokines in inflammatory bowel diseases. *Front Immunol* 2018;9:1700.
31. Lichnog C, Klabunde S, Becker E, et al. Cellular mechanisms of etrolizumab treatment in inflammatory bowel disease. *Front Pharmacol* 2019;10:39.
32. Battat R, Dulai PS, Vande Castele N, et al. Biomarkers are associated with clinical and endoscopic outcomes with vedolizumab treatment in ulcerative colitis. *Inflamm Bowel Dis* 2019;25:410–20.
33. Zundler S, Schillinger D, Fischer A, et al. Blockade of $\alpha E\beta 7$ integrin suppresses accumulation of CD8+ and Th9 lymphocytes from patients with IBD in the inflamed gut in vivo. *Gut* 2017;66:1936–48.
34. Gorfu G, Rivera-Nieves J, Ley K. Role of beta7 integrins in intestinal lymphocyte homing and retention. *Curr Mol Med* 2009;9:836–50.
35. Smids C, Horjus Talabur Horje CS, Drylewicz J, et al. Intestinal T cell profiling in inflammatory bowel disease: linking t cell subsets to disease activity and disease course. *J Crohns Colitis* 2018;12:465–75.
36. Cupi ML, Sarra M, Marafini I, et al. Plasma cells in the mucosa of patients with inflammatory bowel disease produce granzyme B and possess cytotoxic activities. *J Immunol* 2014;192:6083–91.
37. Corraliza AM, Ricart E, López-García A, et al. Differences in peripheral and tissue immune cell populations following haematopoietic stem cell transplantation in Crohn's disease patients. *J Crohns Colitis* 2019;13:634–47.
38. Selby WS, Janossy G, Bofill M, Jewell DP. Intestinal lymphocyte subpopulations in inflammatory bowel disease: an analysis by immunohistological and cell isolation techniques. *Gut* 1984;25:32–40.
39. Coletta M, Paroni M, Alvisi MF, et al. Immunological variables associated with clinical and endoscopic response to vedolizumab in patients with inflammatory bowel diseases. *J Crohns Colitis* 2020;14:1190–201.
40. Zeissig S, Rosati E, Dowds CM, et al. Vedolizumab is associated with changes in innate rather than adaptive immunity in patients with inflammatory bowel disease. *Gut* 2019;68:25–39.
41. Arijs I, De Hertogh G, Lemaire K, et al. Mucosal gene expression of antimicrobial peptides in inflammatory bowel disease before and after first infliximab treatment. *PLoS One* 2009;4:e7984.
42. Arijs I, De Hertogh G, Machiels K, et al. Mucosal gene expression of cell adhesion molecules, chemokines, and chemokine receptors in patients with inflammatory bowel disease before and after infliximab treatment. *Am J Gastroenterol* 2011;106:748–61.
43. Leal RF, Planell N, Kajekar R, et al. Identification of inflammatory mediators in patients with Crohn's disease unresponsive to anti-TNF α therapy. *Gut* 2015;64:233–42.
44. Uzzan M, Tokuyama M, Rosenstein AK, et al. Anti-alpha4beta7 therapy targets lymphoid aggregates in the gastrointestinal tract of hiv-1-infected individuals. *Sci Transl Med* 2018;10. doi: 10.1126/scitranslmed.aau4711.
45. Wyant T, Leach T, Sankoh S, et al. Vedolizumab affects antibody responses to immunisation selectively in the gastrointestinal tract: randomised controlled trial results. *Gut* 2015;64:77–83.
46. Rüegg C, Postigo AA, Sikorski EE, Butcher EC, Pytela R, Erle DJ. Role of integrin alpha 4 beta 7/alpha 4 beta P in lymphocyte adherence to fibronectin and VCAM-1 and in homotypic cell clustering. *J Cell Biol* 1992;117:179–89.
47. Miyamoto S, Katz BZ, Lafrenie RM, Yamada KM. Fibronectin and integrins in cell adhesion, signaling, and morphogenesis. *Ann N Y Acad Sci* 1998;857:119–29.
48. Lamb CA, Mansfield JC, Tew GW, et al. $\alpha E\beta 7$ integrin identifies subsets of pro-inflammatory colonic CD4+ T lymphocytes in ulcerative colitis. *J Crohns Colitis* 2017;11:610–20.
49. Ichikawa R, Lamb CA, Eastham-Anderson J, et al. AlphaE integrin expression is increased in the ileum relative to the colon and unaffected by inflammation. *J Crohns Colitis* 2018;12:1191–9.
50. Andrew DP, Rott LS, Kilshaw PJ, Butcher EC. Distribution of alpha 4 beta 7 and alpha E beta 7 integrins on thymocytes, intestinal epithelial lymphocytes and peripheral lymphocytes. *Eur J Immunol* 1996;26:897–905.
51. Shibahara T, Si-Tahar M, Shaw SK, Madara JL. Adhesion molecules expressed on homing lymphocytes in model intestinal epithelia. *Gastroenterology* 2000;118:289–98.
52. Sathaliyawala T, Kubota M, Yudanin N, et al. Distribution and compartmentalization of human circulating and tissue-resident memory T cell subsets. *Immunity* 2013;38:187–97.
53. Brew R, West DC, Burthem J, Christmas SE. Expression of the human mucosal lymphocyte antigen, HML-1, by T cells activated with mitogen or specific antigen in vitro. *Scand J Immunol* 1995;41:553–62.
54. Verstockt B, Verstockt S, Veny M, et al. Expression levels of 4 genes in colon tissue might be used to predict which patients will enter endoscopic remission after vedolizumab therapy for inflammatory bowel diseases. *Clin Gastroenterol Hepatol* 2020;18:1142–1151.e10.