

The role of gender in a large international OCD sample: A Report from the International College of Obsessive-Compulsive Spectrum Disorders (ICOCS) Network

Beatrice Benatti^{a,b,*}, Nicolaja Girone^a, Laura Celebre^a, Matteo Vismara^{a,b}, Eric Hollander^c, Naomi A. Fineberg^d, Dan J. Stein^e, Humberto Nicolini^{f,g}, Nuria Lanzagorta^g, Donatella Marazziti^h, Stefano Pallantiⁱ, Michael van Ameringen^j, Christine Lochner^k, Oguz Karamustafalioglu^l, Luchezar Hranov^m, Martin Figeeⁿ, Lynne M. Drummond^o, Jon E. Grant^p, Damiaan Denysⁿ, Leonardo F. Fontenelle^{q,v}, Jose M. Menchon^r, Joseph Zohar^s, Carolyn I. Rodriguez^{t,u}, Bernardo Dell'Osso^{a,b,t}

^a Luigi Sacco University Hospital, Psychiatry 2 Unit, University of Milan, Milan, Italy

^b CRC 'Aldo Ravelli' for Neuro-technology and Experimental Brain Therapeutics, University of Milan, Milan, Italy

^c Department of Psychiatry and Behavioral Sciences, Albert Einstein College of Medicine, Montefiore Medical Center, New York, NY, USA

^d University of Hertfordshire, School of Life and Medical Sciences and Hertfordshire Partnership University NHS Foundation Trust, Rosanne House, Welwyn Garden City, United Kingdom

^e SA MRC Unit on Risk & Resilience in Mental Disorders, Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

^f Genomics of Psychiatric and Neurodegenerative Diseases Laboratory, National Institute of Genomic Medicine (INMEGEN), Mexico City, Mexico

^g Carracci Medical Group, Mexico City, Mexico

^h Department of Clinical and Experimental Medicine, Section of Psychiatry, University of Pisa, Unicamillus-Saint Camillus International University of Health Sciences, Rome, Italy

ⁱ Department of Psychiatry, University of Florence, Institute of Neurosciences, Florence, Italy

^j Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, Ontario, Canada

^k SA MRC Unit on Risk & Resilience in Mental Disorders, Department of Psychiatry, Stellenbosch University, South Africa

^l Department of Psychiatry, Sisli Eftal Teaching and Research Hospital, Istanbul, Turkey

^m University Multiprofile Hospital for Active Treatment in Neurology and Psychiatry Sveti Naum, Sofia, Bulgaria

ⁿ Department of Psychiatry, Academic Medical Center, University of Amsterdam, Amsterdam, the Netherlands

^o National Services for OCD/BDD, St. George's, National Health Service Trust, London, United Kingdom

^p Department of Psychiatry and Behavioral Neuroscience, University of Chicago, IL, USA

^q Turner Institute for Brain and Mental Health, Monash University, VIC, Australia

^r Psychiatry Unit at the Hospital Universitat de Bellvitge-IDIBELL, University of Barcelona, Cibersam, Barcelona, Spain

^s Post-Trauma Center, Research Foundation by the Sheba Medical Center, Tel Aviv University, Sackler School of Medicine, Israel

^t Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA, USA

^u Veterans Affairs Palo Alto Health Care System, Palo Alto, CA, USA

^v Obsessive, Compulsive, and Anxiety Spectrum Research Program, Institute of Psychiatry of the Federal University of Rio de Janeiro (UFRJ), D'Oro Institute for Research and Education (IDOR), Rio de Janeiro, Brazil

ARTICLE INFO

Keywords:

OCD
Gender differences
Age at onset
Education

ABSTRACT

Introduction: Obsessive-compulsive disorder (OCD) is characterized by a range of phenotypic expressions. Gender may be a relevant factor in mediating the disorder's heterogeneity. The aim of the present report was to explore a large multisite clinical sample of OCD patients, hypothesizing existing demographic, geographical and clinical differences between male and female patients with OCD.

Methods: Socio-demographic and clinical variables of 491 adult OCD outpatients recruited in the International College of Obsessive-Compulsive Spectrum Disorders (ICOCS) network were investigated with a retrospective analysis on a previously gathered set of data from eleven countries worldwide. Patients were assessed through

* Corresponding author at: Luigi Sacco University Hospital, University of Milan, Via GB Grassi 74, 20154 Milan, Italy.

E-mail address: beatrice.benatti@unimi.it (B. Benatti).

<https://doi.org/10.1016/j.comppsy.2022.152315>

Received 10 July 2021; Received in revised form 8 March 2022; Accepted 9 April 2022

Available online 13 April 2022

0010-440X/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

structured clinical interviews, the Yale–Brown Obsessive-Compulsive Scale (Y-BOCS), the Montgomery-Asberg Depression Rating Scale (MADRS) and the Self-rating Depression Scale (SDS).

Results: Among females, adult onset (>18 years old) was significantly over-represented (67% vs. 33%, $p < 0.005$), and females showed a significantly older age at illness onset compared with males (20.85 ± 10.76 vs. 17.71 ± 8.96 years, $p < 0.005$). Females also had a significantly lower education level than males (13.09 ± 4.02 vs. 13.98 ± 3.85 years; $p < 0.05$), a significantly higher rate of being married (50.8% vs. 33.5%; $p < 0.001$) and a higher rate of living with a partner (47.5% vs. 37.6%; $p < 0.001$) than males. Nonetheless, no significant gender differences emerged in terms of the severity of OCD symptoms nor in the severity of comorbid depressive symptoms. No predictive effect of gender was found for Y-BOCS, MADRS and SDS severity.

Discussion/Conclusions: Our findings showed significant differences between genders in OCD. A sexually dimorphic pattern of genetic susceptibility may have a crucial role to OCD clinical heterogeneity, potentially requiring different specific therapeutic strategies. Further research is warranted to validate gender as an important determinant of the heterogeneity in OCD.

1. Introduction

Obsessive-compulsive disorder (OCD) is characterized by largely variable phenotypic expressions. Moreover, it often occurs with comorbid conditions, with the most common being anxiety, mood and other OC-spectrum disorders [1,2]. Previous findings suggested that gender may be a relevant factor in mediating this heterogeneity [3].

Despite more than 50% of subjects reporting the onset of the disease in late adolescence/early adulthood [4,5] when considering the role of gender, a younger age of onset with greater symptom severity was found in males [3,6,7]. In terms of clinical differences between male and female patients with OCD, a significantly different expression of obsessions and compulsions phenotype were found across genders [3,7]. For example, females were more likely to report obsessions associated with contamination belief or aggressive concerns, while males usually reported blasphemous thoughts [7]. An early study by Drummond [8] found that for patients hospitalized with profound, refractory OCD, being female predicted a better response. However, a subsequent paper from the same Centre failed to show any advantage for women [9]. Besides, gender seems to play a role in the onset of OCD comorbidities, although the available data are discordant. Males with OCD seem to be more vulnerable to social phobia, tic disorders, alcohol use disorders, compulsive Internet use and sexual disorders [10] whereas females with OCD seem more likely to present with eating disorders, specific phobias, trichotillomania, skin picking and compulsive buying [7]. On the other hand, some results showed an association between female gender and alcohol use disorders, anxiety and mood disorders [4]. In conclusion, conflicting data regarding OCD patterns and gender exist. As a complement to those findings and to evaluate the extent of gender differences for those with OCD at a global level, we conducted a post-hoc analysis of international multisite cross-sectional data from the ICOCS network. We aimed to examine the correlates of demographic, geographical and clinical differences between male and female patients with OCD.

2. Methods

The sample included 491 adult outpatients with a DSM-IV-TR OCD diagnosis [11] at different stages of treatment were recruited in the International College of Obsessive-Compulsive Spectrum Disorders (ICOCS) network between 2008 and 2010 [5,12–14]. The network included a multisite collaborative teams of OCD experts spanning academic and tertiary clinics from across the globe, including the Americas (Canada, the United States, and Mexico), Africa (Libya and South Africa), Europe (Spain, Italy, Turkey, Bulgaria, the United Kingdom) and the Middle East (Israel). Local institutional review boards approved the protocol. Patients provided written informed consent to use their anonymized data.

The sociodemographic measures included: age, gender, years of education, marital status, professional status, age of OCD onset, psychopharmacological therapy, presence of monotherapy vs augmentation.

Comorbidity of associated mood, tic and anxiety disorders (e.g., major depression, generalized anxiety disorder) was assessed using a Structured Clinical Interview based on the DSM-IV criteria (SCID-I and -II) [15,16]. Medical comorbidities were elicited by clinician interview.

OCD illness severity was assessed by the Yale-Brown Obsessive Compulsive Scale (Y-BOCS); severity of OCD was also assessed by means of a qualitative analysis on Y-BOCS scores, differentiating “mild” OCD (range 9–15), “moderate” OCD (range 16–23), “severe” OCD (range 24–32) and “extreme” OCD (range 33–40) [17,18]. Severity of comorbid depressive symptoms was assessed using the Montgomery-Asberg Depression Rating Scale (MADRS) [19]. Interference with work/school, social life/leisure activities, and family life/home responsibilities was measured with the Sheehan Disability Scale (SDS) [20].

Statistical analyses were performed with Pearson's chi-squared test for categorical variables and ANOVA test for the continuous variables to explore gender differences. A linear regression was used to test whether Y-BOCS, MADRS and SDS severity could be predicted by gender. All analyses were performed using Statistical Package for the Social Sciences (SPSS) 25.0 software for Windows (SPSS Inc., Chicago, IL, USA). Statistical significance was set at $p < 0.05$.

3. Results

The sample included 60.5% females, mean age 38.56 ± 12.86 years. Patients were geographically distributed as follows: 217 (44.2%) from Europe (22% from Italy, 8.1% from Spain, 6.7% from United Kingdom, 4.9% from Turkey, 2.4% from Bulgaria); 115 (23.4%) from the Americas (14.1% from Canada, 7.7% from Mexico and 1.6% from USA); 100 (20.4%) from Africa (20.2% from South Africa and 0.2% from Libya) and 59 (12%) from Middle East (Israel).

For the purpose of the study the whole sample was divided into two subgroups based on sex. Comparison of socio-demographic and clinical variables between female and male samples are provided in [Tables 1 and 2](#).

With respect to geographical differences among clinics involved in the ICOCS network, a significant difference by gender distribution emerged (see [Table 3](#)). Specifically, Americas showed significantly higher rates of female OCD patients compared to male OCDs (30.3% vs. 12.9%; $p < 0.001$, [Fig. 1](#)). However, no significant differences in gender rates emerged for Europe, Africa and the Middle East.

In the female group adult onset (>18 years old) was found to be significantly more represented (67% vs. 33%, $p < 0.005$; [Fig. 2](#)), showing a significantly older age at onset when compared with male patients (20.85 ± 10.76 vs. 17.71 ± 8.96 years, $p < 0.005$). No significant difference emerged neither in the severity of symptoms assessed through the Y-BOCS (22.32 ± 7.36 vs. 21.9 ± 7.26) nor in the severity of comorbid depressive symptoms, assessed through the MADRS and the SDS ([Table 2](#)). Furthermore, no predictive effect of gender was found for Y-BOCS, MADRS and SDS severity with linear regression ($b = 0.416$, $t = 0.577$ $p = 0.564$; $b = -1.220$, $t = -1.300$ $p = 0.194$; $b = 0.300$, $t = 0.363$).

Table 1

Comparison of socio-demographic variables between female and male samples.

	Females	Males	Total sample
Gender	297 (60.5%)	194 (39.5%)	491
Age	40.28 ± 13.39**	36.18 ± 11.72	38.56 ± 12.86
Years of education	13.09 ± 4.02*	13.98 ± 3.85	13.46 ± 3.97
Marital status			
Single	103 (40.2%)	110 (61.5%)	213 (49%)
Married	130 (50.8%)*	60 (33.5%)	190 (43.7%)
Divorced	18 (7%)	7 (3.9%)	25 (5.7%)
Widow/er	5 (2%)	1 (0.6%)	6 (1.4%)
Professional status			
Worker	130 (50.4%)	101 (54%)	231 (51.9%)
Unemployed	73 (28.3%)	38 (20.3%)	111 (24.9%)
Student	37 (14.3%)	35 (18.7%)	72 (16.2%)
Retired	18 (7%)	13 (7%)	31 (7%)
Living			
Alone	40 (13.5%)	40 (20.6%)	80 (16.3%)
With spouse	141 (47.5%)*	63 (37.6%)	204 (41.5%)
With parents	63 (21.2%)	73 (37.6%)	136 (27.7%)
Nursing home	53 (17.8%)	18 (9.3%)	71 (14.5%)

Notes: Values for categorical and continuous variables are expressed in percentages and mean ± SD, respectively. Reported variables had a percentage of missing data ranging from 2% to 12%. Boldface indicates parameters with statistically significant differences between the two subgroups; ** $p < 0.005$ * $p < 0.05$.

Table 2

Comparison of clinical variables between female and male samples.

	Females	Males	Total sample
Age at OCD onset (years)	20.85 ± 10.76**	17.71 ± 8.96	19.53 ± 10.15
Late onset (>18 years)	126 (67%)*	62 (33%)	188 (45.7%)
Medical Comorbidity	50 (54.9%)	41 (45.1%)	91 (20.8%)
Psychiatric Comorbidity	140 (57.6%)	103 (42.4%)	243 (55.9%)
Polycomorbidities	38 (56.7%)	29 (43.3%)	67 (14.9%)
Tic disorder	36 (49.3%)	37(50.7%)	73 (17.6%)
Psychopharmacological therapy	182 (58.7%)	128 (41.3%)	310 (87.8%)
Monotherapy vs augmentation	97 (60.6%) vs 85 (56.7%)	63 (39.4%) vs 65 (43.3%)	160 (51.6%) vs 150 (48.4%)
Psychiatric hospitalization	46 (54.1%)	39 (45.9%)	85 (18.8%)
CBT	137 (60.1%)	91 (39.9%)	228 (46.4%)
Y-BOCS total scores	22.32 ± 7.36	21.9 ± 7.26	22.16 ± 7.32
Mild (9–15)	36 (53.7%)	31 (46.3%)	67 (13.6%)
Moderate (16–23)	110 (65.1%)	59 (34.9%)	169 (34.4%)
Severe (24–32)	88 (58.7%)	62 (41.3%)	150 (30.5%)
Extreme (33–40)	28 (68.3%)	13 (31.7%)	41 (8.4%)
MADRS score	13.13 ± 8.56	14.35 ± 8.37	13.57 ± 8.50
SDS score	16.98 ± 7.79	16.68 ± 7.96	16.87 ± 7.85

Notes: Values for categorical and continuous variables are expressed in percentages and mean ± SD, respectively. OCD: Obsessive-Compulsive Disorder; Polycomorbidities: co-occurrence of at least two comorbid psychiatric conditions other than OCD; CBT: Cognitive behavioral therapy; Y-BOCS: Yale-Brown Obsessive Compulsive Scale; MADRS: Montgomery-Asberg Depression Rating Scale; SDS: Self-rating Depression Scale.

Reported variables had a percentage of missing data ranging from 2% to 12%. Boldface indicates parameters with statistically significant differences between the two subgroups; ** $p < 0.005$ * $p < 0.05$.

$p = 0.717$, respectively; see Table 4). Lastly, no significant differences in terms of psychiatric comorbidity and presence of poly-comorbidities, pharmacological or psychological treatment and presence of tic disorder were found. Regarding years of education, the female group showed a significantly lower duration of education compared to male subgroup (13.09 ± 4.02 vs. 13.98 ± 3.85 years; $p < 0.05$). Moreover, in the female group a significantly higher rate of being married (50.8% vs- 33.5%; $p < 0.001$) and a higher presence of living with spouse (47.5% vs. 37.6%; $p < 0.001$) emerged compared to males. No significant difference emerged regarding professional status between groups.

4. Discussion

In the present report, clinical correlates of gender differences in OCD disorder were examined. To the authors' knowledge, this is the first study analyzing gender differences in an international multisite OCD adult outpatient population.

We found that compared to males, females with OCD 1) had a lower level of education, 2) were more commonly married or living with partners, and 3) had a later age of onset of OCD. And finally, we found a higher female:male ratio in the Americas compared to other study sites (Europe, Africa and Israel).

Consistently with previous findings [3,5,21], OCD female patients showed a significantly higher age of psychopathological onset when compared to the males. In particular, in the female group the adult onset (>18 years old) was found to be significantly more represented. Previous studies showed a younger age at OCD onset with greater symptom severity in males compared to females [3,6,7]. Nonetheless, in our study no significant differences in the severity of OCD symptoms assessed by the Yale-Brown obsessive compulsive scale was found. In addition, no difference in the severity of comorbid depressive symptoms and no predictive effect of gender for Y-BOCS, MADRS and SDS severity were found.

No significant differences in terms of psychiatric comorbidities, including tic disorders, and in comorbid depression severity emerged in the present sample. Previously other large multicentric studies did not show differences in terms of mood disorders and any anxiety disorder rates between genders [22]. Moreover, when comparing a sample of OCD patients in terms of gender differences, also Lochner and colleagues did not find significantly different tic disorders prevalence between males and females groups [23]. Previous reports from the ICOCs group highlighted lower general psychiatric comorbidity rates in the sample compared to other studies, potentially explaining the lack of significant associations [24]. Lastly no significant gender differences occurred in terms of pharmacological or psychological treatment, as also previously highlighted by the Spanish study of Labad and colleagues in 2008 [25]. These specific features need to be further investigated.

Consistently with several other authors [26–28], our results showed that women with OCD were more commonly married than men. In addition, we found a higher frequency of female patients living with partners compared to males, as previously highlighted by Sobin and colleagues [29]. One possible reason could be related to the earlier age at OCD onset among men, preventing stable relationships in those patients.

Regarding years of education, the female group showed a significantly lower duration of education compared to male subgroup. In this perspective, the current literature presented mixed results. Vander Stoep and colleagues attempted to quantify the societal burden of mental disorders in decreased educational attainment, estimating that as many as 46% of high school dropouts might be attributable to the negative effects of prior mental disorders [30]. Another study using data from the Australian National Survey of Mental Health and Wellbeing showed, contrary to our results, that male children with early-onset OCD had higher school dropout rates compared to OCD females and controls [31]. A possible explanation in the role of OCD gender differences and education levels may lie on age at OCD onset and DUI.

Lastly, our multisite results showed a significantly higher prevalence of females with OCD in Americas compared to males and no significantly gender differences in Europe, Africa and Israel.

When considering the role of gender, several studies with children OCD samples have identified a greater proportion of males [32–34], whereas some adult OCD samples find both equal distributions [35,36] or a greater proportion of females [37,38] across centres.


The above-mentioned results should be interpreted considering some methodological limitations. First, the cross-sectional nature of the study allowed only one time assessment. Secondly, some variables were obtained retrospectively, being susceptible to recall bias.

Table 3


Gender distribution of OCD sample recruited in the ICOCS network across the globe.

	Females		Males		Total sample
Geographical area	Europe 118 (39.7%)		Europe 99 (51%)		Europe 217(44.2%)
	Italy 59 (19.9%)		Italy 49 (25.3%)		Italy 108 (22%)
	Spain 19 (6.4%)		Spain 21 (10.8%)		Spain 40 (8.1%)
	Turkey 19 (6.4%)		Turkey 5 (2.6%)		Turkey 24 (4.9%)
	UK 13(4.4%)		UK 20 (10.3%)		UK 33 (6.7%)
	Bulgaria 8 (2,7%)		Bulgaria 4 (2,1%)		Bulgaria 12 (2.4%)
	Americas 90 (30.3%)**		Americas 25 (12.9%)**		Americas 115 (23.4%)
	Mexico 32 (10.8%)		Mexico 6 (3.1%)		Mexico 38 (7.7%)
	Canada 54 (18.2%)		Canada 15 (7.7%)		Canada 69 (14.1%)
	USA 4 3(1.3%)		USA 4 (2.1%)		USA 8 (1.6%)
	Africa 59 (19.9%)		Africa 41 (21.1%)		Africa 100 (20.4%)
	South Africa 58 (19,5%)		South Africa 41(21.1%)		South Africa 99(20.2%)
	Libya 1 (0.3%)		Libya 0 (0%)		Libya 1(0.2%)
	Middle East		Middle East		Middle East
	Israel 30 (10.1%)		Israel 29 (14.9%)		Israel 59 (12%)

Notes: Values for categorical variables are expressed in percentages. Boldface indicates parameters with statistically significant differences between the two subgroups; **p < 0.001 *p < 0.05.

**Fig. 1.** Gender distribution of OCD sample recruited in the ICOCS network across the globe.

Notes: Boldface indicates parameters with statistically significant differences between the two subgroups; **p < 0.001.

**Fig. 2.** Comparison of psychopathological onset between female and male OCD patients.

Notes: Boldface indicates parameters with statistically significant differences between the two subgroups; **p < 0.005.

Nevertheless, our findings suggest that gender is a fundamental contributor to OCD clinical heterogeneity potentially necessitating the establishment of specific therapeutic strategies. Future investigation may, therefore, benefit from a focus on gender diversified severity variables, such as suicidality, stressful life events or history of trauma and from a wider sample size and longer follow-up.

Table 4

Linear regression analysis of predictive effect of gender for Y-BOCS, MADRS and SDS severity.

Subscales	Predictor	Unstandardized Coefficients		Standardized Coefficients	t	p
		b	SE	β		
Y-BOCS	Constant	21.489	1.216		17.672	
	Gender	0.416	0.721	0.028	0.577	0.564
MADRS	Constant	15.572	1.604		9.707	
	Gender	-1.220	0.939	-0.060	-1.300	0.194
SDS	Constant	16.387	1.395		11.750	
	Gender	0.300	0.827	0.019	0.363	0.717

Notes: b: Regression coefficient; SE: standard error; Y-BOCS: Yale-Brown Obsessive Compulsive Scale; MADRS: Montgomery-Asberg Depression Rating Scale; SDS: Self-rating Depression Scale.

Role of funding source

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of Competing Interest

- BB has received lecture honoraria from Lundbeck and Janssen.

- BDO has received lecture honoraria from Angelini, Janssen, Lundbeck, Livanova, Arcapharma, and Neuraxpharm.
- CR in the last 3 years, has served as a consultant for Epiodyne and received research grant support from Biohaven Pharmaceuticals and a stipend from APA Publishing for her role as Deputy Editor at The American Journal of Psychiatry.
- MVA has been on the advisory board of Allergan, Almatica, Brainsway, Lundbeck, Myriad Neuroscience, Otsuka, Purdue Pharma. He has received research support from Janssen-Ortho Inc., Purdue Pharma (Canada), the Canadian Foundation for Innovation, and Hamilton Academic Health Sciences Organization (HAHSO).
- NF declares that in the past 3 years she has held research or networking grants from the ECNP, UK NIHR, EU H2020 (COST Action), University of Hertfordshire; she has accepted travel and/or hospitality expenses from the BAP, ECNP, RCPsych, CINP, International Forum of Mood and Anxiety Disorders, World Psychiatric Association, Indian Association for Biological Psychiatry, Sun; she has received payment from Taylor and Francis and Elsevier for editorial duties. In the past 3 years, she has accepted a paid speaking engagement in a webinar sponsored by Abbott. Previously, she has accepted paid speaking engagements in various industry-supported symposia and has recruited patients for various industry-sponsored studies in the field of OCD treatment. She leads an NHS treatment service for OCD. She holds Board membership for various registered charities linked to OCD. She gives expert advice on psychopharmacology to the UK MHRA.

Acknowledgements

This article is based upon work from COST Action CA16207 “European Network for Problematic Usage of the Internet”, supported by COST (European Cooperation in Science and Technology). www.cost.eu

References

- [1] Hofmeijer-Sevink MK, van Oppen P, van Megen HJ, et al. Clinical relevance of comorbidity in obsessive compulsive disorder: the Netherlands OCD Association study. *J Affect Disord* 2013;150:847–54.
- [2] Lochner C, Fineberg NA, Zohar J, et al. Comorbidity in obsessive-compulsive disorder (OCD): A report from the International College of Obsessive-Compulsive Spectrum Disorders (ICOCs). *Compr Psychiatry* 2014;55:1513–9.
- [3] Benatti B, Celebre L, Gironi N, et al. Clinical characteristics and comorbidity associated with female gender in obsessive-compulsive disorder. *J Psychiatr Res* 2020;131:209–14.
- [4] Fineberg NA, Hengartner MP, Bergbaum, et al. A prospective population-based cohort study of the prevalence, incidence and impact of obsessive-compulsive symptomatology. *Int J Psychiatry Clin Pract* 2013;2013(17):170–8.
- [5] Dell’Osso B, Benatti B, Hollander E, et al. Childhood, adolescent and adult age at onset and related clinical correlates in obsessive-compulsive disorder: A report from the International College of Obsessive-Compulsive Spectrum Disorders (ICOCs). *Int J Psychiatry Clin Pract* 2016;20:210–7.
- [6] Stewart SE, Geller DA, Jenike M, et al. Long-term outcome of pediatric obsessive-compulsive disorder: a meta-analysis and qualitative review of the literature. *Acta Psychiatr Scand* 2004;110:4–13.
- [7] Torresan RC, Ramos-Cerqueira ATA, Shavitt RG. Symptom dimensions, clinical course and comorbidity in men and women with obsessive-compulsive disorder. *Psychiatry Res* 2013;209:186–95.
- [8] Drummond LM. The treatment of severe, chronic, resistant obsessive-compulsive disorder: an evaluation of an inpatient programme using behavioural psychotherapy in combination with other treatments. *Br J Psychiatry* 1993;163:223–9.
- [9] Boschén MJ, Drummond LM, Pillaya A, Morton K. Predicting outcome of treatment for severe, treatment resistant OCD in inpatient and community settings. *J Behav Ther Exp Psychiatry* 2010;41:90–5.
- [10] Bogetto F, Venturello S, Albert U, Maina G, Ravizza L. Gender-related clinical differences in obsessive-compulsive disorder. *Eur Psychiatry* 1999;14:434–41.
- [11] American Psychiatric Association. Diagnostic and statistical manual of mental disorders. In: Text revision. 4th ed. Washington, DC: American Psychiatric Association; 2000.
- [12] Dell’Osso B, Benatti B, Buoli M, et al. The influence of age at onset and duration of illness on long-term outcome in patients with obsessive-compulsive disorder: a report from the International College of Obsessive Compulsive Spectrum Disorders (ICOCs). *Eur Neuropsychopharmacol* 2013;23:865–71.
- [13] Lochner C, Fineberg NA, Zohar J, et al. Comorbidity in obsessive-compulsive disorder (OCD): a report from the international college of obsessive-compulsive spectrum disorders (ICOCs). *Compr Psychiatry* 2014;55:1513–9.
- [14] Van Ameringen M, Simpson W, Patterson B, et al. Pharmacological treatment strategies in obsessive compulsive disorder: a cross-sectional view in nine international OCD centers. *J Psychopharmacol* 2014;28:596–602.
- [15] First MB, Gibbon M, Spitzer R, Williams J, Benjamin L. Structured Clinical Interview for DSM-IV Axis II Personality Disorders. SCID-II. 1997 [Washington DC].
- [16] First MB, Spitzer RL, Gibbon M, Williams JB. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition. SCID-I/P. 2002 [New York].
- [17] Goodman WK, Price LH, Rasmussen SA, et al. The Yale-Brown Obsessive Compulsive Scale. I development, use, and reliability. *Arch Gen Psychiatry* 1989;46:1006–11.
- [18] Rapp AM, Bergman RL, Piacentini J, McGuire JF. Evidence-based assessment of obsessive-compulsive disorder. *J Cent Nerv Syst Dis* 2016;8:13–29.
- [19] Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry* 1979;134:382–9.
- [20] Sheehan DV. The Anxiety Disease. New York: Charles Scribners Sons; 1983.
- [21] Fontenelle LF, Mendlowicz MV, Marques C, Versiani M. Early- and late- onset obsessive-compulsive disorder in adult patients: an exploratory clinical and therapeutic study. *J Psychiatr Res* 2003;37:127–33.
- [22] Tripathi A, Avasthi A, Grover, et al. Gender differences in obsessive-compulsive disorder: findings from a multicentric study from northern India. *Asian J Psychiatr* 2018;37:3–9.
- [23] Lochner C, Hemmings SM, Kinnear CJ, Moolman-Smook JC, Corfield VA, Knowles JA, et al. Gender in obsessive-compulsive disorder: clinical and genetic findings. *Eur Neuropsychopharmacol* 2004;14(5):437–45.
- [24] Dell’Osso B, Nicolini H, Lanzagorta N, et al. Cigarette smoking in patients with obsessive compulsive disorder: a report from the International College of Obsessive Compulsive Spectrum Disorders (ICOCs). *CNS Spectr* 2014;20:469–73.
- [25] Labad J, Menchon JM, Alonso P, et al. Gender differences in obsessive-compulsive symptom dimensions. *Depress Anxiety* 2008;25:832–8.
- [26] Castle DJ, Deale A, Marks IM. Gender differences in obsessive compulsive disorder. *Aust N Z J Psychiatry* 1995;29:114–7.
- [27] Tükel R, Polat A, Genç A, Bozkurt O, Atlı H. Gender-related differences among Turkish patients with obsessive-compulsive disorder. *Compr Psychiatry* 2004;45:362–6.
- [28] Jaisorya T, Janardhan R, Srinath S, Thennarasu K. Sex differences in Indian patients with obsessive-compulsive disorder. *Compr Psychiatry* 2009;50:70–5.
- [29] Sobin C, Blundell M, Weiller F, Gavigan C, Haiman C, Karayiorgou M. Phenotypic characteristics of obsessive-compulsive disorder ascertained in adulthood. *J Psychiatr Res* 1999;33:265–73.
- [30] Vander Stoep A, Weiss NS, Kuo ES, Cheney D, Cohen P. What proportion of failure to complete secondary school in the US population is attributable to adolescent psychiatric disorder? *J Behav Health Serv Res* 2003;30:119–24.
- [31] Leach LS, Butterworth P. The effect of early onset common mental disorders on educational attainment in Australia. *Psychiatry Res* 2012;199:51–7.
- [32] Geller DA. Obsessive-compulsive and spectrum disorders in children and adolescents. *Psychiatr Clin* 2006;29:353–70.
- [33] Mancebo MC, Garcia AM, Pinto A, et al. Juvenile-onset OCD: clinical features in children, adolescents and adults. *Acta Psychiatr Scand* 2008;118:149–59.
- [34] Mathis MA, Alvarenga PD, Funaro G, et al. Gender differences in obsessive-compulsive disorder: a literature review. *Braz J Psychiatry* 2011;33:390–9.
- [35] Fullana MA, Mataix-Cols D, Caspi A, et al. Obsessions and compulsions in the community: prevalence, interference, help-seeking, developmental stability, and co-occurring psychiatric conditions. *Am J Psychiatry* 2009;166:329–36.
- [36] Canals J, Hernández-Martínez C, Cosi S, Voltas N. The epidemiology of obsessive-compulsive disorder in Spanish school children. *J Anxiety Disord* 2012;26:746–52.
- [37] Rasmussen SA, Eisen JL. Epidemiology of obsessive compulsive disorder. *J Clin Psychiatry* 1990;51:10–3.
- [38] Fineberg NA, Hengartner MP, Bergbaum CE, et al. A prospective population-based cohort study of the prevalence, incidence and impact of obsessive-compulsive symptomatology. *Int J Psychiatry Clin Pract* 2013;17:170–8.