

Breakdown of heteroclinic connections in the analytic Hopf-Zero singularity: Rigorous computation of the Stokes constant

Inmaculada Baldomá^{1,2,3}, Maciej J. Capiński⁴, Marcel Guardia^{1,2,3}, Tere M. Seara^{1,2,3}

¹ Departament de Matemàtiques, Universitat Politècnica de Catalunya (UPC), Barcelona, Spain.

² IMTECH, Universitat Politècnica de Catalunya (UPC), Barcelona, Spain.

³ Centre de Recerca Matemàtica, Barcelona, Spain.

⁴AGH University of Science and Technology, Faculty of Applied Mathematics
al. Mickiewicza 30, 30-059 Kraków, Poland

June 29, 2022

Abstract

Consider analytic generic unfoldings of the three dimensional conservative Hopf-Zero singularity. Under open conditions on the parameters determining the singularity, the unfolding possesses two saddle-foci when the unfolding parameter is small enough. One of them has one dimensional stable manifold and two dimensional unstable manifold whereas the other one has one dimensional unstable manifold and two dimensional stable manifold. Baldomá, Castejón and Seara [BCS13] gave an asymptotic formula for the distance between the one dimensional invariant manifolds in a suitable transverse section. This distance is exponentially small with respect to the perturbative parameter, and it depends on what is usually called a Stokes constant. The non-vanishing of this constant implies that the distance between the invariant manifolds at the section is not zero. However, up to now there do not exist analytic techniques to check that condition. In this paper we provide a method for obtaining accurate rigorous computer assisted bounds for the Stokes constant. We apply it to two concrete unfoldings of the Hopf-Zero singularity, obtaining a computer assisted proof that the constant is non-zero

Contents

| | | |
|----------|---|----------|
| 1 | Introduction | 2 |
| 1.1 | Hopf-zero singularity and its unfoldings | 4 |
| 1.2 | The inner equation | 6 |
| 2 | Rigorous computation of the Stokes constant | 8 |
| 2.1 | Scheme of the method. Theoretical approach | 9 |
| 2.1.1 | Existence domain of the solutions of the inner equation | 9 |
| 2.1.2 | Rigorous computation of the Stokes constant | 11 |
| 2.2 | Computing the Stokes constant: method | 13 |

| | | |
|----------|---|-----------|
| 2.2.1 | The linear operator \mathcal{B}^- | 13 |
| 2.2.2 | Explicit constants for the inner equation | 14 |
| 2.2.3 | Bounds for the norm of the first iteration | 15 |
| 2.2.4 | The Lipschitz constant of \mathcal{S} | 16 |
| 2.2.5 | The Lipschitz constant of \mathcal{F}^- | 20 |
| 2.2.6 | Setting ρ_* for the existence result | 21 |
| 2.2.7 | Explicit bounds for the norm of the linear operator \mathcal{G} . . | 21 |
| 2.2.8 | Computation of the Stokes constant | 22 |
| 2.3 | Computing the Stokes constant: algorithm | 23 |
| 3 | Examples | 24 |
| 3.1 | The first example | 24 |
| 3.2 | The second example | 25 |
| 4 | Improving the computation of the Stokes constant | 27 |
| 4.1 | Proof of Lemma 4.1 | 29 |
| A | Proof of Lemma 2.1 | 34 |
| B | Computing the bound on $\Delta\psi(-i\rho^*)$ | 35 |

1 Introduction

One of the fundamental questions in dynamical systems is to assess whether a given model possesses chaotic dynamics or not. In particular, one would like to prove whether the model has a hyperbolic invariant set whose dynamics is conjugated to the symbolic dynamics of the usual Bernoulli shift by means of the construction of a Smale horseshoe. Since the pioneering works by Smale and Shilnikov, it is well known that the construction of such invariant sets may be attained by analyzing the stable and unstable invariant manifolds of hyperbolic invariant objects (critical points, periodic orbits, invariant tori) and their intersections.

Such analysis can be done by classical perturbative techniques such as (suitable versions of) Melnikov Theory [Mel63] or by means of Computer Assisted Proofs [CZ17, CZ18]. However, there are settings where Melnikov Theory nor “direct” Computer Assisted Proofs (that is, rigorous computation of the invariant manifolds) cannot be applied. For instance, in the so-called exponentially small splitting of separatrices setting. That is, on models which depend on a small parameter and where the distance between the stable and unstable invariant manifolds is exponentially small with respect to this parameter.

This phenomenon of exponentially small splitting of separatrices often appears in analytic systems with different time scales, which couple fast rotation with slow hyperbolic motion. Example of such settings are nearly integrable Hamiltonian systems at resonances, near the identity area preserving maps or local bifurcations in Hamiltonian, reversible or volume preserving settings. In such settings, one needs more sophisticated techniques rather than Melnikov Theory to analyze the distance between the stable and unstable invariant manifolds. Most of the results in the area follow the seminal approach proposed by Lazutkin in [Laz03] (there are though other approaches such as [Tre97]). Using

these techniques, one can provide an asymptotic formula for the distance between the invariant manifolds, with respect to the perturbation parameter. If we denote by ε the small parameter, the distance is usually of the form

$$d = d(\varepsilon) \sim \Theta \varepsilon^\alpha e^{\frac{a}{\varepsilon^\beta}} \quad \text{as} \quad \varepsilon \rightarrow 0$$

for some constants Θ , α , a and β . In most of the settings, the constants α , a and β have explicit formulas and can be “easily” computed for given models. However, the constant Θ is of radically different nature and much harder to compute. Indeed, the constants α , a and β depend on certain first order terms of the model whereas Θ , which we refer to as the Stokes constant, depends in a non-trivial way on the “whole jet” of the considered model. Note that it is crucial to know whether Θ vanishes or not, since its vanishing makes the whole first order between the invariant manifolds vanish and, consequently, chaos can not be guaranteed in the system.

The purpose of this paper is to provide (computer assisted) methods to check, in given models, that the Stokes constant does not vanish. Moreover, our method provides a rigorous accurate computation of this constant. To show the main ideas of the method and avoid technicalities, we focus on the simplest setting where this method can be implemented: the breakdown of a one-dimensional heteroclinic connection for generic analytic unfoldings of the volume preserving Hopf-zero singularity.

This problem was analyzed in [BS08, BCS13]. In these papers and the companions [BCS18a, BCS18b, BIS20], the authors prove that, in generic unfoldings of an open set of Hopf-zero singularities, one can encounter Shilnikov chaos [Sn70]. The fundamental difficulty in these models is to prove that the one dimensional and two-dimensional heteroclinic manifolds connecting two saddle-foci in a suitable truncated normal form of the unfolding, break down when one considers the whole vector field. These breakdowns, which are exponentially small, plus some additional generic conditions lead to existence of chaotic motions.

Remark 1.1. *A bifurcation with very similar behavior to that of the conservative Hopf-zero singularity is the Hamiltonian Hopf-zero singularity where a critical point of a 2 degree of freedom Hamiltonian system has a pair of elliptic eigenvalues and a pair of 0 eigenvalues forming a Jordan block (see for instance [GL14]). In generic unfoldings, the 0 eigenvalues become a pair of small real eigenvalues and therefore the critical point becomes a saddle-center. In this setting, one can analyze the one dimensional invariant manifolds of the critical point and obtain an asymptotic formula for their distance (in a suitable section). This distance is exponentially with respect to the perturbative parameter. Then, to prove that they indeed do not intersect, one has to show that a certain Stokes constant is not zero as in the Hopf-zero conservative singularity. The methods presented in this paper can be adapted to this other setting. The Hamiltonian Hopf-zero singularity appears in many physical models, for instance in the Restricted Planar 3 Body Problem (see [BGG21a, BGG21b]). It also plays an important role in the breakdown of small amplitude breathers for the Klein-Gordon equation (albeit in an infinite dimensional setting), see [SK87, GGSZ21]. We plan to provide a computer assisted proof of the Stokes constant to guarantee the non-existence of small breathers in given Klein-Gordon equations in a future work.*

In this paper, we provide a method to compute the Stokes constant associated to the breakdown of the one-dimensional heteroclinic connection in analytic unfoldings of the conservative Hopf-zero singularity.

Let us first explain the Hopf-zero singularity and state the main results about the breakdown of its one-dimensional heteroclinic connection obtained in [BS08, BCS13].

1.1 Hopf-zero singularity and its unfoldings

The Hopf-zero singularity takes place on a vector field $X^* : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, which has the origin as a critical point, and such that the eigenvalues of the linear part at this point are $0, \pm i\alpha^*$, for some $\alpha^* \neq 0$. Hence, after a linear change of variables, we can assume that the linear part of this vector field at the origin is

$$DX^*(0, 0, 0) = \begin{pmatrix} 0 & \alpha^* & 0 \\ -\alpha^* & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

We assume that X^* is analytic. Since $DX^*(0, 0, 0)$ has zero trace, it is reasonable to study it in the context of analytic conservative vector fields (see [BV84] for the analysis of this singularity in the \mathcal{C}^∞ class). In this case, the generic singularity can be met by a generic linear family depending on one parameter, and so it has codimension one.

We study generic analytic families X_μ of conservative vector fields on \mathbb{R}^3 depending on a parameter $\mu \in \mathbb{R}$, such that $X_0 = X^*$, the vector field described above.

Following [Guc81] and [GH90], after some changes of variables, we can write X_μ in its normal form up to order two, namely

$$\begin{aligned} \frac{d\bar{x}}{dt} &= -\beta_1 \bar{x} \bar{z} + \bar{y} (\alpha^* + \alpha_2 \mu + \alpha_3 \bar{z}) + \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu), \\ \frac{d\bar{y}}{dt} &= -\bar{x} (\alpha^* + \alpha_2 \mu + \alpha_3 \bar{z}) - \beta_1 \bar{y} \bar{z} + \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu), \\ \frac{d\bar{z}}{dt} &= -\gamma_0 \mu + \beta_1 \bar{z}^2 + \gamma_2 (\bar{x}^2 + \bar{y}^2) + \gamma_3 \mu^2 + \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu). \end{aligned} \tag{1}$$

Note that the coefficients β_1 , γ_2 and α_3 depend exclusively on the vector field X^* .

From now on, we will assume that X^* and its unfolding X_μ satisfy the following generic conditions:

$$\beta_1 \neq 0, \quad \gamma_0 \neq 0. \tag{2}$$

Depending on the other coefficients α_i and γ_i , one obtains different qualitative behaviors for the orbits of the vector field X_μ . We consider μ satisfying

$$\beta_1 \gamma_0 \mu > 0. \tag{3}$$

In fact, redefining the parameters μ and the variable \bar{z} , one can achieve

$$\beta_1 > 0, \quad \gamma_0 = 1, \tag{4}$$

and consequently the open set defined by (3) is now

$$\mu > 0. \quad (5)$$

Moreover, dividing the variables \bar{x}, \bar{y} and \bar{z} by $\sqrt{\beta_1}$, and scaling time by $\sqrt{\beta_1}$, redefining the coefficients and denoting $\alpha_0 = \alpha^*/\sqrt{\beta_1}$, we can assume that $\beta_1 = 1$, and therefore system (1) becomes

$$\begin{aligned} \frac{d\bar{x}}{dt} &= -\bar{x}\bar{z} + \bar{y}(\alpha_0 + \alpha_2\mu + \alpha_3\bar{z}) + \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu), \\ \frac{d\bar{y}}{dt} &= -\bar{x}(\alpha_0 + \alpha_2\mu + \alpha_3\bar{z}) - \bar{y}\bar{z} + \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu), \\ \frac{d\bar{z}}{dt} &= -\mu + \bar{z}^2 + \gamma_2(\bar{x}^2 + \bar{y}^2) + \gamma_3\mu^2 + \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu). \end{aligned} \quad (6)$$

We denote by X_μ^2 , usually called the normal form of second order, the vector field obtained considering the terms of (6) up to order two. Therefore, one has

$$X_\mu = X_\mu^2 + F_\mu^2, \quad \text{where } F_\mu^2(\bar{x}, \bar{y}, \bar{z}) = \mathcal{O}_3(\bar{x}, \bar{y}, \bar{z}, \mu).$$

It can be easily seen that system (6) has two critical points at distance $\mathcal{O}(\sqrt{\mu})$ to the origin. Therefore, we scale the variables and parameters so that the critical points are $\mathcal{O}(1)$ and not $\mathcal{O}(\sqrt{\mu})$. That is, we define the new parameter $\delta = \sqrt{\mu}$, and the new variables $x = \delta^{-1}\bar{x}$, $y = \delta^{-1}\bar{y}$, $z = \delta^{-1}\bar{z}$ and $t = \delta\bar{t}$. Then, renaming the coefficients $b = \gamma_2$, $c = \alpha_3$, system (6) becomes

$$\begin{aligned} \frac{dx}{dt} &= -xz + \left(\frac{\alpha(\delta^2)}{\delta} + cz \right) y + \delta^{-2}f(\delta x, \delta y, \delta z, \delta), \\ \frac{dy}{dt} &= - \left(\frac{\alpha(\delta^2)}{\delta} + cz \right) x - yz + \delta^{-2}g(\delta x, \delta y, \delta z, \delta), \\ \frac{dz}{dt} &= -1 + b(x^2 + y^2) + z^2 + \delta^{-2}h(\delta x, \delta y, \delta z, \delta), \end{aligned} \quad (7)$$

where f , g and h are real analytic functions of order three in all their variables, $\delta > 0$ is a small parameter and $\alpha(\delta^2) = \alpha_0 + \alpha_2\delta^2$.

Remark 1.2. *Without loss of generality, we can assume that α_0 and c are both positive constants. In particular, for δ small enough, $\alpha(\delta^2)$ will be also positive.*

Observe that, if we do not consider the higher order terms (that is, $f = g = h = 0$), we obtain the unperturbed system

$$\begin{aligned} \frac{dx}{dt} &= -xz + \left(\frac{\alpha(\delta^2)}{\delta} + cz \right) y, \\ \frac{dy}{dt} &= - \left(\frac{\alpha(\delta^2)}{\delta} + cz \right) x - yz, \\ \frac{dz}{dt} &= -1 + b(x^2 + y^2) + z^2. \end{aligned} \quad (8)$$

The next lemma gives the main properties of this system.

Lemma 1.3 ([BCS13]). *For any value of $\delta > 0$, the unperturbed system (8) has the following properties:*

1. It possesses two hyperbolic fixed points $S_{\pm}^0 = (0, 0, \pm 1)$ which are of saddle-focus type with eigenvalues $\mp 1 + |\frac{\alpha}{\delta} \pm c|i$, $\mp 1 - |\frac{\alpha}{\delta} \pm c|i$, and ± 2 .
2. The one-dimensional unstable manifold of S_+^0 and the one-dimensional stable manifold of S_-^0 coincide along the heteroclinic connection $\{(0, 0, z) : -1 < z < 1\}$. The time parameterization of this heteroclinic connection is given by

$$\Upsilon_0(t) = (0, 0, z_0(t)) = (0, 0, -\tanh t),$$

if we require $\Upsilon_0(0) = (0, 0, 0)$.

Their 2-dimensional stable/unstable manifolds also coincide, but we will not deal with this problem in this paper.

The critical points given in Lemma 1.3 are persistent for system (7) for small values of $\delta > 0$. Below we summarize some properties of system (7).

Lemma 1.4 ([BCS13]). *If $\delta > 0$ is small enough, system (7) has two fixed points $S_{\pm}(\delta)$ of saddle-focus type,*

$$S_{\pm}(\delta) = (x_{\pm}(\delta), y_{\pm}(\delta), z_{\pm}(\delta)),$$

with

$$x_{\pm}(\delta) = \mathcal{O}(\delta^2), \quad y_{\pm}(\delta) = \mathcal{O}(\delta^2), \quad z_{\pm}(\delta) = \pm 1 + \mathcal{O}(\delta).$$

The point $S_+(\delta)$ has a one-dimensional unstable manifold and a two-dimensional stable one. Conversely, $S_-(\delta)$ has a one-dimensional stable manifold and a two-dimensional unstable one.

Moreover, there are no other fixed points of (7) in the closed ball $B(\delta^{-1/3})$.

The theorem proven in [BCS13] is the following.

Theorem 1.5 ([BCS13]). *Consider system (7), with $\delta > 0$ small enough. Then, there exists a constant C^* , such that the distance $d^{\text{u,s}}$ between the one-dimensional stable manifold of $S_-(\delta)$ and the one-dimensional unstable manifold of $S_+(\delta)$, when they meet the plane $z = 0$, is given by*

$$d^{\text{u,s}} = \delta^{-2} e^{-\frac{\alpha_0 \pi}{2\delta}} e^{\frac{\pi}{2}(\alpha_0 h_0 + c)} \left(C^* + \mathcal{O}\left(\frac{1}{\log(1/\delta)}\right) \right),$$

where $\alpha_0 = \alpha(0)$, and $h_0 = -\lim_{z \rightarrow 0} z^{-3} h(0, 0, z, 0, 0)$.

In [BCS13] was proven that the constant C^* comes from the so called *inner equation* and that, generically, it does not vanish. However, for a given model is usually very hard to prove analytically whether the associated C^* vanishes or not. In this paper we provide a rigorous (computer assisted) method to check whether it vanishes and to compute its value.

1.2 The inner equation

One of the key parts of the proof of Theorem 1.5 is to analyze an inner equation. This equation provides the Stokes constant C^* and it was obtained and analyzed in [BS08]. To obtain it, we perform the change of coordinates $(\phi, \varphi, \eta) = C_{\delta}(x, y, z)$ given by

$$\phi = \delta(x + iy), \quad \varphi = \delta(x - iy), \quad \eta = \delta z, \quad \tau = \frac{t - i\pi/2}{\delta}.$$

Applying this change to system (7), one obtains

$$\begin{aligned}
\frac{d\phi}{d\tau} &= (-\alpha i - \eta)\phi + \tilde{F}_1(\phi, \varphi, \eta, \delta) \\
\frac{d\varphi}{d\tau} &= (\alpha i - \eta)\varphi + \tilde{F}_2(\phi, \varphi, \eta, \delta) \\
\frac{d\eta}{d\tau} &= -\delta^2 + b\phi\varphi + \eta^2 + \tilde{H}(\phi, \varphi, \eta, \delta)
\end{aligned} \tag{9}$$

where

$$\begin{aligned}
\tilde{F}_1(\phi, \varphi, \eta, \delta) &= f(C_\delta^{-1}(\phi, \varphi, \eta), \delta) + ig(C_\delta^{-1}(\phi, \varphi, \eta), \delta), \\
\tilde{F}_2(\phi, \varphi, \eta, \delta) &= f(C_\delta^{-1}(\phi, \varphi, \eta), \delta) - ig(C_\delta^{-1}(\phi, \varphi, \eta), \delta), \\
\tilde{H}(\phi, \varphi, \eta, \delta) &= h(C_\delta^{-1}(\phi, \varphi, \eta), \delta).
\end{aligned}$$

The inner equation comes from (9) taking $\delta = 0$. Defining $F_i(\phi, \varphi, \eta) = \tilde{F}_i(\phi, \varphi, \eta, 0)$ and $H(\phi, \varphi, \eta) = \tilde{H}(\phi, \varphi, \eta, 0)$ and, for technical reasons, performing the change $\eta = -s^{-1}$, we get

$$\begin{aligned}
\frac{d\phi}{d\tau} &= -\left(\alpha i - \frac{1}{s}\right) + F_1(\phi, \varphi, -s^{-1}) \\
\frac{d\varphi}{d\tau} &= \left(\alpha i + \frac{1}{s}\right) + F_2(\phi, \varphi, -s^{-1}) \\
\frac{ds}{d\tau} &= 1 + s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1})).
\end{aligned} \tag{10}$$

We reparameterize time so that equation (10) becomes a non-autonomous 2-dimensional equation with time s ,

$$\begin{aligned}
\phi' &= \frac{-(\alpha i - \frac{1}{s})\phi + F_1(\phi, \varphi, -s^{-1})}{1 + s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1}))} \\
\varphi' &= \frac{(\alpha i + \frac{1}{s})\varphi + F_2(\phi, \varphi, -s^{-1})}{1 + s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1}))}
\end{aligned} \tag{11}$$

with $' = \frac{d}{ds}$.

To analyze this system, we separate its linear terms from the nonlinear ones. Indeed, defining

$$\mathcal{A}(s) = \begin{pmatrix} -i\alpha + \frac{1}{s} & 0 \\ 0 & i\alpha + \frac{1}{s} \end{pmatrix} \tag{12}$$

and

$$\mathcal{S}(\phi, \varphi, s) = \begin{pmatrix} \frac{(\alpha i - \frac{1}{s})\phi s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1})) + F_1(\phi, \varphi, -s^{-1})}{1 + s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1}))} \\ -\frac{(\alpha i + \frac{1}{s})\varphi s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1})) + F_2(\phi, \varphi, -s^{-1})}{1 + s^2(b\phi\varphi + H(\phi, \varphi, -s^{-1}))} \end{pmatrix}, \tag{13}$$

equation (11) can be expressed as

$$\begin{pmatrix} \frac{d\phi}{ds} \\ \frac{d\varphi}{ds} \end{pmatrix} = \mathcal{A}(s) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} + \mathcal{S}(\phi, \varphi, s). \tag{14}$$

From now on we will refer to (14) as the inner equation.

For $s \in \mathbb{C}$ we shall write $\Re s$ and $\Im s$ for its real and imaginary part, respectively. Following [BS08], we define the *inner domains* as

$$\mathcal{D}_\rho^- = \{s \in \mathbb{C} : |\Im s| \geq -\tan \beta \Re s - \rho, \Re s \leq 0\}, \quad \mathcal{D}_\rho^+ = \{s : -s \in \mathcal{D}_\rho^-\} \quad (15)$$

for some $\rho > 0$.

Theorem 1.6 ([BS08]). *If ρ is big enough, the inner equation has two solutions $\psi^\pm = (\phi^\pm, \varphi^\pm)$ defined in \mathcal{D}_ρ^\pm satisfying the asymptotic condition*

$$\lim_{\Re s \rightarrow \pm\infty} \psi^\pm(s) = 0. \quad (16)$$

Moreover its difference satisfies that, for $s \in \mathcal{D}_\rho^+ \cap \mathcal{D}_\rho^- \cap \{\Im s < 0\}$

$$\Delta\psi(s) = \psi^+(s) - \psi^-(s) = se^{-i\alpha(s-h_0 \log s)} \left[\begin{pmatrix} \Theta \\ 0 \end{pmatrix} + \mathcal{O}\left(\frac{1}{|s|}\right) \right]. \quad (17)$$

In addition $\Theta \neq 0$ if and only if $\Delta\psi \neq 0$.

Note that the Stokes constant $\Theta \in \mathbb{C}$ can be defined as

$$\Theta = \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha(s-h_0 \log s)} \Delta\phi(s). \quad (18)$$

Later, in [BCS13] the authors prove that, if C^* is the constant introduced in Theorem 1.5, then

$$C^* = |\Theta|.$$

However there is no closed formula for Θ , which depends on the full jet of the nonlinear terms in (14). Our strategy to compute Θ is to perform a computer assisted proof.

2 Rigorous computation of the Stokes constant

We propose a method to compute the Stokes constant Θ relying on rigorous, computer assisted, interval arithmetic based validation. The method takes advantage from the constructive method for proving Theorem 1.6 based on fixed point arguments and we strongly believe that it can be applied to other settings as, for instance, the classical rapidly forced pendulum and close to the identity area preserving maps.

The method we propose to compute the Stokes constant Θ is divided into two parts.

- Part 1: We provide an algorithm to give an explicit $\rho_* > 0$ such that the existence of the solutions ψ^\pm of the inner equation, in the domain $\{\Re s = 0, \Im s \leq -\rho_*\}$ is guaranteed.

The algorithm is based in giving explicit bounds (which depend on the nonlinear terms \mathcal{S} of the inner equation, see (13)) of all the constants involved in the fixed point argument. We believe that this algorithm can be generalized to other situations where the proof of the existence of the corresponding solutions of the inner equation relies on fixed point arguments.

In the case of the Hopf-zero singularity, by Theorem 1.6, if one can check (using rigorous computer computations) that $\Delta\psi(-i\rho^*) = \psi^+(-i\rho^*) - \psi^-(-i\rho^*) \neq 0$ one can ensure that $\Theta \neq 0$.

- Part 2: Using that $\Delta\psi(s)$ is defined for $s \in \{\Re s = 0, \Im s \leq -\rho_*\}$ with ρ_* given in Part 1, we give a method which provides rigorous accurate estimates for Θ .

We give an algorithm to compute $\rho_0 \geq \rho_*$ such that, for all $\rho \geq \rho_0$, the Stokes constant and $\Delta\phi(-i\rho)$ satisfies the relation

$$\Theta = i\rho^{-1}\Delta\phi(-i\rho)e^{\alpha(\rho - ih_0 \log \rho - h_0 \frac{\pi}{2})}(1 + g(\rho)) \quad (19)$$

with $|g(\rho)| < 1$. By (17), we know that $|g(\rho)|$ is of order $\mathcal{O}(\rho^{-1})$. We provide explicit upper bounds for it.

Part 2 also relies on evaluating $\Delta\psi(-i\rho)$ but takes more advantage on the fixed point argument techniques used to prove formula (17) in Theorem 1.6.

A similar formula to (19) for Θ can be deduced in other settings such as the rapidly forced pendulum and close to the identity area preserving maps. We should be able to adapt our method to a plethora of different situations.

In Section 2.1 we show the theoretical framework we use to design the method. In particular, the functional setting needed for the fixed point argument. It is divided in Sections 2.1.1 and 2.1.2 which deal with Part 1 and Part 2 respectively. In Section 2.2, we follow the theoretical approach given in the previous sections and compute all the necessary constants to implement the method. After that, in Section 2.3 we write the precise algorithm, pointing out all the constants that need to be computed to find Θ . In Section 3, we apply our method to two examples. Finally, in Section 4, we explain how to improve the accuracy in the computation of the Stokes constant in one of the examples considered in Section 3.

2.1 Scheme of the method. Theoretical approach

2.1.1 Existence domain of the solutions of the inner equation

We analyze the solutions $\psi^\pm = (\phi^\pm, \varphi^\pm)$ of equation (14) in the inner domains \mathcal{D}_ρ^\pm introduced in (15). To prove the existence of the solutions ψ^\pm , we set up a fixed point argument. From now on we use subindices 1 and 2 to denote the two components of all vectors and operators.

Note that the right hand side of equation (14) has a linear part plus higher order terms (which will be treated as perturbation). We consider a fundamental matrix $M(s)$ associated to the matrix \mathcal{A} in (12) given by

$$M(s) = s \begin{pmatrix} e^{-i\alpha s} & 0 \\ 0 & e^{i\alpha s} \end{pmatrix} \quad (20)$$

and we define also the integral operators

$$\mathcal{B}^\pm(h) = \begin{pmatrix} \mathcal{B}_1^\pm(h) \\ \mathcal{B}_2^\pm(h) \end{pmatrix} = M(s) \int_{\pm\infty}^0 M(s+t)^{-1} h(s+t) dt. \quad (21)$$

Then, the solutions ψ^\pm of equation (14) satisfying the asymptotic conditions (16) must be also solutions of the integral equation

$$\psi^\pm = \mathcal{B}^\pm(\mathcal{S}(\psi, s)).$$

Therefore, we look for fixed points of the operators

$$\mathcal{F}^\pm(\psi) = \mathcal{B}^\pm(\mathcal{S}(\psi, s)). \quad (22)$$

We define the Banach spaces

$$\mathcal{X}_\nu^\pm = \{h : \mathcal{D}_\rho^\pm \rightarrow \mathbb{C} : \text{analytic}, \|h\|_\nu < \infty\} \quad \text{where} \quad \|h\|_\nu = \sup_{s \in \mathcal{D}_\rho^\pm} |s^\nu h(s)|. \quad (23)$$

Then, we obtain fixed points of the operators \mathcal{F}^\pm in the Banach spaces $\mathcal{X}_\nu \times \mathcal{X}_\nu$ with the norm

$$\|(\phi, \varphi)\|_\nu = \max\{\|\phi\|_\nu, \|\varphi\|_\nu\},$$

for some ν to be chosen.

In [BS08], it is proven that the operators \mathcal{F}^\pm are contractive operators in some ball of $\mathcal{X}_3 \times \mathcal{X}_3$ if $\rho \geq \rho^*$ is big enough. Consequently the existence of solutions ψ^\pm of equation (14) in the domains \mathcal{D}_ρ^\pm is guaranteed. However, we want to be explicit in the estimates to compute the smallest ρ_* such that one can prove that \mathcal{F}^\pm are contractive operators.

To this end, we need to control the dependence on ρ of the Lipschitz constant of the operators \mathcal{F}^\pm . Let us explain briefly the procedure, which is performed only for the $-$ case being the $+$ case analogous.

- In Section 2.2.1, we provide explicit bounds for the norm of the linear operator \mathcal{B}^- in (21).
- In Section 2.2.2, we define a set of constants depending on the nonlinear terms \mathcal{S} (see (13)) of the inner equation.
- We deal with the bounds of the first iteration, $\mathcal{F}^+(0)$ in Section 2.2.3. We conclude that it belongs to a closed ball of $\mathcal{X}_3 \times \mathcal{X}_3$ if $\rho \geq \rho_*^1$ where ρ_*^1 is determined by the constants in the previous step. The radius of the ball, $M_0(\rho)/2$ is fully determined also by the previous constants.
- In Section 2.2.4, we provide explicit bounds of the derivative of the nonlinear operator \mathcal{S} and consequently of its Lipschitz constant, which depends on ρ . These computations hold true for values of $\rho \geq \rho_*^2 \geq \rho_*^1$ with ρ_*^2 satisfying some explicit conditions.
- In Section 2.2.5, for $\rho \geq \rho_*^2$, we compute the Lipschitz constant $L(\rho)$ of \mathcal{F}^- in the closed ball of $\mathcal{X}_3 \times \mathcal{X}_3$ of radius $M_0(\rho)$.
- In Section 2.2.6, we set $\rho_* \geq \rho_*^2$ for the existence result. We choose ρ^* such that $L(\rho_*) \leq \frac{1}{2}$. Then, since

$$\|\psi^-\|_3 \leq \|\mathcal{F}^-(0)\|_3 + \|\mathcal{F}^-(\psi^-) - \mathcal{F}^-(0)\|_3 \leq \frac{M_0(\rho)}{2} + L(\rho)\|\psi^-\|_3 \leq M_0(\rho)$$

the fixed point theorem ensures the existence of a fixed point ψ^- satisfying $\|\psi^-\|_3 \leq M_0(\rho)$ for $\rho \geq \rho^*$.

- Finally, we compute $\Delta\phi(-i\rho_*)$ by computed assisted proofs techniques. This completes the Part 1 of the algorithm since $\Delta\phi(-i\rho_*) \neq 0$ implies $\Theta \neq 0$.

All the steps described above are written with all the detailed constants in Section 2.3.

2.1.2 Rigorous computation of the Stokes constant

In this section we describe a method to compute rigorously the Stokes constant Θ defined in (18) (Part 2 of the algorithm). The method is based in the alternative formula for Θ proposed in (19):

$$\Theta = i \frac{e^{\alpha(\rho - ih_0 \log \rho - h_0 \frac{\pi}{2})} \Delta\phi(-i\rho)}{\rho} (1 + g(\rho)), \quad \lim_{\rho \rightarrow \infty} g(\rho) = 0. \quad (24)$$

Let us to explain how this formula is derived. The key point is to analyze the difference

$$\Delta\psi(s) = \psi^+(s) - \psi^-(s)$$

as a solution of a linear equation on the vertical axis $\Im s \in (-\infty, -\rho_*)$ where ρ_* is provided by the method explained in Section 2.1.1. Indeed $\Delta\psi = (\Delta\phi, \Delta\varphi)$ satisfies the equation

$$\begin{pmatrix} \Delta\phi' \\ \Delta\varphi' \end{pmatrix} = (\mathcal{A}(s) + \mathcal{K}(s)) \begin{pmatrix} \Delta\phi \\ \Delta\varphi \end{pmatrix}$$

where

$$\mathcal{K}(s) = \int_0^1 DS(\psi^-(s) + t(\psi^+(s) - \psi^-(s)), s) dt \quad (25)$$

and \mathcal{S} is given in (13). We look for the linear terms of lower order in s^{-1} of \mathcal{S} . Indeed, we have that

$$\mathcal{S}(\psi, s) = \frac{1}{s} \begin{pmatrix} -\alpha ih_0 \phi \\ \alpha ih_0 \varphi \end{pmatrix} + \tilde{\mathcal{S}}(\psi, s)$$

with $\tilde{\mathcal{S}}(\psi, s) = \mathcal{O}(|s|^{-2})$ when $\psi \in \mathcal{X}_3$ (see (23)). Then, $\Delta\psi$ satisfies the equation

$$\begin{pmatrix} \Delta\phi' \\ \Delta\varphi' \end{pmatrix} = (\tilde{\mathcal{A}}(s) + \tilde{\mathcal{K}}(s)) \begin{pmatrix} \Delta\phi \\ \Delta\varphi \end{pmatrix} \quad (26)$$

where

$$\tilde{\mathcal{A}}(s) = \begin{pmatrix} -i\alpha + \frac{1}{s} - i\alpha \frac{h_0}{s} & 0 \\ 0 & i\alpha + \frac{1}{s} + i\alpha \frac{h_0}{s} \end{pmatrix} \quad (27)$$

and

$$\tilde{\mathcal{K}}(s) = \int_0^1 D\tilde{\mathcal{S}}(\psi^-(s) + t(\psi^+(s) - \psi^-(s)), s) dt. \quad (28)$$

A fundamental matrix for the linear system $z' = \tilde{\mathcal{A}}(s)z$ is

$$\begin{pmatrix} se^{-i\alpha(s+h_0 \log s)} & 0 \\ 0 & se^{i\alpha(s+h_0 \log s)} \end{pmatrix}.$$

Therefore, any solution of system (26) can be expressed as

$$\begin{pmatrix} \Delta\phi \\ \Delta\varphi \end{pmatrix} = \begin{pmatrix} se^{-i\alpha(s+h_0 \log s)} \left[\kappa_0 + \int_{-i\rho}^s \frac{e^{i\alpha(t+h_0 \log t)}}{t} (\tilde{\mathcal{K}}_{11}\Delta\phi + \tilde{\mathcal{K}}_{12}\Delta\varphi) dt \right] \\ se^{i\alpha(s+h_0 \log s)} \left[\kappa_1 + \int_{-i\rho}^s \frac{e^{-i\alpha(t+h_0 \log t)}}{t} (\tilde{\mathcal{K}}_{21}\Delta\phi + \tilde{\mathcal{K}}_{22}\Delta\varphi) dt \right] \end{pmatrix}$$

with κ_0, κ_1 two constants.

Since $|\psi^\pm| \leq M_0(\rho)|s|^{-3}$, $\Delta\psi$ goes to 0 as $\Im s \rightarrow -\infty$ and, therefore,

$$\kappa_1 = - \int_{-i\rho}^{-i\infty} \frac{e^{-i\alpha(t+h_0 \log t)}}{t} (\tilde{\mathcal{K}}_{21}\Delta\phi + \tilde{\mathcal{K}}_{22}\Delta\varphi) dt.$$

Then, we deduce that the difference $\Delta\psi(s)$ is a fixed point of the equation

$$\begin{aligned} \Delta\psi(s) &= \Delta\psi^0(s) + \mathcal{G}(\Delta\psi)(s), \quad \text{where} \\ \Delta\psi^0(s) &= \begin{pmatrix} se^{-i\alpha(s+h_0 \log s)} \kappa_0 \\ 0 \end{pmatrix} \end{aligned} \quad (29)$$

with κ_0 a constant depending on ρ and \mathcal{G} is the linear operator

$$\mathcal{G}(\Delta\psi) = \begin{pmatrix} se^{-i\alpha(s+h_0 \log s)} \int_{-i\rho}^s \frac{e^{i\alpha(t+h_0 \log t)}}{t} (\tilde{\mathcal{K}}_{11}\Delta\phi + \tilde{\mathcal{K}}_{12}\Delta\varphi) dt \\ se^{i\alpha(s+h_0 \log s)} \int_{-i\infty}^s \frac{e^{-i\alpha(t+h_0 \log t)}}{t} (\tilde{\mathcal{K}}_{21}\Delta\phi + \tilde{\mathcal{K}}_{22}\Delta\varphi) dt \end{pmatrix}. \quad (30)$$

By construction, κ_0 is defined as

$$\kappa_0 = \kappa_0(\rho) = i \frac{e^{\alpha(\rho - ih_0 \log \rho - h_0 \frac{\pi}{2})} \Delta\phi(-i\rho)}{\rho}. \quad (31)$$

Using (18) and (29), we have

$$\begin{aligned} \Theta &= \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha(s+h_0 \log s)} \Delta\phi(s) \\ &= \kappa_0 \left(1 + \kappa_0^{-1} \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha(s+h_0 \log s)} \mathcal{G}_1(\Delta\psi(s)) \right). \end{aligned} \quad (32)$$

We use equality (32) to obtain formula (24) of Θ . To bound $|g(\rho)|$ in formula (24), we need to control the linear operator $s^{-1} e^{i\alpha(s+h_0 \log s)} \mathcal{G}_1$. To this end, we consider a norm with exponential weights,

$$\|\psi\| = \max \left\{ \max_{s \in E} \left| s^{-1} e^{i\alpha(s+h_0 \log s)} \phi \right|, \max_{s \in E} \left| e^{i\alpha(s+h_0 \log s)} \varphi \right| \right\} \quad (33)$$

with $E = \{\Re s = 0, \Im s \in (-\rho_*, -\infty)\}$.

Observe that (29) can be rewritten

$$(\text{Id} - \mathcal{G})(\Delta\psi) = \Delta\psi^0.$$

Now we see that $\text{Id} - \mathcal{G}$ is an invertible operator. Indeed, in [BS08], it was proven that for ρ big enough

$$\|\mathcal{G}_1(\Phi)\| \leq A_1(\rho)\|\Phi\|, \quad \|\mathcal{G}_2(\Phi)\| \leq A_2(\rho)\|\Phi\| \quad (34)$$

with $0 < A(\rho) := \max\{A_1(\rho), A_2(\rho)\} < 1$. Moreover, [BS08] also shows that

$$\lim_{\rho \rightarrow \infty} A(\rho) = 0.$$

These estimates imply that, if ρ is big enough, $\text{Id} - \mathcal{G}$ is invertible and therefore $\Delta\psi = (\text{Id} - \mathcal{G})^{-1}(\Delta\psi^0)$. Moreover,

$$\|\Delta\psi\| \leq \frac{1}{1 - A(\rho)} \|\Delta\psi^0\| = \frac{|\kappa_0(\rho)|}{1 - A(\rho)} \quad (35)$$

and this inequality directly gives

$$\left| \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha(s+h_0 \log s)} \mathcal{G}_1(\Delta\psi(s)) \right| \leq A_1(\rho) \frac{|\kappa_0(\rho)|}{1 - A(\rho)}.$$

Therefore, from (32), we can conclude that the Stokes constant Θ , which is independent of ρ , can be computed as

$$\Theta = \kappa_0(\rho)(1 + g(\rho)),$$

for any ρ big enough, where κ_0 is given in (31) and g satisfies

$$|g(\rho)| \leq \overline{M}(\rho) := \frac{A_1(\rho)}{1 - A(\rho)}. \quad (36)$$

Since $A(\rho), A_1(\rho)$ go to zero as $\rho \rightarrow \infty$, the same happens for $\overline{M}(\rho)$. Then (24) is proven.

Notice that the relative error to approximate Θ by κ_0 is

$$\frac{|\Theta - \kappa_0(\rho)|}{|\kappa_0(\rho)|} \leq \overline{M}(\rho).$$

As a consequence,

$$|\Theta| \in [|\kappa_0(\rho)|(1 - \overline{M}(\rho)), |\kappa_0(\rho)|(1 + \overline{M}(\rho))].$$

In Section 2.2 the procedure described above is implemented:

- Following the fixed point argument in [BS08], in Section 2.2.7 we give an explicit formula for $A(\rho) = \max\{A_1(\rho), A_2(\rho)\}$ in (34) for $\rho \geq \rho_*$, where ρ_* is the constant given by Part 1.
- In Section 2.2.8, we set $\rho_0 \geq \rho_*$ such that $\overline{M}(\rho) < 1$ for $\rho \geq \rho_0$.

2.2 Computing the Stokes constant: method

In this section we are going to give explicit expressions for all the constant involved in the method explained in the previous section.

2.2.1 The linear operator \mathcal{B}^-

Lemma 2.1. *Consider the linear operator \mathcal{B}^- defined in (21).*

1. When $\nu > 1$, the linear operator $\mathcal{B}^- : \mathcal{X}_\nu \times \mathcal{X}_\nu \rightarrow \mathcal{X}_{\nu-1} \times \mathcal{X}_{\nu-1}$ is continuous and

$$\|\mathcal{B}^-(\psi)\|_{\nu-1} \leq B_{\nu+1} \|\psi\|_\nu$$

where

$$\begin{aligned} B_m &= \frac{\pi (m-3)!!}{2 (m-2)!!} && \text{if } m \text{ is even} \\ B_m &= \frac{(m-3)!!}{(m-2)!!} && \text{if } m \text{ is odd.} \end{aligned} \quad (37)$$

2. When $\nu > 0$, the linear operator $\mathcal{B}^- : \mathcal{X}_\nu \times \mathcal{X}_\nu \rightarrow \mathcal{X}_\nu \times \mathcal{X}_\nu$ is continuous and, for all $0 < \gamma \leq \beta$ (see (15)),

$$\|\mathcal{B}^-(\psi)\|_\nu \leq \frac{1}{\alpha \sin \gamma (\cos \gamma)^{\nu+1}} \|\psi\|_\nu.$$

Define $\gamma_* \in (0, \frac{\pi}{2})$ such that $\sin^2 \gamma_* = \frac{1}{\nu+2}$. If $\gamma_* \leq \beta$,

$$\|\mathcal{B}^-(\psi)\|_\nu \leq C_\nu \|\psi\|_\nu \quad \text{where} \quad C_\nu = \frac{(\nu+2)^{\frac{\nu+2}{2}}}{\alpha(\nu+1)^{\frac{\nu+1}{2}}}. \quad (38)$$

This lemma is proven in Appendix A.

From now on we choose β , the angle in the definition (15) of \mathcal{D}_ρ^- , be such that $6 \sin \beta^2 = 1$. Then for all $\nu \geq 4$, the optimal value γ_* in second item of Lemma 2.1 satisfies that $\gamma_* \leq \beta$ and the optimal bound (38) will be used throughout the paper.

We emphasize that, if $s \in \mathcal{D}_\rho^-$, one has that $|s| \geq \rho$. Recall that we are looking for ρ_* the minimum value for ρ to ensure that the inner equation has a solution ψ^- defined in \mathcal{D}_ρ^- . Since we need ρ_*^{-1} to be small, we start by assuming that $\rho_* \geq 2$. We will change this value along the proof.

2.2.2 Explicit constants for the inner equation

We consider the max norm $|(x, y, z)| = \max\{|x|, |y|, |z|\}$. Let $a_3 = \lim_{z \rightarrow 0} z^{-3} F_1(0, 0, z)$, $h_0 = \lim_{z \rightarrow 0} z^{-3} H(0, 0, z)$ and $C_F^0, C_H^0, \bar{C}_H^0$ be such that for $|z| \leq \frac{1}{2}$,

$$\begin{aligned} |\Delta F_1(z)| &= |F_1(0, 0, z) + a_3 z^3| \leq C_F^0 |z|^4, \\ |\Delta F_2(z)| &= |F_2(0, 0, z) + \bar{a}_3 z^3| \leq C_F^0 |z|^4, \\ |\Delta H(z)| &= |H(0, 0, z) + h_0 z^3| \leq C_H^0 |z|^4, \\ |H(0, 0, z)| &\leq \bar{C}_H^0 |z|^3. \end{aligned} \quad (39)$$

We also introduce $C_F, C_F^{\phi, \varphi}, C_H, C_H^{\phi, \varphi}$ such that, for $|(x, y)| \leq |z|$,

$$\begin{aligned} |H(x, y, z)| &\leq C_H |(x, y, z)|^3 \leq C_H |z|^3 \\ |F_{1,2}(x, y, z)| &\leq C_F |(x, y, z)|^3 \leq C_F |z|^3, \\ |\partial_x F_{1,2}(x, y, z)| &\leq C_F^\phi |(x, y, z)|^2 \leq C_F^\phi |z|^2, \\ |\partial_y F_{1,2}(x, y, z)| &\leq C_F^\varphi |(x, y, z)|^2 \leq C_F^\varphi |z|^2, \\ |\partial_x H(x, y, z)| &\leq C_H^\phi |(x, y, z)|^2 \leq C_H^\phi |z|^2, \\ |\partial_y H(x, y, z)| &\leq C_H^\varphi |(x, y, z)|^2 \leq C_H^\varphi |z|^2. \end{aligned} \quad (40)$$

As a consequence, setting

$$\overline{C}_H = C_H^\phi + C_H^\varphi \quad \text{and} \quad \overline{C}_F = C_F^\phi + C_F^\varphi \quad (41)$$

we have

$$\begin{aligned} |H(x, y, z) + h_0 z^3| &\leq C_H^0 |z|^4 + \overline{C}_H |(x, y)| |z|^2 \\ |F_1(x, y, z) + a_3 z^3| &\leq C_F^0 |z|^4 + \overline{C}_F |(x, y)| |z|^2 \\ |F_2(x, y, z) + \overline{a}_3 z^3| &\leq C_F^0 |z|^4 + \overline{C}_F |(x, y)| |z|^2. \end{aligned}$$

2.2.3 Bounds for the norm of the first iteration

The second step in the proof consists on studying $\mathcal{F}^-(0)(s) = \mathcal{B}^-(\mathcal{S}(0, s))$, where \mathcal{F}^- is the operator introduced in (22).

Lemma 2.2. *Chose any $\rho_*^1 > \max\{2, \overline{C}_H^0\}$, take $\rho \geq \rho_*^1$ and define*

$$\mathcal{C}_0(\rho) = \frac{C_F^0 + |a_3| \overline{C}_H^0}{1 - \frac{|\overline{C}_H^0|}{\rho}}.$$

Then $\mathcal{F}^-(0) \in \mathcal{X}_3 \times \mathcal{X}_3$ and

$$\|\mathcal{F}^-(0)\|_3 \leq \frac{11|a_3|}{3\alpha} + B_5 \mathcal{C}_0(\rho).$$

Proof. By (13), it is clear that

$$\mathcal{S}_1(0, s) - \frac{a_3}{s^3} = \frac{F_1(0, 0, -s^{-1})}{1 + s^2 H(0, 0, -s^{-1})} - \frac{a_3}{s^3} = \frac{\Delta F_1(-s^{-1}) - \frac{a_3 H(0, 0, -s^{-1})}{s}}{1 + s^2 H(0, 0, -s^{-1})}$$

and therefore

$$\left| \mathcal{S}_1(0, s) - \frac{a_3}{s^3} \right| \leq \frac{1}{|s|^4} \frac{C_F^0 + |a_3| \overline{C}_H^0}{1 - \frac{\overline{C}_H^0}{\rho}}.$$

An analogous bound works for $\mathcal{S}_2(0, s)$ and therefore

$$\|\mathcal{S}(0, s) - s^{-3}(a_3, \overline{a}_3)\|_4 \leq \frac{C_F^0 + |a_3| \overline{C}_H^0}{1 - \frac{|\overline{C}_H^0|}{\rho}} = \mathcal{C}_0(\rho). \quad (42)$$

We introduce $\mathcal{S}_0(s) = s^{-3}(a_3, \overline{a}_3)$. We have that

$$\mathcal{B}_1^-(\mathcal{S}_0(s)) = a_3 s \int_{-\infty}^0 \frac{e^{iat}}{(s+t)^4} dt = \frac{a_3}{i\alpha s^3} + \frac{4sa_3}{i\alpha} \int_{-\infty}^0 \frac{e^{iat}}{(s+t)^5} dt. \quad (43)$$

Notice that, for $s \in \mathcal{D}_\rho^-$ and $t \in \mathbb{R}$, $|s+t|^2 \geq |s|^2 + t^2$. Then, using also Lemma A.1 (see Appendix A),

$$\left| s \int_{-\infty}^0 \frac{e^{iat}}{(s+t)^5} dt \right| \leq \frac{1}{|s|^3} \int_{-\infty}^0 \frac{1}{(t^2+1)^{5/2}} dt = \frac{2}{3|s|^3}.$$

Using this last bound and formula (43) we obtain

$$|\mathcal{B}_1^-(\mathcal{S}_0(s))| \leq \frac{1}{|s|^3} \left(\frac{|a_3|}{\alpha} + \frac{8|a_3|}{3\alpha} \right) \leq \frac{11|a_3|}{3\alpha|s|^3}.$$

To finish we notice that, from (42) and the first item of Corollary 2.1,

$$\|\mathcal{F}_1^-(0)\|_3 \leq \|\mathcal{B}_1^-(\mathcal{S}_0)\|_3 + \|\mathcal{B}_1^-(\mathcal{S}(0, \cdot) - \mathcal{S}_0)\|_3 \leq \frac{11|a_3|}{3\alpha} + \mathcal{C}_0(\rho)B_5.$$

Analogous computations lead to the same estimate for $\|\mathcal{F}_2^-(0)\|_3$. □

2.2.4 The Lipschitz constant of \mathcal{S}

Let

$$M_0(\rho) = \frac{22|a_3|}{3\alpha} + 2B_5\mathcal{C}_0(\rho) \quad (44)$$

in such a way that $2\|\mathcal{F}^-(0)\|_3 \leq M_0(\rho)$.

Lemma 2.3. *Assume that $\|\phi\|_3, \|\varphi\|_3 \leq M_0(\rho)$ and take $\rho \geq \rho_*^2$ being $\rho_*^2 \geq \rho_*^1$ such that*

$$\min \left\{ 1 - \frac{bM_0^2(\rho_*^2)}{(\rho_*^2)^4} - \frac{C_H}{(\rho_*^2)}, 1 - \frac{M_0(\rho_*^2)}{(\rho_*^2)^2} \right\} > 0. \quad (45)$$

Then

$$\begin{aligned} |\partial_\phi \mathcal{S}_1(\psi, s)|, |\partial_\varphi \mathcal{S}_2(\psi, s)| &\leq \frac{M_{11}^1(\rho)}{|s|} + \frac{M_{11}^2(\rho)}{|s|^2} + \frac{M_{11}^3(\rho)}{|s|^3} + \frac{M_{11}^4(\rho)}{|s|^4} \\ |\partial_\varphi \mathcal{S}_1(\psi, s)|, |\partial_\phi \mathcal{S}_2(\psi, s)| &\leq \frac{M_{12}^2(\rho)}{|s|^2} + \frac{M_{12}^3(\rho)}{|s|^3} + \frac{M_{12}^4(\rho)}{|s|^4} \end{aligned}$$

with

$$\begin{aligned}
M_{11}^1(\rho) &= \frac{\alpha|h_0|}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}}, \\
M_{11}^2(\rho) &= \frac{|h_0| + \alpha C_H^0 + C_F^\phi}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}}, \\
M_{11}^3(\rho) &= \frac{1}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}\right)^2} \left[M_0 \alpha C_H^\phi + C_F C_H^\phi + (\alpha \bar{C}_H M_0 + C_H^0) \left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}\right) \right], \\
M_{11}^4(\rho) &= \frac{M_0}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}\right)^2} \\
&\quad \cdot \left[b(C_F + \alpha M_0) + C_H^\phi + \left(\alpha b M_0 + \bar{C}_H + \frac{bM_0}{\rho} \right) \left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}\right) + \frac{bM_0}{\rho} \right] \\
M_{12}^2(\rho) &= \frac{C_F^\varphi}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}}, \\
M_{12}^3(\rho) &= \frac{M_0 \alpha C_H^\varphi + C_F C_H^\varphi}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}\right)^2}, \\
M_{12}^4(\rho) &= \frac{M_0}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}\right)^2} \left(b(C_F + \alpha M_0) + C_H^\varphi + \frac{bM_0}{\rho} \right).
\end{aligned} \tag{46}$$

Proof. Notice that $\rho_*^2 \geq \rho_*^1$ and therefore, Lemma 2.2 can be applied for $\rho \geq \rho_*^2$. Moreover, if $s \in \mathcal{D}_\rho^-$,

$$|\psi(s)| \leq \frac{M_0(\rho)}{|s|^3} \leq \frac{1}{|s|}$$

so that the bounds in (40) can also be used.

We start with $\partial_\phi \mathcal{S}$. We introduce

$$\begin{aligned}
S_1(\psi, s) &= \frac{\partial_\phi F_1(\psi, -s^{-1})}{1 + s^2(b\phi\varphi + H(\psi, -s^{-1}))} + \frac{F_1(\psi, -s^{-1})s^2(b\varphi + \partial_\phi H(\psi, -s^{-1}))}{(1 + s^2(b\phi\varphi + H(\psi, -s^{-1})))^2} \\
S_2(\psi, s) &= \frac{(\alpha i - \frac{1}{s})s^2(b\phi\varphi + H(\psi, -s^{-1}))}{1 + s^2(b\phi\varphi + H(\psi, -s^{-1}))} - \frac{(\alpha i - \frac{1}{s})\phi s^2(b\varphi + \partial_\phi H(\psi, -s^{-1}))}{(1 + s^2(b\phi\varphi + H(\psi, -s^{-1})))^2}.
\end{aligned}$$

Straightforward computations, lead us to

$$\partial_\phi \mathcal{S}_1 = S_1 + S_2.$$

When $\|\phi\|_3, \|\varphi\|_3 \leq M_0(\rho)$,

$$\begin{aligned}
|S_1(\psi, s)| &\leq \frac{1}{|s|^2} \frac{C_F^\phi}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}} + \frac{1}{|s|^3} \frac{C_F \left(C_H^\phi + b\frac{M_0}{|s|} \right)}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2} \\
&\leq \frac{1}{|s|^2} \frac{C_F^\phi}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}} + \frac{1}{|s|^3} \frac{C_F C_H^\phi}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2} + \frac{1}{|s|^4} \frac{bM_0 C_F}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2}, \\
|S_2(\psi, s)| &\leq \frac{1}{|s|} \frac{\left(\alpha + \frac{1}{|s|} \right) \left(|h_0| + \frac{C_H^0}{|s|} + \frac{\bar{C}_H M_0}{|s|^2} + b\frac{M_0^2}{|s|^3} \right)}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}} + \frac{M_0 \left(\alpha + \frac{1}{|s|} \right) \left(\frac{C_H^\phi}{|s|^2} + \frac{bM_0}{|s|^3} \right)}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2} \\
&= \frac{1}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}} \left(\frac{\alpha|h_0|}{|s|} + \frac{|h_0| + \alpha C_H^0}{|s|^2} + \frac{\alpha \bar{C}_H M_0 + C_H^0}{|s|^3} + \frac{\alpha b M_0^2 + \bar{C}_H M_0}{|s|^4} + \frac{b M_0^2}{|s|^5} \right) \\
&\quad + \frac{1}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2} \left(\frac{M_0 \alpha C_H^\phi}{|s|^3} + \frac{\alpha b M_0^2 + M_0 C_H^\phi}{|s|^4} + \frac{b M_0^2}{|s|^5} \right).
\end{aligned}$$

Therefore we have that

$$|\partial_\phi \mathcal{S}_1(\psi, s)| \leq \frac{M_{11}^1(\rho)}{|s|} + \frac{M_{11}^2(\rho)}{|s|^2} + \frac{M_{11}^3(\rho)}{|s|^3} + \frac{M_{11}^4(\rho)}{|s|^4}$$

where M_{11}^k are the constants introduced in the lemma.

We now compute a bound for $\partial_\varphi \mathcal{S}$. As for $\partial_\phi \mathcal{S}$, we define

$$\begin{aligned}
S_1(\psi, s) &= \frac{\partial_\varphi F_1(\psi, -s^{-1})}{1 + s^2(b\phi\varphi + H(\psi, -s^{-1}))} + \frac{F_1(\psi, -s^{-1})s^2(b\phi + \partial_\varphi H(\psi, -s^{-1}))}{(1 + s^2(b\phi\varphi + H(\psi, -s^{-1})))^2} \\
S_2(\psi, s) &= -\frac{(\alpha i - \frac{1}{s})\phi s^2(b\phi + \partial_\varphi H(\psi, -s^{-1}))}{(1 + s^2(b\phi\varphi + H(\psi, -s^{-1})))^2}.
\end{aligned}$$

and we notice that

$$\partial_\varphi \mathcal{S}_1 = S_1 + S_2.$$

We have that, if $\|\phi\|_3, \|\varphi\|_3 \leq M_0(\rho)$,

$$\begin{aligned}
|S_1(\psi, s)| &\leq \frac{1}{|s|^2} \frac{C_F^\varphi}{1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho}} + \frac{1}{|s|^3} \frac{C_F C_H^\varphi}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2} + \frac{1}{|s|^4} \frac{bM_0 C_F}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2}, \\
|S_2(\psi, s)| &\leq \frac{1}{\left(1 - \frac{bM_0^2}{\rho^4} - \frac{C_H}{\rho} \right)^2} \left(\frac{M_0 \alpha C_H^\varphi}{|s|^3} + \frac{\alpha b M_0^2 + M_0 C_H^\varphi}{|s|^4} + \frac{b M_0^2}{|s|^5} \right).
\end{aligned}$$

Then

$$|\partial_\varphi \mathcal{S}_1(\psi, s)| \leq \frac{M_{12}^2(\rho)}{|s|^2} + \frac{M_{12}^3(\rho)}{|s|^3} + \frac{M_{12}^4(\rho)}{|s|^4}$$

with the constants M_{12}^k defined in the lemma.

Since the bounds for $F_1, \partial_\phi F_1, \partial_\varphi F_1$ are the same as for $F_2, \partial_\phi F_2, \partial_\varphi F_2$ and using the symmetry in the definition of \mathcal{S} , we have that

$$\begin{aligned} |\partial_\phi \mathcal{S}_2(\psi, s)| &\leq \frac{M_{12}^2(\rho)}{|s|^2} + \frac{M_{12}^3(\rho)}{|s|^3} + \frac{M_{12}^4(\rho)}{|s|^4} \\ |\partial_\varphi \mathcal{S}_2(\psi, s)| &\leq \frac{M_{11}^1(\rho)}{|s|} + \frac{M_{11}^2(\rho)}{|s|^2} + \frac{M_{11}^3(\rho)}{|s|^3} + \frac{M_{11}^4(\rho)}{|s|^4}. \end{aligned}$$

□

As a corollary we obtain the following.

Corollary 2.4. *If $\psi, \psi' \in B(M_0(\rho))$ with $\rho \geq \rho_*^2$ as in Lemma 2.3. Then, there exist functions $\Delta \mathcal{S}_j$, $j = 1 \dots 4$, such that*

$$\mathcal{S}(\psi, s) - \mathcal{S}(\psi', s) = \sum_{j=1}^4 \Delta \mathcal{S}_j(\psi, s) - \Delta \mathcal{S}_j(\psi', s)$$

and

$$\begin{aligned} \|\Delta \mathcal{S}_1(\psi, s) - \Delta \mathcal{S}_1(\psi', s)\|_4 &\leq M_{11}^1(\rho) \|\psi - \psi'\|_3 \\ \|\Delta \mathcal{S}_j(\psi, s) - \Delta \mathcal{S}_j(\psi', s)\|_{3+j} &\leq (M_{11}^j(\rho) + M_{12}^j(\rho)) \|\psi - \psi'\|_3. \end{aligned}$$

As a consequence

$$\begin{aligned} |\mathcal{S}(\psi, s) - \mathcal{S}(\psi', s)| &\leq |\psi(s) - \psi'(s)| \left(\frac{M_{11}^1(\rho)}{|s|} + \frac{M_{11}^2(\rho) + M_{12}^2(\rho)}{|s|^2} \right. \\ &\quad \left. + \frac{M_{11}^3(\rho) + M_{12}^3(\rho)}{|s|^3} + \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{|s|^4} \right). \end{aligned}$$

Proof. Indeed:

$$\begin{aligned} \mathcal{S}_1(\psi, s) - \mathcal{S}_1(\psi', s) &= (\phi - \phi') \int_0^1 \partial_\phi \mathcal{S}_1(\psi' + \lambda(\psi - \psi')) d\lambda \\ &\quad + (\varphi - \varphi') \int_0^1 \partial_\varphi \mathcal{S}_1(\psi' + \lambda(\psi - \psi')) d\lambda \\ &= (\phi - \phi')(S_1^\phi + S_2^\phi + S_3^\phi + S_4^\phi) + (\varphi - \varphi')(S_2^\varphi + S_3^\varphi + S_4^\varphi) \end{aligned}$$

with $S_j^{\phi, \varphi} \in \mathcal{X}_j$ and

$$\|S_j^\phi\|_j \leq M_{11}^j(\rho), \quad \|S_j^\varphi\|_j \leq M_{12}^j(\rho).$$

In analogous way we decompose $\mathcal{S}_2(\psi) - \mathcal{S}_2(\psi')$ and by symmetry we obtain that

$$\mathcal{S}(\psi, s) - \mathcal{S}(\psi', s) = \begin{pmatrix} S_{11}(s) & S_{12}(s) \\ S_{21}(s) & S_{22}(s) \end{pmatrix} \begin{pmatrix} \phi - \phi' \\ \varphi - \varphi' \end{pmatrix} \quad (47)$$

with

$$\begin{aligned} |S_{11}(s)|, |S_{22}(s)| &\leq \frac{M_{11}^1(\rho)}{|s|} + \frac{M_{11}^2(\rho)}{|s|^2} + \frac{M_{11}^3(\rho)}{|s|^3} + \frac{M_{11}^4(\rho)}{|s|^4} \\ |S_{12}(s)|, |S_{21}(s)| &\leq \frac{M_{12}^2(\rho)}{|s|^2} + \frac{M_{12}^3(\rho)}{|s|^3} + \frac{M_{12}^4(\rho)}{|s|^4}. \end{aligned}$$

Namely, S_{ij} can be decomposed as a sum of functions belonging to the adequate \mathcal{X}_k . Therefore, taking the supremum norm in (47) we get the result. □

Remark 2.5. Notice that, for some concrete functions $F_{1,2}$ and H the general bounds in that we have used for them and their derivatives (see (40)) may not be sharp. To improve the estimates for $|\partial_{\phi,\varphi}\mathcal{S}|$ in Lemma 2.3, we need to track some terms of the functions $F_{1,2}, H$. Indeed, instead of (40) we can use bounds of the derivatives of the form

$$\begin{aligned} |\partial_x F_{1,2}(x, y, z)| &\leq c_F^\phi |z|^2 + K_F^\phi(x, y) |z| \\ |\partial_y F_{1,2}(x, y, z)| &\leq c_F^\varphi |z|^2 + K_F^\varphi(x, y) |z| \\ |\partial_x H(x, y, z)| &\leq c_H^\phi |z|^2 + K_H^\phi(x, y) |z| \\ |\partial_y H(x, y, z)| &\leq c_H^\varphi |z|^2 + K_H^\varphi(x, y) |z| \end{aligned}$$

which, together with (39), imply

$$\begin{aligned} |F_{1,2}(x, y, z) + a_3 z^3| &\leq C_F^0 |z|^4 + \bar{C}_F |(x, y)| |z|^2 + \bar{K}_F |(x, y)|^2 |z| \\ |H(x, y, z) + h_0 z^3| &\leq C_H^0 |z|^4 + \bar{C}_H |(x, y)| |z|^2 + \bar{K}_H |(x, y)|^2 |z| \end{aligned}$$

with $\bar{C}_F = c_F^\phi + c_F^\varphi$, $\bar{C}_H = c_H^\phi + c_H^\varphi$, $\bar{K}_F = K_F^\phi + K_F^\varphi$, $\bar{K}_H = K_H^\phi + K_H^\varphi$. If necessary, since $|(x, y)| \leq |z|$, we can also use

$$\begin{aligned} |F_{1,2}(x, y, z) + a_3 z^3| &\leq C_F^0 |z|^4 + \tilde{C}_F |(x, y)| |z|^2 \\ |H(x, y, z) + h_0 z^3 + a_4 z^4 + a_5 z^5| &\leq \tilde{C}_H |z|^6 + \tilde{C}_H |(x, y)| |z|^2. \end{aligned}$$

where

$$\tilde{C}_F = \bar{C}_F + \bar{K}_F, \quad \tilde{C}_H = \bar{C}_H + \bar{K}_H.$$

It is clear that, taking

$$\begin{aligned} C_F &= |a_3| + \frac{C_F^0}{\rho} + \frac{\tilde{C}_F M_0}{\rho^2}, & C_F^{\phi,\varphi} &= c_F^{\phi,\varphi} + \frac{K_F^{\phi,\varphi} M_0}{\rho^2} \\ C_H &= |a_4| + \frac{|a_5|}{\rho} + \frac{\tilde{C}_H^0}{\rho^2} + \frac{\tilde{C}_H M_0}{\rho}, & C_H^{\phi,\varphi} &= c_H^{\phi,\varphi} + \frac{K_H^{\phi,\varphi} M_0}{\rho^2}, \end{aligned}$$

we can get a more accurate bound for $|\partial_{\phi,\varphi}\mathcal{S}|$. In fact, we can just change the definition of M_{ij}^k by changing the value of $C_F, C_F^{\phi,\varphi}, C_H^{\phi,\varphi}$ by their new value.

2.2.5 The Lipschitz constant of \mathcal{F}^-

Now we are going to compute the Lipschitz constant of the operator \mathcal{F}^- in (22).

Lemma 2.6. Take $\rho \geq \rho_*^2$ as in Lemma 2.3. The operator $\mathcal{F} : B(M_0) \rightarrow \mathcal{X}_3 \times \mathcal{X}_3$ is Lipschitz with Lipschitz constant $L(\rho) = \min\{L_1(\rho), L_2(\rho)\}$ with

$$\begin{aligned} L_1(\rho) &= C_4 \frac{M_{11}^1(\rho)}{\rho} + B_6 \frac{M_{11}^2(\rho) + M_{12}^2(\rho)}{\rho} + B_7 \frac{M_{11}^3(\rho) + M_{12}^3(\rho)}{\rho^2} \\ &\quad + B_8 \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{\rho^3} \\ L_2(\rho) &= C_4 \frac{M_{11}^1(\rho)}{\rho} + C_5 \frac{M_{11}^2(\rho) + M_{12}^2(\rho)}{\rho^2} + C_6 \frac{M_{11}^3(\rho) + M_{12}^3(\rho)}{\rho^3} \\ &\quad + C_7 \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{\rho^4}. \end{aligned} \tag{48}$$

where B_ν and C_ν are the constants introduced in (37) and (38) respectively.

Proof. We apply the second item of Lemma 2.1 to $\Delta\mathcal{S}_1(\psi, s) - \Delta\mathcal{S}_1(\psi', s)$ and we obtain that

$$\|\mathcal{B}^-(\Delta\mathcal{S}_1(\psi, s) - \Delta\mathcal{S}_1(\psi', s))\|_4 \leq C_4 M_{11}^1(\rho) \|\phi - \phi'\|_3. \quad (49)$$

Now we apply the first item of Lemma 2.1 to $\Delta\mathcal{S}_j(\psi, s) - \Delta\mathcal{S}_j(\psi', s)$ and we obtain

$$\|\mathcal{B}^-(\Delta\mathcal{S}_j(\psi, s) - \Delta\mathcal{S}_j(\psi', s))\|_{2+j} \leq B_{j+4}(M_{11}^j(\rho) + M_{12}^j(\rho)) \|\phi - \phi'\|_3. \quad (50)$$

Then, we get $L_1(\rho)$ adding the results in (49) and (50). Furthermore, applying the second item of Corollary 2.1, we obtain $L_2(\rho)$ using that

$$\|\mathcal{B}^-(\Delta\mathcal{S}_j(\psi, s) - \Delta\mathcal{S}_j(\psi', s))\|_{3+j} \leq C_{j+2}(M_{11}^j(\rho) + M_{12}^j(\rho)) \|\phi - \phi'\|_3,$$

□

Remark 2.7. Notice that B_m is decreasing with respect to m , but C_m is increasing. It is not difficult to check that when $\rho \geq C_7/B_8 \geq 3^9 \cdot 32/(2^{12} \cdot 5\pi) \sim 9.7895$ then $L_1(\rho) \geq L_2(\rho)$. This fact will be used in Section 3.1 and 3.2.

2.2.6 Setting ρ_* for the existence result

We choose $\rho_* \geq \rho_*^2$ satisfying

$$L(\rho_*) \leq \frac{1}{2},$$

so that Lemma 2.3 can be applied for $\rho \geq \rho_*$. Then, the operator $\mathcal{F}^- : B(M_0) \rightarrow B(M_0)$ is contractive. Indeed,

$$\|\mathcal{F}^-(\psi)\|_3 \leq \|\mathcal{F}^-(0)\|_3 + \|\mathcal{F}^-(\psi) - \mathcal{F}^-(0)\|_3 = \frac{M_0}{2} + LM_0 \leq M_0$$

provided $L \leq \frac{1}{2}$. Therefore, the operator has a fixed point ψ^- defined in $\mathcal{D}_{\rho_*}^-$ (see (15)) and therefore satisfies

$$|\phi^-(s)|, |\varphi^-(s)| \leq \frac{M_0}{|s|^3}. \quad (51)$$

2.2.7 Explicit bounds for the norm of the linear operator \mathcal{G}

The next lemma gives estimates for the linear operator \mathcal{G} defined in (30) with respect to the norm introduced in (33).

Lemma 2.8. Take $\rho \geq \rho_*$ and let

$$\begin{aligned} A_1(\rho) &= \frac{M_{11}^2}{\rho} + \frac{M_{11}^3 + M_{12}^2}{2\rho^2} + \frac{M_{11}^4 + M_{12}^3}{3\rho^3} + \frac{M_{12}^4}{4\rho^4} \\ A_2(\rho) &= \frac{M_{12}^2}{2\alpha\rho^2} + \frac{M_{11}^2 + M_{12}^3}{2\alpha\rho^3} + \frac{M_{11}^3 + M_{12}^4}{2\alpha\rho^4} + \frac{M_{11}^4}{2\alpha\rho^5} \end{aligned} \quad (52)$$

where $M_{ij} = M_{ij}(\rho)$ are the constants introduced in (46). Then, we have that, for s with $\Re s = 0$ and $\Im s \leq -\rho$,

$$\begin{aligned} \left| s^{-1} e^{i\alpha(s+h_0 \log s)} \mathcal{G}_1(\Delta\psi) \right| &\leq A_1(\rho) \|\Delta\psi\| \\ \left| e^{i\alpha(s+h_0 \log s)} \mathcal{G}_2(\Delta\psi) \right| &\leq A_2(\rho) \|\Delta\psi\|. \end{aligned} \quad (53)$$

In particular,

$$\|\mathcal{G}(\Delta\psi)\| \leq A(\rho)\|\Delta\psi\|, \quad (54)$$

with

$$A = A(\rho) = \max\{A_1(\rho), A_2(\rho)\}. \quad (55)$$

Proof. In this proof we omit the dependence on ρ of $M_{i,j}^k$. We use Lemma 2.3 to bound $\tilde{\mathcal{K}}_{ij}$, the components of the matrix $\tilde{\mathcal{K}}$ in (28). By construction, if $\psi \in B(M_0)$,

$$\begin{aligned} \left| \tilde{\mathcal{K}}_{11}(\psi, s) \right|, \left| \tilde{\mathcal{K}}_{22}(\psi, s) \right| &\leq \frac{M_{11}^2}{|s|^2} + \frac{M_{11}^3}{|s|^3} + \frac{M_{11}^4}{|s|^4} \\ \left| \tilde{\mathcal{K}}_{12}(\psi, s) \right|, \left| \tilde{\mathcal{K}}_{21}(\psi, s) \right| &\leq \frac{M_{12}^2}{|s|^2} + \frac{M_{12}^3}{|s|^3} + \frac{M_{12}^4}{|s|^4}. \end{aligned}$$

Then, for the first component,

$$\begin{aligned} \left| s^{-1} e^{i\alpha(s+h_0 \log s)} \mathcal{G}_1(\Delta\psi) \right| &\leq \left| \int_{-i\rho}^s \frac{e^{i\alpha(t+h_0 \log t)}}{t} \left(\tilde{\mathcal{K}}_{11} \Delta\phi + \tilde{\mathcal{K}}_{12} \Delta\varphi \right) dt \right| \\ &\leq \int_{-i\rho}^s \left(\frac{M_{11}^2}{|t|^2} + \frac{M_{11}^3}{|t|^3} + \frac{M_{11}^4}{|t|^4} \right) \|\Delta\phi\| dt \\ &\quad + \int_{-i\rho}^s \left(\frac{M_{12}^2}{|t|^3} + \frac{M_{12}^3}{|t|^4} + \frac{M_{12}^4}{|t|^5} \right) \|\Delta\varphi\| dt \\ &\leq \left[\frac{M_{11}^2}{\rho} + \frac{M_{11}^3}{2\rho^2} + \frac{M_{11}^4}{3\rho^3} \right] \|\Delta\phi\| + \left[\frac{M_{12}^2}{2\rho^2} + \frac{M_{12}^3}{3\rho^3} + \frac{M_{12}^4}{4\rho^4} \right] \|\Delta\varphi\| \\ &\leq \left(\frac{M_{11}^2}{\rho} + \frac{M_{11}^3}{2\rho^2} + \frac{M_{12}^2}{2\rho^2} + \frac{M_{11}^4}{3\rho^3} + \frac{M_{12}^3}{3\rho^3} + \frac{M_{12}^4}{4\rho^4} \right) \|\Delta\psi\|. \end{aligned} \quad (56)$$

For the second component, using that $|e^{i\alpha h_0 \log t}| = e^{\alpha h_0 \pi/2}$,

$$\begin{aligned} \left| e^{i\alpha(s+h_0 \log s)} \mathcal{G}_2(\Delta\psi) \right| &\leq \left| s e^{2i\alpha(s+h_0 \log s)} \int_{-\infty}^s \frac{e^{-i\alpha(t+h_0 \log t)}}{t} \left(\tilde{\mathcal{K}}_{21} \Delta\phi + \tilde{\mathcal{K}}_{22} \Delta\varphi \right) dt \right| \\ &\leq |s| e^{-2\alpha \Im s} \int_{-\infty}^s e^{2\alpha \Re t} \left(\left[\frac{M_{12}^2}{|t|^2} + \frac{M_{12}^3}{|t|^3} + \frac{M_{12}^4}{|t|^4} \right] \|\Delta\phi\| \right) dt \\ &\quad + |s| e^{-2\alpha \Im s} \int_{-\infty}^s e^{2\alpha \Re t} \left(\frac{M_{11}^2}{|t|^3} + \frac{M_{11}^3}{|t|^4} + \frac{M_{11}^4}{|t|^5} \|\Delta\varphi\| \right) dt \\ &\leq \left[\frac{M_{12}^2}{2\alpha\rho^2} + \frac{M_{12}^3}{2\alpha\rho^3} + \frac{M_{12}^4}{2\alpha\rho^4} \right] \|\Delta\phi\| + \left[\frac{M_{11}^2}{2\alpha\rho^3} + \frac{M_{11}^3}{2\alpha\rho^4} + \frac{M_{11}^4}{2\alpha\rho^5} \right] \|\Delta\varphi\| \\ &\leq \left(\frac{M_{12}^2}{2\alpha\rho^2} + \frac{M_{11}^2}{2\alpha\rho^3} + \frac{M_{12}^3}{2\alpha\rho^3} + \frac{M_{11}^3}{2\alpha\rho^4} + \frac{M_{12}^4}{2\alpha\rho^4} + \frac{M_{11}^4}{2\alpha\rho^5} \right) \|\Delta\psi\|. \end{aligned}$$

and the result is proven. \square

2.2.8 Computation of the Stokes constant

Using the estimates of the operator \mathcal{G} given in (54) we can provide a rigorous computation of the Stokes constant.

Let $\rho_0 \geq \rho_*$ be such that

$$A(\rho_0) < \frac{1}{2}.$$

Then, the constant $\overline{M}(\rho_0)$ defined in (36) satisfies

$$\overline{M}(\rho_0) = \frac{A_1(\rho_0)}{1 - A(\rho_0)} < \frac{A(\rho_0)}{1 - A(\rho_0)} < 1 \quad (57)$$

and, as a consequence,

$$\Theta \in [\kappa_0(\rho_0)(1 - \overline{M}(\rho_0)), \kappa_0(\rho_0)(1 + \overline{M}(\rho_0))].$$

In the next section we give the precise algorithm which allows, by means of computer rigorous computations, to compute Θ with a previous known accuracy. This algorithm is applied to two concrete examples in Sections 3.1 and 3.2.

2.3 Computing the Stokes constant: algorithm

We describe the steps needed to obtain the values of ρ_* and ρ_0 which guarantees that ψ^+, ψ^- are defined for $s \in (-i\rho_*, -i\infty)$ and a good accuracy of Θ .

- Step 1: Compute the constants a_3, h_0 and $C_F^0, C_H^0, \overline{C}_H^0, C_F^{\phi, \varphi}, C_H^{\phi, \varphi}$ which satisfy (39) and (40) and $\overline{C}_H, \overline{C}_F$ given in (41).
- Step 2: Take $\rho_*^1 \geq \max\{2, \overline{C}_H^0\}$ and compute, for $\rho \geq \rho_*^1$, the constants $\mathcal{C}_0(\rho)$ given in Lemma 2.2 and $M_0(\rho)$ defined in (44).
- Step 3: Choose $\rho_*^2 \geq \rho_*^1$ such that (45) is satisfied. Compute also the constants $M_{11}^j(\rho), j = 1, 2, 3, 4$ and $M_{12}^j(\rho), j = 2, 3, 4$, in (46), for $\rho \geq \rho_*^2$.
- Step 4: Compute the constants $L_1(\rho)$ and $L_2(\rho)$ in (48) for $\rho \geq \rho_*^2$.
- Step 5: Choose $\rho_* \geq \rho_*^2$ satisfying

$$L(\rho_*) = \min\{L_1(\rho_*), L_2(\rho_*)\} \leq \frac{1}{2}.$$

- Step 6: Take ρ_* and check that the difference

$$\psi^+(-i\rho_*) - \psi^-(-i\rho_*) \neq 0.$$

- Step 7: For $\rho \geq \rho_*$, compute the constants $A_1(\rho)$ and $A_2(\rho)$ in (52) and $A(\rho)$ in (55).
- Step 8: Compute $\rho_0 \geq \rho_*$ such that $A(\rho_0) \leq 1/2$. Then, compute $\kappa_0(\rho_0)$ in (31) and $\overline{M}(\rho_0)$ in (57).

Therefore, the Stokes constant satisfies

$$\Theta \in [\kappa_0(\rho_0)(1 - \overline{M}(\rho_0)), \kappa_0(\rho_0)(1 + \overline{M}(\rho_0))]. \quad (58)$$

Remark 2.9. *By Theorem 1.6, the first 6 steps allows us to check whether $\Theta \neq 0$ or not.*

3 Examples

To illustrate the algorithm we consider two concrete examples of analytic unfoldings of a Hopf-Zero singularity (7) whose corresponding inner equation can be found in (9). In both cases, we prove that the associated constants Θ does not vanish and give rigorous estimates for them.

3.1 The first example

As first example we take

$$\alpha = 1, \quad b = 1, \quad g = h = 0 \quad \text{and} \quad f(X, Y, Z, \delta) = Z^3. \quad (59)$$

This corresponds to $F_1(\phi, \varphi, s) = -s^{-3}$, $F_2 = F_1$ and $H = 0$. The inner equation (14) associated to this model is the following

$$\begin{pmatrix} \frac{d\phi}{ds} \\ \frac{d\varphi}{ds} \end{pmatrix} = \mathcal{A}(s) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} + \mathcal{S}(\phi, \varphi, s) \quad (60)$$

with

$$\mathcal{S}(\phi, \varphi, s) = \begin{pmatrix} \mathcal{S}_1(\phi, \varphi, s) \\ \mathcal{S}_2(\phi, \varphi, s) \end{pmatrix} = \begin{pmatrix} \frac{(i - \frac{1}{s}) \varphi \phi^2 s^2 - \frac{1}{s^3}}{1 + \varphi \phi s^2} \\ \frac{(i + \frac{1}{s}) \varphi^2 \phi s^2 - \frac{1}{s^3}}{1 + \varphi \phi s^2} \end{pmatrix} \quad (61)$$

Now we follow the steps of the algorithm in Section 2.3.

Step 1. In this case $h_0 = 0$ and moreover, among all the constants defined in Step 1, the only one that is different from 0 is $a_3 = 1$.

Step 2. In this case we have that $\rho_*^1 = 2$ and $\mathcal{C}_0 = 0$ so that $M_0 = \frac{22}{3}$ is independent on ρ .

Step 3. We have that ρ_*^2 has to be such that

$$\sqrt{M_0} = \sqrt{\frac{22}{3}} < \rho_*^2.$$

In addition $M_{ij}^{1,2,3} = 0$ and

$$\begin{aligned} M_{11}^4(\rho) &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(1 + M_0 + \left(M_0 + \frac{M_0}{\rho}\right) \left(1 - \frac{M_0^2}{\rho^4}\right) + \frac{M_0}{\rho}\right) \\ &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(1 + M_0 \left(1 + \frac{1}{\rho}\right) \left(2 - \frac{M_0^2}{\rho^4}\right)\right) \\ M_{12}^4(\rho) &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(1 + M_0 \left(1 + \frac{1}{\rho}\right)\right). \end{aligned}$$

Step 4 and Step 5. One can check that

$$L_1(\rho) = B_8 \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{\rho^3} \leq \frac{1}{2}$$

for $\rho \geq \rho_* = 9.7895$. Under this condition, as we claimed in Remark 2.7,

$$L(\rho) \leq \frac{1}{2}.$$

Therefore we can guarantee the existence of ψ^\pm for $\rho \geq \rho_* = 9.7895$.

Step 6. Now it only remains to compute

$$\psi^+(-i\rho_*) - \psi^-(-i\rho_*) \neq 0.$$

By means of rigorous computer computations, which are discussed in more detail in Appendix B, we obtain that there exists a

$$\rho_* \in [15.99999965, 16.00000035] \quad (62)$$

for which

$$\begin{aligned} & \psi^+(-i\rho_*) - \psi^-(-i\rho_*) \quad (63) \\ &= \begin{pmatrix} \Delta\phi(-i\rho_*) \\ \Delta\varphi(-i\rho_*) \end{pmatrix} \\ &\in \begin{pmatrix} [-4.50096 \cdot 10^{-10}, 4.50096 \cdot 10^{-10}] - [1.88812 \cdot 10^{-6}, 1.88897 \cdot 10^{-6}]i \\ [-3.85539 \cdot 10^{-10}, 3.85539 \cdot 10^{-10}] + [-4.01544 \cdot 10^{-10}, 3.4832 \cdot 10^{-10}]i \end{pmatrix}. \end{aligned}$$

Therefore, the Stokes constant associated to the first example (59) does not vanish.

Now we follow **Step 7.** and **Step 8.** to provide rigorous accurate estimates for it.

Step 7. The constants A_1 and A_2 in (52) are

$$A_1(\rho) = \frac{M_{11}^4(\rho)}{3\rho^3} + \frac{M_{12}^4(\rho)}{4\rho^4}, \quad A_2(\rho) = \frac{M_{12}^4(\rho)}{2\rho^4} + \frac{M_{11}^4(\rho)}{2\rho^5} \quad (64)$$

which give the constant $A(\rho) = \max\{A_1(\rho), A_2(\rho)\}$ in (55). We obtain

$$\begin{aligned} A_1(\rho_*) &\in [0.010155523, 0.010155525], \\ A_2(\rho_*) &\in [0.0009597786, 0.0009597788], \\ A(\rho_*) &= A_1(\rho_*) < 1/2. \end{aligned}$$

Step 8. One can choose $\rho_0 = \rho_*$. Then, by (31), (57) and (58), one obtains

$$\Theta \in [1.0378681, 1.0598665] + [-0.000253, 0.000253]i \quad (65)$$

We can see that the accuracy of the computation is roughly $2 \cdot 10^{-2}$.

3.2 The second example

The second example, breaks the reversibility. It consists in taking $\alpha = b = 1$, $g = h = 0$ and $f(X, Y, Z, \delta) = Z^3 + 2XYZ$ which corresponds to

$$F_1(\phi, \varphi, s) = F_2(\phi, \varphi, s) = -\frac{1}{s^3} + \frac{i}{s}(\phi^2 - \varphi^2), \quad H = 0$$

and then, the inner equation associated to this unfolding is:

$$\begin{pmatrix} \frac{d\phi}{ds} \\ \frac{d\varphi}{ds} \end{pmatrix} = \mathcal{A}(s) \begin{pmatrix} \phi \\ \varphi \end{pmatrix} + \mathcal{S}(\phi, \varphi, s) \quad (66)$$

with \mathcal{S} defined as

$$\mathcal{S}(\phi, \varphi, s) = \begin{pmatrix} \mathcal{S}_1(\phi, \varphi, s) \\ \mathcal{S}_2(\phi, \varphi, s) \end{pmatrix} = \begin{pmatrix} \frac{(i - \frac{1}{s})\varphi\phi^2s^2 - \frac{1}{s^3} + \frac{i}{2s}(\phi^2 - \varphi^2)}{1 + \varphi\phi s^2} \\ \frac{(i + \frac{1}{s})\varphi^2\phi s^2 - \frac{1}{s^3} + \frac{i}{2s}(\phi^2 - \varphi^2)}{1 + \varphi\phi s^2} \end{pmatrix}.$$

Step 1. We have that all the constants are zero except $a_3 = 1$, $C_F = 2$, $C_F^\phi = C_F^\varphi = 1$.

Step 2. As for the first example $\rho_*^1 = 2$ and $\mathcal{C}_0 = 0$ so that $M_0 = \frac{22}{3}$.

Step 3. We have that ρ_*^2 has to be such that

$$\sqrt{M_0} = \sqrt{\frac{22}{3}} < \rho_*^2.$$

In addition $M_{ij}^{1,3}(\rho) = 0$ being

$$\begin{aligned} M_{11}^2(\rho) &= M_{12}^2(\rho) = \frac{2}{1 - \frac{M_0^2}{\rho^4}} \\ M_{11}^4(\rho) &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(2 + M_0 + \left(M_0 + \frac{M_0}{\rho}\right) \left(1 - \frac{M_0^2}{\rho^4}\right) + \frac{M_0}{\rho}\right) \\ &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(2 + M_0 \left(1 + \frac{1}{\rho}\right) \left(2 - \frac{M_0^2}{\rho^4}\right)\right) \\ M_{12}^4(\rho) &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(2 + M_0 \left(1 + \frac{1}{\rho}\right)\right). \end{aligned}$$

Step 4 and Step 5. One can check that

$$L_1(\rho) = B_6 \frac{M_{11}^2(\rho) + M_{12}^2(\rho)}{\rho} + B_8 \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{\rho^3} \leq \frac{1}{2}$$

for $\rho \geq \rho_* \geq 9.7895$. Therefore, under this condition, using Remark 2.7 as for Example 1, we can guarantee that $L(\rho) \geq 1/2$ and, then, the existence of ψ^\pm .

Remark 3.1. For Example 2, we can obtain more accurate estimates by computing directly the derivatives $\partial_{\phi, \varphi} \mathcal{S}$. Indeed, performing straightforward computations we obtain that $M_{ij}^1(\rho) = M_{ij}^2(\rho) = M_{ij}^3(\rho) = 0$ and

$$\begin{aligned} M_{11}^4(\rho) &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(3 + M_0 \left(1 + \frac{1}{\rho}\right) \left(2 - \frac{M_0^2}{\rho^4}\right)\right) \\ M_{12}^4(\rho) &= \frac{M_0}{\left(1 - \frac{M_0^2}{\rho^4}\right)^2} \left(3 + M_0 \left(1 + \frac{1}{\rho}\right)\right). \end{aligned} \quad (67)$$

Therefore

$$L_1(\rho) = B_8 \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{\rho^3} = \frac{5\pi}{32} \frac{M_{11}^4(\rho) + M_{12}^4(\rho)}{\rho^3}.$$

Since we need to be as precise as possible, we use the constants $M_{ij}^k(\rho)$ defined in Remark 3.1 instead of the constants provided by the general method.

Step 6. We compute $\psi^+(-i\rho_*) - \psi^-(-i\rho_*)$. By means of rigorous computer computations we obtain that there exists a ρ_*

$$\rho_* \in [15.99999965, 16.00000035], \quad (68)$$

for which

$$\begin{aligned} & \psi^+(-i\rho_*) - \psi^-(-i\rho_*) \quad (69) \\ &= \begin{pmatrix} \Delta\phi(-i\rho_*) \\ \Delta\varphi(-i\rho_*) \end{pmatrix} \\ & \left(\begin{array}{l} [8.63066 \cdot 10^{-9}, 9.53086 \cdot 10^{-9}] - [1.88812 \cdot 10^{-6}, 1.88897 \cdot 10^{-6}]i \\ [-4.20777 \cdot 10^{-10}, 3.50313 \cdot 10^{-10}] + [-4.01721 \cdot 10^{-10}, 3.48156 \cdot 10^{-10}]i \end{array} \right). \end{aligned}$$

This implies that $\Theta \neq 0$. Now we perform the last two steps in the algorithm.

Step 7. For Example 2, we have that

$$A_1(\rho) = \frac{M_{11}^4(\rho)}{3\rho^3} + \frac{M_{12}^4(\rho)}{4\rho^4}, \quad A_2(\rho) = \frac{M_{12}^4(\rho)}{2\rho^4} + \frac{M_{11}^4(\rho)}{2\rho^5},$$

and $A(\rho) = \max\{A_1(\rho), A_2(\rho)\}$, with $M_{11}^4(\rho), M_{12}^4(\rho)$ defined in (67). We obtain

$$\begin{aligned} A_1(\rho_*) &\in [0.0114071016, 0.0114071033], \\ A_2(\rho_*) &\in [0.00066984056, 0.00066984069], \\ A(\rho_*) &= A_1(\rho_*) < 1/2. \end{aligned}$$

Step 8. By means of rigorous computer validation, for $\rho_0 = \rho_*$ using (31), (57) and (58) we obtain

$$\Theta \in [1.036525, 1.0612062] + [0.004738, 0.005355]i.$$

We can see that the accuracy of the computation is roughly $2.5 \cdot 10^{-1}$.

4 Improving the computation of the Stokes constant

In this section we give an improvement of the Steps 7 and 8 in Section 2.3 to obtain accurate estimates for the Stokes constant Θ . We explain this improvement for the Example 1 given in Section 3.1 but the method we present is general and can be applied to any system.

Recall that, using (18) and (29),

$$\begin{aligned} \Theta &= \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha s} \Delta\phi(s) = \kappa_0 + \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha s} \mathcal{G}_1(\Delta\psi(s)) \\ &= \kappa_0 + \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha s} \mathcal{G}_1(\Delta\psi_0(s)) + \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha s} \mathcal{G}_1(\mathcal{G}(\Delta\psi)(s)). \end{aligned}$$

Therefore, by (53), (54) and (35), the remainder

$$\mathcal{E}_\Theta = \Theta - \kappa_0 - \lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha s} \mathcal{G}_1(\Delta\psi_0(s))$$

satisfies

$$\begin{aligned} |\mathcal{E}_\Theta| &\leq \sup_s |s^{-1} e^{i\alpha s} \mathcal{G}_1(\mathcal{G}(\Delta\psi)(s))| \leq A_1 \|\mathcal{G}(\Delta\psi)\| \leq A_1 A \|\Delta\psi\| \\ &\leq A_1 A \frac{|\kappa_0|}{1-A}. \end{aligned}$$

where $A = A(\rho) = \max\{A_1, A_2\}$ and A_1, A_2 are given in (64).

Using (30) and the fact that in Example 1, $h_0 = 0$ and therefore $\tilde{\mathcal{K}} = \mathcal{K}$,

$$\lim_{\Im s \rightarrow -\infty} s^{-1} e^{i\alpha s} \mathcal{G}_1(\Delta\psi_0(s)) = -\kappa_0 \int_{-i\infty}^{i\rho} \mathcal{K}_{11}(t) dt = -i\kappa_0 \int_{-\infty}^{\rho} \mathcal{K}_{11}(ir) dr, \quad (70)$$

which implies

$$\Theta = \kappa_0 - i\kappa_0 \int_{-\infty}^{\rho} \mathcal{K}_{11}(ir) dr + \mathcal{E}_\Theta. \quad (71)$$

To obtain this integral we need an approximation of the coefficient \mathcal{K}_{11} .

Lemma 4.1. *The function \mathcal{K}_{11} introduced in (25) associated to equation (60) satisfies*

$$\begin{aligned} \mathcal{K}_{11}(s) &= \frac{i\beta_1}{s^4} + \frac{\beta_2}{s^5} + \frac{i\beta_3}{s^6} + \frac{\beta_4}{s^7} + \mathcal{E}\mathcal{K}\mathcal{T}_{11} \\ \beta_1 &= 3, \quad \beta_2 = -6, \quad \beta_3 = -68, \quad \beta_4 = 48 \end{aligned} \quad (72)$$

and $\mathcal{E}\mathcal{K}\mathcal{T}_{11}$ satisfies

$$|\mathcal{E}\mathcal{K}\mathcal{T}_{11}| \leq \frac{B\mathcal{R}}{|s|^7} + \frac{B_{11} + B_{12} + B_{13} + B_{14}}{|s|^8} \quad (73)$$

where

$$\begin{aligned} B &= \frac{5\pi}{32} (M_{11}^4(\rho) + M_{12}^4(\rho)) M_0 + \frac{225\pi}{2} \\ \mathcal{R} &= \frac{1 + 4M_0(1 + \frac{1}{\rho}) + 3\frac{M_0^2}{\rho^4}}{\left(1 - \frac{M_0^2}{\rho^4}\right)^3} \\ B_{11} &= \frac{M_*^4(1 + \frac{1}{\rho})}{\left(1 - \frac{M_*^2}{\rho^4}\right)^2} \\ B_{12} &= M_*^5 \frac{(2M_*(1 + \frac{1}{\rho}) + 1)(3 + \frac{2M_*^2}{\rho^4})}{\rho^4 \left(1 - \frac{M_*^2}{\rho^4}\right)^2} \\ B_{13} &= 2M_*^3 \left(2M_* \left(1 + \frac{1}{\rho}\right) + 1\right) \\ B_{14} &= 800 \left(1 - \frac{1}{\rho}\right) \end{aligned}$$

and

$$M_*(\rho) = 1 + \frac{4}{\rho} + \frac{20}{\rho^2} + \frac{120}{\rho^3}.$$

This lemma is proven in Section 4.1.

Now take ρ_\diamond be such that $M_*(\rho) \leq M_0(\rho)$ for $\rho \geq \rho_\diamond$. Using the expression of \mathcal{K}_{11} given in (72) to compute (70) and using the remainder estimates in (73) we obtain:

$$\begin{aligned} -i\kappa_0 \int_{-\infty}^{\rho} \mathcal{K}_{11}(ir)dr &= \kappa_0 \int_{-\infty}^{\rho} \frac{\beta_1}{r^4} - \frac{\beta_2}{r^5} - \frac{\beta_3}{r^6} + \frac{\beta_4}{r^7} dr + \mathcal{E}\mathcal{S} \\ &= \kappa_0 \left(-\frac{\beta_1}{3\rho^3} + \frac{\beta_2}{4\rho^4} - \frac{\beta_3}{5\rho^5} - \frac{\beta_4}{6\rho^6} \right) + \mathcal{E}\mathcal{S} \end{aligned} \quad (74)$$

and

$$\begin{aligned} |\mathcal{E}\mathcal{S}| = |\mathcal{E}\mathcal{S}(\rho)| &\leq \kappa_0 \int_{\rho}^{\infty} \frac{B\mathcal{R}}{r^7} + \frac{B_{11} + B_{12} + B_{13} + B_{14}}{r^8} \\ &= \frac{B\mathcal{R}}{6\rho^6} + \frac{B_{11} + B_{12} + B_{13} + B_{14}}{7\rho^7}. \end{aligned}$$

Finally, using the expression (71) for Θ we obtain

$$\begin{aligned} \Theta &= \kappa_0 \left(1 - \frac{\beta_1}{3\rho^3} + \frac{\beta_2}{4\rho^4} - \frac{\beta_3}{5\rho^5} - \frac{\beta_4}{6\rho^6} \right) + \mathcal{E}\mathcal{T}, \\ \mathcal{E}\mathcal{T} &= \mathcal{E}\mathcal{T}(\rho) = \mathcal{E}\mathcal{S} + \mathcal{E}_\Theta. \end{aligned}$$

As we know the constants β_i , one can use this formula to improve the computation of Θ .

Indeed, using the above approach one obtains

$$\Theta \in [1.047906, 1.049289] + [-0.00070294, 0.00070294]i. \quad (75)$$

We can see that the accuracy of the computation is roughly 10^{-3} , which is an improvement when compared to the accuracy $2 \cdot 10^{-1}$ from (65). (For (75) we have used the same ρ_* and the computed value of $\Delta\psi(-i\rho_*)$ as for (65).)

4.1 Proof of Lemma 4.1

Lets call $\mathcal{K}_{ij}(s)$ the 4 elements of the matrix \mathcal{K} . To obtain expansions for these coefficients we compute first an expansion for ψ^\pm associated to Example 1 in (59).

Lemma 4.2. *The functions ψ^\pm can be written as*

$$\psi^\pm = \psi_* + \mathcal{E}^\pm$$

where $\psi_* = (\phi_*, \varphi_*)$ with

$$\begin{aligned} \phi_* &= -\frac{i}{s^3} - \frac{4}{s^4} + \frac{20i}{s^5} + \frac{120}{s^6} \\ \varphi_* &= \frac{i}{s^3} - \frac{4}{s^4} - \frac{20i}{s^5} + \frac{120}{s^6}, \end{aligned}$$

which satisfy

$$|\phi_*(s)|, |\varphi_*(s)| \leq \frac{M_*(\rho)}{|s|^3} \quad (76)$$

for

$$M_*(\rho) = 1 + \frac{4}{\rho} + \frac{20}{\rho^2} + \frac{120}{\rho^3} \quad (77)$$

and the remainders $\mathcal{E}^\pm = (\mathcal{E}_\phi^\pm, \mathcal{E}_\varphi^\pm)$ satisfy

$$|\mathcal{E}_\phi^\pm|, |\mathcal{E}_\varphi^\pm| \leq \frac{B}{|s|^6}. \quad (78)$$

with

$$B = \frac{5\pi}{32} (M_{11}^4(\rho) + M_{12}^4(\rho)) M_0 + \frac{225\pi}{2}.$$

Proof. By (61), the first iteration $\mathcal{F}^-(0)$ analyzed in Lemma 2.2 for Example 1 is given by

$$\begin{aligned} \mathcal{F}_1^-(0) &= s \int_{-\infty}^0 \frac{1}{(s+t)^4} e^{it} dt \\ \mathcal{F}_2^-(0) &= s \int_{-\infty}^0 \frac{1}{(s+t)^4} e^{-it} dt \end{aligned}$$

Integrating by parts, we obtain

$$\begin{aligned} \mathcal{F}_1^-(0) &= -\frac{i}{s^3} + \frac{4s}{i} \int_{-\infty}^0 \frac{1}{(s+t)^5} e^{it} dt \\ &= -\frac{i}{s^3} - \frac{4}{s^4} + \frac{20i}{s^5} + \frac{120}{s^6} + E_1^-(s) = \phi_*(s) + E_1^-(s), \end{aligned}$$

where

$$|E_1^-(s)| = \left| 720s \int_{-\infty}^0 \frac{1}{(s+t)^8} e^{it} dt \right| \leq \frac{720}{|s|^6} \int_0^\infty \frac{1}{(t^2+1)^4} dr = \frac{720}{|s|^6} \frac{5\pi}{32} = \frac{225\pi}{2|s|^6}.$$

Analogously, for the second component

$$\mathcal{F}_2^-(0) = \frac{i}{s^3} - \frac{4}{s^4} + \frac{20i}{s^5} + \frac{120}{s^6} + E_2^-(s) = \varphi_*(s) + E_2^-(s),$$

where E_2^- has the same bounds as E_1^- :

$$|E_2^-(s)| \leq \frac{225\pi}{2|s|^6}.$$

Let us call $\psi_* = (\phi_*, \varphi_*)$ and $E^- = (E_1^-, E_2^-)$. Observe that by Lemma 2.1 and Corollary 2.4 and recalling that, for Example 1, one has $M_{11}^j = M_{12}^j = 0$ for $j = 1, 2, 3$,

$$\begin{aligned} \|\mathcal{F}^-(\psi) - \mathcal{F}^-(0)\|_6 &\leq \frac{5\pi}{32} \|\mathcal{S}(\psi, s) - \mathcal{S}(0, s)\|_7 \\ &\leq \frac{5\pi}{32} (M_{11}^4(\rho) + M_{12}^4(\rho)) \|\psi\|_3. \end{aligned}$$

Now we use that ψ^- is a fixed point of operator \mathcal{F}^- and therefore

$$\begin{aligned} \|\psi^- - \psi_*\|_6 &\leq \|\psi^- - \mathcal{F}^-(0)\|_6 + \|E^-(0)\|_6 \\ &\leq \frac{5\pi}{32} (M_{11}^4(\rho) + M_{12}^4(\rho)) M_0 + \frac{225\pi}{2} = B. \end{aligned}$$

We conclude

$$\begin{aligned} \left| \phi^- - \left(-\frac{i}{s^3} - \frac{4}{s^4} - \frac{20i}{s^5} + \frac{120}{s^6} \right) \right| &\leq \frac{B}{|s|^6} \\ \left| \varphi^- - \left(\frac{i}{s^3} - \frac{4}{s^4} + \frac{20i}{s^5} + \frac{120}{s^6} \right) \right| &\leq \frac{B}{|s|^6}. \end{aligned}$$

□

Using the previous result, we can compute a better asymptotic expansion of \mathcal{K}_{11} . We rely on the expression

$$\mathcal{K}_{11}(s) = \int_0^1 \partial_\phi \mathcal{S}_1(\psi^-(s) + t(\psi^+(s) - \psi^-(s))) dt.$$

Note that, for Example 1,

$$\partial_\phi \mathcal{S}_1 = \frac{2A\varphi\phi + B\varphi^2\phi^2 + C\varphi}{(1 + D\varphi\phi)^2}$$

where

$$A = \left(i - \frac{1}{s} \right) s^2, \quad B = s^4 \left(i - \frac{1}{s} \right), \quad C = \frac{1}{s}, \quad D = s^2.$$

We define the function

$$g(r) = \partial_\phi \mathcal{S}_1(\psi_*(s) + r(\mathcal{E}^-(s) + t(\mathcal{E}^+(s) - \mathcal{E}^-(s))))$$

and using the fundamental theorem of calculus, $g(1) = g(0) + \int_0^1 g'(r) dr$, we have that

$$\begin{aligned} \partial_\phi \mathcal{S}_1(\psi^-(s) + t(\psi^+(s) - \psi^-(s))) &= \partial_\phi \mathcal{S}_1(\psi_*(s)) \\ &+ \left(\mathcal{E}_\phi^-(s) + t(\mathcal{E}_\phi^+(s) - \mathcal{E}_\phi^-(s)) \right) \int_0^1 \partial_{\phi\phi} \mathcal{S}_1(\psi_*(s) + r(\mathcal{E}^-(s) + t(\mathcal{E}^+(s) - \mathcal{E}^-(s)))) dr \\ &+ \left(\mathcal{E}_\varphi^-(s) + t(\mathcal{E}_\varphi^+(s) - \mathcal{E}_\varphi^-(s)) \right) \int_0^1 \partial_{\phi\varphi} \mathcal{S}_1(\psi_*(s) + r(\mathcal{E}^-(s) + t(\mathcal{E}^+(s) - \mathcal{E}^-(s)))) dr. \end{aligned}$$

Therefore,

$$\begin{aligned} \mathcal{K}_{11} &= \partial_\phi \mathcal{S}_1(\psi_*(s)) \\ &+ \int_0^1 \left[\left(\mathcal{E}_\phi^-(s) + t(\mathcal{E}_\phi^+(s) - \mathcal{E}_\phi^-(s)) \right) \int_0^1 \partial_{\phi\phi} \mathcal{S}_1(\psi_*(s) + r(\mathcal{E}^-(s) + t(\mathcal{E}^+(s) - \mathcal{E}^-(s)))) dr \right] dt \\ &+ \int_0^1 \left[\left(\mathcal{E}_\varphi^-(s) + t(\mathcal{E}_\varphi^+(s) - \mathcal{E}_\varphi^-(s)) \right) \int_0^1 \partial_{\phi\varphi} \mathcal{S}_1(\psi_*(s) + r(\mathcal{E}^-(s) + t(\mathcal{E}^+(s) - \mathcal{E}^-(s)))) dr \right] dt. \end{aligned} \tag{79}$$

One can easily check that

$$\partial_{\phi\phi} \mathcal{S}_1(\psi) = \frac{2A\varphi - 2DC\varphi^2}{(1 + D\varphi\phi)^3}, \quad \partial_{\phi\varphi} \mathcal{S}_1(\psi) = \frac{C + 2A\phi - DC\varphi\phi}{(1 + D\varphi\phi)^3}.$$

Moreover, by (51) and (76), we know that

$$|\psi_*(s) + r(\mathcal{E}^-(s) + t(\mathcal{E}^+(s) - \mathcal{E}^-(s)))| \leq \frac{M_0}{|s|^3}.$$

Then, taking into account the definitions of $\mathbf{A}, \mathbf{C}, \mathbf{D}$, one can obtain the following bounds for $|s| \geq \rho$,

$$\begin{aligned} |\partial_{\phi\phi}\mathcal{S}_1(\psi)| &\leq \frac{1}{s} \frac{2M_0(1 + \frac{1}{\rho}) + 2\frac{M_0^2}{\rho^4}}{\left(1 - \frac{M_0^2}{\rho^4}\right)^3} \\ |\partial_{\phi\varphi}\mathcal{S}_1(\psi)| &\leq \frac{1}{s} \frac{1 + 2M_0(1 + \frac{1}{\rho}) + \frac{M_0^2}{\rho^4}}{\left(1 - \frac{M_0^2}{\rho^4}\right)^3}. \end{aligned}$$

Using (79) and the bounds (78), we obtain that \mathcal{K}_{11} satisfies

$$\mathcal{K}_{11}(s) = \partial_{\phi}\mathcal{S}_1(\psi_*(s)) + \mathcal{E}\mathcal{K}_{11} \quad (80)$$

with

$$|\mathcal{E}\mathcal{K}_{11}| \leq \frac{B}{|s|^7} \mathcal{R} \quad \text{where} \quad \mathcal{R} = \frac{1 + 4M_0(1 + \frac{1}{\rho}) + 3\frac{M_0^2}{\rho^4}}{\left(1 - \frac{M_0^2}{\rho^4}\right)^3}. \quad (81)$$

Last step is to compute $\partial_{\phi}\mathcal{S}_1(\psi_*)$ using the formula of ψ_* in Lemma 4.2. We recall that

$$\partial_{\phi}\mathcal{S}_1(\psi_*) = \frac{\varphi_*\phi_*s^2\left(i - \frac{1}{s}\right)\left(2 + \varphi_*\phi_*s^2\right) + \frac{1}{s}\varphi_*}{\left(1 + \varphi_*\phi_*s^2\right)^2}$$

and we write

$$\partial_{\phi}\mathcal{S}_1(\psi_*) = \frac{2\varphi_*\phi_*s^2\left(i - \frac{1}{s}\right) + \frac{1}{s}\varphi_*}{\left(1 + \varphi_*\phi_*s^2\right)^2} + \mathcal{E}\mathcal{R}_1$$

with

$$|\mathcal{E}\mathcal{R}_1| = \left| \frac{\varphi_*\phi_*s^2\left(i - \frac{1}{s}\right)\varphi_*\phi_*s^2}{\left(1 + \varphi_*\phi_*s^2\right)^2} \right| \leq \frac{M_*^4(1 + \frac{1}{\rho})}{\left(1 - \frac{M_*^2}{\rho^4}\right)^2} \frac{1}{|s|^8} = \frac{B_{11}}{|s|^8}. \quad (82)$$

Now, using that

$$\frac{1}{(1+x)^2} = 1 - 2x + \frac{x^2(3+2x)}{1+x^2},$$

we write

$$\begin{aligned} \partial_{\phi}\mathcal{S}_1(\psi_*) &= \left(2\varphi_*\phi_*s^2\left(i - \frac{1}{s}\right) + \frac{1}{s}\varphi_*\right) (1 - 2\varphi_*\phi_*s^2) + \mathcal{E}\mathcal{R}_1 + \mathcal{E}\mathcal{R}_2 \\ &= 2\varphi_*\phi_*s^2\left(i - \frac{1}{s}\right) + \frac{1}{s}\varphi_* + \mathcal{E}\mathcal{R}_1 + \mathcal{E}\mathcal{R}_2 + \mathcal{E}\mathcal{R}_3 \end{aligned}$$

with

$$\begin{aligned}
|\mathcal{ER}_2| &= \left| 2\varphi_*\phi_*s^2 \left(i - \frac{1}{s} \right) + \frac{1}{s}\varphi_* \right| \left| \frac{\varphi_*^2\phi_*^2s^4 (3 + 2\varphi_*\phi_*s^2)}{(1 + \varphi_*\phi_*s^2)^2} \right| \\
&\leq \frac{M_*^5}{|s|^{12}} \frac{(2M_*(1 + \frac{1}{\rho}) + 1)(3 + \frac{2M_*^2}{\rho^4})}{\left(1 - \frac{M_*^2}{\rho^4}\right)^2} \\
&\leq \frac{M_*^5}{|s|^8} \frac{(2M_*(1 + \frac{1}{\rho}) + 1)(3 + \frac{2M_*^2}{\rho^4})}{\rho^4 \left(1 - \frac{M_*^2}{\rho^4}\right)^2} = \frac{B_{12}}{|s|^8} \tag{83} \\
|\mathcal{ER}_3| &= \left| \left(2\varphi_*\phi_*s^2 \left(i - \frac{1}{s} \right) + \frac{1}{s}\varphi_* \right) (-2\varphi_*\phi_*s^2) \right| \\
&\leq \frac{2M_*^3}{|s|^8} \left(2M_* \left(1 + \frac{1}{\rho} \right) + 1 \right) = \frac{B_{13}}{|s|^8}.
\end{aligned}$$

We now substitute the expressions ψ_* in Lemma 4.2 which give

$$\begin{aligned}
\phi_*\varphi_* &= \frac{1}{s^6} - \frac{24}{s^8} + \frac{400}{s^{10}} \\
2\varphi_*\phi_*s^2 \left(i - \frac{1}{s} \right) + \frac{1}{s}\varphi_* &= \frac{3i}{s^4} - \frac{6}{s^5} - \frac{68i}{s^6} + \frac{48}{s^7} + \frac{800i}{s^8} - \frac{800}{s^9}
\end{aligned}$$

which gives

$$\partial_\phi \mathcal{S}_1(\psi_*) = \frac{3i}{s^4} - \frac{6}{s^5} - \frac{68i}{s^6} + \frac{48}{s^7} + \mathcal{ER}_1 + \mathcal{ER}_2 + \mathcal{ER}_3 + \mathcal{ER}_4$$

and

$$|\mathcal{ER}_4| \leq \frac{800}{|s|^8} \left(1 - \frac{1}{\rho} \right) = \frac{B_{14}}{|s|^8}. \tag{84}$$

Using these approximations in (80), we obtain the statement of the lemma taking

$$\mathcal{EK}\mathcal{T}_{11} = \mathcal{EK}_{11} + \mathcal{ER}_1 + \mathcal{ER}_2 + \mathcal{ER}_3 + \mathcal{ER}_4.$$

Using the bounds (81), (82), (83), (84) we get

$$|\mathcal{EK}\mathcal{T}_{11}| \leq \frac{B\mathcal{R}}{|s|^7} + \frac{B_{11} + B_{12} + B_{13} + B_{14}}{|s|^8} \tag{85}$$

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 757802). This work is part of the grant PGC2018-098676-B-100 funded by MCIN/AEI/10.13039/501100011033 and "ERDF A way of making Europe". I.B and T. M. S. has also been partly supported by the and the Catalan grant 2017SGR1049. M. G. and T. M. S. are supported by the Catalan Institution for Research and Advanced Studies via an ICREA Academia Prize

2019. M.C. has been partially supported by the Polish National Science Center (NCN) grants 2019/35/B/ST1/00655 and 2021/41/B/ST1/00407. This work is also supported by the Spanish State Research Agency, through the Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M).

A Proof of Lemma 2.1

To prove Lemma 2.1, we need first the following lemma.

Lemma A.1. *If m is even*

$$\int_{-\infty}^0 \frac{1}{(t^2 + 1)^{m/2}} dt = \frac{\pi (m-3)!!}{2 (m-2)!!} =: B_m$$

and, for m odd

$$\int_{-\infty}^0 \frac{1}{(t^2 + 1)^{m/2}} dt = \frac{(m-3)!!}{(m-2)!!} =: B_m$$

Proof. Integrating by parts,

$$\begin{aligned} I_k &:= \int_{-\infty}^0 \frac{1}{(t^2 + 1)^{k/2}} dt = k \int_{-\infty}^0 \frac{t^2}{(t^2 + 1)^{\frac{k}{2}+1}} \\ &= kI_k - kI_{k+2}. \end{aligned}$$

Therefore $I_{k+2} = \frac{k-1}{k}I_k$ which implies $I_k = \frac{k-3}{k-2}I_{k-2}$ and therefore

$$I_k = \frac{(k-3)!!}{(k-2)!!} J,$$

with $J = I_3$ if k is odd and $J = I_2$ if k is even. Since $I_3 = 1$ and $I_2 = \frac{\pi}{2}$ we are done. \square

We use this lemma to prove Lemma 2.1.

Proof of Lemma 2.1. Let $\psi = (\phi, \varphi) \in \mathcal{X}_\nu$. The first component of $\mathcal{B}^-(\psi)$ is

$$\mathcal{B}_1^-(\psi) = s \int_{-\infty}^0 \frac{e^{i\alpha t}}{s+t} \phi(s+t) dt. \quad (86)$$

We first prove the first item. Indeed, using that $|s+t|^2 \geq |s|^2 + t^2$ for $s \in \mathcal{D}_\rho^-$, one can prove that

$$|\mathcal{B}_1^-(\psi)| \leq |s| \|\phi\|_\nu \int_{-\infty}^0 \frac{1}{|s+t|^{\nu+1}} dt \leq \frac{\|\phi\|_\nu}{|s|^{\nu-1}} \int_{-\infty}^0 \frac{1}{(t^2 + 1)^{\frac{\nu+1}{2}}} = B_{\nu+1} \frac{\|\phi\|_\nu}{|s|^{\nu-1}}.$$

Analogously we deal with $\mathcal{B}_2^-(\psi)$ and we obtain the result in the first item, taking into account that the product norm is the supremum norm.

Now we deal with the second item. By the geometry of \mathcal{D}_ρ^- , and using the Cauchy's theorem, we can change the path of integration in the integral (86) defining $\mathcal{B}_1^-(\psi)$ to $te^{i\gamma}$, $t \in (-\infty, 0]$, with $0 \leq \gamma \leq \beta$. We obtain then

$$\mathcal{B}_1^-(\psi)(s) = s \int_{-\infty}^0 \frac{e^{i\alpha te^{i\gamma}}}{s + te^{i\gamma}} \phi(s + te^{i\gamma}) e^{i\gamma} dt.$$

Notice that $s + te^{i\gamma} \in \mathcal{D}_\rho^-$ and

$$|s + te^{i\gamma}| \geq |s| \sin\left(\frac{\pi}{2} - \gamma\right) = |s| \cos \gamma.$$

Therefore

$$|\mathcal{B}_1^-(\psi)| \leq \frac{\|\phi\|_\nu}{|s|^\nu (\cos \gamma)^{\nu+1}} \int_{-\infty}^0 e^{\alpha \sin \gamma t} dt = \frac{\|\phi\|_\nu}{|s|^\nu \alpha (\cos \gamma)^{\nu+1} \sin \gamma}.$$

The function $(\cos \gamma)^{\nu+1} \sin \gamma$ has only a maximum in $(0, \frac{\pi}{2})$ in γ_* such that $(\nu + 1) \sin^2 \gamma_* = 1$.

As in the first item, $\mathcal{B}_2^-(\psi)$ can be treated in the same way, changing here the integration path to $te^{-i\gamma}$. \square

B Computing the bound on $\Delta\psi(-i\rho^*)$

Here we provide an explicit rigorous estimate, using interval arithmetic bounds, for the distance between $\psi^+(-i\rho)$ and $\psi^-(-i\rho)$ for a given $\rho > 0$ which we have used for our Examples 1 and 2 (discussed in Sections 3.1, 3.2 and 4).

We start with Example 1. We work with the system (10) with $F_1 = -s^{-3}$, $F_2 = 0$, $H = 0$.

By writing

$$\phi = x_1 + iy_1, \quad \varphi = x_2 + iy_2, \quad s = s_1 + is_2,$$

we can rewrite (10) as

$$\begin{aligned} x_1' &= \left(1 + \frac{s_2}{s_1^2 + s_2^2}\right) y_1 + \frac{s_1 x_1}{s_1^2 + s_2^2} - \frac{s_1^3 - 3s_1 s_2^2}{(s_1^2 + s_2^2)^3}, \\ y_1' &= -\left(1 + \frac{s_2}{s_1^2 + s_2^2}\right) x_1 + \frac{s_1 y_1}{s_1^2 + s_2^2} - \frac{s_2^3 - 3s_1^2 s_2}{(s_1^2 + s_2^2)^3}, \\ x_2' &= -\left(1 - \frac{s_2}{s_1^2 + s_2^2}\right) y_2 + \frac{s_1 x_2}{s_1^2 + s_2^2} - \frac{s_1^3 - 3s_1 s_2^2}{(s_1^2 + s_2^2)^3}, \\ y_2' &= \left(1 - \frac{s_2}{s_1^2 + s_2^2}\right) x_2 + \frac{s_1 y_2}{s_1^2 + s_2^2} - \frac{s_2^3 - 3s_1^2 s_2}{(s_1^2 + s_2^2)^3}, \\ s_1' &= -2s_1 s_2 (x_1 y_2 + x_2 y_1) + (x_1 x_2 - y_1 y_2) (s_1^2 - s_2^2) + 1, \\ s_2' &= 2s_1 s_2 (x_1 x_2 - y_1 y_2) + (x_1 y_2 + x_2 y_1) (s_1^2 - s_2^2). \end{aligned} \tag{87}$$

This is an ODE in \mathbb{R}^6 , with real time τ . We shall write Φ_τ for the flow induced by (87).

Note that $(s_1, s_2) = (0, -\rho)$ corresponds to the complex $s = -i\rho$.

For $x \in \mathbb{R}^6$ we shall write

$$\begin{aligned} \tau^+(x) &= \sup \{\tau < 0 : \pi_{s_1} \Phi_\tau(x) = 0\}, \\ \tau^-(x) &= \inf \{\tau > 0 : \pi_{s_1} \Phi_\tau(x) = 0\}, \end{aligned}$$




Figure 1: Illustration for Lemma B.1. Here we depict the projection of the flow onto the $s = (s_1, s_2)$ coordinates. The sets $\mathcal{P}^+(\mathbf{p}^+)$ and $\mathcal{P}^-(\mathbf{p}^-)$ which lead to the bound (91) for $\Delta\psi$ lie on the section $\{s_1 = 0\}$.

and define

$$\begin{aligned} \mathcal{P}^+, \mathcal{P}^- : \mathbb{R}^6 &\rightarrow \{s_1 = 0\}, \\ \mathcal{P}^+(x) &:= \Phi_{\tau^+(x)}(x), \\ \mathcal{P}^-(x) &:= \Phi_{\tau^-(x)}(x). \end{aligned}$$

We do not assume that \mathcal{P}^+ and \mathcal{P}^- are globally defined. Whenever we write $\mathcal{P}^+(x)$ or $\mathcal{P}^-(x)$ we will always validate that the considered point x lies in the domain of the map.

We know that the two solutions ψ^\pm of (11) (with $F_1 = -s^{-3}$, $F_2 = 0$, $H = 0$) satisfy

$$\|\psi^-(s)\| \leq |s|^{-3} M_0 \leq |\Re s|^{-3} M_0 \quad \text{for } \Re s < 0, \quad (88)$$

$$\|\psi^+(s)\| \leq |s|^{-3} M_0 \leq |\Re s|^{-3} M_0 \quad \text{for } \Re s > 0. \quad (89)$$

Lemma B.1. Consider $s^-, s_l^+, s_u^+ \in \mathbb{R}^2$ of the form $s^- = (s_1^-, s_2^-)$, $s_l^+ = (s_1^+, s_{2,l}^+)$, $s_u^+ = (s_1^+, s_{2,u}^+)$, where $s_1^- < 0 < s_1^+$ and $s_{2,l}^+ < s_{2,u}^+$. (The subscripts l and u stand for ‘lower’ and ‘upper’; see Figure 1.) Let $\mathbf{s}^+ \subset \mathbb{R}^2$ be the vertical interval joining s_l^+ and s_u^+ and let

$$\begin{aligned} \mathbf{p}^- &:= \left\{ p \in \mathbb{R}^6 : \pi_s p = s^-, \|\pi_{x,y} p\| \leq |s_1^-|^{-3} M_0 \right\}, \\ \mathbf{p}^+ &:= \left\{ p \in \mathbb{R}^6 : \pi_s p \in \mathbf{s}^+, \|\pi_{x,y} p\| \leq |s_1^+|^{-3} M_0 \right\}, \\ \mathbf{p}_u^+ &:= \left\{ p \in \mathbb{R}^6 : \pi_s p = s_u^+, \|\pi_{x,y} p\| \leq |s_1^+|^{-3} M_0 \right\}, \\ \mathbf{p}_l^+ &:= \left\{ p \in \mathbb{R}^6 : \pi_s p = s_l^+, \|\pi_{x,y} p\| \leq |s_1^+|^{-3} M_0 \right\}. \end{aligned}$$

If

$$\pi_{s_2} \mathcal{P}^+(\mathbf{p}_l^+) < \pi_{s_2} \mathcal{P}^-(\mathbf{p}^-) < \pi_{s_2} \mathcal{P}^+(\mathbf{p}_u^+) < 0 \quad (90)$$

then there exists a $\rho^* \in -\pi_{s_2} \mathcal{P}^-(\mathbf{p}^-)$ such that

$$\Delta\psi(-i\rho^*) = \psi^+(-i\rho^*) - \psi^-(-i\rho^*) \in \pi_{x,y}(\mathcal{P}^+(\mathbf{p}^+) - \mathcal{P}^-(\mathbf{p}^-)). \quad (91)$$

Proof. From (88–89) we see that

$$\begin{aligned}\psi^-(s^-) &\in \pi_{x,y}\mathbf{p}^-, \\ \psi^+(s^+) &\in \pi_{x,y}\mathbf{p}^+.\end{aligned}$$

By the Bolzano theorem applied to

$$\mathbf{s}^+ \ni s^+ \mapsto \pi_{s_2}(\mathcal{P}^+(\psi^-(s^+), s^+) - \mathcal{P}^-(\psi^-(s^-), s^-))$$

from (90) we see that there exists a $s_*^+ \in \mathbf{s}^+$ such that

$$\mathcal{P}^+(\psi^-(s_*^+), s_*^+) = \mathcal{P}^-(\psi^-(s^-), s^-).$$

By definition of \mathcal{P}^\pm we know that $\pi_s \mathcal{P}^\pm(q) \in \{0\} \times \mathbb{R}$, so

$$\pi_s \mathcal{P}^+(\psi^-(s_*^+), s_*^+) = \pi_s \mathcal{P}^-(\psi^-(s^-), s^-) = (0, -\rho^*),$$

for some $\rho^* > 0$ (the sign follows from (90)) and hence

$$\begin{aligned}\psi^-(-i\rho^*) &= \pi_{x,y} \mathcal{P}^-(\psi^-(s^-), s^-) \in \pi_{x,y} \mathcal{P}^-(\mathbf{p}^-), \\ \psi^+(-i\rho^*) &= \pi_{x,y} \mathcal{P}^-(\psi^-(s_*^+), s_*^+) \in \pi_{x,y} \mathcal{P}^+(\mathbf{p}^+),\end{aligned}$$

which implies (91), as required. \square

In our computer assisted proof we have taken $\bar{\rho} := 16.00008679$ and

$$s^- = (-10^3, -\bar{\rho}), \tag{92}$$

$$s_u^+ = (10^3, -\bar{\rho} + 10^{-6}), \tag{93}$$

$$s_l^+ = (10^3, -\bar{\rho} - 10^{-6}). \tag{94}$$

(the choice of $\bar{\rho}$ is dictated by the fact that then $\rho^* \approx 16$; see (62)). Then, we have validated that, with such choice of s^-, s_u^+, s_l^+ , Lemma B.1 leads to the bound (63). The computation of $\mathcal{P}^+(\mathbf{p}^+), \mathcal{P}^-(\mathbf{p}^-)$ required a long integration time, due to the number 10^3 in our choice of s_1^-, s_1^+ . The benefit of such large value in $\Re s$ is that then $|\Re s|^{-3}$ is a very small number, leading to small sets $\mathbf{p}^\pm, \mathbf{p}_l^+, \mathbf{p}_u^+$. This results in good bounds on $\Delta\psi$. Such choice of $\Re s$ was reached by trial and error.

The computer assisted validation of (63) took under 20 seconds, running on a single thread of a standard laptop.

The computation of $\Delta\psi$ in the second example also follows from Lemma

B.1. The only difference is the formula for the vector field, which is

$$\begin{aligned}
x'_1 &= \frac{s_1}{s_1^2 + s_2^2} x_1 + \left(\alpha + \frac{s_2}{s_1^2 + s_2^2} \right) y_1 - \frac{s_1^3 - 3s_1 s_2^2}{(s_1^2 + s_2^2)^3} \\
&\quad + \frac{1}{s_1^2 + s_2^2} (s_2 (x_1^2 - y_1^2 - x_2^2 + y_2^2) + 2s_1 (x_2 y_2 - x_1 y_1)), \\
y'_1 &= - \left(\alpha + \frac{s_2}{s_1^2 + s_2^2} \right) x_1 + \frac{s_1}{s_1^2 + s_2^2} y_1 - \frac{s_2^3 - 3s_1^2 s_2}{(s_1^2 + s_2^2)^3} \\
&\quad + \frac{1}{s_1^2 + s_2^2} (s_1 (x_1^2 - x_2^2 - y_1^2 + y_2^2) + 2s_2 (x_1 y_1 - x_2 y_2)), \\
x'_2 &= \frac{s_1}{s_1^2 + s_2^2} x_2 - \left(\alpha - \frac{s_2}{s_1^2 + s_2^2} \right) y_2 - \frac{s_1^3 - 3s_1 s_2^2}{(s_1^2 + s_2^2)^3} \\
&\quad + \frac{1}{s_1^2 + s_2^2} (s_2 (x_1^2 - y_1^2 - x_2^2 + y_2^2) + 2s_1 (x_2 y_2 - x_1 y_1)), \\
y'_2 &= \left(\alpha - \frac{s_2}{s_1^2 + s_2^2} \right) x_2 + \frac{s_1}{s_1^2 + s_2^2} y_2 - \frac{s_2^3 - 3s_1^2 s_2}{(s_1^2 + s_2^2)^3} \\
&\quad + \frac{1}{s_1^2 + s_2^2} (s_1 (x_1^2 - x_2^2 - y_1^2 + y_2^2) + 2s_2 (x_1 y_1 - x_2 y_2)), \\
s'_1 &= 1 - 2bs_1 s_2 (x_1 y_2 + x_2 y_1) + b (s_1^2 - s_2^2) (x_1 x_2 - y_1 y_2), \\
s'_2 &= 2bs_1 s_2 (x_1 x_2 - y_1 y_2) + b (s_1^2 - s_2^2) (x_1 y_2 + x_2 y_1).
\end{aligned}$$

We take the same s^- , s_l^+ , s_u^+ as in (92–94) which, with the aid of Lemma B.1 and interval arithmetic integration, leads to the bounds (68–69).

The computer assisted validation of (69) took under 25 seconds, running on a single thread of a standard laptop.

References

- [BCS13] I. Baldomá, O. Castejón, and T. M. Seara. Exponentially small heteroclinic breakdown in the generic Hopf-zero singularity. *J. Dynam. Differential Equations*, 25(2):335–392, 2013.
- [BCS18a] I. Baldomá, O. Castejón, and T. M. Seara. Breakdown of a 2D heteroclinic connection in the Hopf-zero singularity (I). *J. Nonlinear Sci.*, 28(5):1551–1627, 2018.
- [BCS18b] I. Baldomá, O. Castejón, and T. M. Seara. Breakdown of a 2D heteroclinic connection in the Hopf-zero singularity (II): the generic case. *J. Nonlinear Sci.*, 28(4):1489–1549, 2018.
- [BGG21a] I. Baldomá, M. Giralt, and M. Guardia. Breakdown of homoclinic orbits to L3 in the RPC3BP (I). complex singularities and the inner equation. Available at <https://arxiv.org/abs/2107.09942>, 2021.
- [BGG21b] I. Baldomá, M. Giralt, and M. Guardia. Breakdown of homoclinic orbits to L3 in the RPC3BP (II). an asymptotic formula. Available at <https://arxiv.org/abs/2107.09941>, 2021.

- [BIS20] I. Baldomá, S. Ibáñez, and T. M. Seara. Hopf-zero singularities truly unfold chaos. *Commun. Nonlinear Sci. Numer. Simul.*, 84:105162, 19, 2020.
- [BS08] I. Baldomá and T.M. Seara. The inner equation for generic analytic unfoldings of the Hopf-zero singularity. *Discrete Contin. Dyn. Syst., Ser. B*, 10(2-3):323–347, 2008.
- [BV84] H.W. Broer and G. Vegter. Subordinate Šil’nikov bifurcations near some singularities of vector fields having low codimension. *Ergodic Theory Dyn. Syst.*, 4:509–525, 1984.
- [CZ17] M. J. Capiński and P. Zgliczyński. Beyond the Melnikov method: a computer assisted approach. *J. Differential Equations*, 262(1):365–417, 2017.
- [CZ18] M. J. Capiński and P. Zgliczyński. Beyond the Melnikov method II: Multidimensional setting. *J. Differential Equations*, 265(9):3988–4015, 2018.
- [GGSZ21] O. Gomide, M. Guardia, T.M. Seara, and C. Zeng. On small breathers of nonlinear Klein-Gordon equations via exponentially small homoclinic splitting. Available at <https://arxiv.org/abs/2107.14566>, 2021.
- [GH90] J. Guckenheimer and P. Holmes. *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields*, volume 42. Springer Verlag, 1990.
- [GL14] V. Gelfreich and L. Lerman. Separatrix splitting at a Hamiltonian $0^2i\omega$ bifurcation. *Regul. Chaotic Dyn.*, 19(6):635–655, 2014.
- [Guc81] J. Guckenheimer. On a codimension two bifurcation. *Dynamical Systems and Turbulence, Warwick 1980*, pages 99–142, 1981.
- [Laz03] V. F. Lazutkin. Splitting of separatrices for the Chirikov standard map. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 300(Teor. Predst. Din. Sist. Spets. Vyp. 8):25–55, 285, 2003.
- [Mel63] V. K. Melnikov. On the stability of the center for time periodic perturbations. *Trans. Moscow Math. Soc.*, 12:1–57, 1963.
- [SK87] H. Segur and M. D. Kruskal. Nonexistence of small-amplitude breather solutions in ϕ^4 theory. *Phys. Rev. Lett.*, 58:747–750, Feb 1987.
- [Sn70] L. P. Sil’nikov. On the question of the structure of an extended neighborhood of a structurally stable state of equilibrium of saddle-focus type. *Mat. Sb. (N.S.)*, 81 (123):92–103, 1970.
- [Tre97] D. Treschev. Separatrix splitting for a pendulum with rapidly oscillating suspension point. *Russ. J. Math. Phys.*, 5(1):63–98, 1997.