

## Protocol

## Urban and social determinants of alcohol and tobacco consumption among adolescents in Madrid

Maitane Berasaluce<sup>a</sup>, Irene Martín-Turrero<sup>a</sup>, Roberto Valiente<sup>a,b,c</sup>, Lucía Martínez-Manrique<sup>a,d</sup>, María Sandín-Vázquez<sup>a,e</sup>, Xisca Sureda<sup>a,f,g,h,\*</sup><sup>a</sup> Public Health and Epidemiology Research Group, School of Medicine, University of Alcalá, Alcalá de Henares, Madrid, Spain<sup>b</sup> Centre for Research on Environment, Society and Health (CRESH), School of GeoSciences, University of Edinburgh, Edinburgh, United Kingdom<sup>c</sup> SPECTRUM Consortium, United Kingdom<sup>d</sup> Department of Preventive Medicine, Hospital Universitario de Móstoles, Móstoles, Madrid, Spain<sup>e</sup> Department of Community Health and Social Sciences, Graduate School of Public Health & Health Policy, City University of New York, New York, United States of America<sup>f</sup> Department of Epidemiology & Biostatistics, Graduate School of Public Health & Health Policy, City University of New York, New York, United States of America<sup>g</sup> Tobacco Control Research Group, Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain<sup>h</sup> CIBER de Enfermedades Respiratorias (CIBERES), Spain

## ARTICLE INFO

## Article history:

Received 28 June 2023

Accepted 25 September 2023

Available online 25 November 2023

## Keywords:

Alcohol drinking

Tobacco use

Adolescent

Urban health

Social determinants of health

## ABSTRACT

**Objective:** This study aims to describe the accessibility to and promotion of alcohol and tobacco around secondary schools in Madrid and its distribution in relation with area-level socioeconomic deprivation; analyze the relationship between this exposure and individual consumption characteristics of students between 14 and 18 years old; and explore other facilitators of this consumption.

**Method:** Mixed-methods study conducted in three phases: 1) we collected data on accessibility to and promotion of alcohol and tobacco in the environment using systematic social observation around 55 secondary schools; 2) we administered 2287 questionnaires among the students in these centers to gather information about characteristics and determinants of consumption; and 3) we conducted 20 semi-structured interviews and one discussion group to deepen in the results obtained in surveys and systematic social observation. We will use Geographic Information Systems to integrate and analyze the data from a spatial perspective.

© 2023 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Determinantes urbanos y sociales del consumo de alcohol y tabaco en adolescentes en Madrid

## RESUMEN

**Objetivo:** Este estudio tiene como objetivo describir la accesibilidad y la promoción de alcohol y tabaco alrededor de los centros de enseñanza secundaria en Madrid y su distribución en relación con la privación socioeconómica del área; analizar la relación entre esta exposición y las características individuales de consumo en estudiantes de entre 14 y 18 años; y explorar otros facilitadores del consumo.

**Método:** Estudio de métodos mixtos en tres fases: 1) recogimos datos sobre accesibilidad y promoción de estas sustancias en el entorno mediante observación social sistemática alrededor de 55 centros de secundaria; 2) administramos 2287 cuestionarios a estudiantes de estos centros sobre características y determinantes de consumo; y 3) realizamos 20 entrevistas semiestructuradas y un grupo de discusión en estos centros para profundizar en los resultados obtenidos en las encuestas y la observación social sistemática. Utilizaremos Sistemas de Información Geográfica para integrar y analizar los datos desde una perspectiva espacial.

© 2023 SESPAS. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Palabras clave:

Consumo de alcohol

Consumo de tabaco

Adolescentes

Salud urbana

Determinantes sociales de la salud

## Introduction

Adolescents and young people are especially vulnerable to alcohol and tobacco consumption, affecting neurodevelopment during these stages.<sup>1,2</sup> In addition to health consequences, it has been

shown that alcohol and tobacco consumption also have a high economic and social impact on the populations.<sup>3–5</sup> In Spain, 70.5% and 30.7% of adolescents aged between 14 and 18-years old reported alcohol and tobacco use within the last year, respectively.<sup>6</sup>

In the last decade there has been increasing interest in how the urban, social, and cultural environments influence people's behaviors related to health, such as substance use.<sup>7</sup> Evidence suggests that greater accessibility to alcohol and tobacco products and higher exposure to their promotion may increase alcohol and

\* Corresponding author.

E-mail address: [francisca.sureda@uah.es](mailto:francisca.sureda@uah.es) (X. Sureda).





**Figure 2.** Field collection map in a buffer of 500 meters by street network around the educational center.

We used systematic social observation to gather data on alcohol and tobacco points of sale, promotion elements/banners and signs of consumption within a predefined street network buffer of 500 meters around schools that accepted to participate in the study. Data were collected between August and October 2021, excluding school holiday weeks. An updated version of the OHCITIES questionnaire<sup>18,19</sup> was used to record the alcohol items. We modified an existing questionnaire used elsewhere to collect tobacco exposure.<sup>20–22</sup>

We provided theoretical-practical training to fieldworkers for data collection of alcohol and tobacco environments. We prepared maps of the buffer observation area around each secondary school (Fig. 2).

#### Phase 2: individual data collection

We conducted a survey among students between 14–18 years, enrolled in secondary schools in the city of Madrid between February and April 2022. An initial sample size of 1,089 students was calculated, 363 participants for each socioeconomic level stratum (assuming power of 80% and losses of 20%). This would allow us to detect changes in alcohol and tobacco prevalence of 7% between different socioeconomic strata based on the results obtained in a previous study.<sup>23</sup> Forty nine of the 55 centers involved in this study agreed to participate in the collection of individual consumption data, obtaining a total sample of 2287 students (with a loss of six centers in this phase, of which four were from strata representing high SES and high or medium alcohol and tobacco outlet density areas).

The questionnaire included information related to sociodemographic data, alcohol and tobacco consumption patterns and

determinants of their consumption. Alcohol and tobacco related questions were based on the Survey on Drug Use in Secondary Education in Spain (ESTUDES).<sup>24</sup> The questionnaires were provided and filled in paper format, except for seven centers that claimed for an online format. In these cases, we used a free online survey tool ([www.surveymonkey.com](http://www.surveymonkey.com)).

#### Phase 3: Qualitative data collection

A theoretical, intentional, non-probabilistic purposive sampling was carried out.<sup>25</sup> Inclusion criteria were being a student between 14 and 18 years of age and having ever consumed alcohol or tobacco. We recruited informants through the dissemination of a flyer in schools. We performed twenty semi-structured interviews (six from schools located in high and medium SES neighborhood, respectively, and eight from low SES neighborhood schools) and a discussion group (organized with five students from a medium SES neighborhood school) between April and September 2021. The topics covered in the interviews and discussion group tried to complement the information collected by the questionnaire in relation to alcohol and tobacco consumption patterns and factors influencing its consumption. Moreover, a section to explore health promotion strategies to prevent alcohol and tobacco consumption was included.

#### Statistical and geographic analyses

We will obtain prevalence ratios of consumption and consumption patterns. We will carry out qualitative and thematic descriptive analysis using ATLAS-ti software<sup>26</sup> and we will use sequential explanatory method<sup>27</sup> to relate quantitative and qualitative results. Transversally, we will use Geographic Information Systems to geocode exposure and consumption data. Geographic Information Systems will allow us to create measurements to characterize alcohol and tobacco environment (i.e., outlet proximity and density, density and visibility of promotion elements and signs of consumption) and analyze spatial inequalities in the alcohol and tobacco accessibility, promotion and consumption among different geographies in the city of Madrid. We will use Poisson regression models with multilevel robust variance, with the individual at level 1, the educational center at level 2 and the neighborhood at level 3 to assess associations between ecological data and individual characteristics.

#### Ethical considerations

This study was conducted according to the principles promulgated by the Declaration of Helsinki and the Ethics Committee of the University of Alcalá (CEI/HU/2020/48, CEIP2021/5/128) granted ethical approval. We informed participants and their legal representatives about the study and we requested a written informed consent from both parties, including also the permission to be contacted in the future. Confidentiality and anonymity of all data is guaranteed.

#### Discussion

The quantitative measurement of environmental exposure may be subject to discrepancies in the definition of operational variables trying to address the complexities of reality, as well as depending on the quality of the data. Likewise, information collected through questionnaires may be subject to reporting biases. To try to avoid these limitations, we will carry out a quality control of the data and we will contrast the quantitative information from the individual questionnaires with the semi-structured interviews and discussion group. The main strength of this project is the use of a mixed

methodology that combines qualitative with quantitative and geographic approaches. This will allow us to conduct an in-depth analysis of the relationships between the accessibility and promotion of alcohol and tobacco around schools and alcohol and tobacco consumption and its determinants in secondary school students. In addition, it will allow us to learn about other facilitators of the onset of alcohol and tobacco consumption in this population.

### Editor in charge

Miguel Ángel Negrín Hernández.

### Authors contributions

M. Berasaluce: writing original draft, methodology. I. Martín-Turrero: writing, review and editing, methodology. R. Valiente: conceptualization, writing, review and editing, methodology. L. Martínez-Manrique: review and editing, methodology. M. Sandín Vázquez: conceptualization, review and editing, methodology. X. Sureda: conceptualization, writing, review and editing, methodology, project administration, funding acquisition.

### Funding

This study is funded by two projects of the Community de Madrid and the Universidad de Alcalá (CM/JIN/2019-021 and CM/JIN/2021-030) and by the Fondo de Investigaciones Sanitarias (PI19/01626).

RV is a member of SPECTRUM a UK Prevention Research Partnership Consortium. UKPRP is an initiative funded by the UK Research and Innovation Councils, the Department of Health and Social Care (England) and the UK devolved administrations, and leading health research charities.

### Conflicts of interest

None.

### Acknowledgments

The authors would like to thank all the secondary schools and students that participated in this research project. We also thanks APLICA INVESTIGACIÓN Y TRASLACION S.COOP.MAD. for conducting the field work.

### References

- Guerra C, Pascual M. Mechanisms involved in the neurotoxic, cognitive, and neurobehavioral effects of alcohol consumption during adolescence. *Alcohol*. 2010;44:15–26.
- Lydon DM, Wilson SJ, Child A, et al. Adolescent brain maturation and smoking: what we know and where we're headed. *Neurosci Biobehav Rev*. 2014;45:323–42.
- Rehm J, Rehm MX, Shield KD, et al. Alcohol consumption, alcohol dependence and related harms in Spain, and the effect of treatment-based interventions on alcohol dependence. *Adicciones*. 2013;25:11–8.
- Sudhinaraset M, Wigglesworth C, Takeuchi DT. Social and cultural contexts of alcohol use: Influences in a social-ecological framework. *Alcohol Res*. 2016;38:35–45.
- Trangenstein PJ, Curriero FC, Webster D, et al. Outlet type, access to alcohol, and violent crime. *Alcohol Clin Exp Res*. 2018;42:2234–45.
- Observatorio Español de las Drogas y las Adicciones. Informe 2022. Alcohol, tabaco y drogas ilegales en España. Madrid: Ministerio de Sanidad, Delegación del Gobierno para el Plan Nacional sobre Drogas; 2022.
- Pearce J, Barnett R, Moon G. Sociospatial inequalities in health-related behaviours: pathways linking place and smoking. *Prog Hum Geogr*. 2011;36:3–24.
- Bryden A, Roberts B, McKee M, et al. A systematic review of the influence on alcohol use of community level availability and marketing of alcohol. *Health Place*. 2012;18:349–57.
- Valiente R, Escobar F, Urtasun M, et al. Tobacco retail environment and smoking: a systematic review of geographic exposure measures and implications for future studies. *Nicotine Tob Res*. 2021;23:1263–73.
- Shareck M, Kestens Y, Vallée J, et al. The added value of accounting for activity space when examining the association between tobacco retailer availability and smoking among young adults. *Tob Control*. 2015;25:406–12.
- Badland H, Mavoa S, Livingston M, et al. Testing spatial measures of alcohol outlet density with self-rated health in the Australian context: implications for policy and practice. *Drug Alcohol Rev*. 2016;35:298–306.
- Myran DT, Chen JT, Bearnot B, et al. Alcohol availability across neighborhoods in Ontario following alcohol sales deregulation, 2013–2017. *Am J Public Health*. 2019;109:899–905.
- Valiente R, Sureda X, Bilal U, et al. Regulating the local availability of tobacco retailing in Madrid, Spain: a GIS study to evaluate compliance. *Tob Control*. 2019;28:325–33.
- Martín-Turrero I, Valiente R, Molina-de la Fuente I, et al. Accessibility and availability of alcohol outlets around schools: an ecological study in the city of Madrid, Spain, according to socioeconomic area-level. *Environ Res*. 2022;204:112323.
- Gullón P, Bilal U, Cebrecos A, et al. Intersection of neighborhood dynamics and socioeconomic status in small-area walkability: the Heart Healthy Hoods project. *Int J Health Geogr*. 2017;16:21.
- Portal de datos abiertos del Ayuntamiento de Madrid. Censo de locales, sus actividades y terrazas de hostelería y restauración. Available at: <https://datos.madrid.es/portal/site/egob/menuitem.c05c1f754a33a9f8e4b2e4b284f1a5a0/?vgnnextoid=66665cde99be2410VgnVCM1000000b205a0aRCRD&vgnnextchannel=374512b9ace9f310VgnVCM1000001715a0aRCRD&vgnnextfmt=default>.
- Ministerio de Hacienda y Función Pública. Expendedurías. Available at: <https://www.hacienda.gob.es/es-ES/Areas%20Tematicas/CMTabacos/Paginas/Red-Expendedurias-y-PVR.aspx>.
- Sureda X, Espelt A, Villalbí JR, et al. Development and evaluation of the OHCTIES instrument: assessing alcohol urban environments in the Heart Healthy Hoods project. *BMJ Open*. 2017;7:1–10.
- Pastor A, Espelt A, Villalbí J, et al. Disponibilidad y promoción de alcohol según la tipología de los locales y las condiciones socioeconómicas del área. *Adicciones*. 2020;34:23–36.
- Sureda X, Bilal U, Fernández E, et al. Second-hand smoke exposure in outdoor hospitality venues: smoking visibility and assessment of airborne markers. *Environ Res*. 2018;165:220–7.
- Fu M, Fernández E, Martínez-Sánchez JM, et al. Second-hand smoke exposure in indoor and outdoor areas of cafés and restaurants: need for extending smoking regulation outdoors? *Environ Res*. 2016;148:421–8.
- Navas-Acien A, Çarkoğlu A, Ergör G, et al. Compliance with smoke-free legislation within public buildings: a cross-sectional study in Turkey. *Bull World Health Organ*. 2016;94:92–102.
- Larsen K, To T, Irving HM, et al. Smoking and binge-drinking among adolescents, Ontario, Canada: does the school neighbourhood matter? *Health Place*. 2017;47:108–14.
- Observatorio Español de las Drogas y las Adicciones. Encuesta sobre alcohol y otras drogas en España, EDADES 2019/20. Madrid: Ministerio de Sanidad, Delegación del Gobierno para el Plan Nacional sobre Drogas; 2021.
- Morse JM, Field PA. Qualitative research methods for health professionals. Thousand Oaks: Sage; 1995.
- ATLAS.ti. El Programa de Investigación y Análisis de Datos Cualitativos. Available at: <https://atlasti.com/es>.
- Ivankova NV, Creswell JW, Stick SL. Using mixed-methods sequential explanatory design: from theory to practice. *Field Meth*. 2006;18:3–20.