

In Search of Radical Transformations from Metal Enolates. Direct Reactions of *N*-Acyl-1,3-oxazolidin-2-ones with TEMPO Catalyzed by Copper(II) Acetate

Eduard Balaguer-Garcia, Marina Pérez-Palau, Cristina Bello, Anna Maria Costa,* Pedro Romea,* and Fèlix Urpí*



Cite This: *J. Org. Chem.* 2025, 90, 7332–7339



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: The direct reactions of a diverse range of *N*-acyl-1,3-oxazolidin-2-ones with TEMPO, catalyzed by copper(II) acetate and 4,7-dimethyl-1,10-phenanthroline without the need for any base, are herein described. These reactions afford the corresponding aminoxylated derivatives with high chemoselectivity and complete regioselectivity, achieving excellent yields under mild conditions. Further treatment of the resulting imides enables access to a variety of formally protected hydroxy compounds, which can be regarded as valuable synthetic intermediates. The efficient formal synthesis of isoproterenol highlights the potential of this methodology and sets the stage for further advancements in the catalytic chemistry of metal enolates.



INTRODUCTION


Metal enolates are widely regarded as some of the most versatile nucleophilic intermediates in organic synthesis, playing a pivotal role in carbon–carbon and carbon–heteroatom bond-forming reactions that proceed through polar mechanisms.^{1,2} These include alkylation, aldol and Mannich reactions, Michael additions, and Davis oxidations, among others.³ Largely due to their extensive utility, however, the potential of metal enolates to undergo alternative transformations through one-electron mechanisms,⁴ via ligand-to-metal charge transfer (LMCT) or valence tautomerism (Scheme 1),⁵ has been relatively overlooked, representing an underexplored area of research.

Given the unpredictable biradical nature of metal enolates, the persistent oxygen-centered radical (2,2,6,6-tetramethylpi-

peridin-1-yl)oxyl (TEMPO) has proven invaluable both as a probe to elucidate radical pathways and as an oxidizing reagent.⁶ Indeed, TEMPO was instrumental in the discovery of the biradical character of titanium enolates derived from chiral imides,⁷ thereby enabling the stereoselective formation of the corresponding α -hydroxy derivatives (eq 1, Scheme 2).^{8,9} Importantly, this mechanistic insight facilitated the development of new radical-based, stereoselective alkylation reactions from such substrates.^{10,11} In turn, Yazaki and Ohshima demonstrated that the anaerobic iron-catalyzed α -oxidation of carboxylic acids with TEMPO proceeds through a radical mechanism (eq 2, Scheme 2),¹² while Liu and Feng utilized TEMPO to confirm the radical pathway in the iron-catalyzed asymmetric alkylation of 2-acylimidazoles (eq 3, Scheme 2).¹³ Furthermore, Maulide showed that treatment of β,γ -unsaturated amides with triflic anhydride and TEMPO induces highly regioselective γ -oxidations via a radical mechanism (eq 4, Scheme 2).¹⁴

Motivated by these insights¹⁵ and building upon our previous work on the radical reactions of titanium enolates derived from *N*-acyl-1,3-oxazolidin-2-ones (eq 1, Scheme 2),

Scheme 1. Nucleophilic- and Radical-like Reactivity of Metal Enolates



Received: February 27, 2025


Revised: May 13, 2025

Accepted: May 16, 2025

Published: May 27, 2025



Scheme 2. Radical-like Reactions of Metal Enolates with TEMPO




we hypothesized that Lewis acids could catalyze direct radical-like reactions from these imides. In this context and given the lack of knowledge of the electronic character of the metal enolates from such imides, we decided to explore the reaction of model *N*-phenylacetyl-1,3-oxazolidin-2-one with TEMPO in the presence of substoichiometric amounts of first-period transition metal salts. Encouragingly, preliminary experiments demonstrated that copper(II) acetate effectively catalyzes their oxidation by TEMPO, provided that relatively acidic substrates are used. This finding prompted further investigation into the optimal reaction conditions and the scope of the overall transformation. Herein, we present our results on the reactions of *N*-(arylacetyl) and *N*-(β,γ -unsaturated acyl) oxazolidinones with TEMPO, catalyzed by copper(II) acetate and 4,7-dimethyl-1,10-phenanthroline leading to the selective formation of α - and γ -oxygenated derivatives respectively in high yields (Scheme 3).

RESULTS AND DISCUSSION

To identify the most appropriate experimental conditions, we first investigated the reaction of *N*-phenylacetyl-1,3-oxazolidin-


Scheme 3. Current Work. Reaction of Imides with TEMPO Catalyzed by Cu(OAc)₂ and 4,7-Dimethyl-1,10-phenanthroline



2-one (**1a**) with TEMPO in the presence of copper(II) acetate as the catalyst (Table 1). Thus, a series of solvents and temperatures were systematically evaluated to optimize the conversion of **1a** using two equivalents of TEMPO and one equivalent of triethylamine at 70 °C. The results, summarized in Table 1, indicated that the choice of solvent was critical. A protic solvent, such as isopropanol, led to a very poor yield of the α oxidized derivative **2a** (entry 1 in Table 1), while nonprotic solvents like THF and DCE significantly improved the conversion of **1a** (entries 2 and 3 in Table 1). Among the solvents tested, polar aprotic solvents, including DMSO, DMF, ethyl acetate, or acetonitrile gave the best results (entries 4–7 in Table 1). Notably, acetonitrile provided a significant 45% conversion, which was increased to completion when three equivalents of TEMPO were used (entry 8 in Table 1). Temperature also had a significant impact on the reaction outcome. Indeed, the conversion of **1a** was less than 30% at room temperature but raised to 60% at 30 °C (entries 8–12 in Table 1), which proved that mild heating was beneficial for the reaction. At this stage, we speculated that ligands known to coordinate to copper(II) might enhance the reaction outcome. To test this hypothesis, we evaluated the effect of 15 mol % of various ligands, including 1,4-diazabicyclo[2.2.2]octane (DABCO, **L1** in Table 1), 2,2'-bipyridine (bipy, **L2** in Table 1), and 1,10-phenanthroline (phen, **L3** in Table 1) at 30 °C for 16 h. These reactions were conducted in the presence of a bulkier tertiary amine such as diisopropylethylamine, to minimize potential coordination with the metal. The presence of DABCO (**L1**) was found to be detrimental and **2a** was obtained with a poor 15% of conversion (compare entries 11 and 13). In contrast, bipyridine (**L2**) and especially 1,10-phenanthroline (**L3**) improved the conversion up to 88% (compare entry 11 with entries 14–15). Surprisingly, we observed that a tertiary amine such as diisopropylethylamine, commonly used to deprotonate carbonyl substrates, was unnecessary. This suggested that copper(II) acetate could effectively catalyze the TEMPO-oxidation without the need for an external base to produce **2a** in up to 77% by simple addition of 15 mol % of 1,10-phenanthroline (**L3**). Varying the equivalents of copper(II) acetate and ligand **L3** led to the anticipated changes in conversion (compare entries 17–19 in Table 1). Finally, extending the reaction time to 24 h resulted in nearly quantitative conversion (entry 20 in Table 1).

Considering these findings, we explored whether other ligands derived from aromatic amines could also enhance the efficiency of such a straightforward transformation. Aiming to identify the most suitable ligand, we assessed the influence of substoichiometric amounts (15 mol %) of **L3**–**L11** on the oxidation of **1a** with two equivalents of TEMPO at 30 °C for a short reaction time (6 h). Bipyridine **L4** and phenanthroline **L5**, both containing strong electron-withdrawing groups, were found to be ineffective (entries 2 and 3 in Table 2). In turn, biquinoline **L6** yielded worse results than phenanthroline **L3** (entries 1 and 4 in Table 2), while 2,9-dimethyl-1,10-phenanthroline **L7** led to a reduced yield, likely due to the steric hindrance from the two methyl groups (entry 5 in Table 2). Interestingly, 4,7-disubstituted-1,10-phenanthrolines **L8**–**L11**, which feature groups with varying electronic properties, gave a wide range of conversion rates (entries 6–10 in Table 2). The methoxy substituted phenanthroline **L9** proved particularly ineffective, affording only a modest 15% yield after 24 h (entries 7 and 8 in Table 2), likely due to the poor solubility of the resulting copper complex. In contrast, 4,7-

Table 1. Preliminary Studies



entry	solvent	TEMPO (equiv)	R_3N	ligand	temperature ($^{\circ}\text{C}$)	conversion (%)
1	<i>i</i> -PrOH	2	Et_3N		70	<10
2	THF	2	Et_3N		70	21
3	DCE	2	Et_3N		70	23
4	DMSO	2	Et_3N		70	34
5	DMF	2	Et_3N		70	31
6	EtOAc	2	Et_3N		70	35
7	CH_3CN	2	Et_3N		70	45 ^a
8	CH_3CN	3	Et_3N		70	>95
9	CH_3CN	3	Et_3N		50	>95
10	CH_3CN	3	Et_3N		40	75
11	CH_3CN	3	Et_3N		30	60
12	CH_3CN	3	Et_3N		20	30
13	CH_3CN	3	<i>i</i> -Pr ₂ NEt	L1	30	15
14	CH_3CN	3	<i>i</i> -Pr ₂ NEt	L2	30	75
15	CH_3CN	3	<i>i</i> -Pr ₂ NEt	L3	30	88
16	CH_3CN	3		L2	30	59
17	CH_3CN	3		L3	30	77
18 ^b	CH_3CN	3		L3	30	65
19 ^c	CH_3CN	3		L3	30	85
20 ^d	CH_3CN	3		L3	30	95

^aTEMPO was completely consumed. ^b5 mol % of $\text{Cu}(\text{OAc})_2$ and 7.5 mol % of **L3** were used. ^c15 mol % of $\text{Cu}(\text{OAc})_2$ and 22.5 mol % of **L3** were used. ^dReaction time of 24 h.

dimethyl-1,10-phenanthroline **L11** emerged as the most promising ligand, delivering **2a** in 47% yield (entry 10 in Table 2). Finally, extending the reaction time to 24 h gave the desired adduct **2a** in an outstanding 99% yield (entry 11 in Table 2).

Having established the optimal experimental conditions to achieve a quantitative yield with model imide **1a**, we next investigated the scope of the reaction. This involved examining the reaction of *N*-arylacetyl-1,3-oxazolidin-2-ones **1a–1r** with TEMPO catalyzed by $\text{Cu}(\text{OAc})_2/\text{L11}$.


The results summarized in Table 3 demonstrate that the reaction exhibits a broad substrate scope, showing tolerance for a variety of aryl groups. Substrates possessing electron-donating groups, such as ethers (OBn or OMe, **1b–1e** in Table 3) and alkyl groups (Me, **1f–1h** in Table 3), consistently afforded the expected adducts (**2b–2h**) in yields of up to 99%, regardless of the relative position of the substituent. Notably, *ortho*-methoxy and *ortho*-methyl imides **1e** and **1h**, respectively, gave the lowest yields of the corresponding derivatives (**2e** and **2h**, 61% and 67% respectively), likely due to significant steric hindrance at the *ortho* position. Otherwise, electron-withdrawing groups had minimal impact on the reaction outcome, with adducts **2i–2k** being isolated in excellent yields. Finally, more complex oxygenated (**1l–1n**) and nitrogenated (**1o–1p**) imides also performed well, providing the corresponding adducts **2l–2p** in high yields.

The only exceptions were imides **1q** and **1r**, which contain an unprotected alcohol or amine, respectively, in the aromatic ring, as they turned out to be completely unreactive.

All in all, these results highlight the broad applicability of the reaction, which efficiently accommodates a wide range of *N*-arylacetyl imides **1**, yielding α -oxygenated adducts **2** with remarkable efficiency.

Next, we investigated the reduction of the N–O bond of the OTMP group to obtain the α hydroxy derivative.^{9a,16} Unfortunately, none of the conditions tested (Zn/AcOH, catalytic hydrogenation, ...) proved effective, so we shifted our focus to the removal of the oxazolidinone scaffold (Scheme 4). Treatment of the model adduct **2l** with LiOOH, aiming to obtain the corresponding carboxylic acid, was unsuccessful, as the oxidative conditions compromised the OTMP moiety. The use of LiOH resulted in the formation of the undesired amide **3l** in 87% yield as the steric bulk of the OTMP group directed the nucleophilic attack of the hydroxyl anion to the endocyclic carbonyl. In contrast, reductive conditions proved much more effective. Indeed, treatment with LiBH_4 at $0\text{ }^{\circ}\text{C}$ gave alcohol **4l** in 97% yield. On the other hand, the stronger reducing agent LiAlH_4 at $-50\text{ }^{\circ}\text{C}$ led to the formation of aldehyde **5l** in 64% yield. Finally, methyl ester **6l** was isolated in an excellent 96% yield by heating a methanol solution of **2l** in the presence 5 mol % of ytterbium triflate.¹⁷

Table 2. Influence of the Ligands



Reaction scheme showing the oxidation of 1a to 2a using 10 mol% Cu(OAc)₂ and 15 mol% of ligands L3–L11 in CH₃CN at 30 °C for 6 h.


Chemical structures of ligands L3–L11 are shown below the reaction scheme:

- L3: R: H
- L8: R: Cl
- L9: R: OMe
- L10: R: C₄H₈N
- L11: R: Me

entry	ligand	yield (%)
1	L3	38
2	L4	—
3	L5	—
4	L6	19
5	L7	21
6	L8	16
7	L9	6
8 ^a	L9	15
9	L10	26
10	L11	47
11 ^a	L11	99

^aReaction time of 24 h.

Scheme 4. Removal of the Oxazolidinone Heterocycle




Once we had demonstrated the broad scope of the reaction and the ability to efficiently remove the heterocycle, we next

applied the reported procedure to the formal synthesis of isoproterenol, an amino alcohol used in the treatment of respiratory diseases.¹⁸ The synthesis commenced with the well-known oxidation of 1m on a large scale (Scheme 5).¹⁹ To our pleasure, the desired adduct 2m was isolated in 97% yield, which was then converted into alcohol 4m with a remarkable 87% yield. Taken together, these results highlight the robustness of the overall method. Activation of the hydroxyl group followed by treatment of the resultant mesylate with isopropylamine afforded amine 7 in a 57% two-step yield. Subsequent reduction of the N–O bond with zinc in acetic acid was feasible following the removal of the oxazolidinone scaffold, and it successfully gave the desired amino alcohol 8 in quantitative yield. In summary, advanced intermediate 8, which provides a direct route to isoproterenol, was obtained from imide 1m in five steps with an overall yield of 47%.


The satisfactory α oxidation of *N*-arylacetyl oxazolidinones 1 prompted us to explore whether similar experimental conditions could be extended to other substrates, aiming to broaden the scope of the oxidation method. With this objective in mind, we examined the reactivity of tertiary arylacetic amides, structurally related thioimides, as well as fully saturated *N*-acyl oxazolidinones, along with their α,β - and β,γ -unsaturated counterparts (Scheme 6). Amide 9a did not show any reactivity under the established reaction conditions. In contrast, thioimides 10a and 11a underwent oxidation more rapidly than 1a, although the yields of the α oxidized adducts, 15a and 16a respectively, were lower. Altogether, these findings suggest that the presence of an imide or thioimide group, capable of chelating the putative metal enolate, is critical for the reaction and highlight the advantage of employing the more robust oxazolidinone scaffold. Among the other substrates tested, *N*-propanoyl oxazolidinone 12a resulted totally unreactive, whereas the α,β -unsaturated imide 13a yielded only tiny amounts of the γ -aminoxylated adduct 17a. However, the slightly more acidic *N*-[3-butenoyl]-1,3-oxazolidin-2-one 14a proved to be highly reactive and exclusively produced the γ -oxidized adduct 17a in 82% yield, along with 15% of the unreactive *N*-(2-butenoyl) oxazolidinone 13a (see Scheme 6 and Table 4).

From such a promising result, experimental parameters, including solvent, ligand, temperature, and time, were reevaluated carefully. The optimized conditions were essentially the same as those used for arylacetyl imides 1, apart from the reaction time, which was safely reduced to 16 h. Next, these conditions were applied to a broad range of β,γ -unsaturated imides 14a–14o to efficiently provide the corresponding oxidized derivatives 17a–17o (Table 4). Importantly, only the γ -aminoxylated adducts 17a–17n were obtained, without observing the formation of the corresponding α adducts, consistent with the results reported by Maulide in a related procedure proceeding through a radical pathway (see eq 4 in Scheme 2).¹⁵ The only exception was the doubly unsaturated imide 14o, which yielded the fully unsaturated $\text{C}\epsilon$ -oxidized adduct 17o in 84% (Table 4).

The data presented in Table 4 demonstrated that the experimental conditions could be applicable to a wide range of substrates with satisfactory yields. Importantly, β,γ -unsaturated imides containing disubstituted olefins ($\text{R}^1 = \text{R}^2 = \text{R}^4 = \text{H}$) produced the corresponding α,β -unsaturated γ -oxidized adducts with complete regioselectivity in yields ranging from 81% to 99%. Indeed, imides bearing simple alkyl groups ($\text{R}^3 = \text{H}$, alkyl in 14a–14d) afforded adducts 17a–17d in high

Table 3. Scope of the Reaction^a^aReactions at 0.3 mmol scale and 0.2 M.

Scheme 5. Formal Synthesis of Isoproterenol




yields, regardless of the steric hindrance of the substituents. Furthermore, the reaction tolerated the presence of a variety of functional groups, including phenyl, alkene, alkyne, chloride, ketone, and ester groups (**14e–14j**), with γ -oxidized adducts **17e–17j** isolated in yields around 90%. Otherwise, imide **14k** possessing a doubly substituted γ position ($R^2 = R^3 = Me$), produced the $\alpha,\beta,\gamma,\delta$ doubly unsaturated imide instead of the desired adduct **17k**. To our pleasure, imides **14l** and **14m**, which contain substituents at R^1 and R^4 that could potentially hinder the reaction, underwent a totally regioselective oxidation, albeit under more stringent conditions. In this context, the high yield obtained from α -methyl- β,γ -unsaturated imide **14m** was particularly rewarding, as the generation of


chelated metal enolates from such substrates is typically challenging. It is also worth mentioning that imide **14n**, which contains an allene group, provided the fully conjugated adduct **17n** in 50% yield. Finally, $\beta,\gamma,\delta,\epsilon$ -doubly unsaturated imide **14o** afforded the ϵ -aminoxylated adduct **17o** with a complete regioselectivity in an 84% yield.

The removal of the oxazolidinone scaffold of adducts **17** was troublesome. The use of a reducing agent, such as $LiBH_4$, with model adduct **17b**, a transformation successfully employed for arylacetyl derivatives **2**, was in this case useless and produced an almost equimolar mixture of the expected alcohol **18b** and the fully saturated alcohol **19b** in a good overall yield (Scheme 7).²⁰ Instead, heating a mixture of **17b** and a catalytic amount

Scheme 6. Substrate Tests




of ytterbium triflate in methanol gave the small but densely functionalized ester **20b** in excellent yield (Scheme 7).


Scheme 7. Methyl Ester from γ -Oxidized Adduct 17b

The results summarized in Table 4 demonstrate the efficiency of the direct and catalytic oxidation of β,γ -unsaturated imides with TEMPO to provide the corresponding γ -aminoxylated adducts in high yields with remarkable chemo- and regioselectivities.

Finally, and in the absence of a more detailed mechanistic study or experimental evidence, our working hypothesis to account for the observed results is based on the reaction of the biradical form of a copper(II) enolate with TEMPO, followed by the subsequent oxidation of the resulting copper(I) complex with a second TEMPO molecule (Scheme 8).²¹ We are fully aware that the complex reactivity of TEMPO may conceal alternative behavior, and a more detailed mechanistic

Table 4. Scope of the Reaction^a

Scheme 8. Mechanistic Working Hypothesis



study is thus necessary to confirm the radical character of the process.

CONCLUSION

In summary, we have developed a direct anaerobic oxidation method for *N*-(arylacetyl) and *N*-(β,γ -unsaturated acyl)-1,3-oxazolidin-2-ones with TEMPO, catalyzed by copper(II) acetate and 4,7-dimethyl-1,10-phenanthroline under mild experimental conditions. This method is applicable to a wide variety of substrates and exhibits high chemo- and regioselectivity, providing exclusively the corresponding α - and γ -OTMP derivatives in high yields. As proof of its synthetic utility, the method has been successfully employed in the formal synthesis of isoproterenol, an amino alcohol used in the treatment of respiratory diseases.

EXPERIMENTAL SECTION

General Procedure for the Synthesis of 2. A round-bottom flask equipped with a magnetic stirring bar was charged with *N*-arylacetyl-1,3-oxazolidin-2-one **1** (0.3 mmol, 1.0 equiv), anhydrous $\text{Cu}(\text{OAc})_2$ (5.45 mg, 30 μmol , 0.10 equiv, 10 mol %), 4,7-dimethyl-1,10-phenanthroline (9.37 mg, 45 μmol , 0.15 equiv, 15 mol %), and TEMPO (93.8 mg, 0.6 mmol, 2.0 equiv), followed by the addition of acetonitrile (1.5 mL) to get a 0.2 M solution. The resultant mixture was stirred under N_2 at 30 $^\circ\text{C}$ for 24 h and then quenched with sat. NH_4Cl (1.5 mL).

The aqueous layer was extracted with EtOAc (3 \times 1.5 mL). The combined organic extracts were dried (MgSO_4) and concentrated in vacuo. The resulting residue was purified by column chromatography to yield the α aminoxylated adduct **2**.

General Procedure for the Synthesis of 14. A round-bottom flask equipped with a magnetic stirring bar was charged with *N*-(β,γ -unsaturated acyl)-1,3-oxazolidin-2-one **14** (0.3 mmol, 1.0 equiv), anhydrous $\text{Cu}(\text{OAc})_2$ (5.45 mg, 30 μmol , 0.10 equiv, 10 mol %), 4,7-dimethyl-1,10-phenanthroline (9.37 mg, 45 μmol , 0.15 equiv, 15 mol %), and TEMPO (93.8 mg, 0.6 mmol, 2.0 equiv), followed by the addition of acetonitrile (1.5 mL) to get a 0.2 M solution. The resultant mixture was stirred under N_2 at 30 $^\circ\text{C}$ for 16 h and then quenched with sat. NH_4Cl (1.5 mL).

The aqueous layer was extracted with EtOAc (3 \times 1.5 mL). The combined organic extracts were dried (MgSO_4) and concentrated in

vacuo. The resulting residue was purified by column chromatography to yield the α,β -unsaturated γ aminoxylated adduct **17**.

ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its Supporting Information.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.5c00457>.

Experimental procedures, characterization data of all new compounds, and NMR spectra (PDF)

AUTHOR INFORMATION

Corresponding Authors

Anna Maria Costa – Department of Inorganic and Organic Chemistry, Section of Organic Chemistry, and Institut de Biomedicina de la Universitat de Barcelona, Universitat de Barcelona, 08028 Barcelona, Catalonia, Spain; orcid.org/0000-0003-4345-4750; Email: amcosta@ub.edu

Pedro Romea – Department of Inorganic and Organic Chemistry, Section of Organic Chemistry, and Institut de Biomedicina de la Universitat de Barcelona, Universitat de Barcelona, 08028 Barcelona, Catalonia, Spain; orcid.org/0000-0002-0259-9155; Email: pedro.romea@ub.edu

Fèlix Urpi – Department of Inorganic and Organic Chemistry, Section of Organic Chemistry, and Institut de Biomedicina de la Universitat de Barcelona, Universitat de Barcelona, 08028 Barcelona, Catalonia, Spain; orcid.org/0000-0003-4289-6506; Email: felix.urpi@ub.edu

Authors

Eduard Balaguer-Garcia – Department of Inorganic and Organic Chemistry, Section of Organic Chemistry, and Institut de Biomedicina de la Universitat de Barcelona, Universitat de Barcelona, 08028 Barcelona, Catalonia, Spain

Marina Pérez-Palau – Department of Inorganic and Organic Chemistry, Section of Organic Chemistry, and Institut de Biomedicina de la Universitat de Barcelona, Universitat de Barcelona, 08028 Barcelona, Catalonia, Spain; Present Address: M.P.-P.: Postdoctoral Researcher at Institut de Química Avançada-CSIC, Barcelona, 08034, Spain

Cristina Bello – Department of Inorganic and Organic Chemistry, Section of Organic Chemistry, and Institut de Biomedicina de la Universitat de Barcelona, Universitat de Barcelona, 08028 Barcelona, Catalonia, Spain; Present Address: C.B.: Almirall, Carretera Martorell 41–61, Sant Andreu de la Barca, 08740, Spain.

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.joc.5c00457>

Author Contributions

A.M.C., P.R., and F.U. conceived the project and wrote the manuscript. E.B.-G. conducted most of the experimental research, wrote, and edited ESI. M.P.-P. carried out initial experiments. C.B. performed experimental work on the reactions of β,γ -unsaturated imides. All the authors revised the manuscript and gave approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support from the Spanish Ministerio de Ciencia, Innovación y Universidades (MICIU/AEI/10.13039/501100011033/FEDER, UE, Grant PID2021-126251NB-I00), and the Generalitat de Catalunya (2021SGR 268), as well as doctorate studentships to E.B.-G. (FI, Generalitat de Catalunya) and M.P.-P. (FPU, MCIU) are gratefully acknowledged.

■ REFERENCES

- (1) Corey, E. J.; Cheng, X.-M. *The Logic of Chemical Synthesis*; John Wiley & Sons: New York, 1989.
- (2) Mekelburger, H. B.; Wilcox, C. S.; *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon Press: Oxford, 1991; Vol. 2.
- (3) Nicolaou, K. C.; Chen, J. S. *Classics in Total Synthesis III*; Wiley: Weinheim, 2011.
- (4) The importance of radical chemistry has grown significantly over the last years. For leading references, see (a) Beeson, T. D.; Mastracchio, A.; Hong, J.-B.; Ashton, K.; MacMillan, D. W. C. Enantioselective Organocatalysis Using SOMO Activation. *Science* **2007**, *316*, 582–585. (b) Studer, A.; Curran, D. P. The electron is a catalyst. *Nat. Chem.* **2014**, *6*, 765–773. (c) Smith, J. M.; Harwood, S. J.; Baran, P. S. Radical Retrosynthesis. *Acc. Chem. Res.* **2018**, *51*, 1807–1817. (d) Zhang, B.; He, J.; Gao, Y.; Levy, L.; Oderinde, M. S.; Palkowitz, M. D.; Dhar, T. G. M.; Mandler, M. D.; Collins, M. R.; Schmitt, D. C.; Bolduc, P. N.; Chen, T. Y.; Clementson, S.; Petersen, N. N.; Laudadio, G.; Bi, C.; Kawamata, Y.; Baran, P. S. Complex molecule synthesis by electrocatalytic decarboxylative cross-coupling. *Nature* **2023**, *623*, 745–751.
- (5) For a review on valence tautomerism in metal complexes, see Tezgerevska, T.; Alley, K. G.; Boskovic, C. Valence tautomerism in metal complexes: Stimulated and reversible intramolecular electron transfer between metal centers and organic ligands. *Coord. Chem. Rev.* **2014**, *268*, 23–40.
- (6) (a) Vogler, T.; Studer, A. Applications of TEMPO in Synthesis. *Synthesis* **2008**, 2008, 1979–1993. (b) Tebben, L.; Studer, A. Nitroxides Applications in Synthesis and in Polymer Chemistry. *Angew. Chem., Int. Ed.* **2011**, *50*, S034–S068.
- (7) Moreira, I.; Bofill, J. M.; Anglada, J. M.; Solsona, J. G.; Nebot, J.; Romea, P.; Urpí, F. Unconventional Biradical Character of Titanium Enolates. *J. Am. Chem. Soc.* **2008**, *130*, 3242–3243.
- (8) Mabe, P. J.; Zakarian, A. Asymmetric Radical Addition of TEMPO to Titanium Enolates. *Org. Lett.* **2014**, *16*, 516–519.
- (9) (a) Gómez-Palomino, A.; Pellicena, M.; Romo, J. M.; Solà, R.; Romea, P.; Urpí, F.; Font-Bardia, M. Stereoselective Aminooxygenation of Biradical Titanium Enolates with TEMPO. *Chem.—Eur. J.* **2014**, *20*, 10153–10159. (b) Heras, C.; Gómez-Palomino, A.; Romea, P.; Urpí, F.; Bofill, J. M.; Moreira, I.; de, P. R. Experimental and Computational Evidence of the Biradical Structure and Reactivity of Titanium(IV) Enolates. *J. Org. Chem.* **2017**, *82*, 8909–8916.
- (10) For pioneering and insightful applications of biradical enolate chemistry in synthesis, see: (a) Beaumont, S.; Ilardi, E. A.; Monroe, L. R.; Zakarian, A. Valence Tautomerism in Titanium Enolates: Catalytic Radical Haloalkylation and Application in the Total Synthesis of Neodysidenin. *J. Am. Chem. Soc.* **2010**, *132*, 1482–1483. (b) Gu, Z.; Herrmann, A. T.; Zakarian, A. Dual Ti–Ru Catalysis in the Direct Radical Haloalkylation of N-Acyl Oxazolidinones. *Angew. Chem., Int. Ed.* **2011**, *50*, 7136–7139.
- (11) (a) Gómez-Palomino, A.; Pérez-Palau, M.; Romea, P.; Urpí, F.; Del Olmo, M.; Hesse, T.; Fleckenstein, S.; Gómez-Bengoa, E.; Sotorrios, L.; Font-Bardia, M. Stereoselective Decarboxylative Alkylation of Titanium(IV) Enolates with Diacyl Peroxides. *Org. Lett.* **2020**, *22*, 199–203. (b) Pérez-Palau, M.; Sanosa, N.; Romea, P.; Urpí, F.; López, R.; Gómez-Bengoa, E.; Font-Bardia, M. Stereoselective Alkylation of Chiral Titanium(IV) Enolates with *tert*-Butyl Peresters. *Org. Lett.* **2021**, *23*, 8852–8856.
- (12) Tanaka, T.; Yazaki, R.; Ohshima, T. Chemoselective Catalytic α -Oxidation of Carboxylic Acids: Iron/Alkali Metal Cooperative Redox Active Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 4517–4524.
- (13) Xu, N.; Pu, M.; Yu, H.; Yang, G.; Liu, X.; Feng, X. Iron-Catalyzed Asymmetric α -Alkylation of 2-Acylimidazoles via Dehydrogenative Radical Cross-Coupling with Alkanes. *Angew. Chem., Int. Ed.* **2024**, *63*, No. e202314256.
- (14) Heindl, S.; Riomet, M.; Matyasovsky, J.; Lemmerer, M.; Malzer, N.; Maulide, N. Chemoselective γ -Oxidation of β,γ -Unsaturated Amides with TEMPO. *Angew. Chem., Int. Ed.* **2021**, *60*, 19123–19127.
- (15) For another copper-catalyzed precedent, see Taninokuchi, S.; Yazaki, R.; Ohshima, T. Catalytic Aerobic Chemoselective α -Oxidation of Acylpyrazoles en Route to α -Hydroxy Acid Derivatives. *Org. Lett.* **2017**, *19*, 3187–3190.
- (16) (a) Boger, D. L.; Garbaccio, R. M.; Jin, Q. Synthesis and Evaluation of CC-1065 and Duocarmycin Analogues Incorporating the Iso-CI and Iso-CBI Alkylation Subunits: Impact of Relocation of the C-4 Carbonyl. *J. Org. Chem.* **1997**, *62*, 8875–8891. (b) Atobe, M.; Yamazaki, N.; Kibayashi, C. Enantioselective Synthesis of Primary 1-(Aryl)alkylamines by Nucleophilic 1,2-Addition of Organolithium Reagents to Hydroxyoxime Ethers and Application to Asymmetric Synthesis of G-Protein-Coupled Receptor Ligands. *J. Org. Chem.* **2004**, *69*, 5595–5607. (c) Dinca, E.; Hartmann, P.; Smrcek, J.; Dix, I.; Jones, P. G.; Jahn, U. General and Efficient α -Oxygenation of Carbonyl Compounds by TEMPO Induced by Single-Electron-Transfer Oxidation of Their Enolates. *Eur. J. Org. Chem.* **2012**, *2012*, 4461–4482. (d) Chen, J.; Xu, Y.; Shao, W.; Ji, J.; Wang, B.; Yang, M.; Mao, G.; Xiao, F.; Deng, G.-J. Pd-Catalyzed C–O Bond Formation Enabling the Synthesis of Congested *N,N,O*-Trisubstituted Hydroxylamines. *Org. Lett.* **2022**, *24*, 8271–8276.
- (17) (a) Guissart, C.; Barros, A.; Rosa Barata, L.; Evano, G. Broadly Applicable Ytterbium-Catalyzed Esterification, Hydrolysis, and Amidation of Imides. *Org. Lett.* **2018**, *20*, 5098–5102. (b) Stevens, J. M.; Parra-Rivera, A. C.; Dixon, D. D.; Beutner, G. L.; DelMonte, A. J.; Frantz, D. E.; Janey, J. M.; Paulson, J.; Talley, M. R. Direct Lewis Acid Catalyzed Conversion of Enantioenriched *N*-Acylloxazolidinones to Chiral Esters, Amides, and Acids. *J. Org. Chem.* **2018**, *83*, 14245–14261.
- (18) Brittain, R. T.; Jack, D.; Ritchie, A. C. In *Advances in Drug Research*; Harper, N. J., Simmons, A. B., Eds.; Academic Press: New York, 1970; Vol. 5.
- (19) For other synthetic approaches, see: (a) Corey, E. J.; Link, J. O. The first enantioselective syntheses of pure *R*- and *S*-isoproterenol. *Tetrahedron Lett.* **1990**, *31*, 601–604. (b) Kumar, P.; Upadhyay, R. K.; Pandey, R. K. Asymmetric dihydroxylation route to (*R*)-isoprenaline, (*R*)-norfluoxetine and (*R*)-fluoxetine. *Tetrahedron: Asymmetry* **2004**, *15*, 3955–3959. (c) Blay, G.; Hernández-Olmos, V.; Pedro, J. R. Synthesis of (*S*)-(+)-sotalol and (*R*)-(–)-isoproterenol via a catalytic enantioselective Henry reaction. *Tetrahedron: Asymmetry* **2010**, *21*, 578–581.
- (20) Penning, T. D.; Djuric, S. W.; Haack, R. A.; Kalish, V. J.; Miyashiro, J. M.; Rowell, B. W.; Yu, S. S. Improved Procedure for the Reduction of *N*-Acylloxazolidinones. *Synth. Commun.* **1990**, *20*, 307–312.
- (21) For recent related examples on the oxidation of copper(I) complexes with peroxides, see (b) Xue, M.; Cui, J.; Zhu, X.; Wang, F.; Lv, D.; Nie, Z.; Li, Y.; Bao, H. Copper-Catalyzed Radical Enantioselective Synthesis of γ -Butyrolactones with Two Non-vicinal Carbon Stereocenters. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202304275. (c) Xiang, H.; Ye, Y. Cu-Catalyzed α -Alkylation of Glycine Derivatives for C(sp³)–H/C(sp³)–H Bond Selective Functionalization. *ACS Catal.* **2024**, *14*, 522–532.