

UNIVERSIDAD DE BARCELONA  
FACULTAD DE BIOLOGIA  
DEPARTAMENTO DE BIOLOGIA CELULAR

---

UNIVERSIDAD DE BARCELONA  
INSTITUTO DE INVESTIGACIÓN BIOMÉDICA  
PARQUE CIENTÍFICO DE BARCELONA

**Función de Netrina1 y Semaforinas  
secretables en la guía neuronal y axonal en el  
hipocampo y el cerebelo**

**Patricia Guijarro Larraz**  
Programa de Doctorado de Biología Celular  
Bienio 1998-2000

**Barcelona, Abril de 2006**

## **BIBLIOGRAFÍA**



## BIBLIOGRAFÍA

- Ackerman, S. L., Kozak, L. P., Przyborski, S. A., Rund, L. A., Boyer, B. B., Knowles, B. B. (1997) The mouse rostral cerebellar malformation gene encodes an UNC-5-like protein. *Nature*, **386**, 838-42.
- Ackerman, S. L., Knowles, B. B. (1998) Cloning and mapping of the UNC5C gene to human chromosome 4q21-q23. *Genomics*, **52**, 205-8.
- Adams, N. C., Tomoda, T., Cooper, M., Dietz, G., Hatten, M. E. (2002) Mice that lack astrotactin have slowed neuronal migration. *Development*, **129**, 965-972.
- Adams, R. H., Lohrum, M., Klostermann, A., Betz, H., Puschel, A. W. (1997) The chemorepulsive activity of secreted semaphorins is regulated by furin-dependent proteolytic processing. *Embo J*, **16**, 6077-86.
- Aizawa, H., Wakatsuki, S., Ishii, A., Moriyama, K., Sasaki, Y., Ohashi, K., Sekine-Aizawa, Y., Sehara-Fujisawa, A., Mizuno, K., Goshima, Y., Yahara, I. (2001) Phosphorylation of cofilin by LIM-kinase is necessary for semaphorin 3A-induced growth cone collapse. *Nat Neurosci*, **4**, 367-73.
- Alabed, Y. Z., Grados-Munro, E., Ferraro, G. B., Hsieh, S. H., Fournier, A. E. (2006) Neuronal responses to myelin are mediated by rho kinase. *J Neurochem*, **96**, 1616-25.
- Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., Watson, J. D. (1992) *Biología Molecular de la Célula*. Ediciones Omega, S.A., Barcelona.
- Alcantara, S., Ruiz, M., D'Arcangelo, G., Ezan, F., de Lecea, L., Curran, T., Sotelo, C., Soriano, E. (1998) Regional and cellular patterns of reelin mRNA expression in the forebrain of the developing and adult mouse. *J Neurosci*, **18**, 7779-99.
- Alcantara, S., Ruiz, M., De Castro, F., Soriano, E., Sotelo, C. (2000) Netrin 1 acts as an attractive or as a repulsive cue for distinct migrating neurons during the development of the cerebellar system. *Development*, **127**, 1359-1372.
- Alvarado-Mallart, R.-M. (2005) The chick/quail transplantation model: Discovery of the isthmic organizer center. *Brain Research Reviews*, **49**, 109-113.
- Amaral, D. G., Witter, M. P. (1995) Hippocampal formation. In Paxinos, G. (ed.), *The rat nervous system*. Academic Press, New York, pp. 443-494.
- Anderson, S. A., Eisenstat, D. D., Shi, L., Rubenstein, J. L. (1997) Interneuron migration from basal forebrain to neocortex: dependence on Dlx genes. *Science*, **278**, 474-6.
- Anderson, S. A., Marin, O., Horn, C., Jennings, K., Rubenstein, J. L. (2001) Distinct cortical migrations from the medial and lateral ganglionic eminences. *Development*, **128**, 353-63.
- Andrews, W., Liapi, A., Plachez, C., Camurri, L., Zhang, J., Mori, S., Murakami, F., Parnavelas, J. G., Sundaresan, V., Richards, L. J. (2006) Robo1 regulates the development of major axon tracts and interneuron migration in the forebrain. *Development*, **133**, 2243-52.
- Ara, J., Bannerman, P., Hahn, A., Ramirez, S., Pleasure, D. (2004) Modulation of Sciatic Nerve Expression of Class 3 Semaphorins by Nerve Injury. *Neurochemical Research*, **29**, 1153-1159.
- Arnaud, L., Ballif, B. A., Forster, E., Cooper, J. A. (2003) Fyn tyrosine kinase is a critical regulator of disabled-1 during brain development. *Curr Biol*, **13**, 9-17.
- Astic, L., Pellier-Monnin, V., Saucier, D., Charrier, C., Mehlen, P. (2002) Expression of netrin-1 and netrin-1 receptor, DCC, in the rat olfactory nerve pathway during development and axonal regeneration. *Neuroscience*, **109**, 643-56.
- Bagnard, D., Vaillant, C., Khuth, S. T., Dufay, N., Lohrum, M., Puschel, A. W., Belin, M. F., Bolz, J., Thomasset, N. (2001) Semaphorin 3A-vascular endothelial growth factor-165 balance mediates migration and apoptosis of neural progenitor cells by the recruitment of shared receptor. *J Neurosci*, **21**, 3332-41.
- Bagri, A., Cheng, H. J., Yaron, A., Pleasure, S. J., Tessier-Lavigne, M. (2003) Stereotyped pruning of long hippocampal axon branches triggered by retraction inducers of the semaphorin family. *Cell*, **113**, 285-99.
- Barallobre, M. J., Del Rio, J. A., Alcantara, S., Borrell, V., Aguado, F., Ruiz, M., Carmona, M. A., Martin, M., Fabre, M., Yuste, R., Tessier-Lavigne, M., Soriano, E. (2000) Aberrant development of hippocampal circuits and altered neural activity in netrin 1-deficient mice. *Development*, **127**, 4797-810.

- Barallobre, M. J., Pascual, M., Del Rio, J. A., Soriano, E. (2005) The Netrin family of guidance factors: emphasis on Netrin-1 signalling. *Brain Res Brain Res Rev*, **49**, 22-47.
- Barberis, D., Artigiani, S., Casazza, A., Corso, S., Giordano, S., Love, C. A., Jones, E. Y., Comoglio, P. M., Tamagnone, L. (2004) Plexin signaling hampers integrin-based adhesion, leading to Rho-kinase independent cell rounding, and inhibiting lamellipodia extension and cell motility. *Faseb J*, **18**, 592-4.
- Bartley, T. D., Hunt, R. W., Welcher, A. A., Boyle, W. J., Parker, V. P., Lindberg, R. A., Lu, H. S., Colombero, A. M., Elliott, R. L., Guthrie, B. A., et al. (1994) B61 is a ligand for the ECK receptor protein-tyrosine kinase. *Nature*, **368**, 558-60.
- Barton, W. A., Liu, B. P., Tzvetkova, D., Jeffrey, P. D., Fournier, A. E., Sah, D., Cate, R., Strittmatter, S. M., Nikolov, D. B. (2003) Structure and axon outgrowth inhibitor binding of the Nogo-66 receptor and related proteins. *Embo J*, **22**, 3291-302.
- Barzik, M., Kotova, T. I., Higgs, H. N., Hazelwood, L., Hanein, D., Gertler, F. B., Schafer, D. A. (2005) Ena/VASP proteins enhance actin polymerization in the presence of barbed end capping proteins. *J Biol Chem*, **280**, 28653-62.
- Bashaw, G. J., Kidd, T., Murray, D., Pawson, T., Goodman, C. S. (2000) Repulsive axon guidance: Abelson and Enabled play opposing roles downstream of the roundabout receptor. *Cell*, **101**, 703-15.
- Bashaw, G. J. (2004) Semaphorin signaling unplugged; a nervy AKAP cAMP(s) out on plexin. *Neuron*, **42**, 363-6.
- Bates, C. A., Stelzner, D. J. (1993) Extension and regeneration of corticospinal axons after early spinal injury and the maintenance of corticospinal topography. *Exp Neurol*, **123**, 106-17.
- Beffert, U., Weeber, E. J., Durudas, A., Qiu, S., Masiulis, I., Sweatt, J. D., Li, W. P., Adelmann, G., Frotscher, M., Hammer, R. E., Herz, J. (2005) Modulation of synaptic plasticity and memory by Reelin involves differential splicing of the lipoprotein receptor Apoer2. *Neuron*, **47**, 567-79.
- Behar, O., Golden, J. A., Mashimo, H., Schoen, F. J., Fishman, M. C. (1996) Semaphorin III is needed for normal patterning and growth of nerves, bones and heart. *Nature*, **383**, 525-8.
- Ben-Ari, Y., Khalilov, I., Represa, A., Gozlan, H. (2004) Interneurons set the tune of developing networks. *Trends in Neurosciences*, **27**, 422-427.
- Benson, M. D., Romero, M. I., Lush, M. E., Lu, Q. R., Henkemeyer, M., Parada, L. F. (2005) Ephrin-B3 is a myelin-based inhibitor of neurite outgrowth. *PNAS*, **102**, 10694-10699.
- Ben-Zvi, A., Yagil, Z., Hagalili, Y., Klein, H., Lerman, O., Behar, O. (2006) Semaphorin 3A and neurotrophins: a balance between apoptosis and survival signaling in embryonic DRG neurons. *J Neurochem*, **96**, 585-97.
- Berthie, B., Axelrad, H. (1994) Granular layer collaterals of the unipolar brush cell axon display rosette-like excrescences. A Golgi study in the rat cerebellar cortex. *Neurosci Lett*, **167**, 161-5.
- Bertrand, J., Winton, M. J., Rodriguez-Hernandez, N., Campenot, R. B., McKerracher, L. (2005) Application of Rho antagonist to neuronal cell bodies promotes neurite growth in compartmented cultures and regeneration of retinal ganglion cell axons in the optic nerve of adult rats. *J Neurosci*, **25**, 1113-21.
- Blanco, M. J., Pena-Melian, A., Angela Nieto, M. (2002) Expression of EphA receptors and ligands during chick cerebellar development. *Mechanisms of Development*, **114**, 225-229.
- Bloch-Gallego, E., Ezan, F., Tessier-Lavigne, M., Sotelo, C. (1999) Floor Plate and Netrin-1 Are Involved in the Migration and Survival of Inferior Olivary Neurons. *J. Neurosci.*, **19**, 4407-4420.
- Bloch-Gallego, E., Causeret, F., Ezan, F., Backer, S., Hidalgo-Sanchez, M. (2005) Development of precerebellar nuclei: instructive factors and intracellular mediators in neuronal migration, survival and axon pathfinding. *Brain Res Brain Res Rev*, **49**, 253-66.
- Bock, H. H., Herz, J. (2003) Reelin activates SRC family tyrosine kinases in neurons. *Curr Biol*, **13**, 18-26.
- Borrell, V., Del Rio, J. A., Alcantara, S., Derer, M., Martinez, A., D'Arcangelo, G., Nakajima, K., Mikoshiba, K., Derer, P., Curran, T., Soriano, E. (1999a) Reelin regulates the development and synaptogenesis of the layer-specific entorhino-hippocampal connections. *J Neurosci*, **19**, 1345-58.
- Borrell, V., Ruiz, M., Del Rio, J. A., Soriano, E. (1999b) Development of commissural connections in the hippocampus of reeler mice: evidence of an inhibitory influence of Cajal-Retzius cells. *Exp Neurol*, **156**, 268-82.

- Braisted, J. E., Catalano, S. M., Stimac, R., Kennedy, T. E., Tessier-Lavigne, M., Shatz, C. J., O'Leary, D. D. (2000) Netrin-1 promotes thalamic axon growth and is required for proper development of the thalamocortical projection. *J Neurosci*, **20**, 5792-801.
- Brose, K., Bland, K. S., Wang, K. H., Arnott, D., Henzel, W., Goodman, C. S., Tessier-Lavigne, M., Kidd, T. (1999) Slit proteins bind Robo receptors and have an evolutionarily conserved role in repulsive axon guidance. *Cell*, **96**, 795-806.
- Brose, K., Tessier-Lavigne, M. (2000) Slit proteins: key regulators of axon guidance, axonal branching, and cell migration. *Curr Opin Neurobiol*, **10**, 95-102.
- Brown, C. B., Feiner, L., Lu, M. M., Li, J., Ma, X., Webber, A. L., Jia, L., Raper, J. A., Epstein, J. A. (2001) PlexinA2 and semaphorin signaling during cardiac neural crest development. *Development*, **128**, 3071-80.
- Brown, M., Jacobs, T., Eickholt, B., Ferrari, G., Teo, M., Monfries, C., Qi, R. Z., Leung, T., Lim, L., Hall, C. (2004) Alpha2-chimaerin, cyclin-dependent Kinase 5/p35, and its target collapsin response mediator protein-2 are essential components in semaphorin 3A-induced growth-cone collapse. *J Neurosci*, **24**, 8994-9004.
- Brownlee, H., Gao, P. P., Frisen, J., Dreyfus, C., Zhou, R., Black, I. B. (2000) Multiple ephrins regulate hippocampal neurite outgrowth. *J Comp Neurol*, **425**, 315-22.
- Bundesen, L. Q., Scheel, T. A., Bregman, B. S., Kromer, L. F. (2003) Ephrin-B2 and EphB2 regulation of astrocyte-meningeal fibroblast interactions in response to spinal cord lesions in adult rats. *J Neurosci*, **23**, 7789-800.
- Cackowski, F. C., Xu, L., Hu, B., Cheng, S. Y. (2004) Identification of two novel alternatively spliced Neuropilin-1 isoforms. *Genomics*, **84**, 82-94.
- Cai, D., Shen, Y., De Bellard, M., Tang, S., Filbin, M. T. (1999) Prior exposure to neurotrophins blocks inhibition of axonal regeneration by MAG and myelin via a cAMP-dependent mechanism. *Neuron*, **22**, 89-101.
- Cai, D., Qiu, J., Cao, Z., McAtee, M., Bregman, B. S., Filbin, M. T. (2001) Neuronal cyclic AMP controls the developmental loss in ability of axons to regenerate. *J Neurosci*, **21**, 4731-9.
- Cai, H., Reed, R. R. (1999) Cloning and Characterization of Neuropilin-1-Interacting Protein: a PSD-95/Dlg/ZO-1 domain-containing protein that interacts with the cytoplasmic domain of Neuropilin-1. *J Neurosci*, **19**, 6519-6527.
- Campbell, D. S., Holt, C. E. (2001) Chemotropic responses of retinal growth cones mediated by rapid local protein synthesis and degradation. *Neuron*, **32**, 1013-26.
- Campbell, D. S., Regan, A. G., Lopez, J. S., Tannahill, D., Harris, W. A., Holt, C. E. (2001) Semaphorin 3A elicits stage-dependent collapse, turning, and branching in *Xenopus* retinal growth cones. *J Neurosci*, **21**, 8538-47.
- Carlier, M. F., Laurent, V., Santolini, J., Melki, R., Didry, D., Xia, G. X., Hong, Y., Chua, N. H., Pantaloni, D. (1997) Actin depolymerizing factor (ADF/cofilin) enhances the rate of filament turnover: implication in actin-based motility. *J Cell Biol*, **136**, 1307-22.
- Castellani, V., Chedotal, A., Schachner, M., Faivre-Sarrailh, C., Rougon, G. (2000) Analysis of the L1-deficient mouse phenotype reveals cross-talk between Sema3A and L1 signaling pathways in axonal guidance. *Neuron*, **27**, 237-49.
- Castellani, V., De Angelis, E., Kenwrick, S., Rougon, G. (2002) Cis and trans interactions of L1 with neuropilin-1 control axonal responses to semaphorin 3A. *Embo J*, **21**, 6348-57.
- Castellani, V., Falk, J., Rougon, G. (2004) Semaphorin3A-induced receptor endocytosis during axon guidance responses is mediated by L1 CAM. *Mol Cell Neurosci*, **26**, 89-100.
- Catalano, S. M., Messersmith, E. K., Goodman, C. S., Shatz, C. J., Chedotal, A. (1998) Many major CNS axon projections develop normally in the absence of semaphorin III. *Mol Cell Neurosci*, **11**, 173-82.
- Cebria, F., Newmark, P. A. (2005) Planarian homologs of netrin and netrin receptor are required for proper regeneration of the central nervous system and the maintenance of nervous system architecture. *Development*, **132**, 3691-703.
- Chedotal, A., Del Rio, J. A., Ruiz, M., He, Z., Borrell, V., de Castro, F., Ezan, F., Goodman, C. S., Tessier-Lavigne, M., Sotelo, C., Soriano, E. (1998) Semaphorins III and IV repel hippocampal axons via two distinct receptors. *Development*, **125**, 4313-23.
- Chedotal, A., Kerjan, G., Moreau-Fauvarque, C. (2005) The brain within the tumor: new roles for axon guidance molecules in cancers. *Cell Death Differ*, **12**, 1044-56.

- Chen, H., Chedotal, A., He, Z., Goodman, C. S., Tessier-Lavigne, M. (1997) Neuropilin-2, a novel member of the neuropilin family, is a high affinity receptor for the semaphorins Sema E and Sema IV but not Sema III. *Neuron*, **19**, 547-59.
- Chen, H., He, Z., Bagri, A., Tessier-Lavigne, M. (1998) Semaphorin-neuropilin interactions underlying sympathetic axon responses to class III semaphorins. *Neuron*, **21**, 1283-90.
- Chen, H., Bagri, A., Zupicich, J. A., Zou, Y., Stoeckli, E., Pleasure, S. J., Lowenstein, D. H., Skarnes, W. C., Chedotal, A., Tessier-Lavigne, M. (2000a) Neuropilin-2 regulates the development of selective cranial and sensory nerves and hippocampal mossy fiber projections. *Neuron*, **25**, 43-56.
- Chen, M. S., Huber, A. B., van der Haar, M. E., Frank, M., Schnell, L., Spillmann, A. A., Christ, F., Schwab, M. E. (2000b) Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. **403**, 434-439.
- Chen, Y. Q., Hsieh, J. T., Yao, F., Fang, B., Pong, R. C., Cipriano, S. C., Krepulat, F. (1999) Induction of apoptosis and G2/M cell cycle arrest by DCC. *Oncogene*, **18**, 2747-54.
- Chen, Z. Y., Sun, C., Reuhl, K., Bergemann, A., Henkemeyer, M., Zhou, R. (2004) Abnormal hippocampal axon bundling in EphB receptor mutant mice. *J Neurosci*, **24**, 2366-74.
- Cheng, H., Cao, Y., Olson, L. (1996) Spinal cord repair in adult paraplegic rats: partial restoration of hind limb function. *Science*, **273**, 510-3.
- Cheng, H. J., Nakamoto, M., Bergemann, A. D., Flanagan, J. G. (1995) Complementary gradients in expression and binding of ELF-1 and Mek4 in development of the topographic retinotectal projection map. *Cell*, **82**, 371-81.
- Cheng, H. J., Bagri, A., Yaron, A., Stein, E., Pleasure, S. J., Tessier-Lavigne, M. (2001) Plexin-A3 mediates semaphorin signaling and regulates the development of hippocampal axonal projections. *Neuron*, **32**, 249-63.
- Colamarino, S. A., Tessier-Lavigne, M. (1995) The axonal chemoattractant netrin-1 is also a chemorepellent for trochlear motor axons. *Cell*, **81**, 621-9.
- Committee, S. N. (1999) Unified nomenclature for the semaphorins/collapsins. *Cell*, **97**, 551-2.
- Condic, M. L. (2001) Adult neuronal regeneration induced by transgenic integrin expression. *J Neurosci*, **21**, 4782-8.
- Corset, V., Nguyen-Ba-Charvet, K. T., Forcet, C., Moyse, E., Chedotal, A., Mehlen, P. (2000) Netrin-1-mediated axon outgrowth and cAMP production requires interaction with adenosine A2b receptor. *Nature*, **407**, 747-750.
- Cramer, L. P. (1997) Molecular mechanism of actin-dependent retrograde flow in lamellipodia of motile cells. *Front Biosci*, **2**, d260-70.
- D'Arcangelo, G., Curran, T. (1998) Reeler: new tales on an old mutant mouse. *BioEssays*, **20**, 235-244.
- D'Arcangelo, G. (2005) Apoer2: a reelin receptor to remember. *Neuron*, **47**, 471-3.
- D'Arcangelo, G. (2006) Reelin mouse mutants as models of cortical development disorders. *Epilepsy Behav*, **8**, 81-90.
- de la Torre, J. R., Hopker, V. H., Ming, G. L., Poo, M. M., Tessier-Lavigne, M., Hemmati-Brivanlou, A., Holt, C. E. (1997) Turning of retinal growth cones in a netrin-1 gradient mediated by the netrin receptor DCC. *Neuron*, **19**, 1211-24.
- De Winter, F., Holtmaat, A. J., Verhaagen, J. (2002a) Neuropilin and class 3 semaphorins in nervous system regeneration. *Adv Exp Med Biol*, **515**, 115-39.
- De Winter, F., Oudega, M., Lankhorst, A. J., Hamers, F. P., Blits, B., Ruitenbergh, M. J., Pasterkamp, R. J., Gispen, W. H., Verhaagen, J. (2002b) Injury-induced class 3 semaphorin expression in the rat spinal cord. *Exp Neurol*, **175**, 61-75.
- de Wit, J., Verhaagen, J. (2003) Role of semaphorins in the adult nervous system. *Prog Neurobiol*, **71**, 249-67.
- Del Rio, J. A., Heimrich, B., Borrell, V., Forster, E., Drakew, A., Alcantara, S., Nakajima, K., Miyata, T., Ogawa, M., Mikoshiba, K., Derer, P., Frotscher, M., Soriano, E. (1997) A role for Cajal-Retzius cells and reelin in the development of hippocampal connections. *Nature*, **385**, 70-4.
- del Rio, J. A., Sole, M., Borrell, V., Martinez, A., Soriano, E. (2002) Involvement of Cajal-Retzius cells in robust and layer-specific regeneration of the entorhino-hippocampal pathways. *Eur J Neurosci*, **15**, 1881-90.
- Del Rio, J. A., Gonzalez-Billault, C., Urena, J. M., Jimenez, E. M., Barallobre, M. J., Pascual, M., Pujadas, L., Simo, S., La Torre, A., Wandosell, F., Avila, J., Soriano, E. (2004) MAP1B is

- required for Netrin 1 signaling in neuronal migration and axonal guidance. *Curr Biol*, **14**, 840-50.
- Delaney, C. L., Brenner, M., Messing, A. (1996) Conditional ablation of cerebellar astrocytes in postnatal transgenic mice. *Journal of Neuroscience*, **16**, 6908-6918.
- Dent, E. W., Barnes, A. M., Tang, F., Kalil, K. (2004) Netrin-1 and semaphorin 3A promote or inhibit cortical axon branching, respectively, by reorganization of the cytoskeleton. *J Neurosci*, **24**, 3002-12.
- Dickson, B. J., Keleman, K. (2002) Netrins. *Curr Biol*, **12**, R154-5.
- Domeniconi, M., Cao, Z., Spencer, T., Sivasankaran, R., Wang, K. C., Nikulina, E., Kimura, N., Cai, H., Deng, K., Gao, Y. (2002) Myelin-Associated Glycoprotein Interacts with the Nogo66 Receptor to Inhibit Neurite Outgrowth. *Neuron*, **35**, 283-290.
- Dou, C. L., Levine, J. M. (1997) Identification of a neuronal cell surface receptor for a growth inhibitory chondroitin sulfate proteoglycan (NG2). *J Neurochem*, **68**, 1021-30.
- Drescher, U., Kremoser, C., Handwerker, C., Loschinger, J., Noda, M., Bonhoeffer, F. (1995) In vitro guidance of retinal ganglion cell axons by RAGS, a 25 kDa tectal protein related to ligands for Eph receptor tyrosine kinases. *Cell*, **82**, 359-70.
- Dubreuil, C. I., Marklund, N., Deschamps, K., McIntosh, T. K., McKerracher, L. (2006) Activation of Rho after traumatic brain injury and seizure in rats. *Exp Neurol*, **198**, 361-9.
- Dupuy, S. T., Houser, C. R. (1996) Prominent Expression of Two Forms of Glutamate Decarboxylase in the Embryonic and Early Postnatal Rat Hippocampal Formation. *J. Neurosci.*, **16**, 6919-6932.
- Eickholt, B. J., Mackenzie, S. L., Graham, A., Walsh, F. S., Doherty, P. (1999) Evidence for collapsin-1 functioning in the control of neural crest migration in both trunk and hindbrain regions. *Development*, **126**, 2181-9.
- Eickholt, B. J., Walsh, F. S., Doherty, P. (2002) An inactive pool of GSK-3 at the leading edge of growth cones is implicated in Semaphorin 3A signaling. *J Cell Biol*, **157**, 211-7.
- Ellezam, B., Selles-Navarro, I., Manitt, C., Kennedy, T. E., McKerracher, L. (2001) Expression of netrin-1 and its receptors DCC and UNC-5H2 after axotomy and during regeneration of adult rat retinal ganglion cells. *Exp Neurol*, **168**, 105-15.
- Ellezam, B., Dubreuil, C., Winton, M., Loy, L., Dergham, P., Selles-Navarro, I., McKerracher, L. (2002) Inactivation of intracellular Rho to stimulate axon growth and regeneration. *Prog Brain Res*, **137**, 371-80.
- Engelkamp, D. (2002) Cloning of three mouse Unc5 genes and their expression patterns at mid-gestation. *Mech Dev*, **118**, 191-7.
- Fan, J., Mansfield, S. G., Redmond, T., Gordon-Weeks, P. R., Raper, J. A. (1993) The organization of F-actin and microtubules in growth cones exposed to a brain-derived collapsing factor. *J Cell Biol*, **121**, 867-78.
- Fatemi, S. H., Earle, J. A., McMenomy, T. (2000) Reduction in Reelin immunoreactivity in hippocampus of subjects with schizophrenia, bipolar disorder and major depression. *Mol Psychiatry*, **5**, 654-63, 571.
- Fazeli, A., Dickinson, S. L., Hermiston, M. L., Tighe, R. V., Steen, R. G., Small, C. G., Stoeckli, E. T., Keino-Masu, K., Masu, M., Rayburn, H., Simons, J., Bronson, R. T., Gordon, J. I., Tessier-Lavigne, M., Weinberg, R. A. (1997) Phenotype of mice lacking functional Deleted in colorectal cancer (Dcc) gene. *Nature*, **386**, 796-804.
- Feiner, L., Koppel, A. M., Kobayashi, H., Raper, J. A. (1997) Secreted chick semaphorins bind recombinant neuropilin with similar affinities but bind different subsets of neurons in situ. *Neuron*, **19**, 539-45.
- Feiner, L., Webber, A. L., Brown, C. B., Lu, M. M., Jia, L., Feinstein, P., Mombaerts, P., Epstein, J. A., Raper, J. A. (2001) Targeted disruption of semaphorin 3C leads to persistent truncus arteriosus and aortic arch interruption. *Development*, **128**, 3061-70.
- Feldheim, D. A., Kim, Y. I., Bergemann, A. D., Frisen, J., Barbacid, M., Flanagan, J. G. (2000) Genetic analysis of ephrin-A2 and ephrin-A5 shows their requirement in multiple aspects of retinocollicular mapping. *Neuron*, **25**, 563-74.
- Feldheim, D. A., Nakamoto, M., Osterfield, M., Gale, N. W., DeChiara, T. M., Rohatgi, R., Yancopoulos, G. D., Flanagan, J. G. (2004) Loss-of-function analysis of EphA receptors in retinotectal mapping. *J Neurosci*, **24**, 2542-50.



- Fernandez-Gonzalez, A., Spada, A. R. L., Treadaway, J., Higdon, J. C., Harris, B. S., Sidman, R. L., Morgan, J. I., Zuo, J. (2002) Purkinje cell degeneration (pcd) Phenotypes Caused by Mutations in the Axotomy-Induced Gene, *Nna1*. *Science*, **295**, 1904-1906.
- Filbin, M. T. (1995) Myelin-associated glycoprotein: a role in myelination and in the inhibition of axonal regeneration? *Curr Opin Neurobiol*, **5**, 588-95.
- Filbin, M. T. (1996) The muddle with MAG. *Mol Cell Neurosci*, **8**, 84-92.
- Finger, J. H., Bronson, R. T., Harris, B., Johnson, K., Przyborski, S. A., Ackerman, S. L. (2002) The netrin 1 receptors *Unc5h3* and *Dcc* are necessary at multiple choice points for the guidance of corticospinal tract axons. *J Neurosci*, **22**, 10346-56.
- Fiore, R., Puschel, A. W. (2003) The function of semaphorins during nervous system development. *Front Biosci*, **8**, s484-99.
- Flace, P., Benagiano, V., Lorusso, L., Girolamo, F., Rizzi, A., Virgintino, D., Roncali, L., Ambrosi, G. (2004) Glutamic acid decarboxylase immunoreactive large neuron types in the granular layer of the human cerebellar cortex. *Anat Embryol (Berl)*, **208**, 55-64.
- Forcet, C., Ye, X., Granger, L., Corset, V., Shin, H., Bredesen, D. E., Mehlen, P. (2001) The dependence receptor DCC (deleted in colorectal cancer) defines an alternative mechanism for caspase activation. *Proc Natl Acad Sci U S A*, **98**, 3416-21.
- Forcet, C., Stein, E., Pays, L., Corset, V., Llambi, F., Tessier-Lavigne, M., Mehlen, P. (2002) Netrin-1-mediated axon outgrowth requires deleted in colorectal cancer-dependent MAPK activation. *Nature*, **417**, 443-7.
- Fournier, A. E., Nakamura, F., Kawamoto, S., Goshima, Y., Kalb, R. G., Strittmatter, S. M. (2000) Semaphorin3A enhances endocytosis at sites of receptor-F-actin colocalization during growth cone collapse. *J Cell Biol*, **149**, 411-22.
- Fournier, A. E., GrandPre, T., Strittmatter, S. M. (2001) Identification of a receptor mediating Nogo-66 inhibition of axonal regeneration. **409**, 341-346.
- Fournier, A. E., Takizawa, B. T., Strittmatter, S. M. (2003) Rho kinase inhibition enhances axonal regeneration in the injured CNS. *J Neurosci*, **23**, 1416-23.
- Francis, S. H., Corbin, J. D. (1999) Cyclic nucleotide-dependent protein kinases: intracellular receptors for cAMP and cGMP action. *Crit Rev Clin Lab Sci*, **36**, 275-328.
- Freund, T. F., Buzsaki, G. (1996) Interneurons of the hippocampus. *Hippocampus*, **6**, 347-470.
- Frotscher, M., Seress, L., Schwedtfeger, W. K., Buhl, E. (1991). The mossy cells of the fascia dentata: a comparative study of their fine structure and synaptic connections in rodents and primates. *J Comp Neurol*, **312**, 145-63.
- Frotscher, M., Heimrich, B. (1993) Formation of layer-specific fiber projections to the hippocampus in vitro. *Proc Natl Acad Sci U S A*, **90**, 10400-3.
- Fuh, G., Garcia, K. C., de Vos, A. M. (2000) The interaction of neuropilin-1 with vascular endothelial growth factor and its receptor flt-1. *J Biol Chem*, **275**, 26690-5.
- Fujisawa, H., Kitsukawa, T., Kawakami, A., Takagi, S., Shimizu, M., Hirata, T. (1997) Roles of a neuronal cell-surface molecule, neuropilin, in nerve fiber fasciculation and guidance. *Cell Tissue Res*, **290**, 465-70.
- Fujisawa, H. (2004) Discovery of Semaphorin receptors, Neuropilin and Plexin, and their functions in neural development. *J Neurobiol*, **59**, 24-33.
- Fujitani, M., Kawai, H., Proia, R. L., Kashiwagi, A., Yasuda, H., Yamashita, T. (2005) Binding of soluble myelin-associated glycoprotein to specific gangliosides induces the association of p75NTR to lipid rafts and signal transduction. *J Neurochem*, **94**, 15-21.
- Fukata, Y., Itoh, T. J., Kimura, T., Menager, C., Nishimura, T., Shimizu, T., Watanabe, H., Inagaki, N., Iwamatsu, A., Hotani, H., Kaibuchi, K. (2002) CRMP-2 binds to tubulin heterodimers to promote microtubule assembly. *Nat Cell Biol*, **4**, 583-91.
- Gagliardini, V., Fankhauser, C. (1999) Semaphorin III can induce death in sensory neurons. *Mol Cell Neurosci*, **14**, 301-16.
- Gagnon, M. L., Bielenberg, D. R., Gechtman, Z., Miao, H. Q., Takashima, S., Soker, S., Klagsbrun, M. (2000) Identification of a natural soluble neuropilin-1 that binds vascular endothelial growth factor: In vivo expression and antitumor activity. *Proc Natl Acad Sci U S A*, **97**, 2573-8.
- Gallagher, E., Howell, B. W., Soriano, P., Cooper, J. A., Hawkes, R. (1998) Cerebellar abnormalities in the disabled (*mdab1-1*) mouse. *J Comp Neurol*, **402**, 238-51.

- Gammill, L. S., Gonzalez, C., Gu, C., Bronner-Fraser, M. (2006) Guidance of trunk neural crest migration requires neuropilin 2/semaphorin 3F signaling. *Development*, **133**, 99-106.
- Geisbrecht, B. V., Dowd, K. A., Barfield, R. W., Longo, P. A., Leahy, D. J. (2003) Netrin binds discrete subdomains of DCC and UNC5 and mediates interactions between DCC and heparin. *J Biol Chem*, **278**, 32561-8.
- Germroth, P., Schwerdtfeger, W. K., Buhl, E. H. (1989) GABAergic neurons in the entorhinal cortex project to the hippocampus. *Brain Res*, **494**, 187-92.
- Gherardi, E., Love, C. A., Esnouf, R. M., Jones, E. Y. (2004) The sema domain. *Curr Opin Struct Biol*, **14**, 669-78.
- Giger, R. J., Wolfner, D. P., De Wit, G. M., Verhaagen, J. (1996) Anatomy of rat semaphorin III/collapsin-1 mRNA expression and relationship to developing nerve tracts during neuroembryogenesis. *J Comp Neurol*, **375**, 378-92.
- Giger, R. J., Pasterkamp, R. J., Heijnen, S., Holtmaat, A. J., Verhaagen, J. (1998a) Anatomical distribution of the chemorepellent semaphorin III/collapsin-1 in the adult rat and human brain: predominant expression in structures of the olfactory-hippocampal pathway and the motor system. *J Neurosci Res*, **52**, 27-42.
- Giger, R. J., Urquhart, E. R., Gillespie, S. K., Levensgood, D. V., Ginty, D. D., Kolodkin, A. L. (1998b) Neuropilin-2 is a receptor for semaphorin IV: insight into the structural basis of receptor function and specificity. *Neuron*, **21**, 1079-92.
- Giger, R. J., Cloutier, J. F., Sahay, A., Prinjha, R. K., Levensgood, D. V., Moore, S. E., Pickering, S., Simmons, D., Rastan, S., Walsh, F. S., Kolodkin, A. L., Ginty, D. D., Geppert, M. (2000) Neuropilin-2 is required in vivo for selective axon guidance responses to secreted semaphorins. *Neuron*, **25**, 29-41.
- Gitler, A. D., Lu, M. M., Epstein, J. A. (2004) PlexinD1 and semaphorin signaling are required in endothelial cells for cardiovascular development. *Dev Cell*, **7**, 107-16.
- Gluzman-Poltorak, Z., Cohen, T., Herzog, Y., Neufeld, G. (2000) Neuropilin-2 is a receptor for the vascular endothelial growth factor (VEGF) forms VEGF-145 and VEGF-165 [corrected]. *J Biol Chem*, **275**, 18040-5.
- Gluzman-Poltorak, Z., Cohen, T., Shibuya, M., Neufeld, G. (2001) Vascular endothelial growth factor receptor-1 and neuropilin-2 form complexes. *J Biol Chem*, **276**, 18688-94.
- Goldberg, J. L., Vargas, M. E., Wang, J. T., Mandemakers, W., Oster, S. F., Sretavan, D. W., Barres, B. A. (2004) An Oligodendrocyte Lineage-Specific Semaphorin, Sema5A, Inhibits Axon Growth by Retinal Ganglion Cells. *J Neurosci.*, **24**, 4989-4999.
- Goldowitz, D., Hamre, K. M., Przyborski, S. A., Ackerman, S. L. (2000) Granule Cells and Cerebellar Boundaries: Analysis of Unc5h3 Mutant Chimeras. *J Neurosci.*, **20**, 4129-4137.
- Gomez, T. (2005) Neurobiology: channels for pathfinding. *Nature*, **434**, 835-8.
- Goshima, Y., Nakamura, F., Strittmatter, P., Strittmatter, S. M. (1995) Collapsin-induced growth cone collapse mediated by an intracellular protein related to UNC-33. *Nature*, **376**, 509-14.
- Gozlan, H., Ben-Ari, Y. (2003) Interneurons are the Source and the Targets of the First Synapses Formed in the Rat Developing Hippocampal Circuit. *Cereb. Cortex*, **13**, 684-692.
- Graef, I. A., Wang, F., Charron, F., Chen, L., Neilson, J., Tessier-Lavigne, M., Crabtree, G. R. (2003) Neurotrophins and netrins require calcineurin/NFAT signaling to stimulate outgrowth of embryonic axons. *Cell*, **113**, 657-70.
- GrandPre, T., Nakamura, F., Vartanian, T., Strittmatter, S. M. (2000) Identification of the Nogo inhibitor of axon regeneration as a Reticulon protein. **403**, 439-444.
- Groves, M. L., McKeon, R., Werner, E., Nagarsheth, M., Meador, W., English, A. W. (2005) Axon regeneration in peripheral nerves is enhanced by proteoglycan degradation. *Exp Neurol*, **195**, 278-92.
- Gu, C., Limberg, B. J., Whitaker, G. B., Perman, B., Leahy, D. J., Rosenbaum, J. S., Ginty, D. D., Kolodkin, A. L. (2002) Characterization of neuropilin-1 structural features that confer binding to semaphorin 3A and vascular endothelial growth factor 165. *J Biol Chem*, **277**, 18069-76.
- Gu, C., Rodriguez, E. R., Reimert, D. V., Shu, T., Fritzsche, B., Richards, L. J., Kolodkin, A. L., Ginty, D. D. (2003) Neuropilin-1 conveys semaphorin and VEGF signaling during neural and cardiovascular development. *Dev Cell*, **5**, 45-57.

- Gu, C., Yoshida, Y., Livet, J., Reimert, D. V., Mann, F., Merte, J., Henderson, C. E., Jessell, T. M., Kolodkin, A. L., Ginty, D. D. (2005) Semaphorin 3E and plexin-D1 control vascular pattern independently of neuropilins. *Science*, **307**, 265-8.
- Guirland, C., Suzuki, S., Kojima, M., Lu, B., Zheng, J. Q. (2004) Lipid Rafts Mediate Chemotropic Guidance of Nerve Growth Cones. *Neuron*, **42**, 51-62.
- Guthrie, S. (2004) Axon guidance: mice and men need Rig and Robo. *Curr Biol*, **14**, R632-4.
- Guttmann-Raviv, N., Kessler, O., Shraga-Heled, N., Lange, T., Herzog, Y., Neufeld, G. (2006) The neuropilins and their role in tumorigenesis and tumor progression. *Cancer Lett*, **231**, 1-11.
- Hagino, S., Iseki, K., Mori, T., Zhang, Y., Hikake, T., Yokoya, S., Takeuchi, M., Hasimoto, H., Kikuchi, S., Wanaka, A. (2003) Slit and glypican-1 mRNAs are coexpressed in the reactive astrocytes of the injured adult brain. *Glia*, **42**, 130-8.
- Hamasaki, T., Goto, S., Nishikawa, S., Ushio, Y. (2001) A Role of Netrin-1 in the Formation of the Subcortical Structure Striatum: Repulsive Action on the Migration of Late-Born Striatal Neurons. *J. Neurosci.*, **21**, 4272-4280.
- Hansen, M. J., Dallal, G. E., Flanagan, J. G. (2004) Retinal axon response to ephrin-as shows a graded, concentration-dependent transition from growth promotion to inhibition. *Neuron*, **42**, 717-30.
- Hasan, S. J., Keirstead, H. S., Muir, G. D., Steeves, J. D. (1993) Axonal regeneration contributes to repair of injured brainstem-spinal neurons in embryonic chick. *J Neurosci*, **13**, 492-507.
- Hasegawa, Y., Fujitani, M., Hata, K., Tohyama, M., Yamagishi, S., Yamashita, T. (2004) Promotion of axon regeneration by myelin-associated glycoprotein and Nogo through divergent signals downstream of Gi/G. *J Neurosci*, **24**, 6826-32.
- Hatten, M. E., Heintz, N. (1995) Mechanisms of neural patterning and specification in the developing cerebellum. *Annu Rev Neurosci*, **18**, 385-408.
- Hatten, M. E. (1999) Central nervous system neuronal migration. *Annu Rev Neurosci*, **22**, 511-39.
- He, Z., Tessier-Lavigne, M. (1997) Neuropilin is a receptor for the axonal chemorepellent Semaphorin III. *Cell*, **90**, 739-51.
- He, Z., Koprivica, V. (2004) The Nogo signaling pathway for regeneration block. *Annual Review of Neuroscience*, **27**, 341-368.
- Hedgecock, E. M., Culotti, J. G., Hall, D. H. (1990) The unc-5, unc-6, and unc-40 genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in *C. elegans*. *Neuron*, **4**, 61-85.
- Hedrick, L., Cho, K. R., Fearon, E. R., Wu, T. C., Kinzler, K. W., Vogelstein, B. (1994) The DCC gene product in cellular differentiation and colorectal tumorigenesis. *Genes Dev*, **8**, 1174-83.
- Hennou, S., Khalilov, I., Diabira, D., Ben-Ari, Y., Gozlan, H. (2002) Early sequential formation of functional GABA<sub>A</sub> and glutamatergic synapses on CA1 interneurons of the rat foetal hippocampus. *Eur J Neurosci*, **16**, 197-208.
- Herincs, Z., Corset, V., Cahuzac, N., Furne, C., Castellani, V., Hueber, A. O., Mehlen, P. (2005) DCC association with lipid rafts is required for netrin-1-mediated axon guidance. *J Cell Sci*, **118**, 1687-92.
- Herrick, T. M., Cooper, J. A. (2002) A hypomorphic allele of *dab1* reveals regional differences in reelin-Dab1 signaling during brain development. *Development*, **129**, 787-96.
- Herrup, K., Kuemerle, B. (1997) The compartmentalization of the cerebellum. *Annu Rev Neurosci*, **20**, 61-90.
- Hirai, H., Maru, Y., Hagiwara, K., Nishida, J., Takaku, F. (1987) A novel putative tyrosine kinase receptor encoded by the *eph* gene. *Science*, **238**, 1717-20.
- Holtmaat, A. J., De Winter, F., De Wit, J., Gorter, J. A., da Silva, F. H., Verhaagen, J. (2002) Semaphorins: contributors to structural stability of hippocampal networks? *Prog Brain Res*, **138**, 17-38.
- Holtmaat, A. J., Gorter, J. A., De Wit, J., Tolner, E. A., Spijker, S., Giger, R. J., Lopes da Silva, F. H., Verhaagen, J. (2003) Transient downregulation of Sema3A mRNA in a rat model for temporal lobe epilepsy. A novel molecular event potentially contributing to mossy fiber sprouting. *Exp Neurol*, **182**, 142-50.
- Homayouni, R., Rice, D. S., Curran, T. (2001) Disabled-1 interacts with a novel developmentally regulated protocadherin. *Biochem Biophys Res Commun*, **289**, 539-47.

- Hong, K., Hinck, L., Nishiyama, M., Poo, M. M., Tessier-Lavigne, M., Stein, E. (1999) A ligand-gated association between cytoplasmic domains of UNC5 and DCC family receptors converts netrin-induced growth cone attraction to repulsion. *Cell*, **97**, 927-41.
- Hong, K., Nishiyama, M., Henley, J., Tessier-Lavigne, M., Poo, M. (2000a) Calcium signalling in the guidance of nerve growth by netrin-1. *Nature*, **403**, 93-8.
- Hong, S. E., Shugart, Y. Y., Huang, D. T., Shahwan, S. A., Grant, P. E., Hourihane, J. O., Martin, N. D., Walsh, C. A. (2000b) Autosomal recessive lissencephaly with cerebellar hypoplasia is associated with human RELN mutations. *Nat Genet*, **26**, 93-6.
- Hopker, V. H., Shewan, D., Tessier-Lavigne, M., Poo, M., Holt, C. (1999) Growth-cone attraction to netrin-1 is converted to repulsion by laminin-1. *Nature*, **401**, 69-73.
- Hoshino, M., Nakamura, S., Mori, K., Kawauchi, T., Terao, M., Nishimura, Y. V., Fukuda, A., Fuse, T., Matsuo, N., Sone, M., Watanabe, M., Bito, H., Terashima, T., Wright, C. V., Kawaguchi, Y., Nakao, K., Nabeshima, Y. (2005) Ptf1a, a bHLH transcriptional gene, defines GABAergic neuronal fates in cerebellum. *Neuron*, **47**, 201-13.
- Howell, B. W., Gertler, F. B., Cooper, J. A. (1997) Mouse disabled (mDab1): a Src binding protein implicated in neuronal development. *Embo J*, **16**, 121-32.
- Hu, H., Marton, T. F., Goodman, C. S. (2001) Plexin B mediates axon guidance in *Drosophila* by simultaneously inhibiting active Rac and enhancing RhoA signaling. *Neuron*, **32**, 39-51.
- Hu, H., Li, M., Labrador, J. P., McEwen, J., Lai, E. C., Goodman, C. S., Bashaw, G. J. (2005) Cross GTPase-activating protein (CrossGAP)/Vilse links the Roundabout receptor to Rac to regulate midline repulsion. *Proc Natl Acad Sci U S A*, **102**, 4613-8.
- Huang, X., Cheng, H. J., Tessier-Lavigne, M., Jin, Y. (2002) MAX-1, a novel PH/MyTH4/FERM domain cytoplasmic protein implicated in netrin-mediated axon repulsion. *Neuron*, **34**, 563-76.
- Huber, A. B., Kolodkin, A. L., Ginty, D. D., Cloutier, J. F. (2003) Signaling at the growth cone: ligand-receptor complexes and the control of axon growth and guidance. *Annu Rev Neurosci*, **26**, 509-63.
- Husmann, K., Faissner, A., Schachner, M. (1992) Tenascin promotes cerebellar granule cell migration and neurite outgrowth by different domains in the fibronectin type III repeats. *J. Cell Biol.*, **116**, 1475-1486.
- Ice, R. J., Wildonger, J., Mann, R. S., Hiebert, S. W. (2005) Comment on "Nervy links protein kinase A to plexin-mediated semaphorin repulsion". *Science*, **309**, 558; author reply 558.
- Ishii, N., Wadsworth, W. G., Stern, B. D., Culotti, J. G., Hedgecock, E. M. (1992) UNC-6, a laminin-related protein, guides cell and pioneer axon migrations in *C. elegans*. *Neuron*, **9**, 873-81.
- Ishizuka, N., Weber, J., Amaral, D. G. (1990) Organization of intrahippocampal projections originating from CA3 pyramidal cells in the rat. *J Comp Neurol*, **295**, 580-623.
- Jain, A., Brady-Kalnay, S. M., Bellamkonda, R. V. (2004) Modulation of Rho GTPase activity alleviates chondroitin sulfate proteoglycan-dependent inhibition of neurite extension. *J Neurosci Res*, **77**, 299-307.
- Jarjour, A. A., Manitt, C., Moore, S. W., Thompson, K. M., Yuh, S.-J., Kennedy, T. E. (2003) Netrin-1 Is a Chemorepellent for Oligodendrocyte Precursor Cells in the Embryonic Spinal Cord. *J. Neurosci.*, **23**, 3735-3744.
- Jensen, P., Zoghbi, H. Y., Goldowitz, D. (2002) Dissection of the Cellular and Molecular Events that Position Cerebellar Purkinje Cells: A Study of the math1 Null-Mutant Mouse. *J. Neurosci.*, **22**, 8110-8116.
- Julien, F., Bechara, A., Fiore, R., Nawabi, H., Zhou, H., Hoyo-Becerra, C., Bozon, M., Rougon, G., Grumet, M., Puschel, A. W., Sanes, J. R., Castellani, V. (2005) Dual functional activity of semaphorin 3B is required for positioning the anterior commissure. *Neuron*, **48**, 63-75.
- Kandel, E. R., Schwartz, J. H., Jessell, T. M. (1993) *Principles of Neural Science*. Prentice-Hall International.
- Karam, S. D., Burrows, R. C., Logan, C., Koblar, S., Pasquale, E. B., Bothwell, M. (2000) Eph Receptors and Ephrins in the Developing Chick Cerebellum: Relationship to Sagittal Patterning and Granule Cell Migration. *J. Neurosci.*, **20**, 6488-6500.
- Karam, S. D., Dottori, M., Ogawa, K., Henderson, J. T., Boyd, A. W., Pasquale, E. B., Bothwell, M. (2002) EphA4 is not required for Purkinje cell compartmentation. *Developmental Brain Research*, **135**, 29-38.

- Kawaguchi, Y., Cooper, B., Gannon, M., Ray, M., MacDonald, R. J., Wright, C. V. (2002) The role of the transcriptional regulator Ptf1a in converting intestinal to pancreatic progenitors. *Nat Genet*, **32**, 128-34.
- Kawakami, A., Kitsukawa, T., Takagi, S., Fujisawa, H. (1996) Developmentally regulated expression of a cell surface protein, neuropilin, in the mouse nervous system. *J Neurobiol*, **29**, 1-17.
- Kawasaki, T., Kitsukawa, T., Bekku, Y., Matsuda, Y., Sanbo, M., Yagi, T., Fujisawa, H. (1999) A requirement for neuropilin-1 in embryonic vessel formation. *Development*, **126**, 4895-902.
- Keino-Masu, K., Masu, M., Hinck, L., Leonardo, E. D., Chan, S. S., Culotti, J. G., Tessier-Lavigne, M. (1996) Deleted in Colorectal Cancer (DCC) encodes a netrin receptor. *Cell*, **87**, 175-85.
- Keleman, K., Dickson, B. J. (2001) Short- and long-range repulsion by the Drosophila Unc5 netrin receptor. *Neuron*, **32**, 605-17.
- Keleman, K., Rajagopalan, S., Cleppien, D., Teis, D., Paiha, K., Huber, L. A., Technau, G. M., Dickson, B. J. (2002) Comm sorts robo to control axon guidance at the Drosophila midline. *Cell*, **110**, 415-27.
- Keleman, K., Ribeiro, C., Dickson, B. J. (2005) Comm function in commissural axon guidance: cell-autonomous sorting of Robo in vivo. *Nat Neurosci*, **8**, 156-63.
- Kennedy, T. E., Serafini, T., de la Torre, J. R., Tessier-Lavigne, M. (1994) Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. *Cell*, **78**, 425-35.
- Kerjan, G., Dolan, J., Haumaitre, C., Schneider-Maunoury, S., Fujisawa, H., Mitchell, K. J., Chedotal, A. (2005) The transmembrane semaphorin Sema6A controls cerebellar granule cell migration. *Nat Neurosci*, **8**, 1516-24.
- Kidd, T., Brose, K., Mitchell, K. J., Fetter, R. D., Tessier-Lavigne, M., Goodman, C. S., Tear, G. (1998) Roundabout controls axon crossing of the CNS midline and defines a novel subfamily of evolutionarily conserved guidance receptors. *Cell*, **92**, 205-15.
- Kidd, T., Bland, K. S., Goodman, C. S. (1999) Slit is the midline repellent for the robo receptor in Drosophila. *Cell*, **96**, 785-94.
- Kikuchi, K., Kishino, A., Konishi, O., Kumagai, K., Hosotani, N., Saji, I., Nakayama, C., Kimura, T. (2003) In Vitro and in Vivo Characterization of a Novel Semaphorin 3A Inhibitor, SM-216289 or Xanthofulvin. *J. Biol. Chem.*, **278**, 42985-42991.
- Killeen, M., Tong, J., Krizus, A., Steven, R., Scott, I., Pawson, T., Culotti, J. (2002) UNC-5 function requires phosphorylation of cytoplasmic tyrosine 482, but its UNC-40-independent functions also require a region between the ZU-5 and death domains. *Dev Biol*, **251**, 348-66.
- Kitsukawa, T., Shimono, A., Kawakami, A., Kondoh, H., Fujisawa, H. (1995) Overexpression of a membrane protein, neuropilin, in chimeric mice causes anomalies in the cardiovascular system, nervous system and limbs. *Development*, **121**, 4309-18.
- Kitsukawa, T., Shimizu, M., Sanbo, M., Hirata, T., Taniguchi, M., Bekku, Y., Yagi, T., Fujisawa, H. (1997) Neuropilin-semaphorin III/D-mediated chemorepulsive signals play a crucial role in peripheral nerve projection in mice. *Neuron*, **19**, 995-1005.
- Klein, R. (2004) Eph/ephrin signaling in morphogenesis, neural development and plasticity. *Curr Opin Cell Biol*, **16**, 580-9.
- Klostermann, A., Lohrum, M., Adams, R. H., Puschel, A. W. (1998) The chemorepulsive activity of the axonal guidance signal semaphorin D requires dimerization. *J Biol Chem*, **273**, 7326-31.
- Koeberle, P. D., Bahr, M. (2004) Growth and guidance cues for regenerating axons: where have they gone? *J Neurobiol*, **59**, 162-80.
- Kolodkin, A. L., Matthes, D. J., Goodman, C. S. (1993) The semaphorin genes encode a family of transmembrane and secreted growth cone guidance molecules. *Cell*, **75**, 1389-99.
- Kolodkin, A. L., Levengood, D. V., Rowe, E. G., Tai, Y. T., Giger, R. J., Ginty, D. D. (1997) Neuropilin is a semaphorin III receptor. *Cell*, **90**, 753-62.
- Kolodziej, P. A., Timpe, L. C., Mitchell, K. J., Fried, S. R., Goodman, C. S., Jan, L. Y., Jan, Y. N. (1996) frazzled encodes a Drosophila member of the DCC immunoglobulin subfamily and is required for CNS and motor axon guidance. *Cell*, **87**, 197-204.
- Komatsuzaki, K., Dalvin, S., Kinane, T. B. (2002) Modulation of G(ialpha(2)) signaling by the axonal guidance molecule UNC5H2. *Biochem Biophys Res Commun*, **297**, 898-905.
- Koppel, A. M., Feiner, L., Kobayashi, H., Raper, J. A. (1997) A 70 amino acid region within the semaphorin domain activates specific cellular response of semaphorin family members. *Neuron*, **19**, 531-7.

- Koppel, A. M., Raper, J. A. (1998) Collapsin-1 covalently dimerizes, and dimerization is necessary for collapsing activity. *J Biol Chem*, **273**, 15708-13.
- Koscheck, T., Weyer, A., Schilling, R. L., Schilling, K. (2003) Morphological development and neurochemical differentiation of cerebellar inhibitory interneurons in microexplant cultures. *Neuroscience*, **116**, 973-984.
- Kramer, S. G., Kidd, T., Simpson, J. H., Goodman, C. S. (2001) Switching repulsion to attraction: changing responses to slit during transition in mesoderm migration. *Science*, **292**, 737-40.
- Kruger, R. P., Aurandt, J., Guan, K.-L. (2005) Semaphorins command cells to move. *Nat Rev Mol Cell Biol*, **6**, 789-800.
- Kuramoto, T., Kuwamura, M., Serikawa, T. (2004) Rat neurological mutations cerebellar vermis defect and hobble are caused by mutations in the netrin-1 receptor gene *Unc5h3*. *Molecular Brain Research*, **122**, 103-108.
- Kuwamura, M., Ishida, A., Yamate, J., Kato, K., Kotani, T., Sakuma, S. (1997) Chronological and immunohistochemical observations of cerebellar dysplasia and vermis defect in the hereditary cerebellar vermis defect (CVD) rat. *Acta Neuropathol (Berl)*, **94**, 549-56.
- Laforest, S., Milanini, J., Parat, F., Thimonier, J., Lehmann, M. (2005) Evidences that [beta]1 integrin and Rac1 are involved in the overriding effect of laminin on myelin-associated glycoprotein inhibitory activity on neuronal cells. *Molecular and Cellular Neuroscience*, **30**, 418-428.
- Laine, J., Axelrad, H. (1994) The candelabrum cell: a new interneuron in the cerebellar cortex. *J Comp Neurol*, **339**, 159-73.
- Laine, J., Axelrad, H. (1998) Lugaro cells target basket and stellate cells in the cerebellar cortex. *Neuroreport*, **9**, 2399-403.
- Laine, J., Axelrad, H. (2002) Extending the cerebellar Lugaro cell class. *Neuroscience*, **115**, 363-74.
- Lambert de Rouvroit, C., Goffinet, A. M. (2001) Neuronal migration. *Mech Dev*, **105**, 47-56.
- Lebrand, C., Dent, E. W., Strasser, G. A., Lanier, L. M., Krause, M., Svitkina, T. M., Borisy, G. G., Gertler, F. B. (2004) Critical role of Ena/VASP proteins for filopodia formation in neurons and in function downstream of netrin-1. *Neuron*, **42**, 37-49.
- Lee, J., Li, W., Guan, K. L. (2005) SRC-1 Mediates UNC-5 Signaling in *Caenorhabditis elegans*. *Mol Cell Biol*, **25**, 6485-95.
- Lehmann, M., Fournier, A., Selles-Navarro, I., Dergham, P., Sebok, A., Leclerc, N., Tigyi, G., McKerracher, L. (1999) Inactivation of Rho signaling pathway promotes CNS axon regeneration. *J Neurosci*, **19**, 7537-47.
- Leonardo, E. D., Hinck, L., Masu, M., Keino-Masu, K., Ackerman, S. L., Tessier-Lavigne, M. (1997) Vertebrate homologues of *C. elegans* UNC-5 are candidate netrin receptors. *Nature*, **386**, 833-8.
- Li, D., Field, P. M., Yoshioka, N., Raisman, G. (1994) Axons regenerate with correct specificity in horizontal slice culture of the postnatal rat entorhino-hippocampal system. *Eur J Neurosci*, **6**, 1026-37.
- Li, S., Liu, B. P., Budel, S., Li, M., Ji, B., Walus, L., Li, W., Jirik, A., Rabacchi, S., Choi, E., Worley, D., Sah, D. W., Pepinsky, B., Lee, D., Relton, J., Strittmatter, S. M. (2004) Blockade of Nogo-66, myelin-associated glycoprotein, and oligodendrocyte myelin glycoprotein by soluble Nogo-66 receptor promotes axonal sprouting and recovery after spinal injury. *J Neurosci*, **24**, 10511-20.
- Li, X., Meriane, M., Triki, I., Shekarabi, M., Kennedy, T. E., Larose, L., Lamarche-Vane, N. (2002a) The adaptor protein Nck-1 couples the netrin-1 receptor DCC (deleted in colorectal cancer) to the activation of the small GTPase Rac1 through an atypical mechanism. *J Biol Chem*, **277**, 37788-97.
- Li, X., Saint-Cyr-Proulx, E., Aktories, K., Lamarche-Vane, N. (2002b) Rac1 and Cdc42 but not RhoA or Rho kinase activities are required for neurite outgrowth induced by the Netrin-1 receptor DCC (deleted in colorectal cancer) in N1E-115 neuroblastoma cells. *J Biol Chem*, **277**, 15207-14.
- Liebscher, T., Schnell, L., Schnell, D., Scholl, J., Schneider, R., Gullo, M., Fouad, K., Mir, A., Rausch, M., Kindler, D., Hamers, F. P., Schwab, M. E. (2005) Nogo-A antibody improves regeneration and locomotion of spinal cord-injured rats. *Ann Neurol*, **58**, 706-19.
- Lim, Y. S., Wadsworth, W. G. (2002) Identification of domains of netrin UNC-6 that mediate attractive and repulsive guidance and responses from cells and growth cones. *J Neurosci*, **22**, 7080-7.
- Lin, C. H., Espreafico, E. M., Mooseker, M. S., Forscher, P. (1996) Myosin drives retrograde F-actin flow in neuronal growth cones. *Neuron*, **16**, 769-82.

- Lin, L., Rao, Y., Isacson, O. (2005) Netrin-1 and slit-2 regulate and direct neurite growth of ventral midbrain dopaminergic neurons. *Mol Cell Neurosci*, **28**, 547-55.
- Lindholm, T., Skold, M. K., Suneson, A., Carlstedt, T., Cullheim, S., Risling, M. (2004) Semaphorin and neuropilin expression in motoneurons after intraspinal motoneuron axotomy. *Neuroreport*, **15**, 649-54.
- Liu, G., Beggs, H., Jurgensen, C., Park, H. T., Tang, H., Gorski, J., Jones, K. R., Reichardt, L. F., Wu, J., Rao, Y. (2004) Netrin requires focal adhesion kinase and Src family kinases for axon outgrowth and attraction. *Nat Neurosci*, **7**, 1222-32.
- Liu, X. B., Low, L. K., Jones, E. G., Cheng, H. J. (2005) Stereotyped axon pruning via plexin signaling is associated with synaptic complex elimination in the hippocampus. *J Neurosci*, **25**, 9124-34.
- Livesey, F. J., Hunt, S. P. (1997) Netrin and Netrin Receptor Expression in the Embryonic Mammalian Nervous System Suggests Roles in Retinal, Striatal, Nigral, and Cerebellar Development. *Molecular and Cellular Neuroscience*, **8**, 417-429.
- Llambi, F., Causeret, F., Bloch-Gallego, E., Mehlen, P. (2001) Netrin-1 acts as a survival factor via its receptors UNC5H and DCC. *Embo J*, **20**, 2715-22.
- Llambi, F., Lourenco, F. C., Gozuacik, D., Guix, C., Pays, L., Del Rio, G., Kimchi, A., Mehlen, P. (2005) The dependence receptor UNC5H2 mediates apoptosis through DAP-kinase. *Embo J*, **24**, 1192-201.
- Logan, A., Ahmed, Z., Baird, A., Gonzalez, A. M., Berry, M. (2006) Neurotrophic factor synergy is required for neuronal survival and disinhibited axon regeneration after CNS injury. *Brain*, **129**, 490-502.
- Long, H., Sabatier, C., Ma, L., Plump, A., Yuan, W., Ornitz, D. M., Tamada, A., Murakami, F., Goodman, C. S., Tessier-Lavigne, M. (2004) Conserved roles for Slit and Robo proteins in midline commissural axon guidance. *Neuron*, **42**, 213-23.
- Lu, P., Yang, H., Jones, L. L., Filbin, M. T., Tuszynski, M. H. (2004) Combinatorial therapy with neurotrophins and cAMP promotes axonal regeneration beyond sites of spinal cord injury. *J Neurosci*, **24**, 6402-9.
- Lumsden, A. G., Davies, A. M. (1986) Chemotropic effect of specific target epithelium in the developing mammalian nervous system. *Nature*, **323**, 538-9.
- Luo, Y., Raible, D., Raper, J. A. (1993) Collapsin: a protein in brain that induces the collapse and paralysis of neuronal growth cones. *Cell*, **75**, 217-27.
- Machold, R., Fishell, G. (2005) Math1 is expressed in temporally discrete pools of cerebellar rhombic-lip neural progenitors. *Neuron*, **48**, 17-24.
- Madison, R. D., Zomorodi, A., Robinson, G. A. (2000) Netrin-1 and peripheral nerve regeneration in the adult rat. *Exp Neurol*, **161**, 563-70.
- Madura, T., Yamashita, T., Kubo, T., Fujitani, M., Hosokawa, K., Tohyama, M. (2004) Activation of Rho in the injured axons following spinal cord injury. *EMBO Rep*, **5**, 412-7.
- Magdaleno, S., Keshvara, L., Curran, T. (2002) Rescue of ataxia and preplate splitting by ectopic expression of Reelin in reeler mice. *Neuron*, **33**, 573-86.
- Magyar-Lehmann, S., Suter, C. S., Stahel, W., Schachner, M. (1995) Behaviour of small inhibitory interneurons in early postnatal mouse cerebellar microexplant cultures: a video time-lapse analysis. *Eur J Neurosci*, **7**, 1449-59.
- Maricich, S. M., Herrup, K. (1999) Pax-2 expression defines a subset of GABAergic interneurons and their precursors in the developing murine cerebellum. *J Neurobiol*, **41**, 281-94.
- Marillat, V., Sabatier, C., Failli, V., Matsunaga, E., Sotelo, C., Tessier-Lavigne, M., Chedotal, A. (2004) The slit receptor Rig-1/Robo3 controls midline crossing by hindbrain precerebellar neurons and axons. *Neuron*, **43**, 69-79.
- Marin, O., Rubenstein, J. L. (2001) A long, remarkable journey: tangential migration in the telencephalon. *Nat Rev Neurosci*, **2**, 780-90.
- Marin, O., Yaron, A., Bagri, A., Tessier-Lavigne, M., Rubenstein, J. L. (2001) Sorting of striatal and cortical interneurons regulated by semaphorin-neuropilin interactions. *Science*, **293**, 872-5.
- Marin, O., Plump, A. S., Flames, N., Sanchez-Camacho, C., Tessier-Lavigne, M., Rubenstein, J. L. (2003) Directional guidance of interneuron migration to the cerebral cortex relies on subcortical Slit1/2-independent repulsion and cortical attraction. *Development*, **130**, 1889-901.
- Marin, O., Rubenstein, J. L. (2003) Cell migration in the forebrain. *Annu Rev Neurosci*, **26**, 441-83.

- Martinez, A., Otal, R., Sieber, B. A., Ibanez, C., Soriano, E. (2005) Disruption of ephrin-A/EphA binding alters synaptogenesis and neural connectivity in the hippocampus. *Neuroscience*, **135**, 451-61.
- Martinez, A., Soriano, E. (2005) Functions of ephrin/Eph interactions in the development of the nervous system: emphasis on the hippocampal system. *Brain Res Brain Res Rev*, **49**, 211-26.
- Mathis, C., Collin, L., Borrelli, E. (2003) Oligodendrocyte ablation impairs cerebellum development. *Development*, **130**, 4709-4718.
- McKerracher, L. (2002) Ganglioside rafts as MAG receptors that mediate blockade of axon growth. *PNAS*, **99**, 7811-7813.
- McLaughlin, T., Hindges, R., Yates, P. A., O'Leary, D. D. (2003) Bifunctional action of ephrin-B1 as a repellent and attractant to control bidirectional branch extension in dorsal-ventral retinotopic mapping. *Development*, **130**, 2407-18.
- Mehlen, P., Mazelin, L. (2003) The dependence receptors DCC and UNC5H as a link between neuronal guidance and survival. *Biology of the Cell*, **95**, 425-436.
- Mellitzer, G., Xu, Q., Wilkinson, D. G. (1999) Eph receptors and ephrins restrict cell intermingling and communication. *Nature*, **400**, 77-81.
- Meriane, M., Tcherkezian, J., Webber, C. A., Danek, E. I., Triki, I., McFarlane, S., Bloch-Gallego, E., Lamarche-Vane, N. (2004) Phosphorylation of DCC by Fyn mediates Netrin-1 signaling in growth cone guidance. *J Cell Biol*, **167**, 687-98.
- Merz, D. C., Zheng, H., Killeen, M. T., Krizus, A., Culotti, J. G. (2001) Multiple signaling mechanisms of the UNC-6/netrin receptors UNC-5 and UNC-40/DCC in vivo. *Genetics*, **158**, 1071-80.
- Messersmith, E. K., Leonardo, E. D., Shatz, C. J., Tessier-Lavigne, M., Goodman, C. S., Kolodkin, A. L. (1995) Semaphorin III can function as a selective chemorepellent to pattern sensory projections in the spinal cord. *Neuron*, **14**, 949-59.
- Metin, C., Deleglise, D., Serafini, T., Kennedy, T. E., Tessier-Lavigne, M. (1997) A role for netrin-1 in the guidance of cortical efferents. *Development*, **124**, 5063-74.
- Meyer, G., Feldman, E. L. (2002) Signaling mechanisms that regulate actin-based motility processes in the nervous system. *J Neurochem*, **83**, 490-503.
- Mi, S., Lee, X., Shao, Z., Thill, G., Ji, B., Relton, J., Levesque, M., Allaire, N., Perrin, S., Sands, B., Crowell, T., Cate, R. L., McCoy, J. M., Pepinsky, R. B. (2004) LINGO-1 is a component of the Nogo-66 receptor/p75 signaling complex. *Nat Neurosci*, **7**, 221-8.
- Mi, S., Miller, R. H., Lee, X., Scott, M. L., Shulag-Morskaya, S., Shao, Z., Chang, J., Thill, G., Levesque, M., Zhang, M., Hession, C., Sah, D., Trapp, B., He, Z., Jung, V., McCoy, J. M., Pepinsky, R. B. (2005) LINGO-1 negatively regulates myelination by oligodendrocytes. *Nat Neurosci*, **8**, 745-51.
- Miao, H. Q., Soker, S., Feiner, L., Alonso, J. L., Raper, J. A., Klagsbrun, M. (1999) Neuropilin-1 mediates collapsin-1/semaphorin III inhibition of endothelial cell motility: functional competition of collapsin-1 and vascular endothelial growth factor-165. *J Cell Biol*, **146**, 233-42.
- Mikule, K., Gatlin, J. C., de la Houssaye, B. A., Pfenninger, K. H. (2002) Growth cone collapse induced by semaphorin 3A requires 12/15-lipoxygenase. *J Neurosci*, **22**, 4932-41.
- Ming, G., Song, H., Berninger, B., Inagaki, N., Tessier-Lavigne, M., Poo, M. (1999) Phospholipase C-gamma and phosphoinositide 3-kinase mediate cytoplasmic signaling in nerve growth cone guidance. *Neuron*, **23**, 139-48.
- Ming, G. L., Song, H. J., Berninger, B., Holt, C. E., Tessier-Lavigne, M., Poo, M. M. (1997) cAMP-dependent growth cone guidance by netrin-1. *Neuron*, **19**, 1225-35.
- Ming, G. L., Wong, S. T., Henley, J., Yuan, X. B., Song, H. J., Spitzer, N. C., Poo, M. M. (2002) Adaptation in the chemotactic guidance of nerve growth cones. *Nature*, **417**, 411-8.
- Mingorance, A., Fontana, X., Sole, M., Burgaya, F., Urena, J. M., Teng, F. Y., Tang, B. L., Hunt, D., Anderson, P. N., Bethea, J. R., Schwab, M. E., Soriano, E., del Rio, J. A. (2004) Regulation of Nogo and Nogo receptor during the development of the entorhino-hippocampal pathway and after adult hippocampal lesions. *Mol Cell Neurosci*, **26**, 34-49.
- Mingorance, A., Fontana, X., Soriano, E., Del Rio, J. A. (2005) Overexpression of myelin-associated glycoprotein after axotomy of the perforant pathway. *Mol Cell Neurosci*, **29**, 471-83.
- Mingorance, A., Sole, M., Muneton, V., Martinez, A., Nieto-Sampedro, M., Soriano, E., Del Rio, J. A. (2006) Regeneration of lesioned entorhino-hippocampal axons in vitro by combined degradation of inhibitory proteoglycans and blockade of Nogo-66/NgR signaling. *Faseb J*.



- Mitsui, N., Inatome, R., Takahashi, S., Goshima, Y., Yamamura, H., Yanagi, S. (2002) Involvement of Fes/Fps tyrosine kinase in semaphorin3A signaling. *Embo J*, **21**, 3274-85.
- Miyata, T., Nakajima, K., Aruga, J., Takahashi, S., Ikenaka, K., Mikoshiba, K., Ogawa, M. (1996) Distribution of a reeler gene-related antigen in the developing cerebellum: an immunohistochemical study with an allogeneic antibody CR-50 on normal and reeler mice. *J Comp Neurol*, **372**, 215-28.
- Miyata, T., Nakajima, K., Mikoshiba, K., Ogawa, M. (1997) Regulation of Purkinje cell alignment by reelin as revealed with CR-50 antibody. *J Neurosci*, **17**, 3599-609.
- Mjaatvedt, A. E., Cabin, D. E., Cole, S. E., Long, L. J., Breitwieser, G. E., Reeves, R. H. (1995) Assessment of a mutation in the H5 domain of Girk2 as a candidate for the weaver mutation. *Genome Res*, **5**, 453-63.
- Monnier, P. P., Sierra, A., Schwab, J. M., Henke-Fahle, S., Mueller, B. K. (2003) The Rho/ROCK pathway mediates neurite growth-inhibitory activity associated with the chondroitin sulfate proteoglycans of the CNS glial scar. *Mol Cell Neurosci*, **22**, 319-30.
- Moreau-Fauvarque, C., Kumanogoh, A., Camand, E., Jaillard, C., Barbin, G., Boquet, I., Love, C., Jones, E. Y., Kikutani, H., Lubetzki, C., Dusart, I., Chedotal, A. (2003) The Transmembrane Semaphorin Sema4D/CD100, an Inhibitor of Axonal Growth, Is Expressed on Oligodendrocytes and Upregulated after CNS Lesion. *J. Neurosci.*, **23**, 9229-9239.
- Moreno-Flores, M. T., Martin-Aparicio, E., Avila, J., Diaz-Nido, J., Wandosell, F. (2002) Ephrin-B1 Promotes Dendrite Outgrowth on Cerebellar Granule Neurons. *Molecular and Cellular Neuroscience*, **20**, 429-446.
- Nakajima, K., Mikoshiba, K., Miyata, T., Kudo, C., Ogawa, M. (1997) Disruption of hippocampal development in vivo by CR-50 mAb against reelin. *Proc Natl Acad Sci U S A*, **94**, 8196-201.
- Nakamoto, T., Kain, K. H., Ginsberg, M. H. (2004) Neurobiology: New connections between integrins and axon guidance. *Curr Biol*, **14**, R121-3.
- Nakamura, F., Tanaka, M., Takahashi, T., Kalb, R. G., Strittmatter, S. M. (1998) Neuropilin-1 extracellular domains mediate semaphorin D/III-induced growth cone collapse. *Neuron*, **21**, 1093-100.
- Negishi, M., Oinuma, I., Katoh, H. (2005a) Plexins: axon guidance and signal transduction. *Cell Mol Life Sci*, **62**, 1363-71.
- Negishi, M., Oinuma, I., Katoh, H. (2005b) R-ras as a key player for signaling pathway of plexins. *Mol Neurobiol*, **32**, 217-22.
- Neumann, S., Bradke, F., Tessier-Lavigne, M., Basbaum, A. I. (2002) Regeneration of sensory axons within the injured spinal cord induced by intraganglionic cAMP elevation. *Neuron*, **34**, 885-93.
- Nguyen Ba-Charvet, K. T., Brose, K., Marillat, V., Kidd, T., Goodman, C. S., Tessier-Lavigne, M., Sotelo, C., Chedotal, A. (1999) Slit2-Mediated chemorepulsion and collapse of developing forebrain axons. *Neuron*, **22**, 463-73.
- Niclou, S. P., Franssen, E. H., Ehlert, E. M., Taniguchi, M., Verhaagen, J. (2003) Meningeal cell-derived semaphorin 3A inhibits neurite outgrowth. *Mol Cell Neurosci*, **24**, 902-12.
- Niederost, B., Oertle, T., Fritsche, J., McKinney, R. A., Bandtlow, C. E. (2002) Nogo-A and Myelin-Associated Glycoprotein Mediate Neurite Growth Inhibition by Antagonistic Regulation of RhoA and Rac1. *J. Neurosci.*, **22**, 10368-10376.
- Nieuwenhuys, R., Donkelaar, H. J. T., Nicholson, C. (1998) *The Central Nervous System of Vertebrates*. Springer cop., Berlin-Barcelona.
- Nishihara, E., Yoshida-Komiya, H., Chan, C. S., Liao, L., Davis, R. L., O'Malley, B. W., Xu, J. (2003) SRC-1 null mice exhibit moderate motor dysfunction and delayed development of cerebellar Purkinje cells. *J Neurosci*, **23**, 213-22.
- Nishio, Y., Koda, M., Kitajo, K., Seto, M., Hata, K., Taniguchi, J., Moriya, H., Fujitani, M., Kubo, T., Yamashita, T. (2006) Delayed treatment with Rho-kinase inhibitor does not enhance axonal regeneration or functional recovery after spinal cord injury in rats. *Exp Neurol*.
- Nishiyama, A., Yang, Z., Butt, A. (2005) Astrocytes and NG2-glia: what's in a name? *Journal of Anatomy*, **207**, 687-693.
- Nishiyama, M., Hoshino, A., Tsai, L., Henley, J. R., Goshima, Y., Tessier-Lavigne, M., Poo, M. M., Hong, K. (2003) Cyclic AMP/GMP-dependent modulation of Ca<sup>2+</sup> channels sets the polarity of nerve growth-cone turning. *Nature*, **423**, 990-5.

- Niu, S., Renfro, A., Quattrocchi, C. C., Sheldon, M., D'Arcangelo, G. (2004) Reelin promotes hippocampal dendrite development through the VLDLR/ApoER2-Dab1 pathway. *Neuron*, **41**, 71-84.
- Nobes, C. D., Lauritzen, I., Mattei, M. G., Paris, S., Hall, A., Chardin, P. (1998) A new member of the Rho family, Rnd1, promotes disassembly of actin filament structures and loss of cell adhesion. *J Cell Biol*, **141**, 187-97.
- Ohta, K., Takagi, S., Asou, H., Fujisawa, H. (1992) Involvement of neuronal cell surface molecule B2 in the formation of retinal plexiform layers. *Neuron*, **9**, 151-61.
- Ohta, K., Mizutani, A., Kawakami, A., Murakami, Y., Kasuya, Y., Takagi, S., Tanaka, H., Fujisawa, H. (1995) Plexin: a novel neuronal cell surface molecule that mediates cell adhesion via a homophilic binding mechanism in the presence of calcium ions. *Neuron*, **14**, 1189-99.
- Oinuma, I., Ishikawa, Y., Katoh, H., Negishi, M. (2004) The Semaphorin 4D receptor Plexin-B1 is a GTPase activating protein for R-Ras. *Science*, **305**, 862-5.
- Oiwa, K., Sakakibara, H. (2005) Recent progress in dynein structure and mechanism. *Curr Opin Cell Biol*, **17**, 98-103.
- Okuyama-Yamamoto, A., Yamamoto, T., Miki, A., Terashima, T. (2005) Changes in reelin expression in the mouse olfactory bulb after chemical lesion to the olfactory epithelium. *Eur J Neurosci*, **21**, 2586-92.
- Osborne, N. J., Begbie, J., Chilton, J. K., Schmidt, H., Eickholt, B. J. (2005) Semaphorin/neuropilin signaling influences the positioning of migratory neural crest cells within the hindbrain region of the chick. *Dev Dyn*, **232**, 939-49.
- O'Shea, K., Rheinheimer, J., Dixit, V. (1990) Deposition and role of thrombospondin in the histogenesis of the cerebellar cortex. *J Cell Biol*, **110**, 1275-1283.
- Pantaloni, D., Carlier, M. F. (1993) How profilin promotes actin filament assembly in the presence of thymosin beta 4. *Cell*, **75**, 1007-14.
- Pappas, G. D., Kriho, V., Pesold, C. (2001) Reelin in the extracellular matrix and dendritic spines of the cortex and hippocampus: a comparison between wild type and heterozygous reeler mice by immunoelectron microscopy. *J Neurocytol*, **30**, 413-25.
- Park, J. B., Yiu, G., Kaneko, S., Wang, J., Chang, J., He, X. L., Garcia, K. C., He, Z. (2005) A TNF receptor family member, TROY, is a coreceptor with Nogo receptor in mediating the inhibitory activity of myelin inhibitors. *Neuron*, **45**, 345-51.
- Pascual, M., Perez-Sust, P., Soriano, E. (2004a) The GABAergic septohippocampal pathway in control and reeler mice: target specificity and termination onto Reelin-expressing interneurons. *Mol Cell Neurosci*, **25**, 679-91.
- Pascual, M., Pozas, E., Barallobre, M. J., Tessier-Lavigne, M., Soriano, E. (2004b) Coordinated functions of Netrin-1 and Class 3 secreted Semaphorins in the guidance of reciprocal septohippocampal connections. *Molecular and Cellular Neuroscience*, **26**, 24-33.
- Pascual, M., Pozas, E., Soriano, E. (2005) Role of class 3 semaphorins in the development and maturation of the septohippocampal pathway. *Hippocampus*, **15**, 184-202.
- Pasterkamp, R. J., Giger, R. J., Verhaagen, J. (1998a) Regulation of semaphorin III/collapsin-1 gene expression during peripheral nerve regeneration. *Exp Neurol*, **153**, 313-27.
- Pasterkamp, R. J., De Winter, F., Holtmaat, A. J., Verhaagen, J. (1998b) Evidence for a role of the chemorepellent semaphorin III and its receptor neuropilin-1 in the regeneration of primary olfactory axons. *J Neurosci*, **18**, 9962-76.
- Pasterkamp, R. J., Giger, R. J., Ruitenber, M. J., Holtmaat, A. J., De Wit, J., De Winter, F., Verhaagen, J. (1999) Expression of the gene encoding the chemorepellent semaphorin III is induced in the fibroblast component of neural scar tissue formed following injuries of adult but not neonatal CNS. *Mol Cell Neurosci*, **13**, 143-66.
- Pasterkamp, R. J., Giger, R. J., Baker, R. E., Hermens, W. T., Verhaagen, J. (2000) Ectopic adenoviral vector-directed expression of Sema3A in organotypic spinal cord explants inhibits growth of primary sensory afferents. *Dev Biol*, **220**, 129-41.
- Pasterkamp, R. J., Anderson, P. N., Verhaagen, J. (2001) Peripheral nerve injury fails to induce growth of lesioned ascending dorsal column axons into spinal cord scar tissue expressing the axon repellent Semaphorin3A. *Eur J Neurosci*, **13**, 457-71.
- Pasterkamp, R. J., Kolodkin, A. L. (2003) Semaphorin junction: making tracks toward neural connectivity. *Curr Opin Neurobiol*, **13**, 79-89.

- Pasterkamp, R. J., Peschon, J. J., Spriggs, M. K., Kolodkin, A. L. (2003) Semaphorin 7A promotes axon outgrowth through integrins and MAPKs. *Nature*, **424**, 398-405.
- Pesold, C., Impagnatiello, F., Pisu, M. G., Uzunov, D. P., Costa, E., Guidotti, A., Caruncho, H. J. (1998) Reelin is preferentially expressed in neurons synthesizing gamma-aminobutyric acid in cortex and hippocampus of adult rats. *Proc Natl Acad Sci U S A*, **95**, 3221-6.
- Piper, M., Salih, S., Weinl, C., Holt, C. E., Harris, W. A. (2005) Endocytosis-dependent desensitization and protein synthesis-dependent resensitization in retinal growth cone adaptation. *Nat Neurosci*, **8**, 179-86.
- Pleasure, S. J., Anderson, S., Hevner, R., Bagri, A., Marin, O., Lowenstein, D. H., Rubenstein, J. L. (2000) Cell migration from the ganglionic eminences is required for the development of hippocampal GABAergic interneurons. *Neuron*, **28**, 727-40.
- Polleux, F., Giger, R. J., Ginty, D. D., Kolodkin, A. L., Ghosh, A. (1998) Patterning of cortical efferent projections by semaphorin-neuropilin interactions. *Science*, **282**, 1904-6.
- Polleux, F., Morrow, T., Ghosh, A. (2000) Semaphorin 3A is a chemoattractant for cortical apical dendrites. *Nature*, **404**, 567-73.
- Pozas, E., Pascual, M., Nguyen Ba-Charvet, K. T., Guijarro, P., Sotelo, C., Chedotal, A., Del Rio, J. A., Soriano, E. (2001) Age-dependent effects of secreted Semaphorins 3A, 3F, and 3E on developing hippocampal axons: in vitro effects and phenotype of Semaphorin 3A (-/-) mice. *Mol Cell Neurosci*, **18**, 26-43.
- Prang, P., Del Turco, D., Kapfhammer, J. P. (2001) Regeneration of entorhinal fibers in mouse slice cultures is age dependent and can be stimulated by NT-4, GDNF, and modulators of G-proteins and protein kinase C. *Exp Neurol*, **169**, 135-47.
- Przyborski, S. A., Knowles, B. B., Ackerman, S. L. (1998) Embryonic phenotype of Unc5h3 mutant mice suggests chemorepulsion during the formation of the rostral cerebellar boundary. *Development*, **125**, 41-50.
- Qiu, J., Cai, D., Dai, H., McAtee, M., Hoffman, P. N., Bregman, B. S., Filbin, M. T. (2002) Spinal axon regeneration induced by elevation of cyclic AMP. *Neuron*, **34**, 895-903.
- Qiu, S., Korwek, K. M., Pratt-Davis, A. R., Peters, M., Bergman, M. Y., Weeber, E. J. (2005) Cognitive disruption and altered hippocampus synaptic function in Reelin haploinsufficient mice. *Neurobiol Learn Mem*.
- Rabacchi, S. A., Solowska, J. M., Kruk, B., Luo, Y., Raper, J. A., Baird, D. H. (1999) Collapsin-1/semaphorin-III/D is regulated developmentally in Purkinje cells and collapses pontocerebellar mossy fiber neuronal growth cones. *J Neurosci*, **19**, 4437-48.
- Rajagopalan, S., Vivancos, V., Nicolas, E., Dickson, B. J. (2000) Selecting a longitudinal pathway: Robo receptors specify the lateral position of axons in the Drosophila CNS. *Cell*, **103**, 1033-45.
- Ralevic, V., Burnstock, G. (1998) Receptors for Purines and Pyrimidines. *Pharmacol Rev*, **50**, 413-492.
- Ren, X. R., Ming, G. L., Xie, Y., Hong, Y., Sun, D. M., Zhao, Z. Q., Feng, Z., Wang, Q., Shim, S., Chen, Z. F., Song, H. J., Mei, L., Xiong, W. C. (2004) Focal adhesion kinase in netrin-1 signaling. *Nat Neurosci*, **7**, 1204-12.
- Renzi, M. J., Feiner, L., Koppel, A. M., Raper, J. A. (1999) A dominant negative receptor for specific secreted semaphorins is generated by deleting an extracellular domain from neuropilin-1. *J Neurosci*, **19**, 7870-80.
- Rice, D. S., Curran, T. (2001) Role of the reelin signaling pathway in central nervous system development. *Annu Rev Neurosci*, **24**, 1005-39.
- Rio, C., Rieff, H. I., Qi, P., Khurana, T. S., Corfas, G. (1997) Neuregulin and erbB receptors play a critical role in neuronal migration. *Neuron*, **19**, 39-50.
- Rodger, J., Lindsey, K. A., Leaver, S. G., King, C. E., Dunlop, S. A., Beazley, L. D. (2001) Expression of ephrin-A2 in the superior colliculus and EphA5 in the retina following optic nerve section in adult rat. *Eur J Neurosci*, **14**, 1929-36.
- Rodger, J., Goto, H., Cui, Q., Chen, P. B., Harvey, A. R. (2005) cAMP regulates axon outgrowth and guidance during optic nerve regeneration in goldfish. *Mol Cell Neurosci*, **30**, 452-64.
- Rohm, B., Ottemeyer, A., Lohrum, M., Puschel, A. W. (2000a) Plexin/neuropilin complexes mediate repulsion by the axonal guidance signal semaphorin 3A. *Mech Dev*, **93**, 95-104.
- Rohm, B., Rahim, B., Kleiber, B., Hovatta, I., Puschel, A. W. (2000b) The semaphorin 3A receptor may directly regulate the activity of small GTPases. *FEBS Lett*, **486**, 68-72.

- Rosenstein, J. M., Krum, J. M. (2004) New roles for VEGF in nervous tissue--beyond blood vessels. *Exp Neurol*, **187**, 246-53.
- Rothberg, J. M., Jacobs, J. R., Goodman, C. S., Artavanis-Tsakonas, S. (1990) slit: an extracellular protein necessary for development of midline glia and commissural axon pathways contains both EGF and LRR domains. *Genes Dev*, **4**, 2169-87.
- Sabatier, C., Plump, A. S., Le, M., Brose, K., Tamada, A., Murakami, F., Lee, E. Y., Tessier-Lavigne, M. (2004) The divergent Robo family protein rig-1/Robo3 is a negative regulator of slit responsiveness required for midline crossing by commissural axons. *Cell*, **117**, 157-69.
- Sahay, A., Molliver, M. E., Ginty, D. D., Kolodkin, A. L. (2003) Semaphorin 3F is critical for development of limbic system circuitry and is required in neurons for selective CNS axon guidance events. *J Neurosci*, **23**, 6671-80.
- Salazar, M. A., Kwiatkowski, A. V., Pellegrini, L., Cestra, G., Butler, M. H., Rossman, K. L., Serna, D. M., Sondek, J., Gertler, F. B., De Camilli, P. (2003) Tuba, a novel protein containing bin/amphiphysin/Rvs and Dbl homology domains, links dynamin to regulation of the actin cytoskeleton. *J Biol Chem*, **278**, 49031-43.
- Sandvig, A., Berry, M., Barrett, L. B., Butt, A., Logan, A. (2004) Myelin-, reactive glia-, and scar-derived CNS axon growth inhibitors: expression, receptor signaling, and correlation with axon regeneration. *Glia*, **46**, 225-51.
- Sang, Q., Wu, J., Rao, Y., Hsueh, Y. P., Tan, S. S. (2002) Slit promotes branching and elongation of neurites of interneurons but not projection neurons from the developing telencephalon. *Mol Cell Neurosci*, **21**, 250-65.
- Sasaki, Y., Cheng, C., Uchida, Y., Nakajima, O., Ohshima, T., Yagi, T., Taniguchi, M., Nakayama, T., Kishida, R., Kudo, Y., Ohno, S., Nakamura, F., Goshima, Y. (2002) Fyn and Cdk5 mediate semaphorin-3A signaling, which is involved in regulation of dendrite orientation in cerebral cortex. *Neuron*, **35**, 907-20.
- Satoda, M., Takagi, S., Ohta, K., Hirata, T., Fujisawa, H. (1995) Differential expression of two cell surface proteins, neuropilin and plexin, in *Xenopus* olfactory axon subclasses. *J Neurosci*, **15**, 942-55.
- Scarlato, M., Ara, J., Bannerman, P., Scherer, S., Pleasure, D. (2003) Induction of neuropilins-1 and -2 and their ligands, Sema3A, Sema3F, and VEGF, during Wallerian degeneration in the peripheral nervous system. *Exp Neurol*, **183**, 489-98.
- Schilling, K. (2000) Lineage, development and morphogenesis of cerebellar interneurons. *Prog Brain Res*, **124**, 51-68.
- Schwab, J. M., Tuli, S. K., Failli, V. (2006) The Nogo receptor complex: confining molecules to molecular mechanisms. *Trends Mol Med*.
- Schwab, M. E., Bartholdi, D. (1996) Degeneration and regeneration of axons in the lesioned spinal cord. *Physiol Rev*, **76**, 319-70.
- Schwarz, Q., Gu, C., Fujisawa, H., Sabelko, K., Gertsenstein, M., Nagy, A., Taniguchi, M., Kolodkin, A. L., Ginty, D. D., Shima, D. T., Ruhrberg, C. (2004) Vascular endothelial growth factor controls neuronal migration and cooperates with Sema3A to pattern distinct compartments of the facial nerve. *Genes Dev*, **18**, 2822-34.
- Schweigreiter, R., Walmsley, A. R., Niederost, B., Zimmermann, D. R., Oertle, T., Casademunt, E., Frentzel, S., Dechant, G., Mir, A., Bandtlow, C. E. (2004) Versican V2 and the central inhibitory domain of Nogo-A inhibit neurite growth via p75NTR/NgR-independent pathways that converge at RhoA. *Mol Cell Neurosci*, **27**, 163-74.
- Sechi, A. S., Wehland, J. (2004) ENA/VASP proteins: multifunctional regulators of actin cytoskeleton dynamics. *Front Biosci*, **9**, 1294-310.
- Seeger, M., Tear, G., Ferres-Marco, D., Goodman, C. S. (1993) Mutations affecting growth cone guidance in *Drosophila*: genes necessary for guidance toward or away from the midline. *Neuron*, **10**, 409-26.
- Serafini, T., Kennedy, T. E., Galko, M. J., Mirzayan, C., Jessell, T. M., Tessier-Lavigne, M. (1994) The netrins define a family of axon outgrowth-promoting proteins homologous to *C. elegans* UNC-6. *Cell*, **78**, 409-24.
- Serafini, T., Colamarino, S. A., Leonardo, E. D., Wang, H., Bedington, R., Skarnes, W. C., Tessier-Lavigne, M. (1996) Netrin-1 is required for commissural axon guidance in the developing vertebrate nervous system. *Cell*, **87**, 1001-14.

- Serini, G., Valdembri, D., Zanivan, S., Morterra, G., Burkhardt, C., Caccavari, F., Zammataro, L., Primo, L., Tamagnone, L., Logan, M., Tessier-Lavigne, M., Taniguchi, M., Puschel, A. W., Bussolino, F. (2003) Class 3 semaphorins control vascular morphogenesis by inhibiting integrin function. *Nature*, **424**, 391-7.
- Shearer, M. C., Niclou, S. P., Brown, D., Asher, R. A., Holtmaat, A. J. G. D., Levine, J. M., Verhaagen, J., Fawcett, J. W. (2003) The astrocyte/meningeal cell interface is a barrier to neurite outgrowth which can be overcome by manipulation of inhibitory molecules or axonal signalling pathways. *Molecular and Cellular Neuroscience*, **24**, 913-925.
- Shekarabi, M., Kennedy, T. E. (2002) The netrin-1 receptor DCC promotes filopodia formation and cell spreading by activating Cdc42 and Rac1. *Mol Cell Neurosci*, **19**, 1-17.
- Shekarabi, M., Moore, S. W., Tritsch, N. X., Morris, S. J., Bouchard, J. F., Kennedy, T. E. (2005) Deleted in colorectal cancer binding netrin-1 mediates cell substrate adhesion and recruits Cdc42, Rac1, Pak1, and N-WASP into an intracellular signaling complex that promotes growth cone expansion. *J Neurosci*, **25**, 3132-41.
- Shewan, D., Dwivedy, A., Anderson, R., Holt, C. E. (2002) Age-related changes underlie switch in netrin-1 responsiveness as growth cones advance along visual pathway. *Nat Neurosci*, **5**, 955-62.
- Shim, S., Goh, E. L., Ge, S., Sailor, K., Yuan, J. P., Roderick, H. L., Bootman, M. D., Worley, P. F., Song, H., Ming, G. L. (2005) XTRPC1-dependent chemotropic guidance of neuronal growth cones. *Nat Neurosci*, **8**, 730-5.
- Shimizu, M., Murakami, Y., Suto, F., Fujisawa, H. (2000) Determination of cell adhesion sites of neuropilin-1. *J Cell Biol*, **148**, 1283-93.
- Shirasaki, R., Tamada, A., Katsumata, R., Murakami, F. (1995) Guidance of cerebellofugal axons in the rat embryo: directed growth toward the floor plate and subsequent elongation along the longitudinal axis. *Neuron*, **14**, 961-72.
- Shirasaki, R., Katsumata, R., Murakami, F. (1998) Change in chemoattractant responsiveness of developing axons at an intermediate target. *Science*, **279**, 105-7.
- Shirvan, A., Ziv, I., Fleminger, G., Shina, R., He, Z., Brudo, I., Melamed, E., Barzilai, A. (1999) Semaphorins as mediators of neuronal apoptosis. *J Neurochem*, **73**, 961-71.
- Shirvan, A., Kimron, M., Holdengreber, V., Ziv, I., Ben-Shaul, Y., Melamed, S., Melamed, E., Barzilai, A., Solomon, A. S. (2002) Anti-semaphorin 3A antibodies rescue retinal ganglion cells from cell death following optic nerve axotomy. *J Biol Chem*, **277**, 49799-807.
- Sicotte, M., Tsatas, O., Jeong, S. Y., Cai, C. Q., He, Z., David, S. (2003) Immunization with myelin or recombinant Nogo-66/MAG in alum promotes axon regeneration and sprouting after corticospinal tract lesions in the spinal cord. *Mol Cell Neurosci*, **23**, 251-63.
- Silver, J., Miller, J. H. (2004) Regeneration beyond the glial scar. *Nat Rev Neurosci*, **5**, 146-56.
- Simpson, J. H., Bland, K. S., Fetter, R. D., Goodman, C. S. (2000a) Short-range and long-range guidance by Slit and its Robo receptors: a combinatorial code of Robo receptors controls lateral position. *Cell*, **103**, 1019-32.
- Simpson, J. H., Kidd, T., Bland, K. S., Goodman, C. S. (2000b) Short-range and long-range guidance by slit and its Robo receptors. Robo and Robo2 play distinct roles in midline guidance. *Neuron*, **28**, 753-66.
- Sinagra, M., Verrier, D., Frankova, D., Korwek, K. M., Blahos, J., Weeber, E. J., Manzoni, O. J., Chavis, P. (2005) Reelin, very-low-density lipoprotein receptor, and apolipoprotein E receptor 2 control somatic NMDA receptor composition during hippocampal maturation in vitro. *J Neurosci*, **25**, 6127-36.
- Sivasankaran, R., Pei, J., Wang, K. C., Zhang, Y. P., Shields, C. B., Xu, X. M., He, Z. (2004) PKC mediates inhibitory effects of myelin and chondroitin sulfate proteoglycans on axonal regeneration. *Nat Neurosci*, **7**, 261-8.
- Skaliora, I., Singer, W., Betz, H., Puschel, A. W. (1998) Differential patterns of semaphorin expression in the developing rat brain. *Eur J Neurosci*, **10**, 1215-29.
- Snider, W. D., Zhou, F. Q., Zhong, J., Markus, A. (2002) Signaling the pathway to regeneration. *Neuron*, **35**, 13-6.
- Soker, S., Takashima, S., Miao, H. Q., Neufeld, G., Klagsbrun, M. (1998) Neuropilin-1 is expressed by endothelial and tumor cells as an isoform-specific receptor for vascular endothelial growth factor. *Cell*, **92**, 735-45.

- Soker, S., Miao, H. Q., Nomi, M., Takashima, S., Klagsbrun, M. (2002) VEGF165 mediates formation of complexes containing VEGFR-2 and neuropilin-1 that enhance VEGF165-receptor binding. *J Cell Biochem*, **85**, 357-68.
- Song, H., Ming, G., He, Z., Lehmann, M., McKerracher, L., Tessier-Lavigne, M., Poo, M. (1998) Conversion of neuronal growth cone responses from repulsion to attraction by cyclic nucleotides. *Science*, **281**, 1515-8.
- Song, H., Poo, M. (2001) The cell biology of neuronal navigation. *Nat Cell Biol*, **3**, E81-8.
- Song, H. J., Poo, M. M. (1999) Signal transduction underlying growth cone guidance by diffusible factors. *Curr Opin Neurobiol*, **9**, 355-63.
- Soriano, E., Cobas, A., Fairen, A. (1986) Asynchronism in the neurogenesis of GABAergic and non-GABAergic neurons in the mouse hippocampus. *Brain Res*, **395**, 88-92.
- Soriano, E., Cobas, A., Fairen, A. (1989a) Neurogenesis of glutamic acid decarboxylase immunoreactive cells in the hippocampus of the mouse. I: Regio superior and regio inferior. *J Comp Neurol*, **281**, 586-602.
- Soriano, E., Cobas, A., Fairen, A. (1989b) Neurogenesis of glutamic acid decarboxylase immunoreactive cells in the hippocampus of the mouse. II: Area dentata. *J Comp Neurol*, **281**, 603-11.
- Soriano, E., Del Rio, J. A., Martinez, A., Super, H. (1994) Organization of the embryonic and early postnatal murine hippocampus. I. Immunocytochemical characterization of neuronal populations in the subplate and marginal zone. *J Comp Neurol*, **342**, 571-95.
- Soriano, E., Del Rio, J. A. (2005) The cells of cajal-retzius: still a mystery one century after. *Neuron*, **46**, 389-94.
- Soriano, P. (1999) Generalized lacZ expression with the ROSA26 Cre reporter strain. **21**, 70-71.
- Sotelo, C. (2004) Cellular and genetic regulation of the development of the cerebellar system. *Progress in Neurobiology*, **72**, 295-339.
- Spassky, N., de Castro, F., Le Bras, B., Heydon, K., Queraud-LeSaux, F., Bloch-Gallego, E., Chedotal, A., Zalc, B., Thomas, J. L. (2002) Directional guidance of oligodendroglial migration by class 3 semaphorins and netrin-1. *J Neurosci*, **22**, 5992-6004.
- Sperry, R. W. (1963) Chemoaffinity in the Orderly Growth of Nerve Fiber Patterns and Connections. *Proc Natl Acad Sci U S A*, **50**, 703-10.
- Standfield, B. B., Cowan, W. M. (1988) The Development of the Hippocampal Region. In *Development and Maturation of Cerebral Cortex*. Plenum Press, New York, Vol. 7.
- Stein, E., Savaskan, N. E., Ninnemann, O., Nitsch, R., Zhou, R., Skutella, T. (1999) A role for the Eph ligand ephrin-A3 in entorhino-hippocampal axon targeting. *J Neurosci*, **19**, 8885-93.
- Stein, E., Tessier-Lavigne, M. (2001) Hierarchical organization of guidance receptors: silencing of netrin attraction by slit through a Robo/DCC receptor complex. *Science*, **291**, 1928-38.
- Stein, E., Zou, Y., Poo, M., Tessier-Lavigne, M. (2001) Binding of DCC by netrin-1 to mediate axon guidance independent of adenosine A2B receptor activation. *Science*, **291**, 1976-82.
- Steinmayr, M., Andre, E., Conquet, F., Rondi-Reig, L., Delhaye-Bouchaud, N., Auclair, N., Daniel, H., Crepel, F., Mariani, J., Sotelo, C., Becker-Andre, M. (1998) staggerer phenotype in retinoid-related orphan receptor alpha-deficient mice. *Proc Natl Acad Sci U S A*, **95**, 3960-5.
- Steup, A., Ninnemann, O., Savaskan, N. E., Nitsch, R., Puschel, A. W., Skutella, T. (1999) Semaphorin D acts as a repulsive factor for entorhinal and hippocampal neurons. *Eur J Neurosci*, **11**, 729-34.
- Steup, A., Lohrum, M., Hamscho, N., Savaskan, N. E., Ninnemann, O., Nitsch, R., Fujisawa, H., Puschel, A. W., Skutella, T. (2000) Sema3C and netrin-1 differentially affect axon growth in the hippocampal formation. *Mol Cell Neurosci*, **15**, 141-55.
- Stoppini, L., Buchs, P. A., Muller, D. (1991) A simple method for organotypic cultures of nervous tissue. *J Neurosci Methods*, **37**, 173-82.
- Suchting, S., Bicknell, R., Eichmann, A. (2005) Neuronal clues to vascular guidance. *Exp Cell Res*.
- Sugimoto, Y., Taniguchi, M., Yagi, T., Akagi, Y., Nojyo, Y., Tamamaki, N. (2001) Guidance of glial precursor cell migration by secreted cues in the developing optic nerve. *Development*, **128**, 3321-30.
- Sultan, F., Konig, T., Mock, M., Thier, P. (2002) Quantitative organization of neurotransmitters in the deep cerebellar nuclei of the Lurcher mutant. *J Comp Neurol*, **452**, 311-23.

- Super, H., Soriano, E. (1994) The organization of the embryonic and early postnatal murine hippocampus. II. Development of entorhinal, commissural, and septal connections studied with the lipophilic tracer DiI. *J Comp Neurol*, **344**, 101-20.
- Super, H., Martinez, A., Del Rio, J. A., Soriano, E. (1998a) Involvement of distinct pioneer neurons in the formation of layer-specific connections in the hippocampus. *J Neurosci*, **18**, 4616-26.
- Super, H., Soriano, E., Uylings, H. B. (1998b) The functions of the preplate in development and evolution of the neocortex and hippocampus. *Brain Res Brain Res Rev*, **27**, 40-64.
- Suto, F., Ito, K., Uemura, M., Shimizu, M., Shinkawa, Y., Sanbo, M., Shinoda, T., Tsuboi, M., Takashima, S., Yagi, T., Fujisawa, H. (2005) Plexin-a4 mediates axon-repulsive activities of both secreted and transmembrane semaphorins and plays roles in nerve fiber guidance. *J Neurosci*, **25**, 3628-37.
- Suzuki, T., Nakamoto, T., Ogawa, S., Seo, S., Matsumura, T., Tachibana, K., Morimoto, C., Hirai, H. (2002) MICAL, a novel CasL interacting molecule, associates with vimentin. *J Biol Chem*, **277**, 14933-41.
- Swanson, L. W., Sawchenko, P. E., Cowan, W. M. (1980) Evidence that the commissural, associational and septal projections of the regio inferior of the hippocampus arise from the same neurons. *Brain Res*, **197**, 207-12.
- Swiercz, J. M., Kuner, R., Behrens, J., Offermanns, S. (2002) Plexin-B1 directly interacts with PDZ-RhoGEF/LARG to regulate RhoA and growth cone morphology. *Neuron*, **35**, 51-63.
- Takahashi, T., Nakamura, F., Jin, Z., Kalb, R. G., Strittmatter, S. M. (1998) Semaphorins A and E act as antagonists of neuropilin-1 and agonists of neuropilin-2 receptors. *Nat Neurosci*, **1**, 487-93.
- Takahashi, T., Fournier, A., Nakamura, F., Wang, L. H., Murakami, Y., Kalb, R. G., Fujisawa, H., Strittmatter, S. M. (1999) Plexin-neuropilin-1 complexes form functional semaphorin-3A receptors. *Cell*, **99**, 59-69.
- Takahashi, T., Strittmatter, S. M. (2001) Plexin1 autoinhibition by the plexin sema domain. *Neuron*, **29**, 429-39.
- Takashima, S., Kitakaze, M., Asakura, M., Asanuma, H., Sanada, S., Tashiro, F., Niwa, H., Miyazaki, J., Hirota, S., Kitamura, Y., Kitsukawa, T., Fujisawa, H., Klagsbrun, M., Hori, M. (2002) Targeting of both mouse neuropilin-1 and neuropilin-2 genes severely impairs developmental yolk sac and embryonic angiogenesis. *Proc Natl Acad Sci U S A*, **99**, 3657-62.
- Takayama, C. (1994) Altered distribution of inhibitory synaptic terminals in reeler cerebellum with special reference to malposition of GABAergic neurons. *Neurosci Res*, **20**, 239-50.
- Tamagnone, L., Artigiani, S., Chen, H., He, Z., Ming, G. I., Song, H., Chedotal, A., Winberg, M. L., Goodman, C. S., Poo, M., Tessier-Lavigne, M., Comoglio, P. M. (1999) Plexins are a large family of receptors for transmembrane, secreted, and GPI-anchored semaphorins in vertebrates. *Cell*, **99**, 71-80.
- Tamagnone, L., Comoglio, P. M. (2004) To move or not to move? Semaphorin signalling in cell migration. *EMBO Rep*, **5**, 356-61.
- Tamamaki, N., Watanabe, K., Nojyo, Y. (1984) A whole image of the hippocampal pyramidal neuron revealed by intracellular pressure-injection of horseradish peroxidase. *Brain Res*, **307**, 336-40.
- Tamamaki, N., Fujimori, K., Nojyo, Y., Kaneko, T., Takauji, R. (2003) Evidence that *Sema3A* and *Sema3F* regulate the migration of GABAergic neurons in the developing neocortex. *J Comp Neurol*, **455**, 238-48.
- Tan, A. M., Zhang, W., Levine, J. M. (2005) NG2: a component of the glial scar that inhibits axon growth. *Journal of Anatomy*, **207**, 717-725.
- Tanaka, M., Ohashi, R., Nakamura, R., Shinmura, K., Kamo, T., Sakai, R., Sugimura, H. (2004) Tiam1 mediates neurite outgrowth induced by ephrin-B1 and EphA2. *Embo J*, **23**, 1075-88.
- Taniguchi, M., Yuasa, S., Fujisawa, H., Naruse, I., Saga, S., Mishina, M., Yagi, T. (1997) Disruption of semaphorin III/D gene causes severe abnormality in peripheral nerve projection. *Neuron*, **19**, 519-30.
- Tear, G., Harris, R., Sutaria, S., Kilomanski, K., Goodman, C. S., Seeger, M. A. (1996) commissureless controls growth cone guidance across the CNS midline in *Drosophila* and encodes a novel membrane protein. *Neuron*, **16**, 501-14.
- Terman, J. R., Mao, T., Pasterkamp, R. J., Yu, H. H., Kolodkin, A. L. (2002) MICALs, a family of conserved flavoprotein oxidoreductases, function in plexin-mediated axonal repulsion. *Cell*, **109**, 887-900.

- Terman, J. R., Kolodkin, A. L. (2004) Nervy links protein kinase a to plexin-mediated semaphorin repulsion. *Science*, **303**, 1204-7.
- Tessier-Lavigne, M., Goodman, C. S. (1996) The molecular biology of axon guidance. *Science*, **274**, 1123-33.
- Tissir, F., Goffinet, A. M. (2003) Reelin and brain development. *Nat Rev Neurosci*, **4**, 496-505.
- Tong, J., Killeen, M., Steven, R., Binns, K. L., Culotti, J., Pawson, T. (2001) Netrin stimulates tyrosine phosphorylation of the UNC-5 family of netrin receptors and induces Shp2 binding to the RCM cytodomain. *J Biol Chem*, **276**, 40917-25.
- Torres-Vazquez, J., Gitler, A. D., Fraser, S. D., Berk, J. D., Van, N. P., Fishman, M. C., Childs, S., Epstein, J. A., Weinstein, B. M. (2004) Semaphorin-plexin signaling guides patterning of the developing vasculature. *Dev Cell*, **7**, 117-23.
- Tsai, H.-H., Tessier-Lavigne, M., Miller, R. H. (2003) Netrin 1 mediates spinal cord oligodendrocyte precursor dispersal. *Development*, **130**, 2095-2105.
- Tyzio, R., Represa, A., Jorquera, I., Ben-Ari, Y., Gozlan, H., Aniksztejn, L. (1999) The Establishment of GABAergic and Glutamatergic Synapses on CA1 Pyramidal Neurons is Sequential and Correlates with the Development of the Apical Dendrite. *J. Neurosci.*, **19**, 10372-10382.
- Ughrin, Y. M., Chen, Z. J., Levine, J. M. (2003) Multiple regions of the NG2 proteoglycan inhibit neurite growth and induce growth cone collapse. *J Neurosci*, **23**, 175-86.
- Venkatesh, K., Chivatakarn, O., Lee, H., Joshi, P. S., Kantor, D. B., Newman, B. A., Mage, R., Rader, C., Giger, R. J. (2005) The Nogo-66 Receptor Homolog NgR2 Is a Sialic Acid-Dependent Receptor Selective for Myelin-Associated Glycoprotein. *J. Neurosci.*, **25**, 808-822.
- Ventura, A., Pelicci, P. G. (2002) Semaphorins: green light for redox signaling? *Sci STKE*, **2002**, PE44.
- Vikis, H. G., Li, W., He, Z., Guan, K. L. (2000) The semaphorin receptor plexin-B1 specifically interacts with active Rac in a ligand-dependent manner. *Proc Natl Acad Sci U S A*, **97**, 12457-62.
- Vikis, H. G., Li, W., Guan, K. L. (2002) The plexin-B1/Rac interaction inhibits PAK activation and enhances Sema4D ligand binding. *Genes Dev*, **16**, 836-45.
- Voogd, J., Glickstein, M. (1998) The anatomy of the cerebellum. *Trends in Neuroscience*, **21**, 370-375.
- Vourc'h, P., Andres, C. (2004) Oligodendrocyte myelin glycoprotein (OMgp): evolution, structure and function. *Brain Research Reviews*, **45**, 115-124.
- Vyas, A. A., Patel, H. V., Fromholt, S. E., Heffer-Lauc, M., Vyas, K. A., Dang, J., Schachner, M., Schnaar, R. L. (2002) Gangliosides are functional nerve cell ligands for myelin-associated glycoprotein (MAG), an inhibitor of nerve regeneration. *Proc Natl Acad Sci U S A*, **99**, 8412-7.
- Wang, G. X., Poo, M. M. (2005) Requirement of TRPC channels in netrin-1-induced chemotropic turning of nerve growth cones. *Nature*, **434**, 898-904.
- Wang, H. U., Anderson, D. J. (1997) Eph family transmembrane ligands can mediate repulsive guidance of trunk neural crest migration and motor axon outgrowth. *Neuron*, **18**, 383-96.
- Wang, K. C., Koprivica, V., Kim, J. A., Sivasankaran, R., Guo, Y., Neve, R. L., He, Z. (2002) Oligodendrocyte-myelin glycoprotein is a Nogo receptor ligand that inhibits neurite outgrowth. **417**, 941-944.
- Wang, K. H., Brose, K., Arnott, D., Kidd, T., Goodman, C. S., Henzel, W., Tessier-Lavigne, M. (1999) Biochemical purification of a mammalian slit protein as a positive regulator of sensory axon elongation and branching. *Cell*, **96**, 771-84.
- Wang, L. H., Strittmatter, S. M. (1996) A family of rat CRMP genes is differentially expressed in the nervous system. *J Neurosci*, **16**, 6197-207.
- Wang, V. Y., Zoghbi, H. Y. (2001) Genetic regulation of cerebellar development. *Nat Rev Neurosci*, **2**, 484-91.
- Wang, V. Y., Rose, M. F., Zoghbi, H. Y. (2005) Math1 Expression Redefines the Rhombic Lip Derivatives and Reveals Novel Lineages within the Brainstem and Cerebellum. *Neuron*, **48**, 31-43.
- Weeber, E. J., Beffert, U., Jones, C., Christian, J. M., Forster, E., Sweatt, J. D., Herz, J. (2002) Reelin and ApoE receptors cooperate to enhance hippocampal synaptic plasticity and learning. *J Biol Chem*, **277**, 39944-52.
- Wen, Z., Guirland, C., Ming, G. L., Zheng, J. Q. (2004) A CaMKII/calcineurin switch controls the direction of Ca(2+)-dependent growth cone guidance. *Neuron*, **43**, 835-46.



- Wennerberg, K., Forget, M. A., Ellerbroek, S. M., Arthur, W. T., Burridge, K., Settleman, J., Der, C. J., Hansen, S. H. (2003) Rnd proteins function as RhoA antagonists by activating p190 RhoGAP. *Curr Biol*, **13**, 1106-15.
- Whitaker, G. B., Limberg, B. J., Rosenbaum, J. S. (2001) Vascular endothelial growth factor receptor-2 and neuropilin-1 form a receptor complex that is responsible for the differential signaling potency of VEGF(165) and VEGF(121). *J Biol Chem*, **276**, 25520-31.
- White, F. A., Behar, O. (2000) The development and subsequent elimination of aberrant peripheral axon projections in Semaphorin3A null mutant mice. *Dev Biol*, **225**, 79-86.
- Williams, M. E., Wu, S. C., McKenna, W. L., Hinck, L. (2003) Surface expression of the netrin receptor UNC5H1 is regulated through a protein kinase C-interacting protein/protein kinase-dependent mechanism. *J Neurosci*, **23**, 11279-88.
- Winberg, M. L., Mitchell, K. J., Goodman, C. S. (1998a) Genetic analysis of the mechanisms controlling target selection: complementary and combinatorial functions of netrins, semaphorins, and IgCAMs. *Cell*, **93**, 581-91.
- Winberg, M. L., Noordermeer, J. N., Tamagnone, L., Comoglio, P. M., Spriggs, M. K., Tessier-Lavigne, M., Goodman, C. S. (1998b) Plexin A is a neuronal semaphorin receptor that controls axon guidance. *Cell*, **95**, 903-16.
- Wingate, R. (2005) Math-Map(ic)s. *Neuron*, **48**, 1-4.
- Wingate, R. J. (2001) The rhombic lip and early cerebellar development. *Curr Opin Neurobiol*, **11**, 82-8.
- Wolman, M. A., Liu, Y., Tawarayama, H., Shoji, W., Halloran, M. C. (2004) Repulsion and attraction of axons by semaphorin3D are mediated by different neuropilins in vivo. *J Neurosci*, **24**, 8428-35.
- Wong, K., Ren, X. R., Huang, Y. Z., Xie, Y., Liu, G., Saito, H., Tang, H., Wen, L., Brady-Kalnay, S. M., Mei, L., Wu, J. Y., Xiong, W. C., Rao, Y. (2001) Signal transduction in neuronal migration: roles of GTPase activating proteins and the small GTPase Cdc42 in the Slit-Robo pathway. *Cell*, **107**, 209-21.
- Wu, W., Wong, K., Chen, J., Jiang, Z., Dupuis, S., Wu, J. Y., Rao, Y. (1999) Directional guidance of neuronal migration in the olfactory system by the protein Slit. *Nature*, **400**, 331-6.
- Xu, Q., Mellitzer, G., Robinson, V., Wilkinson, D. G. (1999) In vivo cell sorting in complementary segmental domains mediated by Eph receptors and ephrins. *Nature*, **399**, 267-71.
- Xu, X. X., Yi, T., Tang, B., Lambeth, J. D. (1998) Disabled-2 (Dab2) is an SH3 domain-binding partner of Grb2. *Oncogene*, **16**, 1561-9.
- Yabut, O., Renfro, A., Niu, S., Swann, J. W., Marin, O., D'Arcangelo, G. (2005) Abnormal laminar position and dendrite development of interneurons in the reeler forebrain. *Brain Res Mol Brain Res*.
- Yamada, K., Watanabe, M. (2002) Cytodifferentiation of Bergmann glia and its relationship with Purkinje cells. *Anat Sci Int*, **77**, 94-108.
- Yamanaka, H., Yanagawa, Y., Obata, K. (2004) Development of stellate and basket cells and their apoptosis in mouse cerebellar cortex. *Neuroscience Research*, **50**, 13-22.
- Yamashita, T., Fujitani, M., Yamagishi, S., Hata, K., Mimura, F. (2005) Multiple signals regulate axon regeneration through the nogo receptor complex. *Mol Neurobiol*, **32**, 105-11.
- Yaron, A., Huang, P. H., Cheng, H. J., Tessier-Lavigne, M. (2005) Differential requirement for Plexin-A3 and -A4 in mediating responses of sensory and sympathetic neurons to distinct class 3 Semaphorins. *Neuron*, **45**, 513-23.
- Yebra, M., Montgomery, A. M., Diaferia, G. R., Kaido, T., Silletti, S., Perez, B., Just, M. L., Hildbrand, S., Hurford, R., Florkiewicz, E., Tessier-Lavigne, M., Cirulli, V. (2003) Recognition of the neural chemoattractant Netrin-1 by integrins alpha6beta4 and alpha3beta1 regulates epithelial cell adhesion and migration. *Dev Cell*, **5**, 695-707.
- Yee, K. T., Simon, H. H., Tessier-Lavigne, M., O'Leary, D. D. M. (1999) Extension of Long Leading Processes and Neuronal Migration in the Mammalian Brain Directed by the Chemoattractant Netrin-1. *Neuron*, **24**, 607-622.
- Yuasa, S., Kawamura, K., Kuwano, R., Ono, K. (1996) Neuron-glia interrelations during migration of Purkinje cells in the mouse embryonic cerebellum. *Int J Dev Neurosci*, **14**, 429-38.
- Yue, Y., Chen, Z. Y., Gale, N. W., Blair-Flynn, J., Hu, T. J., Yue, X., Cooper, M., Crockett, D. P., Yancopoulos, G. D., Tessarollo, L., Zhou, R. (2002) Mistargeting hippocampal axons by expression of a truncated Eph receptor. *Proc Natl Acad Sci U S A*, **99**, 10777-82.

- Yurchenco, P. D., Wadsworth, W. G. (2004) Assembly and tissue functions of early embryonic laminins and netrins. *Current Opinion in Cell Biology*, **16**, 572-579.
- Zanata, S. M., Hovatta, I., Rohm, B., Puschel, A. W. (2002) Antagonistic effects of Rnd1 and RhoD GTPases regulate receptor activity in Semaphorin 3A-induced cytoskeletal collapse. *J Neurosci*, **22**, 471-7.
- Zhang, L., Goldman, J. E. (1996a) Generation of cerebellar interneurons from dividing progenitors in white matter. *Neuron*, **16**, 47-54.
- Zhang, L., Goldman, J. E. (1996b) Developmental fates and migratory pathways of dividing progenitors in the postnatal rat cerebellum. *J Comp Neurol*, **370**, 536-50.
- Zheng, B., Ho, C., Li, S., Keirstead, H., Steward, O., Tessier-Lavigne, M. (2003) Lack of enhanced spinal regeneration in Nogo-deficient mice. *Neuron*, **38**, 213-24.
- Zhou, F.-Q., Waterman-Storer, C. M., Cohan, C. S. (2002) Focal loss of actin bundles causes microtubule redistribution and growth cone turning. *J. Cell Biol.*, **157**, 839-849.
- Zhu, Y., Li, H., Zhou, L., Wu, J. Y., Rao, Y. (1999) Cellular and molecular guidance of GABAergic neuronal migration from an extracortical origin to the neocortex. *Neuron*, **23**, 473-85.
- Zhuo, L., Sun, B., Zhang, C. L., Fine, A., Chiu, S. Y., Messing, A. (1997) Live astrocytes visualized by green fluorescent protein in transgenic mice. *Dev Biol*, **187**, 36-42.
- Zuo, J., De Jager, P. L., Takahashi, K. A., Jiang, W., Linden, D. J., Heintz, N. (1997) Neurodegeneration in Lurcher mice caused by mutation in delta2 glutamate receptor gene. *Nature*, **388**, 769-73.