

**DEPARTAMENT DE  
BIOLOGIA CEL·LULAR I ANATOMIA PATOLÒGICA  
FACULTAT DE MEDICINA**



**MECANISMOS INTRACELULARES DE SUPERVIVENCIA  
Y MUERTE NEURONAL EN MODELOS  
EXCITOTÓXICOS Y TRANSGÉNICOS DE LA  
ENFERMEDAD DE HUNTINGTON**

**Tesis presentada por Juan Manuel García Martínez  
para optar al título de Doctor por la Universidad de Barcelona**

## **VI.- BIBLIOGRAFÍA**



---

**A**

- Abe, K., Kurakin, A., Mohseni-Maybodi, M., Kay, B., and Khosravi-Far, R. (2000). "The complexity of TNF-related apoptosis-inducing ligand." *Ann. N. Y. Acad. Sci.*, 926, 52-63.
- Adrain, C., Slee, E. A., Harte, M. T., and Martin, S. J. (1999). "Regulation of apoptotic protease activating factor-1 oligomerization and apoptosis by the WD-40 repeat region." *J. Biol. Chem.*, 274(30), 20855-20860.
- Aebischer, P., and Ridet, J. (2001). "Recombinant proteins for neurodegenerative diseases: the delivery issue." *Trends Neurosci.*, 24(9), 533-540.
- Airaksinen, M. S., and Saarma, M. (2002). "The GDNF family: signalling, biological functions and therapeutic value." *Nat. Rev. Neurosci.*, 3(5), 383-394.
- Akerud, P., Alberch, J., Eketjall, S., Wagner, J., and Arenas, E. (1999). "Differential effects of glial cell line-derived neurotrophic factor and neurturin on developing and adult substantia nigra dopaminergic neurons." *J. Neurochem.*, 73(1), 70-78.
- Alberch, J., Perez-Navarro, E., and Canals, J. M. (2002). "Neuroprotection by neurotrophins and GDNF family members in the excitotoxic model of Huntington's disease." *Brain Res. Bull.*, 57(6), 817-822.
- Alberch, J., Perez-Navarro, E., and Canals, J. M. (2004). "Neurotrophic factors in Huntington's disease." *Prog. Brain Res.*, 146, 195-229.
- Alberch, Perez-Navarro, E., and Canals, J. M. (2007). "Huntington's disease: animal models". Encyclopedia of Neuroscience. Elsevier, en prensa.
- Albin, R. L., and Gilman, S. (1990). "Autoradiographic localization of inhibitory and excitatory amino acid neurotransmitter receptors in human normal and olivopontocerebellar atrophy cerebellar cortex." *Brain Res.*, 522(1), 37-45.
- Albin, R. L., Reiner, A., Anderson, K. D., Dure, L. S., Handelin, B., Balfour, R., Whetsell, W. O., Jr., Penney, J. B., and Young, A. B. (1992). "Preferential loss of striato-external pallidal projection neurons in presymptomatic Huntington's disease." *Ann. Neurol.*, 31(4), 425-430.
- Alessi, D. R., Andjelkovic, M., Caudwell, B., Cron, P., Morrice, N., Cohen, P., and Hemmings, B. A. (1996). "Mechanism of activation of protein kinase B by insulin and IGF-1." *EMBO J.*, 15(23), 6541-6551.
- Alessi, D. R., James, S. R., Downes, C. P., Holmes, A. B., Gaffney, P. R., Reese, C. B., and Cohen, P. (1997). "Characterization of a 3-phosphoinositide-dependent protein kinase which phosphorylates and activates protein kinase Balpha." *Curr. Biol.*, 7(4), 261-269.
- Alexi, T., Venero, J. L., and Hefti, F. (1997). "Protective effects of neurotrophin-4/5 and transforming growth factor-alpha on striatal neuronal phenotypic degeneration after excitotoxic lesioning with quinolinic acid." *Neuroscience*, 78(1), 73-86.

- Alexi, T., Borlongan, C. V., Faull, R. L., Williams, C. E., Clark, R. G., Gluckman, P. D., and Hughes, P. E. (2000). "Neuroprotective strategies for basal ganglia degeneration: Parkinson's and Huntington's diseases." *Prog Neurobiol*, 60(5), 409-470.
- Altar, C. A., Armanini, M., Dugich-Djordjevic, M., Bennett, G. L., Williams, R., Feinglass, S., Anicetti, V., Sinicropi, D., and Bakhit, C. (1992). "Recovery of cholinergic phenotype in the injured rat neostriatum: roles for endogenous and exogenous nerve growth factor." *J. Neurochem.*, 59(6), 2167-2177.
- Altar, C. A., Cai, N., Bliven, T., Juhasz, M., Conner, J. M., Acheson, A. L., Lindsay, R. M., and Wiegand, S. J. (1997). "Anterograde transport of brain-derived neurotrophic factor and its role in the brain." *Nature*, 389(6653), 856-860.
- Altman, J. (1992). "Programmed cell death: the paths to suicide." *Trends Neurosci*, 15(8), 278-280.
- Anderson, K. E., Coadwell, J., Stephens, L. R., and Hawkins, P. T. (1998). "Translocation of PDK-1 to the plasma membrane is important in allowing PDK-1 to activate protein kinase B." *Curr. Biol.*, 8(12), 684-691.
- Andjelkovic, M., Alessi, D. R., Meier, R., Fernandez, A., Lamb, N. J., Frech, M., Cron, P., Cohen, P., Lucocq, J. M., and Hemmings, B. A. (1997). "Role of translocation in the activation and function of protein kinase B." *J. Biol. Chem.*, 272(50), 31515-31524.
- Andreyev, A. Y., Fahy, B., and Fiskum, G. (1998). "Cytochrome c release from brain mitochondria is independent of the mitochondrial permeability transition." *FEBS Lett.*, 439(3), 373-376.
- Anglade, P., Vyas, S., Javoy-Agid, F., Herrero, M. T., Michel, P. P., Marquez, J., Mouatt-Prigent, A., Ruberg, M., Hirsch, E. C., and Agid, Y. (1997). "Apoptosis and autophagy in nigral neurons of patients with Parkinson's disease." *Histol. Histopathol.*, 12(1), 25-31.
- Ankarcrona, M., Dypbukt, J. M., Bonfoco, E., Zhivotovsky, B., Orrenius, S., Lipton, S. A., and Nicotera, P. (1995). "Glutamate-induced neuronal death: a succession of necrosis or apoptosis depending on mitochondrial function." *Neuron*, 15(4), 961-973.
- Araujo, D. M., and Hilt, D. C. (1997). "Glial cell line-derived neurotrophic factor attenuates the excitotoxin-induced behavioral and neurochemical deficits in a rodent model of Huntington's disease." *Neuroscience*, 81(4), 1099-1110.
- Arenas, E., Trupp, M., Akerud, P., and Ibanez, C. F. (1995). "GDNF prevents degeneration and promotes the phenotype of brain noradrenergic neurons in vivo." *Neuron*, 15(6), 1465-1473.
- Arevalo, J. C., Conde, B., Hempstead, B. I., Chao, M. V., Martin-Zanca, D., and Perez, P. (2001). "A novel mutation within the extracellular domain of TrkA causes constitutive receptor activation." *Oncogene*, 20(10), 1229-1234.
- Arevalo, J. C., and Wu, S. H. (2006). "Neurotrophin signaling: many exciting surprises!" *Cell Mol. Life Sci.*, 63(13), 1523-1537.

- Arnoult, D., Parone, P., Martinou, J. C., Antonsson, B., Estaquier, J., and Ameisen, J. C. (2002). "Mitochondrial release of apoptosis-inducing factor occurs downstream of cytochrome c release in response to several proapoptotic stimuli." *J. Cell Biol.*, 159(6), 923-929.
- Arrasate, M., Mitra, S., Schweitzer, E. S., Segal, M. R., and Finkbeiner, S. (2004). "Inclusion body formation reduces levels of mutant huntingtin and the risk of neuronal death." *Nature*, 431(7010), 805-810.
- Ashkenazi, A., and Dixit, V. M. (1998). "Death receptors: signaling and modulation." *Science*, 281(5381), 1305-1308.
- Ashkenazi, A. (2002). "Targeting death and decoy receptors of the tumour-necrosis factor superfamily." *Nat. Rev. Cancer*, 2(6), 420-430.
- Atwal, J. K., Massie, B., Miller, F. D., and Kaplan, D. R. (2000). "The TrkB-Shc site signals neuronal survival and local axon growth via MEK and P13-kinase." *Neuron*, 27(2), 265-277.

---

**B**

- Baehrecke, E. H. (2005). "Autophagy: dual roles in life and death?" *Nat. Rev. Mol. Cell Biol.*, 6(6), 505-510.
- Balendran, A., Casamayor, A., Deak, M., Paterson, A., Gaffney, P., Currie, R., Downes, C. P., and Alessi, D. R. (1999). "PDK1 acquires PDK2 activity in the presence of a synthetic peptide derived from the carboxyl terminus of PRK2." *Curr. Biol.*, 9(8), 393-404.
- Baloh, R. H., Tansey, M. G., Golden, J. P., Creedon, D. J., Heuckeroth, R. O., Keck, C. L., Zimonjic, D. B., Popescu, N. C., Johnson, E. M. J., and Milbrandt, J. (1997). "TrnR2, a novel receptor that mediates neurturin and GDNF signaling through Ret." *Neuron*, 18(5), 793-802.
- Baloh, R. H., Gorodinsky, A., Golden, J. P., Tansey, M. G., Keck, C. L., Popescu, N. C., Johnson, E. M. J., and Milbrandt, J. (1998a). "GFRalpha3 is an orphan member of the GDNF/neurturin/persephin receptor family." *Proc Natl Acad Sci U S A*, 95(10), 5801-5806.
- Baloh, R. H., Tansey, M. G., Lampe, P. A., Fahrner, T. J., Enomoto, H., Simburger, K. S., Leitner, M. L., Araki, T., Johnson, E. M. J., and Milbrandt, J. (1998b). "Artemin, a novel member of the GDNF ligand family, supports peripheral and central neurons and signals through the GFRalpha3-RET receptor complex." *Neuron*, 21(6), 1291-1302.
- Barbacid, M. (1994). "The Trk family of neurotrophin receptors." *J Neurobiol*, 25(11), 1386-1403.
- Barde, Y. A., Edgar, D., and Thoenen, H. (1982). "Purification of a new neurotrophic factor from mammalian brain." *EMBO J*, 1(5), 549-553.
- Barkett, M., and Gilmore, T. D. (1999). "Control of apoptosis by Rel/NF-kappaB transcription factors." *Oncogene*, 18(49), 6910-6924.

- Barthwal, M. K., Sathyanarayana, P., Kundu, C. N., Rana, B., Pradeep, A., Sharma, C., Woodgett, J. R., and Rana, A. (2003). "Negative regulation of mixed lineage kinase 3 by protein kinase B/AKT leads to cell survival." *J. Biol. Chem.*, 278(6), 3897-3902.
- Basu, S., Totty, N. F., Irwin, M. S., Sudol, M., and Downward, J. (2003). "Akt phosphorylates the Yes-associated protein, YAP, to induce interaction with 14-3-3 and attenuation of p73-mediated apoptosis." *Mol. Cell*, 11(1), 11-23.
- Bayascas, J. R., and Alessi, D. R. (2005). "Regulation of Akt/PKB Ser473 phosphorylation." *Mol. Cell*, 18(2), 143-145.
- Beal, M. F., Kowall, N. W., Ellison, D. W., Mazurek, M. F., Swartz, K. J., and Martin, J. B. (1986). "Replication of the neurochemical characteristics of Huntington's disease by quinolinic acid." *Nature*, 321(6066), 168-171.
- Beal, M. F., Kowall, N. W., Swartz, K. J., Ferrante, R. J., and Martin, J. B. (1989). "Differential sparing of somatostatin-neuropeptide Y and cholinergic neurons following striatal excitotoxin lesions." *Synapse*, 3(1), 38-47.
- Beal, M. F., Brouillet, E., Jenkins, B. G., Ferrante, R. J., Kowall, N. W., Miller, J. M., Storey, E., Srivastava, R., Rosen, B. R., and Hyman, B. T. (1993). "Neurochemical and histologic characterization of striatal excitotoxic lesions produced by the mitochondrial toxin 3-nitropropionic acid." *J. Neurosci.*, 13(10), 4181-4192.
- Beal, M. F. (2000). "Energetics in the pathogenesis of neurodegenerative diseases." *Trends Neurosci.*, 23(7), 298-304.
- Becher, M. W., Kotzuk, J. A., Sharp, A. H., Davies, S. W., Bates, G. P., Price, D. L., and Ross, C. A. (1998). "Intranuclear neuronal inclusions in Huntington's disease and dentatorubral and pallidoluysian atrophy: correlation between the density of inclusions and IT15 CAG triplet repeat length." *Neurobiol. Dis.*, 4(6), 387-397.
- Bellacosa, A., Testa, J. R., Staal, S. P., and Tsichlis, P. N. (1991). "A retroviral oncogene, akt, encoding a serine-threonine kinase containing an SH2-like region." *Science*, 254(5029), 274-277.
- Bemelmans, A. P., Horellou, P., Pradier, L., Brunet, I., Colin, P., and Mallet, J. (1999). "Brain-derived neurotrophic factor-mediated protection of striatal neurons in an excitotoxic rat model of Huntington's disease, as demonstrated by adenoviral gene transfer." *Hum. Gene Ther.*, 10(18), 2987-2997.
- Benito-Gutierrez, E., Garcia-Fernandez, J., and Comella, J. X. (2006). "Origin and evolution of the Trk family of neurotrophic receptors." *Mol. Cell Neurosci.*, 31(2), 179-192.
- Benn, S. C., and Woolf, C. J. (2004). "Adult neuron survival strategies--slamming on the brakes." *Nat. Rev. Neurosci.*, 5(9), 686-700.
- Berkemeier, L. R., Winslow, J. W., Kaplan, D. R., Nikolics, K., Goeddel, D. V., and Rosenthal, A. (1991). "Neurotrophin-5: a novel neurotrophic factor that activates trk and trkB." *Neuron*, 7(5), 857-866.

- Bhide, P. G., Day, M., Sapp, E., Schwarz, C., Sheth, A., Kim, J., Young, A. B., Penney, J., Golden, J., Aronin, N., and DiFiglia, M. (1996). "Expression of normal and mutant huntingtin in the developing brain." *J. Neurosci.*, 16(17), 5523-5535.
- Bibel, M., Hoppe, E., and Barde, Y. A. (1999). "Biochemical and functional interactions between the neurotrophin receptors trk and p75NTR." *EMBO J.*, 18(3), 616-622.
- Biswas, S. C., and Greene, L. A. (2002). "Nerve growth factor (NGF) down-regulates the Bcl-2 homology 3 (BH3) domain-only protein Bim and suppresses its proapoptotic activity by phosphorylation." *J. Biol. Chem.*, 277(51), 49511-49516.
- Boatright, K. M., and Salvesen, G. S. (2003). "Caspase activation." *Biochem. Soc. Symp.*, (70), 233-242.
- Bonni, A., Brunet, A., West, A. E., Datta, S. R., Takasu, M. A., and Greenberg, M. E. (1999). "Cell survival promoted by the Ras-MAPK signaling pathway by transcription-dependent and -independent mechanisms." *Science*, 286(5443), 1358-1362.
- Bouillet, P., Metcalf, D., Huang, D. C. S., Tarlinton, D. M., Kay, T. W. H., Kontgen, F., Adams, J. M., and Strasser, A. (1999). "Proapoptotic Bcl-2 relative bim required for certain apoptotic responses, leukocyte homeostasis, and to preclude autoimmunity." *Science*, 286(5445), 1735-1738.
- Bratton, S. B., Walker, G., Srinivasula, S. M., Sun, X. M., Butterworth, M., Alnemri, E. S., and Cohen, G. M. (2001). "Recruitment, activation and retention of caspases-9 and -3 by Apaf-1 apoptosome and associated XIAP complexes." *EMBO J.*, 20(5), 998-1009.
- Brazil, D. P., and Hemmings, B. A. (2001). "Ten years of protein kinase B signalling: a hard Akt to follow." *Trends Biochem. Sci.*, 26(11), 657-664.
- Brodbeck, D., Cron, P., and Hemmings, B. A. (1999). "A human protein kinase Bgamma with regulatory phosphorylation sites in the activation loop and in the C-terminal hydrophobic domain." *J. Biol. Chem.*, 274(14), 9133-9136.
- Broker, L. E., Kruyt, F. A., and Giaccone, G. (2005). "Cell death independent of caspases: a review." *Clin. Cancer Res.*, 11(9), 3155-3162.
- Brouillet, E., Hantraye, P., Ferrante, R. J., Dolan, R., Leroy-Willig, A., Kowall, N. W., and Beal, M. F. (1995). "Chronic mitochondrial energy impairment produces selective striatal degeneration and abnormal choreiform movements in primates." *Proc. Natl. Acad. Sci. U. S. A.*, 92(15), 7105-7109.
- Browne, S. E., Bowling, A. C., MacGarvey, U., Baik, M. J., Berger, S. C., Muqit, M. M., Bird, E. D., and Beal, M. F. (1997). "Oxidative damage and metabolic dysfunction in Huntington's disease: selective vulnerability of the basal ganglia." *Ann. Neurol.*, 41(5), 646-653.
- Brunet, A., Bonni, A., Zigmond, M. J., Lin, M. Z., Juo, P., Hu, L. S., Anderson, M. J., Arden, K. C., Blenis, J., and Greenberg, M. E. (1999). "Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor." *Cell*, 96(6), 857-868.



- Bruning, J. C., Gillette, J. A., Zhao, Y., Bjorbaeck, C., Kotzka, J., Knebel, B., Avci, H., Hanstein, B., Lingohr, P., Moller, D. E., Krone, W., Kahn, C. R., and Muller-Wieland, D. (2000). "Ribosomal subunit kinase-2 is required for growth factor-stimulated transcription of the c-Fos gene." *Proc. Natl. Acad. Sci. U. S. A.*, 97(6), 2462-2467.
- Brunk, U. T., and Terman, A. (2002). "The mitochondrial-lysosomal axis theory of aging: accumulation of damaged mitochondria as a result of imperfect autophagocytosis." *Eur. J. Biochem.*, 269(8), 1996-2002.
- Brustovetsky, N., Brustovetsky, T., Purl, K. J., Capano, M., Crompton, M., and Dubinsky, J. M. (2003). "Increased susceptibility of striatal mitochondria to calcium-induced permeability transition." *J. Neurosci.*, 23(12), 4858-4867.
- Buj-Bello, A., Adu, J., Pinon, L. G., Horton, A., Thompson, J., Rosenthal, A., Chinchetru, M., Buchman, V. L., and Davies, A. M. (1997). "Neurturin responsiveness requires a GPI-linked receptor and the Ret receptor tyrosine kinase." *Nature*, 387(6634), 721-724.
- Burgering, B. M., and Coffey, P. J. (1995). "Protein kinase B (c-Akt) in phosphatidylinositol-3-OH kinase signal transduction." *Nature*, 376(6541), 599-602.
- Bursch, W. (2001). "The autophagosomal-lysosomal compartment in programmed cell death." *Cell Death. Differ.*, 8(6), 569-581.

---

**C**

- Cacalano, G., Farinas, I., Wang, L. C., Hagler, K., Forgie, A., Moore, M., Armanini, M., Phillips, H., Ryan, A. M., Reichardt, L. F., Hynes, M., Davies, A., and Rosenthal, A. (1998). "GFRalpha1 is an essential receptor component for GDNF in the developing nervous system and kidney." *Neuron*, 21(1), 53-62.
- Calabresi, P., Pisani, A., Mercuri, N. B., and Bernardi, G. (1996). "The corticostriatal projection: from synaptic plasticity to dysfunctions of the basal ganglia." *Trends Neurosci.*, 19(1), 19-24.
- Calabresi, P., Gubellini, P., Picconi, B., Centonze, D., Pisani, A., Bonsi, P., Greengard, P., Hipskind, R. A., Borrelli, E., and Bernardi, G. (2001). "Inhibition of mitochondrial complex II induces a long-term potentiation of NMDA-mediated synaptic excitation in the striatum requiring endogenous dopamine." *J. Neurosci.*, 21(14), 5110-5120.
- Canals, J. M., Marco, S., Checa, N., Michels, A., Perez-Navarro, E., Arenas, E., and Alberch, J. (1998). "Differential regulation of the expression of nerve growth factor, brain-derived neurotrophic factor, and neurotrophin-3 after excitotoxicity in a rat model of Huntington's disease." *Neurobiol. Dis.*, 5(5), 357-364.
- Canals, J. M., Checa, N., Marco, S., Michels, A., Perez-Navarro, E., and Alberch, J. (1999). "The neurotrophin receptors trkA, trkB and trkC are differentially regulated after excitotoxic lesion in rat striatum." *Brain Res. Mol. Brain Res.*, 69(2), 242-248.

- Canals, J. M., Pineda, J. R., Torres-Peraza, J. F., Bosch, M., Martin-Ibanez, R., Munoz, M. T., Mengod, G., Ernfors, P., and Alberch, J. (2004). "Brain-derived neurotrophic factor regulates the onset and severity of motor dysfunction associated with enkephalinergic neuronal degeneration in Huntington's disease." *J. Neurosci.*, 24(35), 7727-7739.
- Cande, C., Vahsen, N., Garrido, C., and Kroemer, G. (2004). "Apoptosis-inducing factor (AIF): caspase-independent after all." *Cell Death. Differ.*, 11(6), 591-595.
- Cardone, M. H., Roy, N., Stennicke, H. R., Salvesen, G. S., Franke, T. F., Stanbridge, E., Frisch, S., and Reed, J. C. (1998). "Regulation of cell death protease caspase-9 by phosphorylation." *Science*, 282(5392), 1318-1321.
- Cattaneo, E., Zuccato, C., and Tartari, M. (2005). "Normal huntingtin function: an alternative approach to Huntington's disease." *Nat. Rev. Neurosci.*, 6(12), 919-930.
- Celio, M. R. (1990). "Calbindin-D-28K and Parvalbumin in the Rat Nervous-System." *Neuroscience*, 35(2), 375-475.
- Cepeda, C., Ariano, M. A., Calvert, C. R., Flores-Hernandez, J., Chandler, S. H., Leavitt, B. R., Hayden, M. R., and Levine, M. S. (2001). "NMDA receptor function in mouse models of Huntington disease." *J. Neurosci. Res.*, 66(4), 525-539.
- Chaib-Oukadour, I., Gil, C., and Aguilera, J. (2004). "The C-terminal domain of the heavy chain of tetanus toxin rescues cerebellar granule neurones from apoptotic death: involvement of phosphatidylinositol 3-kinase and mitogen-activated protein kinase pathways." *J. Neurochem.*, 90(5), 1227-1236.
- Chan, E. Y., Luthi-Carter, R., Strand, A., Solano, S. M., Hanson, S. A., DeJohn, M. M., Kooperberg, C., Chase, K. O., DiFiglia, M., Young, A. B., Leavitt, B. R., Cha, J. H., Aronin, N., Hayden, M. R., and Olson, J. M. (2002). "Increased huntingtin protein length reduces the number of polyglutamine-induced gene expression changes in mouse models of Huntington's disease." *Hum. Mol. Genet.*, 11(17), 1939-1951.
- Chan, F. K., Chun, H. J., Zheng, L., Siegel, R. M., Bui, K. L., and Lenardo, M. J. (2000). "A domain in TNF receptors that mediates ligand-independent receptor assembly and signaling." *Science*, 288(5475), 2351-2354.
- Chao, M. V. (2003). "Neurotrophins and their receptors: a convergence point for many signalling pathways." *Nat. Rev. Neurosci.*, 4(4), 299-309.
- Checa, N., Canals, J. M., and Alberch, J. (2000). "Developmental regulation of BDNF and NT-3 expression by quinolinic acid in the striatum and its main connections." *Exp. Neurol.*, 165(1), 118-124.
- Checa, N., Canals, J. M., Gratacos, E., and Alberch, J. (2001). "TrkB and TrkC are differentially regulated by excitotoxicity during development of the basal ganglia." *Exp. Neurol.*, 172(2), 282-292.
- Chen, D., and Zhou, Q. (2004). "Caspase cleavage of BimEL triggers a positive feedback amplification of apoptotic signaling." *Proc. Natl. Acad. Sci. U. S. A.*, 101(5), 1235-1240.

- Chen, H. K., Fernandez-Funez, P., Acevedo, S. F., Lam, Y. C., Kaytor, M. D., Fernandez, M. H., Aitken, A., Skoulakis, E. M., Orr, H. T., Botas, J., and Zoghbi, H. Y. (2003). "Interaction of Akt-phosphorylated ataxin-1 with 14-3-3 mediates neurodegeneration in spinocerebellar ataxia type 1." *Cell*, 113(4), 457-468.
- Chen, L., Willis, S. N., Wei, A., Smith, B. J., Fletcher, J. I., Hinds, M. G., Colman, P. M., Day, C. L., Adams, J. M., and Huang, D. C. (2005). "Differential targeting of prosurvival Bcl-2 proteins by their BH3-only ligands allows complementary apoptotic function." *Mol. Cell*, 17(3), 393-403.
- Chen, M., He, H., Zhan, S., Krajewski, S., Reed, J. C., and Gottlieb, R. A. (2001). "Bid is cleaved by calpain to an active fragment in vitro and during myocardial ischemia/reperfusion." *J. Biol. Chem.*, 276(33), 30724-30728.
- Chen, Q., and Reiner, A. (1996). "Cellular distribution of the NMDA receptor NR2A/2B subunits in the rat striatum." *Brain Res.*, 743(1-2), 346-352.
- Chen, Q., Veenman, C. L., and Reiner, A. (1996). "Cellular expression of ionotropic glutamate receptor subunits on specific striatal neuron types and its implication for striatal vulnerability in glutamate receptor-mediated excitotoxicity." *Neuroscience*, 73(3), 715-731.
- Chen, Q., Surmeier, D. J., and Reiner, A. (1999). "NMDA and non-NMDA receptor-mediated excitotoxicity are potentiated in cultured striatal neurons by prior chronic depolarization." *Exp. Neurol.*, 159(1), 283-296.
- Chen, R. H., Sarnecki, C., and Blenis, J. (1992). "Nuclear localization and regulation of erk- and rsk-encoded protein kinases." *Mol. Cell Biol.*, 12(3), 915-927.
- Chen, R. H., Abate, C., and Blenis, J. (1993). "Phosphorylation of the c-Fos transrepression domain by mitogen-activated protein kinase and 90-kDa ribosomal S6 kinase." *Proc. Natl. Acad. Sci. U. S. A.*, 90(23), 10952-10956.
- Chen, R. H., Juo, P. C., Curran, T., and Blenis, J. (1996). "Phosphorylation of c-Fos at the C-terminus enhances its transforming activity." *Oncogene*, 12(7), 1493-1502.
- Cheng, E. H., Kirsch, D. G., Clem, R. J., Ravi, R., Kastan, M. B., Bedi, A., Ueno, K., and Hardwick, J. M. (1997). "Conversion of Bcl-2 to a Bax-like death effector by caspases." *Science*, 278(5345), 1966-1968.
- Cheng, E. H., Sheiko, T. V., Fisher, J. K., Craigen, W. J., and Korsmeyer, S. J. (2003). "VDAC2 inhibits BAK activation and mitochondrial apoptosis." *Science*, 301(5632), 513-517.
- Cheng, J. Q., Godwin, A. K., Bellacosa, A., Taguchi, T., Franke, T. F., Hamilton, T. C., Tschlis, P. N., and Testa, J. R. (1992). "AKT2, a putative oncogene encoding a member of a subfamily of protein-serine/threonine kinases, is amplified in human ovarian carcinomas." *Proc. Natl. Acad. Sci. U. S. A.*, 89(19), 9267-9271.
- Cheung, N. S., Pascoe, C. J., Giardina, S. F., John, C. A., and Beart, P. M. (1998). "Micromolar L-glutamate induces extensive apoptosis in an apoptotic-necrotic continuum of insult-dependent, excitotoxic injury in cultured cortical neurones." *Neuropharmacology*, 37(10-11), 1419-1429.

- Chiariello, M., Visconti, R., Carlomagno, F., Melillo, R. M., Bucci, C., de, F., V, Fox, G. M., Jing, S., Coso, O. A., Gutkind, J. S., Fusco, A., and Santoro, M. (1998). "Signalling of the Ret receptor tyrosine kinase through the c-Jun NH2-terminal protein kinases (JNKs): evidence for a divergence of the ERKs and JNKs pathways induced by Ret." *Oncogene*, 16(19), 2435-2445.
- Cho, J., Yarygina, O., Oo, T. F., Kholodilov, N. G., and Burke, R. E. (2004). "Glial cell line-derived neurotrophic factor receptor GFRalpha1 is expressed in the rat striatum during postnatal development." *Brain Res. Mol. Brain Res.*, 127(1-2), 96-104.
- Choi, D. W. (1988). "Glutamate neurotoxicity and diseases of the nervous system." *Neuron*, 1(8), 623-634.
- Chong, H., Vikis, H. G., and Guan, K. L. (2003). "Mechanisms of regulating the Raf kinase family." *Cell Signal.*, 15(5), 463-469.
- Chong, Z. Z., Li, F., and Maiese, K. (2005). "Activating Akt and the brain's resources to drive cellular survival and prevent inflammatory injury." *Histol. Histopathol.*, 20(1), 299-315.
- Choo, Y. S., Johnson, G. V., MacDonald, M., Detloff, P. J., and Lesort, M. (2004). "Mutant huntingtin directly increases susceptibility of mitochondria to the calcium-induced permeability transition and cytochrome c release." *Hum. Mol. Genet.*, 13(14), 1407-1420.
- Ciccolini, F., and Svendsen, C. N. (2001). "Neurotrophin responsiveness is differentially regulated in neurons and precursors isolated from the developing striatum." *J. Mol. Neurosci.*, 17(1), 25-33.
- Cirman, T., Oresic, K., Mazovec, G. D., Turk, V., Reed, J. C., Myers, R. M., Salvesen, G. S., and Turk, B. (2004). "Selective disruption of lysosomes in HeLa cells triggers apoptosis mediated by cleavage of Bid by multiple papain-like lysosomal cathepsins." *J. Biol. Chem.*, 279(5), 3578-3587.
- Clodfelder-Miller, B., De, S. P., Zmijewska, A. A., Song, L., and Jope, R. S. (2005). "Physiological and pathological changes in glucose regulate brain Akt and glycogen synthase kinase-3." *J. Biol. Chem.*, 280(48), 39723-39731.
- Coffer, P. J., and Woodgett, J. R. (1991). "Molecular cloning and characterisation of a novel putative protein-serine kinase related to the cAMP-dependent and protein kinase C families." *Eur. J. Biochem.*, 201(2), 475-481.
- Colin, E., Regulier, E., Perrin, V., Durr, A., Brice, A., Aebischer, P., Deglon, N., Humbert, S., and Saudou, F. (2005). "Akt is altered in an animal model of Huntington's disease and in patients." *Eur. J. Neurosci.*, 21(6), 1478-1488.
- Conner, J. M., and Varon, S. (1996). "Maintenance of sympathetic innervation into the hippocampal formation requires a continuous local availability of nerve growth factor." *Neuroscience*, 72(4), 933-945.
- Cooper, J. D., Skepper, J. N., Berzaghi, M. D., Lindholm, D., and Sofroniew, M. V. (1996). "Delayed death of septal cholinergic neurons after excitotoxic ablation of hippocampal neurons during early postnatal development in the rat." *Exp. Neurol.*, 139(1), 143-155.

- Corbit, K. C., Foster, D. A., and Rosner, M. R. (1999). "Protein kinase Cdelta mediates neurogenic but not mitogenic activation of mitogen-activated protein kinase in neuronal cells." *Mol. Cell Biol.*, 19(6), 4209-4218.
- Cory, S., and Adams, J. M. (2002). "The Bcl2 family: regulators of the cellular life-or-death switch." *Nat. Rev. Cancer*, 2(9), 647-656.
- Cosgaya, J. M., Chan, J. R., and Shooter, E. M. (2002). "The neurotrophin receptor p75NTR as a positive modulator of myelination." *Science*, 298(5596), 1245-1248.
- Coyle, J. T., and Schwarcz, R. (1976). "Lesion of striatal neurones with kainic acid provides a model for Huntington's chorea." *Nature*, 263(5574), 244-246.
- Creedon, D. J., Tansey, M. G., Baloh, R. H., Osborne, P. A., Lampe, P. A., Fahrner, T. J., Heuckeroth, R. O., Milbrandt, J., and Johnson, E. M., Jr. (1997). "Neurturin shares receptors and signal transduction pathways with glial cell line-derived neurotrophic factor in sympathetic neurons." *Proc. Natl. Acad. Sci. U. S. A.*, 94(13), 7018-7023.
- Cross, D. A., Alessi, D. R., Cohen, P., Andjelkovich, M., and Hemmings, B. A. (1995). "Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B." *Nature*, 378(6559), 785-789.
- Cudkovicz, M., and Kowall, N. W. (1990). "Degeneration of pyramidal projection neurons in Huntington's disease cortex." *Ann. Neurol.*, 27(2), 200-204.

---

**D**

- Dahlstrand, J., Lardelli, M., and Lendahl, U. (1995). "Nestin Messenger-Rna Expression Correlates with the Central-Nervous-System Progenitor-Cell State in Many, But Not All, Regions of Developing Central-Nervous-System." *Developmental Brain Research*, 84(1), 109-129.
- Datta, S. R., Dudek, H., Tao, X., Masters, S., Fu, H., Gotoh, Y., and Greenberg, M. E. (1997). "Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery." *Cell*, 91(2), 231-241.
- Davies, A. M. (1996). "Paracrine and autocrine actions of neurotrophic factors." *Neurochem. Res.*, 21(7), 749-753.
- Davies, S. W., and Beardsall, K. (1992). "Nerve growth factor selectively prevents excitotoxin induced degeneration of striatal cholinergic neurones." *Neurosci. Lett.*, 140(2), 161-164.
- Davies, S. W., Turmaine, M., Cozens, B. A., DiFiglia, M., Sharp, A. H., Ross, C. A., Scherzinger, E., Wanker, E. E., Mangiarini, L., and Bates, G. P. (1997). "Formation of neuronal intranuclear inclusions underlies the neurological dysfunction in mice transgenic for the HD mutation." *Cell*, 90(3), 537-548.
- Davis, R. J. (1995). "Transcriptional regulation by MAP kinases." *Mol. Reprod. Dev.*, 42(4), 459-467.

- de la Monte, S. M., Vonsattel, J. P., and Richardson, E. P., Jr. (1988). "Morphometric demonstration of atrophic changes in the cerebral cortex, white matter, and neostriatum in Huntington's disease." *J. Neuropathol. Exp. Neurol.*, 47(5), 516-525.
- de, B. S., Canals, S., Casarejos, M. J., Solano, R. M., Menendez, J., and Mena, M. A. (2004). "Role of extracellular signal-regulated protein kinase in neuronal cell death induced by glutathione depletion in neuron/glia mesencephalic cultures." *J. Neurochem.*, 91(3), 667-682.
- Decaudin, D., Marzo, I., Brenner, C., and Kroemer, G. (1998). "Mitochondria in chemotherapy-induced apoptosis: a prospective novel target of cancer therapy (review)." *Int. J. Oncol.*, 12(1), 141-152.
- Deckwerth, T. L., Elliott, J. L., Knudson, C. M., Johnson, E. M., Snider, W. D., and Korsmeyer, S. J. (1996). "BAX is required for neuronal death after trophic factor deprivation and during development." *Neuron*, 17(3), 401-411.
- Degenhardt, K., Sundararajan, R., Lindsten, T., Thompson, C., and White, E. (2002). "Bax and Bak independently promote cytochrome C release from mitochondria." *J. Biol. Chem.*, 277(16), 14127-14134.
- Degterev, A., Boyce, M., and Yuan, J. (2003). "A decade of caspases." *Oncogene*, 22(53), 8543-8567.
- Deng, Y., Ren, X., Yang, L., Lin, Y., and Wu, X. (2003). "A JNK-dependent pathway is required for TNF $\alpha$ -induced apoptosis." *Cell*, 115(1), 61-70.
- Desagher, S., Osen-Sand, A., Nichols, A., Eskes, R., Montessuit, S., Lauper, S., Maundrell, K., Antonsson, B., and Martinou, J. C. (1999). "Bid-induced conformational change of Bax is responsible for mitochondrial cytochrome c release during apoptosis." *J. Cell Biol.*, 144(5), 891-901.
- Diaz-Hernandez, M., Torres-Peraza, J., Salvatori-Abarca, A., Moran, M. A., Gomez-Ramos, P., Alberch, J., and Lucas, J. J. (2005). "Full motor recovery despite striatal neuron loss and formation of irreversible amyloid-like inclusions in a conditional mouse model of Huntington's disease." *J. Neurosci.*, 25(42), 9773-9781.
- DiFiglia, M., Sapp, E., Chase, K. O., Davies, S. W., Bates, G. P., Vonsattel, J. P., and Aronin, N. (1997). "Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain." *Science*, 277(5334), 1990-1993.
- Dijkers, P. F., Medema, R. H., Lammers, J. W. J., Koenderman, L., and Coffey, P. J. (2000). "Expression of the pro-apoptotic Bcl-2 family member Bim is regulated by the forkhead transcription factor FKHR-L1." *Current Biology*, 10(19), 1201-1204.
- Dijkhuizen, P. A., and Ghosh, A. (2005). "BDNF regulates primary dendrite formation in cortical neurons via the PI3-kinase and MAP kinase signaling pathways." *J. Neurobiol.*, 62(2), 278-288.
- DiStefano, P. S., and Johnson, E. M., Jr. (1988). "Identification of a truncated form of the nerve growth factor receptor." *Proc. Natl. Acad. Sci. U. S. A.*, 85(1), 270-274.

- Dolcet, X., Egea, J., Soler, R. M., Martin-Zanca, D., and Comella, J. X. (1999). "Activation of phosphatidylinositol 3-kinase, but not extracellular-regulated kinases, is necessary to mediate brain-derived neurotrophic factor-induced motoneuron survival." *J. Neurochem.*, 73(2), 521-531.
- Donepudi, M., Mac, S. A., Briand, C., and Grutter, M. G. (2003). "Insights into the regulatory mechanism for caspase-8 activation." *Mol. Cell*, 11(2), 543-549.
- Dragatsis, I., Levine, M. S., and Zeitlin, S. (2000). "Inactivation of Hdh in the brain and testis results in progressive neurodegeneration and sterility in mice." *Nat. Genet.*, 26(3), 300-306.
- Du, K., and Montminy, M. (1998). "CREB is a regulatory target for the protein kinase Akt/PKB." *J. Biol. Chem.*, 273(49), 32377-32379.
- Du, Y., Fischer, T. Z., Clinton-Luke, P., Lercher, L. D., and Dreyfus, C. F. (2006). "Distinct effects of p75 in mediating actions of neurotrophins on basal forebrain oligodendrocytes." *Mol. Cell Neurosci.*, 31(2), 366-375.
- Duan, W., Guo, Z., Jiang, H., Ware, M., Li, X. J., and Mattson, M. P. (2003). "Dietary restriction normalizes glucose metabolism and BDNF levels, slows disease progression, and increases survival in huntingtin mutant mice." *Proc. Natl. Acad. Sci. U. S. A.*, 100(5), 2911-2916.
- Duckworth, B. C., and Cantley, L. C. (1997). "Conditional inhibition of the mitogen-activated protein kinase cascade by wortmannin. Dependence on signal strength." *J. Biol. Chem.*, 272(44), 27665-27670.
- Duyao, M., Ambrose, C., Myers, R., Novelletto, A., Persichetti, F., Frontali, M., Folstein, S., Ross, C., Franz, M., Abbott, M., and . (1993). "Trinucleotide repeat length instability and age of onset in Huntington's disease." *Nat. Genet.*, 4(4), 387-392.
- Duyao, M. P., Auerbach, A. B., Ryan, A., Persichetti, F., Barnes, G. T., McNeil, S. M., Ge, P., Vonsattel, J. P., Gusella, J. F., Joyner, A. L., and . (1995). "Inactivation of the mouse Huntington's disease gene homolog Hdh." *Science*, 269(5222), 407-410.

---

**E**

- Edinger, A. L., and Thompson, C. B. (2004). "Death by design: apoptosis, necrosis and autophagy." *Curr. Opin. Cell Biol.*, 16(6), 663-669.
- Ehlers, M. D. (2003). "Activity level controls postsynaptic composition and signaling via the ubiquitin-proteasome system." *Nat. Neurosci.*, 6(3), 231-242.
- Eide, F. F., Vining, E. R., Eide, B. L., Zang, K., Wang, X. Y., and Reichardt, L. F. (1996). "Naturally occurring truncated trkB receptors have dominant inhibitory effects on brain-derived neurotrophic factor signaling." *J Neurosci*, 16(10), 3123-3129.

- Encinas, M., Tansey, M. G., Tsui-Pierchala, B. A., Comella, J. X., Milbrandt, J., and Johnson, E. M., Jr. (2001). "c-Src is required for glial cell line-derived neurotrophic factor (GDNF) family ligand-mediated neuronal survival via a phosphatidylinositol-3 kinase (PI-3K)-dependent pathway." *J. Neurosci.*, 21(5), 1464-1472.
- Enokido, Y., de Sauvage, F., Hongo, J. A., Ninkina, N., Rosenthal, A., Buchman, V. L., and Davies, A. M. (1998). "GFR alpha-4 and the tyrosine kinase Ret form a functional receptor complex for persephin." *Curr Biol*, 8(18), 1019-1022.
- Erin, N., Bronson, S. K., and Billingsley, M. L. (2003). "Calcium-dependent interaction of calcineurin with Bcl-2 in neuronal tissue." *Neuroscience*, 117(3), 541-555.
- Ernfors, P., Ibanez, C. F., Ebendal, T., Olson, L., and Persson, H. (1990). "Molecular cloning and neurotrophic activities of a protein with structural similarities to nerve growth factor: developmental and topographical expression in the brain." *Proc Natl Acad Sci U S A*, 87(14), 5454-5458.
- Esposti, M. D. (2002). "The roles of Bid." *Apoptosis.*, 7(5), 433-440.

---

**F**

- Farkas, L. M., Suter-Crazzolara, C., and Unsicker, K. (1997). "GDNF induces the calretinin phenotype in cultures of embryonic striatal neurons." *J. Neurosci. Res.*, 50(3), 361-372.
- Feng, L., Wang, C. Y., Jiang, H., Oho, C., Dugich-Djordjevic, M., Mei, L., and Lu, B. (1999). "Differential signaling of glial cell line-derived neurotrophic factor and brain-derived neurotrophic factor in cultured ventral mesencephalic neurons." *Neuroscience*, 93(1), 265-273.
- Ferrante, R. J., Kowall, N. W., Beal, M. F., Richardson, E. P., Jr., Bird, E. D., and Martin, J. B. (1985). "Selective sparing of a class of striatal neurons in Huntington's disease." *Science*, 230(4725), 561-563.
- Ferrante, R. J., Kowall, N. W., Beal, M. F., Martin, J. B., Bird, E. D., and Richardson, E. P., Jr. (1987). "Morphologic and histochemical characteristics of a spared subset of striatal neurons in Huntington's disease." *J. Neuropathol. Exp. Neurol.*, 46(1), 12-27.
- Ferrante, R. J., Kowall, N. W., and Richardson, E. P., Jr. (1991). "Proliferative and degenerative changes in striatal spiny neurons in Huntington's disease: a combined study using the section-Golgi method and calbindin D28k immunocytochemistry." *J. Neurosci.*, 11(12), 3877-3887.
- Ferrante, R. J., Kowall, N. W., Cipolloni, P. B., Storey, E., and Beal, M. F. (1993). "Excitotoxin lesions in primates as a model for Huntington's disease: histopathologic and neurochemical characterization." *Exp. Neurol.*, 119(1), 46-71.
- Ferrer, I., Goutan, E., Marin, C., Rey, M. J., and Ribalta, T. (2000). "Brain-derived neurotrophic factor in Huntington disease." *Brain Res.*, 866(1-2), 257-261.



- Filippa, N., Sable, C. L., Filloux, C., Hemmings, B., and Van, O. E. (1999). "Mechanism of protein kinase B activation by cyclic AMP-dependent protein kinase." *Mol. Cell Biol.*, 19(7), 4989-5000.
- Fischer, W., Wictorin, K., Bjorklund, A., Williams, L. R., Varon, S., and Gage, F. H. (1987). "Amelioration of cholinergic neuron atrophy and spatial memory impairment in aged rats by nerve growth factor." *Nature*, 329(6134), 65-68.
- Fishell, G., and van der Kooy, D. (1991). "Pattern formation in the striatum: neurons with early projections to the substantia nigra survive the cell death period." *J. Comp. Neurol.*, 312(1), 33-42.
- Fletcher, J. I., and Huang, D. C. (2006). "BH3-only proteins: orchestrating cell death." *Cell Death. Differ.*, 13(8), 1268-1271.
- Foghsgaard, L., Wissing, D., Mauch, D., Lademann, U., Bastholm, L., Boes, M., Elling, F., Leist, M., and Jaattela, M. (2001). "Cathepsin B acts as a dominant execution protease in tumor cell apoptosis induced by tumor necrosis factor." *J. Cell Biol.*, 153(5), 999-1010.
- Franke, T. F., Yang, S. I., Chan, T. O., Datta, K., Kazlauskas, A., Morrison, D. K., Kaplan, D. R., and Tsichlis, P. N. (1995). "The protein kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated phosphatidylinositol 3-kinase." *Cell*, 81(5), 727-736.
- Fuller, G., Veitch, K., Ho, L. K., Cruise, L., and Morris, B. J. (2001). "Activation of p44/p42 MAP kinase in striatal neurons via kainate receptors and PI3 kinase." *Brain Res. Mol. Brain Res.*, 89(1-2), 126-132.

---

**G**

- Gauthier, L. R., Charrin, B. C., Borrell-Pages, M., Dompierre, J. P., Rangone, H., Cordelieres, F. P., De, M. J., MacDonald, M. E., Lessmann, V., Humbert, S., and Saudou, F. (2004). "Huntingtin controls neurotrophic support and survival of neurons by enhancing BDNF vesicular transport along microtubules." *Cell*, 118(1), 127-138.
- Gavalda, N., Perez-Navarro, E., Gratacos, E., Comella, J. X., and Alberch, J. (2004). "Differential involvement of phosphatidylinositol 3-kinase and p42/p44 mitogen activated protein kinase pathways in brain-derived neurotrophic factor-induced trophic effects on cultured striatal neurons." *Mol. Cell Neurosci.*, 25(3), 460-468.
- Gerfen, C. R. (1992). "The neostriatal mosaic: multiple levels of compartmental organization." *Trends Neurosci.*, 15(4), 133-139.
- Gervais, F. G., Singaraja, R., Xanthoudakis, S., Gutekunst, C. A., Leavitt, B. R., Metzler, M., Hackam, A. S., Tam, J., Vaillancourt, J. P., Houtzager, V., Rasper, D. M., Roy, S., Hayden, M. R., and Nicholson, D. W. (2002). "Recruitment and activation of caspase-8 by the Huntingtin-interacting protein Hip-1 and a novel partner Hippi." *Nat. Cell Biol.*, 4(2), 95-105.
- Ghasemzadeh, M. B., Sharma, S., Surmeier, D. J., Eberwine, J. H., and Chesselet, M. F. (1996). "Multiplicity of glutamate receptor subunits in single striatal neurons: an RNA amplification study." *Mol. Pharmacol.*, 49(5), 852-859.

- Gille, H., Kortenjann, M., Thomae, O., Moomaw, C., Slaughter, C., Cobb, M. H., and Shaw, P. E. (1995). "ERK phosphorylation potentiates Elk-1-mediated ternary complex formation and transactivation." *EMBO J.*, 14(5), 951-962.
- Gines, S., Seong, I. S., Fossale, E., Ivanova, E., Trettel, F., Gusella, J. F., Wheeler, V. C., Persichetti, F., and MacDonald, M. E. (2003a). "Specific progressive cAMP reduction implicates energy deficit in presymptomatic Huntington's disease knock-in mice." *Hum. Mol. Genet.*, 12(5), 497-508.
- Gines, S., Ivanova, E., Seong, I. S., Saura, C. A., and MacDonald, M. E. (2003b). "Enhanced Akt signaling is an early pro-survival response that reflects N-methyl-D-aspartate receptor activation in Huntington's disease knock-in striatal cells." *J. Biol. Chem.*, 278(50), 50514-50522.
- Gines, S., Bosch, M., Marco, S., Gavaldà, N., Az-Hernandez, M., Lucas, J. J., Canals, J. M., and Alberch, J. (2006). "Reduced expression of the TrkB receptor in Huntington's disease mouse models and in human brain." *Eur. J. Neurosci.*, 23(3), 649-658.
- Giovannini, M. G., Blitzer, R. D., Wong, T., Asoma, K., Tsokas, P., Morrison, J. H., Iyengar, R., and Landau, E. M. (2001). "Mitogen-activated protein kinase regulates early phosphorylation and delayed expression of Ca<sup>2+</sup>/calmodulin-dependent protein kinase II in long-term potentiation." *J. Neurosci.*, 21(18), 7053-7062.
- Glass, M., Dragunow, M., and Faull, R. L. (2000). "The pattern of neurodegeneration in Huntington's disease: a comparative study of cannabinoid, dopamine, adenosine and GABA(A) receptor alterations in the human basal ganglia in Huntington's disease." *Neuroscience*, 97(3), 505-519.
- Goldin, M., and Segal, M. (2003). "Protein kinase C and ERK involvement in dendritic spine plasticity in cultured rodent hippocampal neurons." *Eur. J. Neurosci.*, 17(12), 2529-2539.
- Gomez-Santos, C., Ferrer, I., Reiriz, J., Vinals, F., Barrachina, M., and Ambrosio, S. (2002). "MPP<sup>+</sup> increases alpha-synuclein expression and ERK/MAP-kinase phosphorylation in human neuroblastoma SH-SY5Y cells." *Brain Res.*, 935(1-2), 32-39.
- Gonzalez, F. A., Seth, A., Raden, D. L., Bowman, D. S., Fay, F. S., and Davis, R. J. (1993). "Serum-induced translocation of mitogen-activated protein kinase to the cell surface ruffling membrane and the nucleus." *J. Cell Biol.*, 122(5), 1089-1101.
- Gottlieb, T. M., Leal, J. F., Seger, R., Taya, Y., and Oren, M. (2002). "Cross-talk between Akt, p53 and Mdm2: possible implications for the regulation of apoptosis." *Oncogene*, 21(8), 1299-1303.
- Gratacos, E., Checa, N., and Alberch, J. (2001a). "Bone morphogenetic protein-2, but not bone morphogenetic protein-7, promotes dendritic growth and calbindin phenotype in cultured rat striatal neurons." *Neuroscience*, 104(3), 783-790.
- Gratacos, E., Perez-Navarro, E., Tolosa, E., Arenas, E., and Alberch, J. (2001b). "Neuroprotection of striatal neurons against kainate excitotoxicity by neurotrophins and GDNF family members." *J. Neurochem.*, 78(6), 1287-1296.

- Gratacos, E., Checa, N., Perez-Navarro, E., and Alberch, J. (2001c). "Brain-derived neurotrophic factor (BDNF) mediates bone morphogenetic protein-2 (BMP-2) effects on cultured striatal neurones." *J. Neurochem.*, 79(4), 747-755.
- Gratacos, E., Gavalda, N., and Alberch, J. (2002). "Bone morphogenetic protein-6 is a neurotrophic factor for calbindin-positive striatal neurons." *J. Neurosci. Res.*, 70(5), 638-644.
- Green, D. R., and Kroemer, G. (2004). "The pathophysiology of mitochondrial cell death." *Science*, 305(5684), 626-629.
- Greene, J. G., and Greenamyre, J. T. (1995). "Characterization of the excitotoxic potential of the reversible succinate dehydrogenase inhibitor malonate." *J. Neurochem.*, 64(1), 430-436.
- Grewal, S. S., York, R. D., and Stork, P. J. (1999). "Extracellular-signal-regulated kinase signalling in neurons." *Curr. Opin. Neurobiol.*, 9(5), 544-553.
- Griffiths, G. J., Dubrez, L., Morgan, C. P., Jones, N. A., Whitehouse, J., Corfe, B. M., Dive, C., and Hickman, J. A. (1999). "Cell damage-induced conformational changes of the pro-apoptotic protein bak in vivo precede the onset of apoptosis." *Journal of Cell Biology*, 144(5), 903-914.
- Grimes, M. L., Zhou, J., Beattie, E. C., Yuen, E. C., Hall, D. E., Valletta, J. S., Topp, K. S., LaVail, J. H., Bunnett, N. W., and Mobley, W. C. (1996). "Endocytosis of activated TrkA: evidence that nerve growth factor induces formation of signaling endosomes." *J. Neurosci.*, 16(24), 7950-7964.
- Gross, A., Jockel, J., Wei, M. C., and Korsmeyer, S. J. (1998). "Enforced dimerization of BAX results in its translocation, mitochondrial dysfunction and apoptosis." *Embo Journal*, 17(14), 3878-3885.
- Gu, M., Gash, M. T., Mann, V. M., Javoy-Agid, F., Cooper, J. M., and Schapira, A. H. (1996). "Mitochondrial defect in Huntington's disease caudate nucleus." *Ann. Neurol.*, 39(3), 385-389.
- Guo, B., Zhai, D., Cabezas, E., Welsh, K., Nouraini, S., Satterthwait, A. C., and Reed, J. C. (2003). "Humanin peptide suppresses apoptosis by interfering with Bax activation." *Nature*, 423(6938), 456-461.

---

## H

- Hahn, P., Lindsten, T., Lyubarsky, A., Ying, G. S., Pugh, E. N., Jr., Thompson, C. B., and Dunaief, J. L. (2004). "Deficiency of Bax and Bak protects photoreceptors from light damage in vivo." *Cell Death. Differ.*, 11(11), 1192-1197.
- Hallbook, F., Ibanez, C. F., and Persson, H. (1991). "Evolutionary studies of the nerve growth factor family reveal a novel member abundantly expressed in *Xenopus* ovary." *Neuron*, 6(5), 845-858.
- Han, B. H., and Holtzman, D. M. (2000). "BDNF protects the neonatal brain from hypoxic-ischemic injury in vivo via the ERK pathway." *J. Neurosci.*, 20(15), 5775-5781.

- Hara, T., Nakamura, K., Matsui, M., Yamamoto, A., Nakahara, Y., Suzuki-Migishima, R., Yokoyama, M., Mishima, K., Saito, I., Okano, H., and Mizushima, N. (2006). "Suppression of basal autophagy in neural cells causes neurodegenerative disease in mice." *Nature*, 441(7095), 885-889.
- Harjes, P., and Wanker, E. E. (2003). "The hunt for huntingtin function: interaction partners tell many different stories." *Trends Biochem. Sci.*, 28(8), 425-433.
- Hassouna, I., Wickert, H., Zimmermann, M., and Gillardon, F. (1996). "Increase in bax expression in substantia nigra following 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) treatment of mice." *Neurosci. Lett.*, 204(1-2), 85-88.
- Hefti, F. (1994). "Development of effective therapy for Alzheimer's disease based on neurotrophic factors." *Neurobiol. Aging*, 15 Suppl 2, S193-S194.
- Henderson, C. E., Phillips, H. S., Pollock, R. A., Davies, A. M., Lemeulle, C., Armanini, M., Simmons, L., Moffet, B., Vandlen, R. A., Simpson LC [corrected to Simmons, and . (1994). "GDNF: a potent survival factor for motoneurons present in peripheral nerve and muscle." *Science*, 266(5187), 1062-1064.
- Hengartner, M. O. (2000). "The biochemistry of apoptosis." *Nature*, 407(6805), 770-776.
- Henshaw, R., Jenkins, B. G., Schulz, J. B., Ferrante, R. J., Kowall, N. W., Rosen, B. R., and Beal, M. F. (1994). "Malonate produces striatal lesions by indirect NMDA receptor activation." *Brain Res.*, 647(1), 161-166.
- Hetman, M., Kanning, K., Cavanaugh, J. E., and Xia, Z. (1999). "Neuroprotection by brain-derived neurotrophic factor is mediated by extracellular signal-regulated kinase and phosphatidylinositol 3-kinase." *J. Biol. Chem.*, 274(32), 22569-22580.
- Hill, M. M., Andjelkovic, M., Brazil, D. P., Ferrari, S., Fabbro, D., and Hemmings, B. A. (2001). "Insulin-stimulated protein kinase B phosphorylation on Ser-473 is independent of its activity and occurs through a staurosporine-insensitive kinase." *J. Biol. Chem.*, 276(28), 25643-25646.
- Hirsch, T., Marchetti, P., Susin, S. A., Dallaporta, B., Zamzami, N., Marzo, I., Geuskens, M., and Kroemer, G. (1997). "The apoptosis-necrosis paradox. Apoptogenic proteases activated after mitochondrial permeability transition determine the mode of cell death." *Oncogene*, 15(13), 1573-1581.
- Hodgson, J. G., Agopyan, N., Gutekunst, C. A., Leavitt, B. R., LePiane, F., Singaraja, R., Smith, D. J., Bissada, N., McCutcheon, K., Nasir, J., Jamot, L., Li, X. J., Stevens, M. E., Rosemond, E., Roder, J. C., Phillips, A. G., Rubin, E. M., Hersch, S. M., and Hayden, M. R. (1999). "A YAC mouse model for Huntington's disease with full-length mutant huntingtin, cytoplasmic toxicity, and selective striatal neurodegeneration." *Neuron*, 23(1), 181-192.
- Hohn, A., Leibrock, J., Bailey, K., and Barde, Y. A. (1990). "Identification and characterization of a novel member of the nerve growth factor/brain-derived neurotrophic factor family." *Nature*, 344(6264), 339-341.

- Holgado-Madruga, M., Moscatello, D. K., Emler, D. R., Dieterich, R., and Wong, A. J. (1997). "Grb2-associated binder-1 mediates phosphatidylinositol 3-kinase activation and the promotion of cell survival by nerve growth factor." *Proc. Natl. Acad. Sci. U. S. A.*, 94(23), 12419-12424.
- Hollmann, M., and Heinemann, S. (1994). "Cloned glutamate receptors." *Annu. Rev. Neurosci.*, 17, 31-108.
- Holtzman, D. M., Sheldon, R. A., Jaffe, W., Cheng, Y., and Ferriero, D. M. (1996). "Nerve growth factor protects the neonatal brain against hypoxic-ischemic injury." *Ann. Neurol.*, 39(1), 114-122.
- Hornung, J. P., Koppel, H., and Clarke, P. G. (1989). "Endocytosis and autophagy in dying neurons: an ultrastructural study in chick embryos." *J. Comp Neurol.*, 283(3), 425-437.
- Horvitz, H. R. (2003). "Worms, life, and death (Nobel lecture)." *ChemBiochem.*, 4(8), 697-711.
- Hsu, Y. T., Wolter, K. G., and Youle, R. J. (1997). "Cytosol-to-membrane redistribution of Bax and Bcl-X-L during apoptosis." *Proceedings of the National Academy of Sciences of the United States of America*, 94(8), 3668-3672.
- Huang, D. C. S., and Strasser, A. (2000). "BH3-Only proteins - Essential initiators of apoptotic cell death." *Cell*, 103(6), 839-842.
- Huang, E. J., and Reichardt, L. F. (2001). "Neurotrophins: Roles in neuronal development and function." *Annual Review of Neuroscience*, 24, 677-736.
- Humbert, S., Bryson, E. A., Cordelieres, F. P., Connors, N. C., Datta, S. R., Finkbeiner, S., Greenberg, M. E., and Saudou, F. (2002). "The IGF-1/Akt pathway is neuroprotective in Huntington's disease and involves Huntingtin phosphorylation by Akt." *Dev. Cell*, 2(6), 831-837.
- Humpel, C., Marksteiner, J., and Saria, A. (1996). "Glial-cell-line-derived neurotrophic factor enhances biosynthesis of substance P in striatal neurons in vitro." *Cell Tissue Res.*, 286(2), 249-255.
- Huntington G. (1872). "On chorea". *Med. Surg. Reporter* 26, 302-321.
- Hurlbert, M. S., Zhou, W., Wasmeier, C., Kaddis, F. G., Hutton, J. C., and Freed, C. R. (1999). "Mice transgenic for an expanded CAG repeat in the Huntington's disease gene develop diabetes." *Diabetes*, 48(3), 649-651.
- Ichinose, T., and Snider, W. D. (2000). "Differential effects of TrkC isoforms on sensory axon outgrowth." *J. Neurosci. Res.*, 59(3), 365-371.
- Imaizumi, K., Tsuda, M., Wanaka, A., Tohyama, M., and Takagi, T. (1994). "Differential expression of sgk mRNA, a member of the Ser/Thr protein kinase gene family, in rat brain after CNS injury." *Brain Res. Mol. Brain Res.*, 26(1-2), 189-196.
- Imaizumi, K., Morihara, T., Mori, Y., Katayama, T., Tsuda, M., Furuyama, T., Wanaka, A., Takeda, M., and Tohyama, M. (1999). "The cell death-promoting gene DP5, which interacts with the BCL2 family, is induced during neuronal apoptosis following exposure to amyloid beta protein." *J. Biol. Chem.*, 274(12), 7975-7981.

---

Ip, N. Y., Ibanez, C. F., Nye, S. H., McClain, J., Jones, P. F., Gies, D. R., Belluscio, L., Le Beau, M. M., Espinosa, R., and Squinto, S. P. (1992). "Mammalian neurotrophin-4: structure, chromosomal localization, tissue distribution, and receptor specificity." *Proc. Natl. Acad. Sci. U. S. A.*, 89(7), 3060-3064.

Ip, Y. T., and Davis, R. J. (1998). "Signal transduction by the c-Jun N-terminal kinase (JNK)--from inflammation to development." *Curr. Opin. Cell Biol.*, 10(2), 205-219.

Irmeler, M., Thome, M., Hahne, M., Schneider, P., Hofmann, K., Steiner, V., Bodmer, J. L., Schroter, M., Burns, K., Mattmann, C., Rimoldi, D., French, L. E., and Tschopp, J. (1997). "Inhibition of death receptor signals by cellular FLIP." *Nature*, 388(6638), 190-195.

---

Jenkins, B. G., Koroshetz, W. J., Beal, M. F., and Rosen, B. R. (1993). "Evidence for impairment of energy metabolism in vivo in Huntington's disease using localized <sup>1</sup>H NMR spectroscopy." *Neurology*, 43(12), 2689-2695.

Jenkins, B. G., Rosas, H. D., Chen, Y. C., Makabe, T., Myers, R., MacDonald, M., Rosen, B. R., Beal, M. F., and Koroshetz, W. J. (1998). "<sup>1</sup>H NMR spectroscopy studies of Huntington's disease: correlations with CAG repeat numbers." *Neurology*, 50(5), 1357-1365.

Jin, K. L., Mao, X. O., Nagayama, T., Goldsmith, P. C., and Greenberg, D. A. (2000). "Induction of vascular endothelial growth factor receptors and phosphatidylinositol 3'-kinase/Akt signaling by global cerebral ischemia in the rat." *Neuroscience*, 100(4), 713-717.

Jin, Z., and El-Deiry, W. S. (2005). "Overview of cell death signaling pathways." *Cancer Biol. Ther.*, 4(2), 139-163.

Jing, S., Wen, D., Yu, Y., Holst, P. L., Luo, Y., Fang, M., Tamir, R., Antonio, L., Hu, Z., Cupples, R., Louis, J. C., Hu, S., Altrock, B. W., and Fox, G. M. (1996). "GDNF-induced activation of the ret protein tyrosine kinase is mediated by GDNFR-alpha, a novel receptor for GDNF." *Cell*, 85(7), 1113-1124.

Johnson, D. E. (2000). "Noncaspase proteases in apoptosis." *Leukemia*, 14(9), 1695-1703.

Jones, P. F., Jakubowicz, T., Pitossi, F. J., Maurer, F., and Hemmings, B. A. (1991a). "Molecular cloning and identification of a serine/threonine protein kinase of the second-messenger subfamily." *Proc. Natl. Acad. Sci. U. S. A.*, 88(10), 4171-4175.

Jones, P. F., Jakubowicz, T., and Hemmings, B. A. (1991b). "Molecular cloning of a second form of rac protein kinase." *Cell Regul.*, 2(12), 1001-1009.

Jung, K. M., Tan, S., Landman, N., Petrova, K., Murray, S., Lewis, R., Kim, P. K., Kim, D. S., Ryu, S. H., Chao, M. V., and Kim, T. W. (2003). "Regulated intramembrane proteolysis of the p75 neurotrophin receptor modulates its association with the TrkA receptor." *J. Biol. Chem.*, 278(43), 42161-42169.

**K**

Kane, L. P., Shapiro, V. S., Stokoe, D., and Weiss, A. (1999). "Induction of NF-kappaB by the Akt/PKB kinase." *Curr. Biol.*, 9(11), 601-604.

Kawaguchi, Y., Wilson, C. J., Augood, S. J., and Emson, P. C. (1995). "Striatal interneurons: chemical, physiological and morphological characterization." *Trends Neurosci.*, 18(12), 527-535.

Kawasaki, H., Morooka, T., Shimohama, S., Kimura, J., Hirano, T., Gotoh, Y., and Nishida, E. (1997). "Activation and involvement of p38 mitogen-activated protein kinase in glutamate-induced apoptosis in rat cerebellar granule cells." *J. Biol. Chem.*, 272(30), 18518-18521.

Kelleher, R. J., III, Govindarajan, A., Jung, H. Y., Kang, H., and Tonegawa, S. (2004). "Translational control by MAPK signaling in long-term synaptic plasticity and memory." *Cell*, 116(3), 467-479.

Kells, A. P., Fong, D. M., Dragunow, M., During, M. J., Young, D., and Connor, B. (2004). "AAV-mediated gene delivery of BDNF or GDNF is neuroprotective in a model of Huntington disease." *Mol. Ther.*, 9(5), 682-688.

Kerr, J. F. R., Wyllie, A. H., and Currie, A. R. (1972). "Apoptosis - Basic Biological Phenomenon with Wide-Ranging Implications in Tissue Kinetics." *British Journal of Cancer*, 26(4), 239-&.

Kim, A. H., Khursigara, G., Sun, X., Franke, T. F., and Chao, M. V. (2001). "Akt phosphorylates and negatively regulates apoptosis signal-regulating kinase 1." *Mol. Cell Biol.*, 21(3), 893-901.

Kim, M., Lee, H. S., Laforet, G., McIntyre, C., Martin, E. J., Chang, P., Kim, T. W., Williams, M., Reddy, P. H., Tagle, D., Boyce, F. M., Won, L., Heller, A., Aronin, N., and DiFiglia, M. (1999). "Mutant huntingtin expression in clonal striatal cells: dissociation of inclusion formation and neuronal survival by caspase inhibition." *J. Neurosci.*, 19(3), 964-973.

Kim, R. (2005). "Unknotting the roles of Bcl-2 and Bcl-xL in cell death." *Biochem. Biophys. Res. Commun.*, 333(2), 336-343.

Kohn, A. D., Kovacina, K. S., and Roth, R. A. (1995). "Insulin stimulates the kinase activity of RAC-PK, a pleckstrin homology domain containing ser/thr kinase." *EMBO J.*, 14(17), 4288-4295.

Kokaia, Z., Bengzon, J., Metsis, M., Kokaia, M., Persson, H., and Lindvall, O. (1993). "Coexpression of neurotrophins and their receptors in neurons of the central nervous system." *Proc. Natl. Acad. Sci. U. S. A.*, 90(14), 6711-6715.

Kolch, W. (2005). "Coordinating ERK/MAPK signalling through scaffolds and inhibitors." *Nat. Rev. Mol. Cell Biol.*, 6(11), 827-837.

- Komatsu, M., Waguri, S., Chiba, T., Murata, S., Iwata, J., Tanida, I., Ueno, T., Koike, M., Uchiyama, Y., Kominami, E., and Tanaka, K. (2006). "Loss of autophagy in the central nervous system causes neurodegeneration in mice." *Nature*, 441(7095), 880-884.
- Konishi, H., Matsuzaki, H., Tanaka, M., Takemura, Y., Kuroda, S., Ono, Y., and Kikkawa, U. (1997). "Activation of protein kinase B (Akt/RAC-protein kinase) by cellular stress and its association with heat shock protein Hsp27." *FEBS Lett.*, 410(2-3), 493-498.
- Kordower, J. H., Emborg, M. E., Bloch, J., Ma, S. Y., Chu, Y., Leventhal, L., McBride, J., Chen, E. Y., Palfi, S., Roitberg, B. Z., Brown, W. D., Holden, J. E., Pyzalski, R., Taylor, M. D., Carvey, P., Ling, Z., Trono, D., Hantraye, P., Deglon, N., and Aebischer, P. (2000). "Neurodegeneration prevented by lentiviral vector delivery of GDNF in primate models of Parkinson's disease." *Science*, 290(5492), 767-773.
- Kotzbauer, P. T., Lampe, P. A., Heuckeroth, R. O., Golden, J. P., Creedon, D. J., Johnson, E. M. J., and Milbrandt, J. (1996). "Neurturin, a relative of glial-cell-line-derived neurotrophic factor." *Nature*, 384(6608), 467-470.
- Krajewski, S., Tanaka, S., Takayama, S., Schibler, M. J., Fenton, W., and Reed, J. C. (1993). "Investigation of the Subcellular-Distribution of the Bcl-2 Oncoprotein - Residence in the Nuclear-Envelope, Endoplasmic-Reticulum, and Outer Mitochondrial-Membranes." *Cancer Research*, 53(19), 4701-4714.
- Krajewski, S., Mai, J. K., Krajewska, M., Sikorska, M., Mossakowski, M. J., and Reed, J. C. (1995). "Upregulation of bax protein levels in neurons following cerebral ischemia." *J. Neurosci.*, 15(10), 6364-6376.
- Krantic, S., Mechawar, N., Reix, S., and Quirion, R. (2005). "Molecular basis of programmed cell death involved in neurodegeneration." *Trends Neurosci.*, 28(12), 670-676.
- Kuida, K., Zheng, T. S., Na, S., Kuan, C., Yang, D., Karasuyama, H., Rakic, P., and Flavell, R. A. (1996). "Decreased apoptosis in the brain and premature lethality in CPP32-deficient mice." *Nature*, 384(6607), 368-372.
- Kuida, K., Haydar, T. F., Kuan, C. Y., Gu, Y., Taya, C., Karasuyama, H., Su, M. S., Rakic, P., and Flavell, R. A. (1998). "Reduced apoptosis and cytochrome c-mediated caspase activation in mice lacking caspase 9." *Cell*, 94(3), 325-337.
- Kulich, S. M., and Chu, C. T. (2001). "Sustained extracellular signal-regulated kinase activation by 6-hydroxydopamine: implications for Parkinson's disease." *J. Neurochem.*, 77(4), 1058-1066.
- Kuwana, T., Mackey, M. R., Perkins, G., Ellisman, M. H., Latterich, M., Schneiter, R., Green, D. R., and Newmeyer, D. D. (2002). "Bid, Bax, and lipids cooperate to form supramolecular openings in the outer mitochondrial membrane." *Cell*, 111(3), 331-342.



- Landles, C., and Bates, G. P. (2004). "Huntingtin and the molecular pathogenesis of Huntington's disease. Fourth in molecular medicine review series." *EMBO Rep.*, 5(10), 958-963.
- Landwehrmeyer, G. B., Standaert, D. G., Testa, C. M., Penney, J. B., Jr., and Young, A. B. (1995). "NMDA receptor subunit mRNA expression by projection neurons and interneurons in rat striatum." *J. Neurosci.*, 15(7 Pt 2), 5297-5307.
- Lankiewicz, S., Marc, L. C., Truc, B. N., Krohn, A. J., Poppe, M., Cole, G. M., Saido, T. C., and Prehn, J. H. (2000). "Activation of calpain I converts excitotoxic neuron death into a caspase-independent cell death." *J. Biol. Chem.*, 275(22), 17064-17071.
- Leavitt, B. R., Wellington, C. L., and Hayden, M. R. (1999). "Recent insights into the molecular pathogenesis of Huntington disease." *Semin. Neurol.*, 19(4), 385-395.
- LeBlanc, A. C. (2003). "Natural cellular inhibitors of caspases." *Prog. Neuropsychopharmacol. Biol. Psychiatry*, 27(2), 215-229.
- Lee, R., Kermani, P., Teng, K. K., and Hempstead, B. L. (2001). "Regulation of cell survival by secreted proneurotrophins." *Science*, 294(5548), 1945-1948.
- Leeds, P., Leng, Y., Chalecka-Franaszek, E., and Chuang, D. M. (2005). "Neurotrophins protect against cytosine arabinoside-induced apoptosis of immature rat cerebellar neurons." *Neurochem. Int.*, 46(1), 61-72.
- Leibrock, J., Lottspeich, F., Hohn, A., Hofer, M., Hengerer, B., Masiakowski, P., Thoenen, H., and Barde, Y. A. (1989). "Molecular cloning and expression of brain-derived neurotrophic factor." *Nature*, 341(6238), 149-152.
- Lenormand, P., Sardet, C., Pages, G., L'Allemain, G., Brunet, A., and Pouyssegur, J. (1993). "Growth factors induce nuclear translocation of MAP kinases (p42mapk and p44mapk) but not of their activator MAP kinase kinase (p45mapkk) in fibroblasts." *J. Cell Biol.*, 122(5), 1079-1088.
- Leong, M. L., Maiyar, A. C., Kim, B., O'Keeffe, B. A., and Firestone, G. L. (2003). "Expression of the serum- and glucocorticoid-inducible protein kinase, Sgk, is a cell survival response to multiple types of environmental stress stimuli in mammary epithelial cells." *J. Biol. Chem.*, 278(8), 5871-5882.
- Letai, A., Bassik, M. C., Walensky, L. D., Sorcinelli, M. D., Weiler, S., and Korsmeyer, S. J. (2002). "Distinct BH3 domains either sensitize or activate mitochondrial apoptosis, serving as prototype cancer therapeutics." *Cancer Cell*, 2(3), 183-192.
- Levi-Montalcini, R., and Angeletti, P. U. (1968). "Nerve growth factor." *Physiol Rev*, 48(3), 534-569.
- Levine, B., and Klionsky, D. J. (2004). "Development by self-digestion: molecular mechanisms and biological functions of autophagy." *Dev. Cell*, 6(4), 463-477.

- Levine, M. S., Klapstein, G. J., Koppel, A., Gruen, E., Cepeda, C., Vargas, M. E., Jokel, E. S., Carpenter, E. M., Zanjani, H., Hurst, R. S., Efstratiadis, A., Zeitlin, S., and Chesselet, M. F. (1999). "Enhanced sensitivity to N-methyl-D-aspartate receptor activation in transgenic and knockin mouse models of Huntington's disease." *J. Neurosci. Res.*, 58(4), 515-532.
- Ley, R., Balmanno, K., Hadfield, K., Weston, C., and Cook, S. J. (2003). "Activation of the ERK1/2 signaling pathway promotes phosphorylation and proteasome-dependent degradation of the BH3-only protein, Bim." *Journal of Biological Chemistry*, 278(21), 18811-18816.
- Li, H., Zhu, H., Xu, C. J., and Yuan, J. (1998). "Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis." *Cell*, 94(4), 491-501.
- Li, S. H., and Li, X. J. (2004). "Huntingtin-protein interactions and the pathogenesis of Huntington's disease." *Trends Genet.*, 20(3), 146-154.
- Li, Z., Zhang, J., Liu, Z., Woo, C. W., and Thiele, C. J. (2007). "Downregulation of Bim by brain-derived neurotrophic factor activation of TrkB protects neuroblastoma cells from paclitaxel but not etoposide or cisplatin-induced cell death." *Cell Death. Differ.*, 14(2), 318-326.
- Lin, A. (2003). "Activation of the JNK signaling pathway: breaking the brake on apoptosis." *Bioessays*, 25(1), 17-24.
- Lin, L. F., Doherty, D. H., Lile, J. D., Bektesh, S., and Collins, F. (1993). "GDNF: a glial cell line-derived neurotrophic factor for midbrain dopaminergic neurons." *Science*, 260(5111), 1130-1132.
- Lindsten, T., Ross, A. J., King, A., Zong, W. X., Rathmell, J. C., Shiels, H. A., Ulrich, E., Waymire, K. G., Mahar, P., Frauwirth, K., Chen, Y. F., Wei, M., Eng, V. M., Adelman, D. M., Simon, M. C., Ma, A., Golden, J. A., Evan, G., Korsmeyer, S. J., MacGregor, G. R., and Thompson, C. B. (2000). "The combined functions of proapoptotic Bcl-2 family members Bak and Bax are essential for normal development of multiple tissues." *Molecular Cell*, 6(6), 1389-1399.
- Lord, S. J., Rajotte, R. V., Korbitt, G. S., and Bleackley, R. C. (2003). "Granzyme B: a natural born killer." *Immunol. Rev.*, 193, 31-38.
- Lovat, P. E., Oliverio, S., Corazzari, M., Rodolfo, C., Ranalli, M., Goranov, B., Melino, G., Redfern, C. P., and Piacentini, M. (2003). "Bak: a downstream mediator of fenretinide-induced apoptosis of SH-SY5Y neuroblastoma cells." *Cancer Res.*, 63(21), 7310-7313.
- Luciano, F., Jacquet, A., Colosetti, P., Herrant, M., Cagnol, S., Pages, G., and Auberger, P. (2003). "Phosphorylation of Bim-EL by Erk1/2 on serine 69 promotes its degradation via the proteasome pathway and regulates its proapoptotic function." *Oncogene*, 22(43), 6785-6793.
- Luo, X., Budihardjo, I., Zou, H., Slaughter, C., and Wang, X. (1998). "Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors." *Cell*, 94(4), 481-490.

Luthi-Carter, R., Strand, A., Peters, N. L., Solano, S. M., Hollingsworth, Z. R., Menon, A. S., Frey, A. S., Spektor, B. S., Penney, E. B., Schilling, G., Ross, C. A., Borchelt, D. R., Tapscott, S. J., Young, A. B., Cha, J. H., and Olson, J. M. (2000). "Decreased expression of striatal signaling genes in a mouse model of Huntington's disease." *Hum. Mol. Genet.*, 9(9), 1259-1271.

Luthi-Carter, R., Hanson, S. A., Strand, A. D., Bergstrom, D. A., Chun, W., Peters, N. L., Woods, A. M., Chan, E. Y., Kooperberg, C., Krainc, D., Young, A. B., Tapscott, S. J., and Olson, J. M. (2002). "Dysregulation of gene expression in the R6/2 model of polyglutamine disease: parallel changes in muscle and brain." *Hum. Mol. Genet.*, 11(17), 1911-1926.

---

**M**

Macdonald, V., and Halliday, G. (2002). "Pyramidal cell loss in motor cortices in Huntington's disease." *Neurobiol. Dis.*, 10(3), 378-386.

MacEwan, D. J. (2002). "TNF ligands and receptors--a matter of life and death." *Br. J. Pharmacol.*, 135(4), 855-875.

Maisonpierre, P. C., Belluscio, L., Squinto, S., Ip, N. Y., Furth, M. E., Lindsay, R. M., and Yancopoulos, G. D. (1990). "Neurotrophin-3: a neurotrophic factor related to NGF and BDNF." *Science*, 247(4949 Pt 1), 1446-1451.

Mandic, A., Viktorsson, K., Strandberg, L., Heiden, T., Hansson, J., Linder, S., and Shoshan, M. C. (2002). "Calpain-mediated Bid cleavage and calpain-independent Bak modulation: two separate pathways in cisplatin-induced apoptosis." *Mol. Cell Biol.*, 22(9), 3003-3013.

Mangiarini, L., Sathasivam, K., Seller, M., Cozens, B., Harper, A., Hetherington, C., Lawton, M., Trotter, Y., Lehrach, H., Davies, S. W., and Bates, G. P. (1996). "Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice." *Cell*, 87(3), 493-506.

Mann, D. M., Oliver, R., and Snowden, J. S. (1993). "The topographic distribution of brain atrophy in Huntington's disease and progressive supranuclear palsy." *Acta Neuropathol. (Berl)*, 85(5), 553-559.

Mao, L., Tang, Q., Samdani, S., Liu, Z., and Wang, J. Q. (2004). "Regulation of MAPK/ERK phosphorylation via ionotropic glutamate receptors in cultured rat striatal neurons." *Eur. J. Neurosci.*, 19(5), 1207-1216.

Marco, S., Perez-Navarro, E., Tolosa, E., Arenas, E., and Alberch, J. (2002a). "Striatopallidal neurons are selectively protected by neurturin in an excitotoxic model of Huntington's disease." *J. Neurobiol.*, 50(4), 323-332.

Marco, S., Canudas, A. M., Canals, J. M., Gavalda, N., Perez-Navarro, E., and Alberch, J. (2002b). "Excitatory amino acids differentially regulate the expression of GDNF, neurturin, and their receptors in the adult rat striatum." *Exp. Neurol.*, 174(2), 243-252.

- Mariathasan, S., Zakarian, A., Bouchard, D., Michie, A. M., Zuniga-Pflucker, J. C., and Ohashi, P. S. (2001). "Duration and strength of extracellular signal-regulated kinase signals are altered during positive versus negative thymocyte selection." *J. Immunol.*, 167(9), 4966-4973.
- Markus, A., Zhong, J., and Snider, W. D. (2002). "Raf and akt mediate distinct aspects of sensory axon growth." *Neuron*, 35(1), 65-76.
- Marshall, C. J. (1995). "Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation." *Cell*, 80(2), 179-185.
- Marshall, M. S. (1995). "Ras target proteins in eukaryotic cells." *FASEB J.*, 9(13), 1311-1318.
- Martin, J. B., and Gusella, J. F. (1986). "Huntington's disease. Pathogenesis and management." *N. Engl. J. Med.*, 315(20), 1267-1276.
- Martin, L. J. (2001). "Neuronal cell death in nervous system development, disease, and injury (Review)." *Int. J. Mol. Med.*, 7(5), 455-478.
- Martinez-Serrano, A., and Bjorklund, A. (1996). "Protection of the neostriatum against excitotoxic damage by neurotrophin-producing, genetically modified neural stem cells." *J. Neurosci.*, 16(15), 4604-4616.
- Martinez-Vicente, M., Sovak, G., and Cuervo, A. M. (2005). "Protein degradation and aging." *Exp. Gerontol.*, 40(8-9), 622-633.
- Masure, S., Cik, M., Pangalos, M. N., Bonaventure, P., Verhasselt, P., Lesage, A. S., Leysen, J. E., and Gordon, R. D. (1998). "Molecular cloning, expression and tissue distribution of glial-cell-line-derived neurotrophic factor family receptor alpha-3 (GFRalpha-3)." *Eur J Biochem*, 251(3), 622-630.
- Masure, S., Cik, M., Hoefnagel, E., Nosrat, C. A., Van, D. L., I, Scott, R., Van Gompel, P., Lesage, A. S., Verhasselt, P., Ibanez, C. F., and Gordon, R. D. (2000). "Mammalian GFRalpha -4, a divergent member of the GFRalpha family of coreceptors for glial cell line-derived neurotrophic factor family ligands, is a receptor for the neurotrophic factor persephin." *J Biol Chem*, 275(50), 39427-39434.
- Mazzucchelli, C., Vantaggiato, C., Ciamei, A., Fasano, S., Pakhotin, P., Krezel, W., Welzl, H., Wolfer, D. P., Pages, G., Valverde, O., Marowsky, A., Porrazzo, A., Orban, P. C., Maldonado, R., Ehrenguber, M. U., Cestari, V., Lipp, H. P., Chapman, P. F., Pouyssegur, J., and Brambilla, R. (2002). "Knockout of ERK1 MAP kinase enhances synaptic plasticity in the striatum and facilitates striatal-mediated learning and memory." *Neuron*, 34(5), 807-820.
- Mcdonnell, T. J., Deane, N., Platt, F. M., Nunez, G., Jaeger, U., Mckearn, J. P., and Korsmeyer, S. J. (1989). "Bcl-2-Immunoglobulin Transgenic Mice Demonstrate Extended B-Cell Survival and Follicular Lymphoproliferation." *Cell*, 57(1), 79-88.
- Menalled, L. B., Sison, J. D., Wu, Y., Olivieri, M., Li, X. J., Li, H., Zeitlin, S., and Chesselet, M. F. (2002). "Early motor dysfunction and striosomal distribution of huntingtin microaggregates in Huntington's disease knock-in mice." *J. Neurosci.*, 22(18), 8266-8276.

- Metzler, M., Helgason, C. D., Dragatsis, I., Zhang, T., Gan, L., Pineault, N., Zeitlin, S. O., Humphries, R. K., and Hayden, M. R. (2000). "Huntingtin is required for normal hematopoiesis." *Hum. Mol. Genet.*, 9(3), 387-394.
- Michaelidis, T. M., Sendtner, M., Cooper, J. D., Airaksinen, M. S., Holtmann, B., Meyer, M., and Thoenen, H. (1996). "Inactivation of bcl-2 results in progressive degeneration of motoneurons, sympathetic and sensory neurons during early postnatal development." *Neuron*, 17(1), 75-89.
- Mikhailov, V., Mikhailova, M., Pulkrabek, D. J., Dong, Z., Venkatachalam, M. A., and Saikumar, P. (2001). "Bcl-2 prevents Bax oligomerization in the mitochondrial outer membrane." *Journal of Biological Chemistry*, 276(21), 18361-18374.
- Mikhailov, V., Mikhailova, M., Degenhardt, K., Venkatachalam, M. A., White, E., and Saikumar, P. (2003). "Association of Bax and Bak homo-oligomers in mitochondria. Bax requirement for Bak reorganization and cytochrome c release." *J. Biol. Chem.*, 278(7), 5367-5376.
- Milbrandt, J., de Sauvage, F. J., Fahrner, T. J., Baloh, R. H., Leitner, M. L., Tansey, M. G., Lampe, P. A., Heuckeroth, R. O., Kotzbauer, P. T., Simburger, K. S., Golden, J. P., Davies, J. A., Vejsada, R., Kato, A. C., Hynes, M., Sherman, D., Nishimura, M., Wang, L. C., Vandlen, R., Moffat, B., Klein, R. D., Poulsen, K., Gray, C., Garcés, A., and Johnson, E. M. J. (1998). "Persephin, a novel neurotrophic factor related to GDNF and neurturin." *Neuron*, 20(2), 245-253.
- Miller, F. D., and Kaplan, D. R. (2001). "Neurotrophin signalling pathways regulating neuronal apoptosis." *Cell Mol. Life Sci.*, 58(8), 1045-1053.
- Miller, T. M., Tansey, M. G., Johnson, E. M., Jr., and Creedon, D. J. (1997). "Inhibition of phosphatidylinositol 3-kinase activity blocks depolarization- and insulin-like growth factor I-mediated survival of cerebellar granule cells." *J. Biol. Chem.*, 272(15), 9847-9853.
- Miranda, R. C., Sohrabji, F., and Toran-Allerand, C. D. (1993). "Neuronal colocalization of mRNAs for neurotrophins and their receptors in the developing central nervous system suggests a potential for autocrine interactions." *Proc. Natl. Acad. Sci. U. S. A.*, 90(14), 6439-6443.
- Mitchell, I. J., Cooper, A. J., and Griffiths, M. R. (1999). "The selective vulnerability of striatopallidal neurons." *Prog. Neurobiol.*, 59(6), 691-719.
- Miura, M., Zhu, H., Rotello, R., Hartweg, E. A., and Yuan, J. (1993). "Induction of apoptosis in fibroblasts by IL-1 beta-converting enzyme, a mammalian homolog of the *C. elegans* cell death gene *ced-3*." *Cell*, 75(4), 653-660.
- Motoyama, N., Wang, F., Roth, K. A., Sawa, H., Nakayama, K., Nakayama, K., Negishi, I., Senju, S., Zhang, Q., Fujii, S., and . (1995). "Massive cell death of immature hematopoietic cells and neurons in Bcl-x-deficient mice." *Science*, 267(5203), 1506-1510.
- Mowla, S. J., Farhadi, H. F., Pareek, S., Atwal, J. K., Morris, S. J., Seidah, N. G., and Murphy, R. A. (2001). "Biosynthesis and post-translational processing of the precursor to brain-derived neurotrophic factor." *J. Biol. Chem.*, 276(16), 12660-12666.

- Murakami, H., Iwashita, T., Asai, N., Shimono, Y., Iwata, Y., Kawai, K., and Takahashi, M. (1999). "Enhanced phosphatidylinositol 3-kinase activity and high phosphorylation state of its downstream signalling molecules mediated by ret with the MEN 2B mutation." *Biochem. Biophys. Res. Commun.*, 262(1), 68-75.
- Murphy, L. O., Smith, S., Chen, R. H., Fingar, D. C., and Blenis, J. (2002). "Molecular interpretation of ERK signal duration by immediate early gene products." *Nat. Cell Biol.*, 4(8), 556-564.
- Murphy, L. O., MacKeigan, J. P., and Blenis, J. (2004). "A network of immediate early gene products propagates subtle differences in mitogen-activated protein kinase signal amplitude and duration." *Mol. Cell Biol.*, 24(1), 144-153.
- Murphy, L. O., and Blenis, J. (2006). "MAPK signal specificity: the right place at the right time." *Trends Biochem. Sci.*, 31(5), 268-275.
- Murray, B., Alessandrini, A., Cole, A. J., Yee, A. G., and Furshpan, E. J. (1998). "Inhibition of the p44/42 MAP kinase pathway protects hippocampal neurons in a cell-culture model of seizure activity." *Proc. Natl. Acad. Sci. U. S. A.*, 95(20), 11975-11980.
- Myers, R. H., Vonsattel, J. P., Stevens, T. J., Cupples, L. A., Richardson, E. P., Martin, J. B., and Bird, E. D. (1988). "Clinical and neuropathologic assessment of severity in Huntington's disease." *Neurology*, 38(3), 341-347.

---

## N

- Napankangas, U., Lindqvist, N., Lindholm, D., and Hallbook, F. (2003). "Rat retinal ganglion cells upregulate the pro-apoptotic BH3-only protein Bim after optic nerve transection." *Brain Res. Mol. Brain Res.*, 120(1), 30-37.
- Nasir, J., Floresco, S. B., O'Kusky, J. R., Diewert, V. M., Richman, J. M., Zeisler, J., Borowski, A., Marth, J. D., Phillips, A. G., and Hayden, M. R. (1995). "Targeted disruption of the Huntington's disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes." *Cell*, 81(5), 811-823.
- Nicholls, D. G., and Budd, S. L. (2000). "Mitochondria and neuronal survival." *Physiol Rev.*, 80(1), 315-360.
- Nishimune, H., Vasseur, S., Wiese, S., Birling, M. C., Holtmann, B., Sendtner, M., Iovanna, J. L., and Henderson, C. E. (2000). "Reg-2 is a motoneuron neurotrophic factor and a signalling intermediate in the CNTF survival pathway." *Nat. Cell Biol.*, 2(12), 906-914.
- Nixon, R. A., Wegiel, J., Kumar, A., Yu, W. H., Peterhoff, C., Cataldo, A., and Cuervo, A. M. (2005). "Extensive involvement of autophagy in Alzheimer disease: an immuno-electron microscopy study." *J. Neuropathol. Exp. Neurol.*, 64(2), 113-122.
- Nixon, R. A. (2006). "Autophagy in neurodegenerative disease: friend, foe or turncoat?" *Trends Neurosci.*, 29(9), 528-535.

Nonomura, T., Kubo, T., Oka, T., Shimoke, K., Yamada, M., Enokido, Y., and Hatanaka, H. (1996). "Signaling pathways and survival effects of BDNF and NT-3 on cultured cerebellar granule cells." *Brain Res. Dev. Brain Res.*, 97(1), 42-50.

Nykjaer, A., Lee, R., Teng, K. K., Jansen, P., Madsen, P., Nielsen, M. S., Jacobsen, C., Kliemannel, M., Schwarz, E., Willnow, T. E., Hempstead, B. L., and Petersen, C. M. (2004). "Sortilin is essential for proNGF-induced neuronal cell death." *Nature*, 427(6977), 843-848.

**O**

O'Connor, L., Strasser, A., O'Reilly, L. A., Hausmann, G., Adams, J. M., Cory, S., and Huang, D. C. S. (1998). "Bim: a novel member of the Bcl-2 family that promotes apoptosis." *Embo Journal*, 17(2), 384-395.

O'Neill, E., and Kolch, W. (2004). "Conferring specificity on the ubiquitous Raf/MEK signalling pathway." *Br. J. Cancer*, 90(2), 283-288.

Oda, E., Ohki, R., Murasawa, H., Nemoto, J., Shibue, T., Yamashita, T., Tokino, T., Taniguchi, T., and Tanaka, N. (2000). "Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis." *Science*, 288(5468), 1053-1058.

Okada, H., and Mak, T. W. (2004). "Pathways of apoptotic and non-apoptotic death in tumour cells." *Nat. Rev. Cancer*, 4(8), 592-603.

Okuno, S., Saito, A., Hayashi, T., and Chan, P. H. (2004). "The c-Jun N-terminal protein kinase signaling pathway mediates Bax activation and subsequent neuronal apoptosis through interaction with Bim after transient focal cerebral ischemia." *J. Neurosci.*, 24(36), 7879-7887.

Oltvai, Z. N., Milliman, C. L., and Korsmeyer, S. J. (1993). "Bcl-2 Heterodimerizes In-Vivo with A Conserved Homolog, Bax, That Accelerates Programmed Cell-Death." *Cell*, 74(4), 609-619.

Oppenheim, R. W., Houenou, L. J., Johnson, J. E., Lin, L. F., Li, L., Lo, A. C., Newsome, A. L., Preette, D. M., and Wang, S. (1995). "Developing motor neurons rescued from programmed and axotomy-induced cell death by GDNF." *Nature*, 373(6512), 344-346.

Oppenheim, R. W., Flavell, R. A., Vinsant, S., Preette, D., Kuan, C. Y., and Rakic, P. (2001). "Programmed cell death of developing mammalian neurons after genetic deletion of caspases." *J. Neurosci.*, 21(13), 4752-4760.

Ordway, J. M., Tallaksen-Greene, S., Gutekunst, C. A., Bernstein, E. M., Cearley, J. A., Wiener, H. W., Dure, L. S., Lindsey, R., Hersch, S. M., Jope, R. S., Albin, R. L., and Detloff, P. J. (1997). "Ectopically expressed CAG repeats cause intranuclear inclusions and a progressive late onset neurological phenotype in the mouse." *Cell*, 91(6), 753-763.

- Palfi, S., Ferrante, R. J., Brouillet, E., Beal, M. F., Dolan, R., Guyot, M. C., Peschanski, M., and Hantraye, P. (1996). "Chronic 3-nitropropionic acid treatment in baboons replicates the cognitive and motor deficits of Huntington's disease." *J. Neurosci.*, 16(9), 3019-3025.
- Panov, A. V., Gutekunst, C. A., Leavitt, B. R., Hayden, M. R., Burke, J. R., Strittmatter, W. J., and Greenamyre, J. T. (2002). "Early mitochondrial calcium defects in Huntington's disease are a direct effect of polyglutamines." *Nat. Neurosci.*, 5(8), 731-736.
- Panov, A. V., Burke, J. R., Strittmatter, W. J., and Greenamyre, J. T. (2003). "In vitro effects of polyglutamine tracts on Ca<sup>2+</sup>-dependent depolarization of rat and human mitochondria: relevance to Huntington's disease." *Arch. Biochem. Biophys.*, 410(1), 1-6.
- Pap, M., and Cooper, G. M. (1998). "Role of glycogen synthase kinase-3 in the phosphatidylinositol 3-Kinase/Akt cell survival pathway." *J. Biol. Chem.*, 273(32), 19929-19932.
- Paradis, E., Douillard, H., Koutroumanis, M., Goodyer, C., and LeBlanc, A. (1996). "Amyloid beta peptide of Alzheimer's disease downregulates Bcl-2 and upregulates bax expression in human neurons." *J. Neurosci.*, 16(23), 7533-7539.
- Paradis, S., and Ruvkun, G. (1998). "Caenorhabditis elegans Akt/PKB transduces insulin receptor-like signals from AGE-1 PI3 kinase to the DAF-16 transcription factor." *Genes Dev.*, 12(16), 2488-2498.
- Paratcha, G., and Ibanez, C. F. (2002). "Lipid rafts and the control of neurotrophic factor signaling in the nervous system: variations on a theme." *Curr. Opin. Neurobiol.*, 12(5), 542-549.
- Paratcha, G., Ledda, F., and Ibanez, C. F. (2003). "The neural cell adhesion molecule NCAM is an alternative signaling receptor for GDNF family ligands." *Cell*, 113(7), 867-879.
- Parrizas, M., Saltiel, A. R., and LeRoith, D. (1997). "Insulin-like growth factor 1 inhibits apoptosis using the phosphatidylinositol 3'-kinase and mitogen-activated protein kinase pathways." *J. Biol. Chem.*, 272(1), 154-161.
- Perez-Navarro, E., Alberch, J., Arenas, E., Calvo, N., and Marsal, J. (1994). "Nerve growth factor and basic fibroblast growth factor protect cholinergic neurons against quinolinic acid excitotoxicity in rat neostriatum." *Eur. J. Neurosci.*, 6(5), 706-711.
- Perez-Navarro, E., Arenas, E., Reiriz, J., Calvo, N., and Alberch, J. (1996). "Glial cell line-derived neurotrophic factor protects striatal calbindin-immunoreactive neurons from excitotoxic damage." *Neuroscience*, 75(2), 345-352.
- Perez-Navarro, E., Arenas, E., Marco, S., and Alberch, J. (1999a). "Intrastriatal grafting of a GDNF-producing cell line protects striatonigral neurons from quinolinic acid excitotoxicity in vivo." *Eur. J. Neurosci.*, 11(1), 241-249.



- Perez-Navarro, E., Alberch, J., Neveu, I., and Arenas, E. (1999b). "Brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4/5 differentially regulate the phenotype and prevent degenerative changes in striatal projection neurons after excitotoxicity in vivo." *Neuroscience*, 91(4), 1257-1264.
- Perez-Navarro, E., Canudas, A. M., Akerund, P., Alberch, J., and Arenas, E. (2000a). "Brain-derived neurotrophic factor, neurotrophin-3, and neurotrophin-4/5 prevent the death of striatal projection neurons in a rodent model of Huntington's disease." *J. Neurochem.*, 75(5), 2190-2199.
- Perez-Navarro, E., Akerud, P., Marco, S., Canals, J. M., Tolosa, E., Arenas, E., and Alberch, J. (2000b). "Neurturin protects striatal projection neurons but not interneurons in a rat model of Huntington's disease." *Neuroscience*, 98(1), 89-96.
- Perez-Navarro, E., Gavalda, N., Gratacos, E., and Alberch, J. (2005). "Brain-derived neurotrophic factor prevents changes in Bcl-2 family members and caspase-3 activation induced by excitotoxicity in the striatum." *J. Neurochem.*, 92(3), 678-691.
- Perez-Navarro, E., Canals, J. M., Gines, S., and Alberch, J. (2006). "Cellular and molecular mechanisms involved in the selective vulnerability of striatal projection neurons in Huntington's disease." *Histol. Histopathol.*, 21(11), 1217-1232.
- Perkinton, M. S., Sihra, T. S., and Williams, R. J. (1999). "Ca(2+)-permeable AMPA receptors induce phosphorylation of cAMP response element-binding protein through a phosphatidylinositol 3-kinase-dependent stimulation of the mitogen-activated protein kinase signaling cascade in neurons." *J. Neurosci.*, 19(14), 5861-5874.
- Perkinton, M. S., Ip, J. K., Wood, G. L., Crossthwaite, A. J., and Williams, R. J. (2002). "Phosphatidylinositol 3-kinase is a central mediator of NMDA receptor signalling to MAP kinase (Erk1/2), Akt/PKB and CREB in striatal neurones." *J. Neurochem.*, 80(2), 239-254.
- Perron, J. C., and Bixby, J. L. (1999). "Distinct neurite outgrowth signaling pathways converge on ERK activation." *Mol. Cell Neurosci.*, 13(5), 362-378.
- Perry, G., Roder, H., Nunomura, A., Takeda, A., Friedlich, A. L., Zhu, X., Raina, A. K., Holbrook, N., Siedlak, S. L., Harris, P. L., and Smith, M. A. (1999). "Activation of neuronal extracellular receptor kinase (ERK) in Alzheimer disease links oxidative stress to abnormal phosphorylation." *Neuroreport*, 10(11), 2411-2415.
- Persad, S., Attwell, S., Gray, V., Mawji, N., Deng, J. T., Leung, D., Yan, J., Sanghera, J., Walsh, M. P., and Dedhar, S. (2001). "Regulation of protein kinase B/Akt-serine 473 phosphorylation by integrin-linked kinase: critical roles for kinase activity and amino acids arginine 211 and serine 343." *J. Biol. Chem.*, 276(29), 27462-27469.
- Poteryaev, D., Titievsky, A., Sun, Y. F., Thomas-Crusells, J., Lindahl, M., Billaud, M., Arumae, U., and Saarma, M. (1999). "GDNF triggers a novel ret-independent Src kinase family-coupled signaling via a GPI-linked GDNF receptor alpha1." *FEBS Lett.*, 463(1-2), 63-66.

- Proskuryakov, S. Y., Gabai, V. L., and Konoplyannikov, A. G. (2002). "Necrosis is an active and controlled form of programmed cell death." *Biochemistry (Mosc.)*, 67(4), 387-408.
- Proskuryakov, S. Y., Konoplyannikov, A. G., and Gabai, V. L. (2003). "Necrosis: a specific form of programmed cell death?" *Exp. Cell Res.*, 283(1), 1-16.
- Purves, D., Snider, W. D., and Voyvodic, J. T. (1988). "Trophic regulation of nerve cell morphology and innervation in the autonomic nervous system." *Nature*, 336(6195), 123-128.
- Putcha, G. V., Moulder, K. L., Golden, J. P., Bouillet, P., Adams, J. A., Strasser, A., and Johnson, E. M. (2001). "Induction of BIM, a proapoptotic BH3-only BCL-2 family member, is critical for neuronal apoptosis." *Neuron*, 29(3), 615-628.
- Putcha, G. V., Le, S. Y., Frank, S., Besirli, C. G., Clark, K., Chu, B. Y., Alix, S., Youle, R. J., LaMarche, A., Maroney, A. C., and Johnson, E. M. (2003). "JNK-mediated BIM phosphorylation potentiates BAX-dependent apoptosis." *Neuron*, 38(6), 899-914.
- Puthalakath, H., Huang, D. C., O'Reilly, L. A., King, S. M., and Strasser, A. (1999). "The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex." *Mol. Cell*, 3(3), 287-296.
- Puthalakath, H., Villunger, A., O'Reilly, L. A., Beaumont, J. G., Coultas, L., Cheney, R. E., Huang, D. C., and Strasser, A. (2001). "Bmf: a proapoptotic BH3-only protein regulated by interaction with the myosin V actin motor complex, activated by anoikis." *Science*, 293(5536), 1829-1832.

---

**Q**

- Qi, X. J., Wildey, G. M., and Howe, P. H. (2006). "Evidence that Ser87 of BimEL is phosphorylated by Akt and regulates BimEL apoptotic function." *J. Biol. Chem.*, 281(2), 813-823.

---

**R**

- Rane, M. J., Pan, Y., Singh, S., Powell, D. W., Wu, R., Cummins, T., Chen, Q., McLeish, K. R., and Klein, J. B. (2003). "Heat shock protein 27 controls apoptosis by regulating Akt activation." *J. Biol. Chem.*, 278(30), 27828-27835.
- Rangone, H., Poizat, G., Troncoso, J., Ross, C. A., MacDonald, M. E., Saudou, F., and Humbert, S. (2004). "The serum- and glucocorticoid-induced kinase SGK inhibits mutant huntingtin-induced toxicity by phosphorylating serine 421 of huntingtin." *Eur. J. Neurosci.*, 19(2), 273-279.
- Rangone, H., Pardo, R., Colin, E., Girault, J. A., Saudou, F., and Humbert, S. (2005). "Phosphorylation of arfaptin 2 at Ser260 by Akt Inhibits PolyQ-huntingtin-induced toxicity by rescuing proteasome impairment." *J. Biol. Chem.*, 280(23), 22021-22028.

- Rapoport, M., and Ferreira, A. (2000). "PD98059 prevents neurite degeneration induced by fibrillar beta-amyloid in mature hippocampal neurons." *J. Neurochem.*, 74(1), 125-133.
- Ravikumar, B., Vacher, C., Berger, Z., Davies, J. E., Luo, S., Oroz, L. G., Scaravilli, F., Easton, D. F., Duden, R., O'Kane, C. J., and Rubinsztein, D. C. (2004). "Inhibition of mTOR induces autophagy and reduces toxicity of polyglutamine expansions in fly and mouse models of Huntington disease." *Nat. Genet.*, 36(6), 585-595.
- Ravikumar, B., cevedo-Arozena, A., Imarisio, S., Berger, Z., Vacher, C., O'Kane, C. J., Brown, S. D., and Rubinsztein, D. C. (2005). "Dynein mutations impair autophagic clearance of aggregate-prone proteins." *Nat. Genet.*, 37(7), 771-776.
- Reed, J. C. (2006). "Proapoptotic multidomain Bcl-2/Bax-family proteins: mechanisms, physiological roles, and therapeutic opportunities." *Cell Death. Differ.*, 13(8), 1378-1386.
- Reginato, M. J., Mills, K. R., Paulus, J. K., Lynch, D. K., Sgroi, D. C., Debnath, J., Muthuswamy, S. K., and Brugge, J. S. (2003). "Integrins and EGFR coordinately regulate the pro-apoptotic protein Bim to prevent anoikis." *Nature Cell Biology*, 5(8), 733-740.
- Rego, A. C., and Oliveira, C. R. (2003). "Mitochondrial dysfunction and reactive oxygen species in excitotoxicity and apoptosis: implications for the pathogenesis of neurodegenerative diseases." *Neurochem. Res.*, 28(10), 1563-1574.
- Reiner, A., Albin, R. L., Anderson, K. D., D'Amato, C. J., Penney, J. B., and Young, A. B. (1988). "Differential loss of striatal projection neurons in Huntington disease." *Proc. Natl. Acad. Sci. U. S. A.*, 85(15), 5733-5737.
- Richfield, E. K., Maguire-Zeiss, K. A., Vonkeman, H. E., and Voorn, P. (1995). "Preferential loss of preproenkephalin versus preprotachykinin neurons from the striatum of Huntington's disease patients." *Ann. Neurol.*, 38(6), 852-861.
- Rigamonti, D., Bauer, J. H., De-Fraja, C., Conti, L., Sipione, S., Sciorati, C., Clementi, E., Hackam, A., Hayden, M. R., Li, Y., Cooper, J. K., Ross, C. A., Govoni, S., Vincenz, C., and Cattaneo, E. (2000). "Wild-type huntingtin protects from apoptosis upstream of caspase-3." *J. Neurosci.*, 20(10), 3705-3713.
- Rigamonti, D., Sipione, S., Goffredo, D., Zuccato, C., Fossale, E., and Cattaneo, E. (2001). "Huntingtin's neuroprotective activity occurs via inhibition of procaspase-9 processing." *J. Biol. Chem.*, 276(18), 14545-14548.
- Righi, M., Tongiorgi, E., and Cattaneo, A. (2000). "Brain-derived neurotrophic factor (BDNF) induces dendritic targeting of BDNF and tyrosine kinase B mRNAs in hippocampal neurons through a phosphatidylinositol-3 kinase-dependent pathway." *J. Neurosci.*, 20(9), 3165-3174.
- Ross, C. A. (2002). "Polyglutamine pathogenesis: emergence of unifying mechanisms for Huntington's disease and related disorders." *Neuron*, 35(5), 819-822.

- Rosler, O. G., Giehl, K. M., and Thiel, G. (2004). "Neuroprotection of immortalized hippocampal neurones by brain-derived neurotrophic factor and Raf-1 protein kinase: role of extracellular signal-regulated protein kinase and phosphatidylinositol 3-kinase." *J. Neurochem.*, 88(5), 1240-1252.
- Roux, P. P., and Blenis, J. (2004). "ERK and p38 MAPK-activated protein kinases: a family of protein kinases with diverse biological functions." *Microbiol. Mol. Biol. Rev.*, 68(2), 320-344.
- Ruvolo, P. P., Deng, X., Carr, B. K., and May, W. S. (1998). "A functional role for mitochondrial protein kinase Calpha in Bcl2 phosphorylation and suppression of apoptosis." *J. Biol. Chem.*, 273(39), 25436-25442.
- 
- S
- Sable, C. L., Filippa, N., Hemmings, B., and Van, O. E. (1997). "cAMP stimulates protein kinase B in a Wortmannin-insensitive manner." *FEBS Lett.*, 409(2), 253-257.
- Saelens, X., Festjens, N., Vande, W. L., van, G. M., van, L. G., and Vandenabeele, P. (2004). "Toxic proteins released from mitochondria in cell death." *Oncogene*, 23(16), 2861-2874.
- Sakurai, M., Hayashi, T., Abe, K., Itoyama, Y., and Tabayashi, K. (2001). "Induction of phosphatidylinositol 3-kinase and serine-threonine kinase-like immunoreactivity in rabbit spinal cord after transient ischemia." *Neurosci. Lett.*, 302(1), 17-20.
- Salvesen, G. S., and Dixit, V. M. (1999). "Caspase activation: the induced-proximity model." *Proc. Natl. Acad. Sci. U. S. A.*, 96(20), 10964-10967.
- Sarbassov, D. D., Guertin, D. A., Ali, S. M., and Sabatini, D. M. (2005). "Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex." *Science*, 307(5712), 1098-1101.
- Sarig, R., Zaltsman, Y., Marcellus, R. C., Flavell, R., Mak, T. W., and Gross, A. (2003). "BID-D59A is a potent inducer of apoptosis in primary embryonic fibroblasts." *J. Biol. Chem.*, 278(12), 10707-10715.
- Sattler, M., Liang, H., Nettlesheim, D., Meadows, R. P., Harlan, J. E., Eberstadt, M., Yoon, H. S., Shuker, S. B., Chang, B. S., Minn, A. J., Thompson, C. B., and Fesik, S. W. (1997). "Structure of Bcl-xL-Bak peptide complex: recognition between regulators of apoptosis." *Science*, 275(5302), 983-986.
- Saudou, F., Finkbeiner, S., Devys, D., and Greenberg, M. E. (1998). "Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear inclusions." *Cell*, 95(1), 55-66.
- Sawada, M., Sun, W., Hayes, P., Leskov, K., Boothman, D. A., and Matsuyama, S. (2003). "Ku70 suppresses the apoptotic translocation of Bax to mitochondria." *Nat. Cell Biol.*, 5(4), 320-329.
- Schaeffer, H. J., and Weber, M. J. (1999). "Mitogen-activated protein kinases: specific messages from ubiquitous messengers." *Mol. Cell Biol.*, 19(4), 2435-2444.

- Scheid, M. P., Schubert, K. M., and Duronio, V. (1999). "Regulation of bad phosphorylation and association with Bcl-x(L) by the MAPK/Erk kinase." *J. Biol. Chem.*, 274(43), 31108-31113.
- Schoepfer, R., Monyer, H., Sommer, B., Wisden, W., Sprengel, R., Kuner, T., Lomeli, H., Herb, A., Kohler, M., Burnashev, N., and . (1994). "Molecular biology of glutamate receptors." *Prog. Neurobiol.*, 42(2), 353-357.
- Schumacher, J. M., Short, M. P., Hyman, B. T., Breakefield, X. O., and Isacson, O. (1991). "Intracerebral implantation of nerve growth factor-producing fibroblasts protects striatum against neurotoxic levels of excitatory amino acids." *Neuroscience*, 45(3), 561-570.
- Schwarcz, R., and Kohler, C. (1983). "Differential vulnerability of central neurons of the rat to quinolinic acid." *Neurosci. Lett.*, 38(1), 85-90.
- Schwarzschild, M. A., Cole, R. L., and Hyman, S. E. (1997). "Glutamate, but not dopamine, stimulates stress-activated protein kinase and AP-1-mediated transcription in striatal neurons." *J. Neurosci.*, 17(10), 3455-3466.
- Schweichel, J. U., and Merker, H. J. (1973). "The morphology of various types of cell death in prenatal tissues." *Teratology*, 7(3), 253-266.
- Scott, R. P., and Ibanez, C. F. (2001). "Determinants of ligand binding specificity in the glial cell line-derived neurotrophic factor family receptor alpha s." *Journal of Biological Chemistry*, 276(2), 1450-1458.
- Sgambato, V., Pages, C., Rogard, M., Besson, M. J., and Caboche, J. (1998). "Extracellular signal-regulated kinase (ERK) controls immediate early gene induction on corticostriatal stimulation." *J. Neurosci.*, 18(21), 8814-8825.
- Shimizu, S., Kanaseki, T., Mizushima, N., Mizuta, T., rakawa-Kobayashi, S., Thompson, C. B., and Tsujimoto, Y. (2004). "Role of Bcl-2 family proteins in a non-apoptotic programmed cell death dependent on autophagy genes." *Nat. Cell Biol.*, 6(12), 1221-1228.
- Shinjo, T., Kuribara, R., Inukai, T., Hosoi, H., Kinoshita, T., Miyajima, A., Houghton, P. J., Look, A. T., Ozawa, K., and Inaba, T. (2001). "Downregulation of Bim, a proapoptotic relative of Bcl-2, is a pivotal step in cytokine-initiated survival signaling in murine hematopoietic progenitors." *Molecular and Cellular Biology*, 21(3), 854-864.
- Shinoda, S., Schindler, C. K., Meller, R., So, N. K., Araki, T., Yamamoto, A., Lan, J. Q., Taki, W., Simon, R. P., and Henshall, D. C. (2004). "Bim regulation may determine hippocampal vulnerability after injurious seizures and in temporal lobe epilepsy." *J. Clin. Invest*, 113(7), 1059-1068.
- Simons, K., and Toomre, D. (2000). "Lipid rafts and signal transduction." *Nat. Rev. Mol. Cell Biol.*, 1(1), 31-39.
- Simpson, L., and Parsons, R. (2001). "PTEN: life as a tumor suppressor." *Exp. Cell Res.*, 264(1), 29-41.
- Sofroniew, M. V., Galletly, N. P., Isacson, O., and Svendsen, C. N. (1990). "Survival of adult basal forebrain cholinergic neurons after loss of target neurons." *Science*, 247(4940), 338-342.

- Sole, C., Dolcet, X., Segura, M. F., Gutierrez, H., az-Meco, M. T., Gozzelino, R., Sanchis, D., Bayascas, J. R., Gallego, C., Moscat, J., Davies, A. M., and Comella, J. X. (2004). "The death receptor antagonist FAIM promotes neurite outgrowth by a mechanism that depends on ERK and NF-kapp B signaling." *J. Cell Biol.*, 167(3), 479-492.
- Soler, R. M., Dolcet, X., Encinas, M., Egea, J., Bayascas, J. R., and Comella, J. X. (1999). "Receptors of the glial cell line-derived neurotrophic factor family of neurotrophic factors signal cell survival through the phosphatidylinositol 3-kinase pathway in spinal cord motoneurons." *J. Neurosci.*, 19(21), 9160-9169.
- Solit, D. B., Basso, A. D., Olshen, A. B., Scher, H. I., and Rosen, N. (2003). "Inhibition of heat shock protein 90 function down-regulates Akt kinase and sensitizes tumors to Taxol." *Cancer Res.*, 63(9), 2139-2144.
- Song, G., Ouyang, G., and Bao, S. (2005). "The activation of Akt/PKB signaling pathway and cell survival." *J. Cell Mol. Med.*, 9(1), 59-71.
- Spreafico, R., Frassoni, C., Arcelli, P., Selvaggio, M., and De, B. S. (1995). "In situ labeling of apoptotic cell death in the cerebral cortex and thalamus of rats during development." *J. Comp Neurol.*, 363(2), 281-295.
- Springer, J. E., Azbill, R. D., Nottingham, S. A., and Kennedy, S. E. (2000). "Calcineurin-mediated BAD dephosphorylation activates the caspase-3 apoptotic cascade in traumatic spinal cord injury." *J. Neurosci.*, 20(19), 7246-7251.
- Srivastava, R. K., Srivastava, A. R., Korsmeyer, S. J., Nesterova, M., Cho-Chung, Y. S., and Longo, D. L. (1998). "Involvement of microtubules in the regulation of Bcl2 phosphorylation and apoptosis through cyclic AMP-dependent protein kinase." *Mol. Cell Biol.*, 18(6), 3509-3517.
- Stanciu, M., Wang, Y., Kentor, R., Burke, N., Watkins, S., Kress, G., Reynolds, I., Klann, E., Angiolieri, M. R., Johnson, J. W., and DeFranco, D. B. (2000). "Persistent activation of ERK contributes to glutamate-induced oxidative toxicity in a neuronal cell line and primary cortical neuron cultures." *J. Biol. Chem.*, 275(16), 12200-12206.
- Standaert, D. G., Testa, C. M., Young, A. B., and Penney, J. B., Jr. (1994). "Organization of N-methyl-D-aspartate glutamate receptor gene expression in the basal ganglia of the rat." *J. Comp Neurol.*, 343(1), 1-16.
- Stehlik, C., de, M. R., Binder, B. R., and Lipp, J. (1998). "Cytokine induced expression of porcine inhibitor of apoptosis protein (iap) family member is regulated by NF-kappa B." *Biochem. Biophys. Res. Commun.*, 243(3), 827-832.
- Stennicke, H. R., and Salvesen, G. S. (2000). "Caspases - controlling intracellular signals by protease zymogen activation." *Biochim. Biophys. Acta*, 1477(1-2), 299-306.
- Stokoe, D., Stephens, L. R., Copeland, T., Gaffney, P. R., Reese, C. B., Painter, G. F., Holmes, A. B., McCormick, F., and Hawkins, P. T. (1997). "Dual role of phosphatidylinositol-3,4,5-trisphosphate in the activation of protein kinase B." *Science*, 277(5325), 567-570.

Strong, T. V., Tagle, D. A., Valdes, J. M., Elmer, L. W., Boehm, K., Swaroop, M., Kaatz, K. W., Collins, F. S., and Albin, R. L. (1993). "Widespread expression of the human and rat Huntington's disease gene in brain and nonneural tissues." *Nat. Genet.*, 5(3), 259-265.

Stroppolo, A., Guinea, B., Tian, C., Sommer, J., and Ehrlich, M. E. (2001). "Role of phosphatidylinositide 3-kinase in brain-derived neurotrophic factor-induced DARPP-32 expression in medium size spiny neurons in vitro." *J. Neurochem.*, 79(5), 1027-1032.

Subramaniam, S., Zirrgiebel, U., von Bohlen Und, H. O., Strelau, J., Laliberte, C., Kaplan, D. R., and Unsicker, K. (2004). "ERK activation promotes neuronal degeneration predominantly through plasma membrane damage and independently of caspase-3." *J. Cell Biol.*, 165(3), 357-369.

---

## T

Tabrizi, S. J., Cleeter, M. W., Xuereb, J., Taanman, J. W., Cooper, J. M., and Schapira, A. H. (1999). "Biochemical abnormalities and excitotoxicity in Huntington's disease brain." *Ann. Neurol.*, 45(1), 25-32.

Tafari, M., Karpnich, N. O., Hurster, K. A., Pastorino, J. G., Schneider, T., Russo, M. A., and Farber, J. L. (2002). "Cytochrome c release upon Fas receptor activation depends on translocation of full-length bid and the induction of the mitochondrial permeability transition." *J. Biol. Chem.*, 277(12), 10073-10082.

Tamatani, M., Che, Y. H., Matsuzaki, H., Ogawa, S., Okado, H., Miyake, S., Mizuno, T., and Tohyama, M. (1999). "Tumor necrosis factor induces Bcl-2 and Bcl-x expression through NFkappaB activation in primary hippocampal neurons." *J. Biol. Chem.*, 274(13), 8531-8538.

Thandi, S., Blank, J. L., and Challiss, R. A. (2002). "Group-I metabotropic glutamate receptors, mGlu1a and mGlu5a, couple to extracellular signal-regulated kinase (ERK) activation via distinct, but overlapping, signalling pathways." *J. Neurochem.*, 83(5), 1139-1153.

The Huntington's Disease Collaborative Research Group. (1993). "A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes." *Cell*, 72(6), 971-983.

Thoenen, H., Barde, Y. A., Davies, A. M., and Johnson, J. E. (1987). "Neurotrophic factors and neuronal death." *Ciba Found. Symp.*, 126, 82-95.

Thoenen, H., and Sendtner, M. (2002). "Neurotrophins: from enthusiastic expectations through sobering experiences to rational therapeutic approaches." *Nat. Neurosci.*, 5 Suppl, 1046-1050.

Thompson, J., Dolcet, X., Hilton, M., Tolcos, M., and Davies, A. M. (2004). "HGF promotes survival and growth of maturing sympathetic neurons by PI-3 kinase- and MAP kinase-dependent mechanisms." *Mol. Cell Neurosci.*, 27(4), 441-452.

Toker, A., and Newton, A. C. (2000). "Akt/protein kinase B is regulated by autophosphorylation at the hypothetical PDK-2 site." *J. Biol. Chem.*, 275(12), 8271-8274.

- Tonks, N. K., and Neel, B. G. (2001). "Combinatorial control of the specificity of protein tyrosine phosphatases." *Curr. Opin. Cell Biol.*, 13(2), 182-195.
- Treanor, J. J., Goodman, L., de Sauvage, F., Stone, D. M., Poulsen, K. T., Beck, C. D., Gray, C., Armanini, M. P., Pollock, R. A., Hefti, F., Phillips, H. S., Goddard, A., Moore, M. W., Buj-Bello, A., Davies, A. M., Asai, N., Takahashi, M., Vandlen, R., Henderson, C. E., and Rosenthal, A. (1996). "Characterization of a multicomponent receptor for GDNF." *Nature*, 382(6586), 80-83.
- Trupp, M., Arenas, E., Fainzilber, M., Nilsson, A. S., Sieber, B. A., Grigoriou, M., Kilkenny, C., Salazar-Gruoso, E., Pachnis, V., and Arumae, U. (1996). "Functional receptor for GDNF encoded by the c-ret proto-oncogene." *Nature*, 381(6585), 785-789.
- Trupp, M., Scott, R., Whittemore, S. R., and Ibanez, C. F. (1999). "Ret-dependent and -independent mechanisms of glial cell line-derived neurotrophic factor signaling in neuronal cells." *J. Biol. Chem.*, 274(30), 20885-20894.
- Tsui-Pierchala, B., Milbrandt, J., and Johnson, E. M. (2002). "NGF utilizes c-Ret via a novel GFL-independent, inter-RTK signaling mechanism to maintain the trophic status of mature sympathetic neurons." *Neuron*, 33(2), 261-273.
- Tu, V. C., Bahl, J. J., and Chen, Q. M. (2002). "Signals of oxidant-induced cardiomyocyte hypertrophy: key activation of p70 S6 kinase-1 and phosphoinositide 3-kinase." *J. Pharmacol. Exp. Ther.*, 300(3), 1101-1110.
- 
- Vaillant, A. R., Mazzoni, I., Tudan, C., Boudreau, M., Kaplan, D. R., and Miller, F. D. (1999). "Depolarization and neurotrophins converge on the phosphatidylinositol 3-kinase-Akt pathway to synergistically regulate neuronal survival." *J. Cell Biol.*, 146(5), 955-966.
- Valentijn, A. J., and Gilmore, A. P. (2004). "Translocation of full-length Bid to mitochondria during anoikis." *J. Biol. Chem.*, 279(31), 32848-32857.
- Vanhoutte, P., Barnier, J. V., Guibert, B., Pages, C., Besson, M. J., Hipskind, R. A., and Caboche, J. (1999). "Glutamate induces phosphorylation of Elk-1 and CREB, along with c-fos activation, via an extracellular signal-regulated kinase-dependent pathway in brain slices." *Mol. Cell Biol.*, 19(1), 136-146.
- Varfolomeev, E. E., Schuchmann, M., Luria, V., Chiannikulchai, N., Beckmann, J. S., Mett, I. L., Rebrikov, D., Brodianski, V. M., Kemper, O. C., Kollet, O., Lapidot, T., Soffer, D., Sobe, T., Avraham, K. B., Goncharov, T., Holtmann, H., Lonai, P., and Wallach, D. (1998). "Targeted disruption of the mouse Caspase 8 gene ablates cell death induction by the TNF receptors, Fas/Apo1, and DR3 and is lethal prenatally." *Immunity.*, 9(2), 267-276.
- Vaux, D. L., Cory, S., and Adams, J. M. (1988). "Bcl-2 Gene Promotes Hematopoietic-Cell Survival and Cooperates with C-Myc to immortalize Pre-B-Cells." *Nature*, 335(6189), 440-442.
- Vaux, D. L., and Silke, J. (2003). "Mammalian mitochondrial IAP binding proteins." *Biochem. Biophys. Res. Commun.*, 304(3), 499-504.



- Venero, J. L., Beck, K. D., and Hefti, F. (1994). "Intrastriatal infusion of nerve growth factor after quinolinic acid prevents reduction of cellular expression of choline acetyltransferase messenger RNA and trkA messenger RNA, but not glutamate decarboxylase messenger RNA." *Neuroscience*, 61(2), 257-268.
- Ventimiglia, R., Mather, P. E., Jones, B. E., and Lindsay, R. M. (1995). "The neurotrophins BDNF, NT-3 and NT-4/5 promote survival and morphological and biochemical differentiation of striatal neurons in vitro." *Eur. J. Neurosci.*, 7(2), 213-222.
- Verma, S., Zhao, L. J., and Chinnadurai, G. (2001). "Phosphorylation of the proapoptotic protein BIK: mapping of phosphorylation sites and effect on apoptosis." *J. Biol. Chem.*, 276(7), 4671-4676.
- Vis, J. C., Schipper, E., de Boer-van Huizen RT, Verbeek, M. M., de Waal, R. M., Wesseling, P., ten Donkelaar, H. J., and Kremer, B. (2005). "Expression pattern of apoptosis-related markers in Huntington's disease." *Acta Neuropathol. (Berl)*, 109(3), 321-328.
- von Bartheld, C. S., Byers, M. R., Williams, R., and Bothwell, M. (1996). "Anterograde transport of neurotrophins and axodendritic transfer in the developing visual system." *Nature*, 379(6568), 830-833.
- Vonsattel, J. P., Myers, R. H., Stevens, T. J., Ferrante, R. J., Bird, E. D., and Richardson, E. P., Jr. (1985). "Neuropathological classification of Huntington's disease." *J. Neuropathol. Exp. Neurol.*, 44(6), 559-577.

---

## W

- Wagey, R., Pelech, S. L., Duronio, V., and Krieger, C. (1998). "Phosphatidylinositol 3-kinase: increased activity and protein level in amyotrophic lateral sclerosis." *J. Neurochem.*, 71(2), 716-722.
- Wajant, H., and Scheurich, P. (2001). "Tumor necrosis factor receptor-associated factor (TRAF) 2 and its role in TNF signaling." *Int. J. Biochem. Cell Biol.*, 33(1), 19-32.
- Walczak, H., and Krammer, P. H. (2000). "The CD95 (APO-1/Fas) and the TRAIL (APO-2L) apoptosis systems." *Exp. Cell Res.*, 256(1), 58-66.
- Wallach, D., Varfolomeev, E. E., Malinin, N. L., Goltsev, Y. V., Kovalenko, A. V., and Boldin, M. P. (1999). "Tumor necrosis factor receptor and Fas signaling mechanisms." *Annu. Rev. Immunol.*, 17, 331-367.
- Wang, J. M., Chao, J. R., Chen, W., Kuo, M. L., Yen, J. J., and Yang-Yen, H. F. (1999). "The antiapoptotic gene mcl-1 is up-regulated by the phosphatidylinositol 3-kinase/Akt signaling pathway through a transcription factor complex containing CREB." *Mol. Cell Biol.*, 19(9), 6195-6206.
- Wang, J. Q., Tang, Q., Parelkar, N. K., Liu, Z., Samdani, S., Choe, E. S., Yang, L., and Mao, L. (2004). "Glutamate signaling to Ras-MAPK in striatal neurons: mechanisms for inducible gene expression and plasticity." *Mol. Neurobiol.*, 29(1), 1-14.

- Wang, K., Yin, X. M., Chao, D. T., Milliman, C. L., and Korsmeyer, S. J. (1996). "BID: a novel BH3 domain-only death agonist." *Genes Dev.*, 10(22), 2859-2869.
- Ward, M. W., Rego, A. C., Frenguelli, B. G., and Nicholls, D. G. (2000). "Mitochondrial membrane potential and glutamate excitotoxicity in cultured cerebellar granule cells." *J. Neurosci.*, 20(19), 7208-7219.
- Ward, M. W., Kogel, D., and Prehn, J. H. (2004). "Neuronal apoptosis: BH3-only proteins the real killers?" *J. Bioenerg. Biomembr.*, 36(4), 295-298.
- Ward, M. W., Rehm, M., Duesmann, H., Kacmar, S., Concannon, C. G., and Prehn, J. H. (2006). "Real time single cell analysis of Bid cleavage and Bid translocation during caspase-dependent and neuronal caspase-independent apoptosis." *J. Biol. Chem.*, 281(9), 5837-5844.
- Wei, M. C., Lindsten, T., Mootha, V. K., Weiler, S., Gross, A., Ashiya, M., Thompson, C. B., and Korsmeyer, S. J. (2000). "tBID, a membrane-targeted death ligand, oligomerizes BAK to release cytochrome c." *Genes Dev.*, 14(16), 2060-2071.
- Wei, M. C., Zong, W. X., Cheng, E. H. Y., Lindsten, T., Panoutsakopoulou, V., Ross, A. J., Roth, K. A., MacGregor, G. R., Thompson, C. B., and Korsmeyer, S. J. (2001). "Proapoptotic BAX and BAK: A requisite gateway to mitochondrial dysfunction and death." *Science*, 292(5517), 727-730.
- Wellbrock, C., Karasarides, M., and Marais, R. (2004). "The RAF proteins take centre stage." *Nat. Rev. Mol. Cell Biol.*, 5(11), 875-885.
- Wennstrom, S., and Downward, J. (1999). "Role of phosphoinositide 3-kinase in activation of ras and mitogen-activated protein kinase by epidermal growth factor." *Mol. Cell Biol.*, 19(6), 4279-4288.
- Weston, C. R., and Davis, R. J. (2002). "The JNK signal transduction pathway." *Curr. Opin. Genet. Dev.*, 12(1), 14-21.
- Weston, C. R., Balmanno, K., Chalmers, C., Hadfield, K., Molton, S. A., Ley, R., Wagner, E. F., and Cook, S. J. (2003). "Activation of ERK1/2 by Delta Raf-1 : ER\* represses bim expression independently of the JNK or PI3K pathways." *Oncogene*, 22(9), 1281-1293.
- White, J. K., Auerbach, W., Duyao, M. P., Vonsattel, J. P., Gusella, J. F., Joyner, A. L., and MacDonald, M. E. (1997). "Huntingtin is required for neurogenesis and is not impaired by the Huntington's disease CAG expansion." *Nat. Genet.*, 17(4), 404-410.
- Williams, M. R., Arthur, J. S., Balendran, A., van der, K. J., Poli, V., Cohen, P., and Alessi, D. R. (2000). "The role of 3-phosphoinositide-dependent protein kinase 1 in activating AGC kinases defined in embryonic stem cells." *Curr. Biol.*, 10(8), 439-448.
- Willis, S. N., and Adams, J. M. (2005). "Life in the balance: how BH3-only proteins induce apoptosis." *Curr. Opin. Cell Biol.*, 17(6), 617-625.
- Willis, S. N., Chen, L., Dewson, G., Wei, A., Naik, E., Fletcher, J. I., Adams, J. M., and Huang, D. C. (2005). "Proapoptotic Bak is sequestered by Mcl-1 and Bcl-xL, but not Bcl-2, until displaced by BH3-only proteins." *Genes Dev.*, 19(11), 1294-1305.

- Wolf, B. B., and Green, D. R. (1999). "Suicidal tendencies: apoptotic cell death by caspase family proteinases." *J. Biol. Chem.*, 274(29), 20049-20052.
- Woo, M., Hakem, R., Soengas, M. S., Duncan, G. S., Shahinian, A., Kagi, D., Hakem, A., McCurrach, M., Khoo, W., Kaufman, S. A., Senaldi, G., Howard, T., Lowe, S. W., and Mak, T. W. (1998). "Essential contribution of caspase 3/CPP32 to apoptosis and its associated nuclear changes." *Genes Dev.*, 12(6), 806-819.
- Worby, C. A., Vega, Q. C., Zhao, Y., Chao, H. H., Seasholtz, A. F., and Dixon, J. E. (1996). "Glial cell line-derived neurotrophic factor signals through the RET receptor and activates mitogen-activated protein kinase." *J. Biol. Chem.*, 271(39), 23619-23622.
- Wu, C., Lai, C. F., and Mobley, W. C. (2001). "Nerve growth factor activates persistent Rap1 signaling in endosomes." *J. Neurosci.*, 21(15), 5406-5416.
- Wu, G. Y., Deisseroth, K., and Tsien, R. W. (2001). "Spaced stimuli stabilize MAPK pathway activation and its effects on dendritic morphology." *Nat. Neurosci.*, 4(2), 151-158.
- Wu, X., Zhu, D., Jiang, X., Okagaki, P., Mearow, K., Zhu, G., McCall, S., Banaudha, K., Lipsky, R. H., and Marini, A. M. (2004). "AMPA protects cultured neurons against glutamate excitotoxicity through a phosphatidylinositol 3-kinase-dependent activation in extracellular signal-regulated kinase to upregulate BDNF gene expression." *J. Neurochem.*, 90(4), 807-818.

---

**X**

- Xia, Z., Dickens, M., Raingeaud, J., Davis, R. J., and Greenberg, M. E. (1995). "Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis." *Science*, 270(5240), 1326-1331.

---

**Y**

- Yacoubian, T. A., and Lo, D. C. (2000). "Truncated and full-length TrkB receptors regulate distinct modes of dendritic growth." *Nat. Neurosci.*, 3(4), 342-349.
- Yamada, M., Ohnishi, H., Sano, S., Nakatani, A., Ikeuchi, T., and Hatanaka, H. (1997). "Insulin receptor substrate (IRS)-1 and IRS-2 are tyrosine-phosphorylated and associated with phosphatidylinositol 3-kinase in response to brain-derived neurotrophic factor in cultured cerebral cortical neurons." *J. Biol. Chem.*, 272(48), 30334-30339.
- Yamada, M., Tanabe, K., Wada, K., Shimoke, K., Ishikawa, Y., Ikeuchi, T., Koizumi, S., and Hatanaka, H. (2001). "Differences in survival-promoting effects and intracellular signaling properties of BDNF and IGF-1 in cultured cerebral cortical neurons." *J. Neurochem.*, 78(5), 940-951.
- Yamaguchi, H., and Wang, H. G. (2001). "The protein kinase PKB/Akt regulates cell survival and apoptosis by inhibiting Bax conformational change." *Oncogene*, 20(53), 7779-7786.

- Yamaguchi, H., and Wang, H. G. (2002). "Bcl-XL protects BimEL-induced Bax conformational change and cytochrome C release independent of interacting with Bax or BimEL." *J. Biol. Chem.*, 277(44), 41604-41612.
- Yamamoto, A., Lucas, J. J., and Hen, R. (2000). "Reversal of neuropathology and motor dysfunction in a conditional model of Huntington's disease." *Cell*, 101(1), 57-66.
- Yang, D. D., Kuan, C. Y., Whitmarsh, A. J., Rincon, M., Zheng, T. S., Davis, R. J., Rakic, P., and Flavell, R. A. (1997). "Absence of excitotoxicity-induced apoptosis in the hippocampus of mice lacking the Jnk3 gene." *Nature*, 389(6653), 865-870.
- Yang, L., Mao, L., Tang, Q., Samdani, S., Liu, Z., and Wang, J. Q. (2004). "A Novel Ca<sup>2+</sup>-independent signaling pathway to extracellular signal-regulated protein kinase by coactivation of NMDA receptors and metabotropic glutamate receptor 5 in neurons." *J. Neurosci.*, 24(48), 10846-10857.
- Yang, X., Chang, H. Y., and Baltimore, D. (1998). "Autoproteolytic activation of procaspases by oligomerization." *Mol. Cell*, 1(2), 319-325.
- Yao, R., and Cooper, G. M. (1995). "Requirement for phosphatidylinositol-3 kinase in the prevention of apoptosis by nerve growth factor." *Science*, 267(5206), 2003-2006.
- Yin, X. M., Wang, K., Gross, A., Zhao, Y. G., Zinkel, S., Klocke, B., Roth, K. A., and Korsmeyer, S. J. (1999). "Bid-deficient mice are resistant to Fas-induced hepatocellular apoptosis." *Nature*, 400(6747), 886-891.
- Yin, X. M. (2006). "Bid, a BH3-only multi-functional molecule, is at the cross road of life and death." *Gene*, 369, 7-19.
- Ying, S. W., Futter, M., Rosenblum, K., Webber, M. J., Hunt, S. P., Bliss, T. V., and Bramham, C. R. (2002). "Brain-derived neurotrophic factor induces long-term potentiation in intact adult hippocampus: requirement for ERK activation coupled to CREB and upregulation of Arc synthesis." *J. Neurosci.*, 22(5), 1532-1540.
- York, R. D., Yao, H., Dillon, T., Ellig, C. L., Eckert, S. P., McCleskey, E. W., and Stork, P. J. (1998). "Rap1 mediates sustained MAP kinase activation induced by nerve growth factor." *Nature*, 392(6676), 622-626.
- York, R. D., Molliver, D. C., Grewal, S. S., Stenberg, P. E., McCleskey, E. W., and Stork, P. J. (2000). "Role of phosphoinositide 3-kinase and endocytosis in nerve growth factor-induced extracellular signal-regulated kinase activation via Ras and Rap1." *Mol. Cell Biol.*, 20(21), 8069-8083.
- Young, A. B., Greenamyre, J. T., Hollingsworth, Z., Albin, R., D'Amato, C., Shoulson, I., and Penney, J. B. (1988). "NMDA receptor losses in putamen from patients with Huntington's disease." *Science*, 241(4868), 981-983.
- Yu, J., Zhang, L., Hwang, P. M., Kinzler, K. W., and Vogelstein, B. (2001). "PUMA induces the rapid apoptosis of colorectal cancer cells." *Mol. Cell*, 7(3), 673-682.

Yu, W. H., Cuervo, A. M., Kumar, A., Peterhoff, C. M., Schmidt, S. D., Lee, J. H., Mohan, P. S., Mercken, M., Farmery, M. R., Tjernberg, L. O., Jiang, Y., Duff, K., Uchiyama, Y., Naslund, J., Mathews, P. M., Cataldo, A. M., and Nixon, R. A. (2005). "Macroautophagy--a novel Beta-amyloid peptide-generating pathway activated in Alzheimer's disease." *J. Cell Biol.*, 171(1), 87-98.

Yuan, J., Lipinski, M., and Degterev, A. (2003). "Diversity in the mechanisms of neuronal cell death." *Neuron*, 40(2), 401-413.

---

## Z

Zeron, M. M., Chen, N., Moshaver, A., Lee, A. T., Wellington, C. L., Hayden, M. R., and Raymond, L. A. (2001). "Mutant huntingtin enhances excitotoxic cell death." *Mol. Cell Neurosci.*, 17(1), 41-53.

Zeron, M. M., Hansson, O., Chen, N., Wellington, C. L., Leavitt, B. R., Brundin, P., Hayden, M. R., and Raymond, L. A. (2002). "Increased sensitivity to N-methyl-D-aspartate receptor-mediated excitotoxicity in a mouse model of Huntington's disease." *Neuron*, 33(6), 849-860.

Zha, J., Harada, H., Yang, E., Jockel, J., and Korsmeyer, S. J. (1996). "Serine phosphorylation of death agonist BAD in response to survival factor results in binding to 14-3-3 not BCL-X(L)." *Cell*, 87(4), 619-628.

Zhang, Y., Li, M., Drozda, M., Chen, M., Ren, S., Mejia Sanchez, R. O., Leavitt, B. R., Cattaneo, E., Ferrante, R. J., Hayden, M. R., and Friedlander, R. M. (2003a). "Depletion of wild-type huntingtin in mouse models of neurologic diseases." *J. Neurochem.*, 87(1), 101-106.

Zhang, Y., Ona, V. O., Li, M., Drozda, M., Dubois-Dauphin, M., Przedborski, S., Ferrante, R. J., and Friedlander, R. M. (2003b). "Sequential activation of individual caspases, and of alterations in Bcl-2 proapoptotic signals in a mouse model of Huntington's disease." *J. Neurochem.*, 87(5), 1184-1192.

Zheng, T. S., Hunot, S., Kuida, K., Momoi, T., Srinivasan, A., Nicholson, D. W., Lazebnik, Y., and Flavell, R. A. (2000). "Deficiency in caspase-9 or caspase-3 induces compensatory caspase activation." *Nat. Med.*, 6(11), 1241-1247.

Zhu, D., Wu, X., Strauss, K. I., Lipsky, R. H., Qureshi, Z., Terhakopian, A., Novelli, A., Banaudha, K., and Marini, A. M. (2005). "N-methyl-D-aspartate and TrkB receptors protect neurons against glutamate excitotoxicity through an extracellular signal-regulated kinase pathway." *J. Neurosci. Res.*, 80(1), 104-113.

Zhu, J. H., Kulich, S. M., Oury, T. D., and Chu, C. T. (2002). "Cytoplasmic aggregates of phosphorylated extracellular signal-regulated protein kinases in Lewy body diseases." *Am. J. Pathol.*, 161(6), 2087-2098.

Zhu, Y., Swanson, B. J., Wang, M., Hildeman, D. A., Schaefer, B. C., Liu, X., Suzuki, H., Mihara, K., Kappler, J., and Marrack, P. (2004). "Constitutive association of the proapoptotic protein Bim with Bcl-2-related proteins on mitochondria in T cells." *Proc. Natl. Acad. Sci. U. S. A.*, 101(20), 7681-7686.

- Zong, W. X., Lindsten, T., Ross, A. J., MacGregor, G. R., and Thompson, C. B. (2001). "BH3-only proteins that bind pro-survival Bcl-2 family members fail to induce apoptosis in the absence of Bax and Bak." *Genes Dev.*, 15(12), 1481-1486.
- Zubenko, G. S., Stiffler, J. S., Hughes, H. B., and Martinez, A. J. (1999). "Reductions in brain phosphatidylinositol kinase activities in Alzheimer's disease." *Biol. Psychiatry*, 45(6), 731-736.
- Zuccato, C., Ciammola, A., Rigamonti, D., Leavitt, B. R., Goffredo, D., Conti, L., MacDonald, M. E., Friedlander, R. M., Silani, V., Hayden, M. R., Timmusk, T., Sipione, S., and Cattaneo, E. (2001). "Loss of huntingtin-mediated BDNF gene transcription in Huntington's disease." *Science*, 293(5529), 493-498.
- Zuccato, C., Tartari, M., Crotti, A., Goffredo, D., Valenza, M., Conti, L., Cataudella, T., Leavitt, B. R., Hayden, M. R., Timmusk, T., Rigamonti, D., and Cattaneo, E. (2003). "Huntingtin interacts with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes." *Nat. Genet.*, 35(1), 76-83.

