

UNIVERSITAT DE BARCELONA

FACULTAT DE FARMÀCIA

DEPARTAMENT DE BIOQUÍMICA I BIOLOGIA MOLECULAR

REGULACIÓ DEL PROMOTOR DE *Sp3*

ALICIA TAPIAS SOLER
Barcelona 2008

BIBLIOGRAFIA



A

- Abella A, Dubus P, Malumbres M, Rane SG, Kiyokawa H, Sicard A, et al. (2005). Cdk4 promotes adipogenesis through PPARgamma activation. *Cell Metab* **2**: 239-49.
- Abramovitch S, Glaser T, Ouchi T, Werner H (2003). BRCA1-Sp1 interactions in transcriptional regulation of the IGF-IR gene. *FEBS Lett* **541**: 149-54.
- Adachi N, Nomoto M, Kohno K, Koyama H (2000). Cell-cycle regulation of the DNA topoisomerase IIalpha promoter is mediated by proximal CCAAT boxes: possible involvement of acetylation. *Gene* **245**: 49-57.
- Adrian GS, Seto E, Fischbach KS, Rivera EV, Adrian EK, Herbert DC, et al. (1996). YY1 and Sp1 transcription factors bind the human transferrin gene in an age-related manner. *J Gerontol A Biol Sci Med Sci* **51**: B66-75.
- Alevizopoulos A, Dusserre Y, Tsai-Pflugfelder M, von der Weid T, Wahli W, Mermod N (1995). A proline-rich TGF-beta-responsive transcriptional activator interacts with histone H3. *Genes Dev* **9**: 3051-66.
- Alroy I, Soussan L, Seger R, Yarden Y (1999). Neu differentiation factor stimulates phosphorylation and activation of the Sp1 transcription factor. *Mol Cell Biol* **19**: 1961-72.
- Ammanamanchi S, Freeman JW, Brattain MG (2003). Acetylated sp3 is a transcriptional activator. *J Biol Chem* **278**: 35775-80.
- Ammendola R, Mesuraca M, Russo T, Cimino F (1992). Sp1 DNA binding efficiency is highly reduced in nuclear extracts from aged rat tissues. *J Biol Chem* **267**: 17944-8.
- Angel P, Hattori K, Smeal T, Karin M (1988). The jun proto-oncogene is positively autoregulated by its product, Jun/AP-1. *Cell* **55**: 875-85.
- Angel P, Imagawa M, Chiu R, Stein B, Imbra RJ, Rahmsdorf HJ, et al. (1987). Phorbol ester-inducible genes contain a common cis element recognized by a TPA-modulated trans-acting factor. *Cell* **49**: 729-39.
- Apt D, Watts RM, Suske G, Bernard HU (1996). High Sp1/Sp3 ratios in epithelial cells during epithelial differentiation and cellular transformation correlate with the activation of the HPV-16 promoter. *Virology* **224**: 281-91.

BIBLIOGRAFIA

- Araki E, Murakami T, Shirotani T, Kanai F, Shinohara Y, Shimada F, et al. (1991). A cluster of four Sp1 binding sites required for efficient expression of the human insulin receptor gene. *J Biol Chem* **266**: 3944-8.
- Armentero MT, Horwitz M, Mermod N (1994). Targeting of DNA polymerase to the adenovirus origin of DNA replication by interaction with nuclear factor I. *Proc Natl Acad Sci U S A* **91**: 11537-41.
- Armstrong SA, Barry DA, Leggett RW, Mueller CR (1997). Casein kinase II-mediated phosphorylation of the C terminus of Sp1 decreases its DNA binding activity. *J Biol Chem* **272**: 13489-95.
- Arsura M, Introna M, Passerini F, Mantovani A, Golay J (1992). B-myb antisense oligonucleotides inhibit proliferation of human hematopoietic cell lines. *Blood* **79**: 2708-16.
- Athanikar JN, Sanchez HB, Osborne TF (1997). Promoter selective transcriptional synergy mediated by sterol regulatory element binding protein and Sp1: a critical role for the Btd domain of Sp1. *Mol Cell Biol* **17**: 5193-200.
- Aznar S, Valeron PF, del Rincon SV, Perez LF, Perona R, Lacal JC (2001). Simultaneous tyrosine and serine phosphorylation of STAT3 transcription factor is involved in Rho A GTPase oncogenic transformation. *Mol Biol Cell* **12**: 3282-94.

B

- Baker SJ, Markowitz S, Fearon ER, Willson JK, Vogelstein B (1990a). Suppression of human colorectal carcinoma cell growth by wild-type p53. *Science* **249**: 912-5.
- Baker SJ, Preisinger AC, Jessup JM, Paraskeva C, Markowitz S, Willson JK, et al. (1990b). p53 gene mutations occur in combination with 17p allelic deletions as late events in colorectal tumorigenesis. *Cancer Res* **50**: 7717-22.
- Bakovic M, Waite K, Vance DE (2003). Oncogenic Ha-Ras transformation modulates the transcription of the CTP:phosphocholine cytidylyltransferase alpha gene via p42/44MAPK and transcription factor Sp3. *J Biol Chem* **278**: 14753-61.

- Banchio C, Schang LM, Vance DE (2004). Phosphorylation of Sp1 by cyclin-dependent kinase 2 modulates the role of Sp1 in CTP:phosphocholine cytidylyltransferase alpha regulation during the S phase of the cell cycle. *J Biol Chem* **279**: 40220-6.
- Bartusel T, Schubert S, Klempnauer KH (2005). Regulation of the cyclin D1 and cyclin A1 promoters by B-Myb is mediated by Sp1 binding sites. *Gene* **351**: 171-80.
- Bassal S, El-Osta A (2005). DNA damage detection and repair, and the involvement of epigenetic states. *Hum Mutat* **25**: 101-9.
- Bates S, Phillips AC, Clark PA, Stott F, Peters G, Ludwig RL, et al. (1998). p14ARF links the tumour suppressors RB and p53. *Nature* **395**: 124-5.
- Bell SM, Schreiner CM, Waclaw RR, Campbell K, Potter SS, Scott WJ (2003). Sp8 is crucial for limb outgrowth and neuropore closure. *Proc Natl Acad Sci U S A* **100**: 12195-200.
- Ben-Shachar D, Karry R (2007). Sp1 expression is disrupted in schizophrenia; a possible mechanism for the abnormal expression of mitochondrial complex I genes, NDUFV1 and NDUFV2. *PLoS ONE* **2**: e817.
- Benatti P, Basile V, Merico D, Fantoni LI, Tagliafico E, Imbriano C (2008). A balance between NF-Y and p53 governs the pro- and anti-apoptotic transcriptional response. *Nucleic Acids Res* **36**: 1415-28.
- Benoist C, O'Hare K, Breathnach R, Chambon P (1980). The ovalbumin gene-sequence of putative control regions. *Nucleic Acids Res* **8**: 127-42.
- Berger I, Shaul Y (1994). The human junD gene is positively and selectively autoregulated. *DNA Cell Biol* **13**: 249-55.
- Bernier D, Thomassin H, Allard D, Guertin M, Hamel D, Blaquier M, et al. (1993). Functional analysis of developmentally regulated chromatin-hypersensitive domains carrying the alpha 1-fetoprotein gene promoter and the albumin/alpha 1-fetoprotein intergenic enhancer. *Mol Cell Biol* **13**: 1619-33.
- Bhattacharya A, Deng JM, Zhang Z, Behringer R, de Crombrugghe B, Maity SN (2003). The B subunit of the CCAAT box binding transcription factor complex (CBF/NF-Y) is essential for early mouse development and cell proliferation. *Cancer Res* **63**: 8167-72.

BIBLIOGRAFIA

- Black AR, Black JD,Azizkhan-Clifford J (2001). Sp1 and kruppel-like factor family of transcription factors in cell growth regulation and cancer. *J Cell Physiol* **188**: 143-60.
- Black AR, Jensen D, Lin SY,Azizkhan JC (1999). Growth/cell cycle regulation of Sp1 phosphorylation. *J Biol Chem* **274**: 1207-15.
- Bonello MR,Khachigian LM (2004). Fibroblast growth factor-2 represses platelet-derived growth factor receptor-alpha (PDGFR-alpha) transcription via ERK1/2-dependent Sp1 phosphorylation and an atypical cis-acting element in the proximal PDGFR-alpha promoter. *J Biol Chem* **279**: 2377-82.
- Borellini F,Glazer RI (1993). Induction of Sp1-p53 DNA-binding heterocomplexes during granulocyte/macrophage colony-stimulating factor-dependent proliferation in human erythroleukemia cell line TF-1. *J Biol Chem* **268**: 7923-8.
- Bosher J, Robinson EC,Hay RT (1990). Interactions between the adenovirus type 2 DNA polymerase and the DNA binding domain of nuclear factor I. *New Biol* **2**: 1083-90.
- Botz J, Zerfass-Thome K, Spitkovsky D, Delius H, Vogt B, Eilers M, et al. (1996). Cell cycle regulation of the murine cyclin E gene depends on an E2F binding site in the promoter. *Mol Cell Biol* **16**: 3401-9.
- Boutillier S, Lannes B, Buee L, Delacourte A, Rouaux C, Mohr M, et al. (2007). Sp3 and sp4 transcription factor levels are increased in brains of patients with Alzheimer's disease. *Neurodegener Dis* **4**: 413-23.
- Bouwman P, Gollner H, Elsasser HP, Eckhoff G, Karis A, Grosveld F, et al. (2000). Transcription factor Sp3 is essential for post-natal survival and late tooth development. *Embo J* **19**: 655-61.
- Bouwman P,Philipsen S (2002). Regulation of the activity of Sp1-related transcription factors. *Mol Cell Endocrinol* **195**: 27-38.
- Brandt TL,Kroll DJ (1997). NF-M trans-activates the human DNA topoisomerase II alpha promoter independently of c-Myb in HL-60 cells. *Leuk Res* **21**: 711-20.
- Braun H, Koop R, Ertmer A, Nacht S,Suske G (2001). Transcription factor Sp3 is regulated by acetylation. *Nucleic Acids Res* **29**: 4994-5000.

Brodin G, Ahgren A, ten Dijke P, Heldin CH, Heuchel R (2000). Efficient TGF-beta induction of the Smad7 gene requires cooperation between AP-1, Sp1, and Smad proteins on the mouse Smad7 promoter. *J Biol Chem* **275**: 29023-30.

Broude EV, Swift ME, Vivo C, Chang BD, Davis BM, Kalurupalle S, et al. (2007). p21(Waf1/Cip1/Sdi1) mediates retinoblastoma protein degradation. *Oncogene* **26**: 6954-8.

Brusselbach S, Mohle-Steinlein U, Wang ZQ, Schreiber M, Lucibello FC, Muller R, et al. (1995). Cell proliferation and cell cycle progression are not impaired in fibroblasts and ES cells lacking c-Fos. *Oncogene* **10**: 79-86.

Bucher P (1990). Weight matrix descriptions of four eukaryotic RNA polymerase II promoter elements derived from 502 unrelated promoter sequences. *J Mol Biol* **212**: 563-78.

C

Caretti G, Motta MC, Mantovani R (1999). NF-Y associates with H3-H4 tetramers and octamers by multiple mechanisms. *Mol Cell Biol* **19**: 8591-603.

Caretti G, Salsi V, Vecchi C, Imbriano C, Mantovani R (2003). Dynamic recruitment of NF-Y and histone acetyltransferases on cell-cycle promoters. *J Biol Chem* **278**: 30435-40.

Carrano AC, Eytan E, Hershko A, Pagano M (1999). SKP2 is required for ubiquitin-mediated degradation of the CDK inhibitor p27. *Nat Cell Biol* **1**: 193-9.

Casanovas O, Miro F, Estanyol JM, Itarte E, Agell N, Bachs O (2000). Osmotic stress regulates the stability of cyclin D1 in a p38SAPK2-dependent manner. *J Biol Chem* **275**: 35091-7.

Chang BD, Watanabe K, Broude EV, Fang J, Poole JC, Kalinichenko TV, et al. (2000). Effects of p21Waf1/Cip1/Sdi1 on cellular gene expression: implications for carcinogenesis, senescence, and age-related diseases. *Proc Natl Acad Sci U S A* **97**: 4291-6.

BIBLIOGRAFIA

- Chang WC,Chen BK (2005). Transcription factor Sp1 functions as an anchor protein in gene transcription of human 12(S)-lipoxygenase. *Biochem Biophys Res Commun* **338**: 117-21.
- Chellappan SP, Hiebert S, Mudryj M, Horowitz JM,Nevins JR (1991). The E2F transcription factor is a cellular target for the RB protein. *Cell* **65**: 1053-61.
- Chiang CM,Roeder RG (1995). Cloning of an intrinsic human TFIID subunit that interacts with multiple transcriptional activators. *Science* **267**: 531-6.
- Chinenov Y,Kerppola TK (2001). Close encounters of many kinds: Fos-Jun interactions that mediate transcription regulatory specificity. *Oncogene* **20**: 2438-52.
- Chiu R, Angel P,Karin M (1989). Jun-B differs in its biological properties from, and is a negative regulator of, c-Jun. *Cell* **59**: 979-86.
- Christensen MA, Zhou W, Qing H, Lehman A, Philipsen S,Song W (2004). Transcriptional regulation of BACE1, the beta-amyloid precursor protein beta-secretase, by Sp1. *Mol Cell Biol* **24**: 865-74.
- Chu S, Cockrell CA,Ferro TJ (2003). Expression of alpha-ENaC2 is dependent on an upstream Sp1 binding motif and is modulated by protein phosphatase 1 in lung epithelial cells. *Biochem Biophys Res Commun* **303**: 1159-68.
- Chu S,Ferro TJ (2005). Sp1: regulation of gene expression by phosphorylation. *Gene* **348**: 1-11.
- Chun RF, Semmes OJ, Neuveut C,Jeang KT (1998). Modulation of Sp1 phosphorylation by human immunodeficiency virus type 1 Tat. *J Virol* **72**: 2615-29.
- Chupreata S, Du M, Todisco A,Merchant JL (2000). EGF stimulates gastrin promoter through activation of Sp1 kinase activity. *Am J Physiol Cell Physiol* **278**: C697-708.
- Cieslik K, Lee CM, Tang JL,Wu KK (1999). Transcriptional regulation of endothelial nitric-oxide synthase by an interaction between casein kinase 2 and protein phosphatase 2A. *J Biol Chem* **274**: 34669-75.
- Ciudad CJ, Morris AE, Jeng C,Chasin LA (1992). Point mutational analysis of the hamster dihydrofolate reductase minimum promoter. *J Biol Chem* **267**: 3650-6.

Ciudad CJ, Urlaub G, Chasin LA (1988). Deletion analysis of the Chinese hamster dihydrofolate reductase gene promoter. *J Biol Chem* **263**: 16274-82.

Collins FS, Metherall JE, Yamakawa M, Pan J, Weissman SM, Forget BG (1985). A point mutation in the A gamma-globin gene promoter in Greek hereditary persistence of fetal haemoglobin. *Nature* **313**: 325-6.

Conaway RC, Brower CS, Conaway JW (2002). Emerging roles of ubiquitin in transcription regulation. *Science* **296**: 1254-8.

Coqueret O (2003). New roles for p21 and p27 cell-cycle inhibitors: a function for each cell compartment? *Trends Cell Biol* **13**: 65-70.

Courey AJ, Tjian R (1988). Analysis of Sp1 in vivo reveals multiple transcriptional domains, including a novel glutamine-rich activation motif. *Cell* **55**: 887-98.

Cress WD, Nevins JR (1996). A role for a bent DNA structure in E2F-mediated transcription activation. *Mol Cell Biol* **16**: 2119-27.

Crossley M, Whitelaw E, Perkins A, Williams G, Fujiwara Y, Orkin SH (1996). Isolation and characterization of the cDNA encoding BKLF/TEF-2, a major CACCC-box-binding protein in erythroid cells and selected other cells. *Mol Cell Biol* **16**: 1695-705.

Crowe DL, Nguyen DC, Tsang KJ, Kyo S (2001). E2F-1 represses transcription of the human telomerase reverse transcriptase gene. *Nucleic Acids Res* **29**: 2789-94.

Currie RA (1997). Functional interaction between the DNA binding subunit trimerization domain of NF-Y and the high mobility group protein HMG-I(Y). *J Biol Chem* **272**: 30880-8.

D

Daniel S, Kim KH (1996). Sp1 mediates glucose activation of the acetyl-CoA carboxylase promoter. *J Biol Chem* **271**: 1385-92.

Daniel S, Zhang S, DePaoli-Roach AA, Kim KH (1996). Dephosphorylation of Sp1 by protein phosphatase 1 is involved in the glucose-mediated activation of the acetyl-CoA carboxylase gene. *J Biol Chem* **271**: 14692-7.

BIBLIOGRAFIA

- Darnell JE, Jr. (1997). STATs and gene regulation. *Science* **277**: 1630-5.
- das Neves L, Duchala CS, Tolentino-Silva F, Haxhiu MA, Colmenares C, Macklin WB, et al. (1999). Disruption of the murine nuclear factor I-A gene (Nfia) results in perinatal lethality, hydrocephalus, and agenesis of the corpus callosum. *Proc Natl Acad Sci U S A* **96**: 11946-51.
- del Pozo JC, Estelle M (2000). F-box proteins and protein degradation: an emerging theme in cellular regulation. *Plant Mol Biol* **44**: 123-8.
- Dennig J, Beato M, Suske G (1996). An inhibitor domain in Sp3 regulates its glutamine-rich activation domains. *Embo J* **15**: 5659-67.
- Deshaijes RJ (1999). SCF and Cullin/Ring H2-based ubiquitin ligases. *Annu Rev Cell Dev Biol* **15**: 435-67.
- Discher DJ, Bishopric NH, Wu X, Peterson CA, Webster KA (1998). Hypoxia regulates beta-enolase and pyruvate kinase-M promoters by modulating Sp1/Sp3 binding to a conserved GC element. *J Biol Chem* **273**: 26087-93.
- Doetzlhofer A, Rotheneder H, Lagger G, Koranda M, Kurtev V, Brosch G, et al. (1999). Histone deacetylase 1 can repress transcription by binding to Sp1. *Mol Cell Biol* **19**: 5504-11.
- Dorn A, Bollekens J, Staub A, Benoist C, Mathis D (1987). A multiplicity of CCAAT box-binding proteins. *Cell* **50**: 863-72.
- Dou QP, Fridovich-Keil JL, Pardee AB (1991). Inducible proteins binding to the murine thymidine kinase promoter in late G1/S phase. *Proc Natl Acad Sci U S A* **88**: 1157-61.
- Du XL, Edelstein D, Rossetti L, Fantus IG, Goldberg H, Ziyadeh F, et al. (2000). Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc Natl Acad Sci U S A* **97**: 12222-6.

E

Efstratiadis A, Posakony JW, Maniatis T, Lawn RM, O'Connell C, Spritz RA, et al. (1980). The structure and evolution of the human beta-globin gene family. *Cell* **21**: 653-68.

Elkon R, Linhart C, Sharan R, Shamir R, Shiloh Y (2003). Genome-wide in silico identification of transcriptional regulators controlling the cell cycle in human cells. *Genome Res* **13**: 773-80.

Elledge SJ, Winston J, Harper JW (1996). A question of balance: the role of cyclin-kinase inhibitors in development and tumorigenesis. *Trends Cell Biol* **6**: 388-92.

Ellis DJ, Dehm SM, Bonham K (2006). The modification of Sp3 isoforms by SUMOylation has differential effects on the SRC1A promoter. *Gene* **379**: 68-78.

Elrod-Erickson M, Rould MA, Nekludova L, Pabo CO (1996). Zif268 protein-DNA complex refined at 1.6 Å: a model system for understanding zinc finger-DNA interactions. *Structure* **4**: 1171-80.

F

Feng WC, Southwood CM, Bieker JJ (1994). Analyses of beta-thalassemia mutant DNA interactions with erythroid Kruppel-like factor (EKLF), an erythroid cell-specific transcription factor. *J Biol Chem* **269**: 1493-500.

Ferrante RJ, Kubilus JK, Lee J, Ryu H, Beesen A, Zucker B, et al. (2003). Histone deacetylase inhibition by sodium butyrate chemotherapy ameliorates the neurodegenerative phenotype in Huntington's disease mice. *J Neurosci* **23**: 9418-27.

Fisch S, Gray S, Heymans S, Haldar SM, Wang B, Pfister O, et al. (2007). Kruppel-like factor 15 is a regulator of cardiomyocyte hypertrophy. *Proc Natl Acad Sci U S A* **104**: 7074-9.

Fojas de Borja P, Collins NK, Du P, Azizkhan-Clifford J, Mudryj M (2001). Cyclin A-CDK phosphorylates Sp1 and enhances Sp1-mediated transcription. *Embo J* **20**: 5737-47.

Frampton J, Ramqvist T, Graf T (1996). v-Myb of E26 leukemia virus up-regulates bcl-2 and suppresses apoptosis in myeloid cells. *Genes Dev* **10**: 2720-31.

Frit P, Kwon K, Coin F, Auriol J, Dubaele S, Salles B, et al. (2002). Transcriptional activators stimulate DNA repair. *Mol Cell* **10**: 1391-401.

Frontini M, Imbriano C, Manni I, Mantovani R (2004). Cell cycle regulation of NF-YC nuclear localization. *Cell Cycle* **3**: 217-22.

Fu YH, Nishinaka T, Yokoyama K, Chiu R (1998). A retinoblastoma susceptibility gene product, RB, targeting protease is regulated through the cell cycle. *FEBS Lett* **421**: 89-93.

Fukada T, Ohtani T, Yoshida Y, Shirogane T, Nishida K, Nakajima K, et al. (1998). STAT3 orchestrates contradictory signals in cytokine-induced G1 to S cell-cycle transition. *Embo J* **17**: 6670-7.

G

Gao B, Jaffe H, Kunos G (1998). Histone H1 isoforms purified from rat liver bind nonspecifically to the nuclear factor 1 recognition sequence and serve as generalized transcriptional repressors. *Mol Cell Biochem* **178**: 187-96.

Gartel AL, Goufman E, Najmabadi F, Tyner AL (2000). Sp1 and Sp3 activate p21 (WAF1/CIP1) gene transcription in the Caco-2 colon adenocarcinoma cell line. *Oncogene* **19**: 5182-8.

Gartel AL, Ye X, Goufman E, Shianov P, Hay N, Najmabadi F, et al. (2001). Myc represses the p21(WAF1/CIP1) promoter and interacts with Sp1/Sp3. *Proc Natl Acad Sci U S A* **98**: 4510-5.

Gelinas R, Endlich B, Pfeiffer C, Yagi M, Stamatoyannopoulos G (1985). G to A substitution in the distal CCAAT box of the A gamma-globin gene in Greek hereditary persistence of fetal haemoglobin. *Nature* **313**: 323-5.

Gerondakis S, Grumont R, Gugasyan R, Wong L, Isomura I, Ho W, et al. (2006). Unravelling the complexities of the NF-kappaB signalling pathway using mouse knockout and transgenic models. *Oncogene* **25**: 6781-99.

Geurts JM, Schoenmakers EF, Roijer E, Astrom AK, Stenman G, van de Ven WJ (1998). Identification of NF1B as recurrent translocation partner gene of HMGIC in pleomorphic adenomas. *Oncogene* **16**: 865-72.

Gewirtz AM, Anfossi G, Venturelli D, Valpreda S, Sims R, Calabretta B (1989). G1/S transition in normal human T-lymphocytes requires the nuclear protein encoded by c-myb. *Science* **245**: 180-3.

Gewirtz AM, Calabretta B (1988). A c-myb antisense oligodeoxynucleotide inhibits normal human hematopoiesis in vitro. *Science* **242**: 1303-6.

Gill G (2003). Post-translational modification by the small ubiquitin-related modifier SUMO has big effects on transcription factor activity. *Curr Opin Genet Dev* **13**: 108-13.

Gollner H, Bouwman P, Mangold M, Karis A, Braun H, Rohner I, et al. (2001a). Complex phenotype of mice homozygous for a null mutation in the Sp4 transcription factor gene. *Genes Cells* **6**: 689-97.

Gollner H, Dani C, Phillips B, Philipsen S, Suske G (2001b). Impaired ossification in mice lacking the transcription factor Sp3. *Mech Dev* **106**: 77-83.

Gonda TJ, Sheiness DK, Bishop JM (1982). Transcripts from the cellular homologs of retroviral oncogenes: distribution among chicken tissues. *Mol Cell Biol* **2**: 617-24.

Gorospe M, Wang X, Holbrook NJ (1998). p53-dependent elevation of p21Waf1 expression by UV light is mediated through mRNA stabilization and involves a vanadate-sensitive regulatory system. *Mol Cell Biol* **18**: 1400-7.

Grana X, Reddy EP (1995). Cell cycle control in mammalian cells: role of cyclins, cyclin dependent kinases (CDKs), growth suppressor genes and cyclin-dependent kinase inhibitors (CKIs). *Oncogene* **11**: 211-9.

Gregory DJ, Garcia-Wilson E, Poole JC, Snowden AW, Roninson IB, Perkins ND (2002). Induction of transcription through the p300 CRD1 motif by p21WAF1/CIP1 is core promoter specific and cyclin dependent kinase independent. *Cell Cycle* **1**: 343-50.

Grekova MC, Scherer SW, Trabb J, Richert JR (2000). Localization of the human SP3 gene to chromosome 7p14-p15.2. The lack of expression in multiple sclerosis does not reflect abnormal gene organization. *J Neuroimmunol* **106**: 214-9.

Gronostajski RM (2000). Roles of the NFI/CTF gene family in transcription and development. *Gene* **249**: 31-45.

Gurtner A, Manni I, Fuschi P, Mantovani R, Guadagni F, Sacchi A, et al. (2003). Requirement for down-regulation of the CCAAT-binding activity of the NF-Y transcription factor during skeletal muscle differentiation. *Mol Biol Cell* **14**: 2706-15.

H

Hagen G, Muller S, Beato M, Suske G (1992). Cloning by recognition site screening of two novel GT box binding proteins: a family of Sp1 related genes. *Nucleic Acids Res* **20**: 5519-25.

Hagen G, Muller S, Beato M, Suske G (1994). Sp1-mediated transcriptional activation is repressed by Sp3. *Embo J* **13**: 3843-51.

Haidweger E, Novy M, Rotheneder H (2001). Modulation of Sp1 activity by a cyclin A/CDK complex. *J Mol Biol* **306**: 201-12.

Halazonetis TD, Georgopoulos K, Greenberg ME, Leder P (1988). c-Jun dimerizes with itself and with c-Fos, forming complexes of different DNA binding affinities. *Cell* **55**: 917-24.

Han I, Kudlow JE (1997). Reduced O glycosylation of Sp1 is associated with increased proteasome susceptibility. *Mol Cell Biol* **17**: 2550-8.

Harada K, Ogden GR (2000). An overview of the cell cycle arrest protein, p21(WAF1). *Oral Oncol* **36**: 3-7.

Harbour JW, Dean DC (2000). The Rb/E2F pathway: expanding roles and emerging paradigms. *Genes Dev* **14**: 2393-409.

Harper JW, Elledge SJ, Keyomarsi K, Dynlacht B, Tsai LH, Zhang P, et al. (1995). Inhibition of cyclin-dependent kinases by p21. *Mol Biol Cell* **6**: 387-400.

Harrison SM, Houzelstein D, Dunwoodie SL, Beddington RS (2000). Sp5, a new member of the Sp1 family, is dynamically expressed during development and genetically interacts with Brachyury. *Dev Biol* **227**: 358-72.

- Hatamochi A, Paterson B, de Crombrugghe B (1986). Differential binding of a CCAAT DNA binding factor to the promoters of the mouse alpha 2(I) and alpha 1(III) collagen genes. *J Biol Chem* **261**: 11310-4.
- He S, Davie JR (2006). Sp1 and Sp3 foci distribution throughout mitosis. *J Cell Sci* **119**: 1063-70.
- He S, Sun JM, Li L, Davie JR (2005). Differential intranuclear organization of transcription factors Sp1 and Sp3. *Mol Biol Cell* **16**: 4073-83.
- Heicklen-Klein A, Ginzburg I (2000). Tau promoter confers neuronal specificity and binds Sp1 and AP-2. *J Neurochem* **75**: 1408-18.
- Helenius M, Hanninen M, Lehtinen SK, Salminen A (1996). Changes associated with aging and replicative senescence in the regulation of transcription factor nuclear factor-kappa B. *Biochem J* **318 (Pt 2)**: 603-8.
- Hernandez EM, Chan CH, Xu B, Notario V, Richert JR (2004). Role of an internal ribosome entry site in the translational control of the human transcription factor Sp3. *Int J Oncol* **24**: 719-24.
- Hernandez EM, Johnson A, Notario V, Chen A, Richert JR (2002). AUA as a translation initiation site in vitro for the human transcription factor Sp3. *J Biochem Mol Biol* **35**: 273-82.
- Hershko A, Ciechanover A (1998). The ubiquitin system. *Annu Rev Biochem* **67**: 425-79.
- Hooft van Huijsduijnen R, Li XY, Black D, Matthes H, Benoist C, Mathis D (1990). Co-evolution from yeast to mouse: cDNA cloning of the two NF-Y (CP-1/CBF) subunits. *Embo J* **9**: 3119-27.
- Hooft van Huijsduijnen RA, Bollekens J, Dorn A, Benoist C, Mathis D (1987). Properties of a CCAAT box-binding protein. *Nucleic Acids Res* **15**: 7265-82.
- Hsiao KM, McMahon SL, Farnham PJ (1994). Multiple DNA elements are required for the growth regulation of the mouse E2F1 promoter. *Genes Dev* **8**: 1526-37.
- Hung JJ, Wang YT, Chang WC (2006). Sp1 deacetylation induced by phorbol ester recruits p300 to activate 12(S)-lipoxygenase gene transcription. *Mol Cell Biol* **26**: 1770-85.

BIBLIOGRAFIA

Husmann M, Dragneva Y, Romahn E, Jehnichen P (2000). Nuclear receptors modulate the interaction of Sp1 and GC-rich DNA via ternary complex formation. *Biochem J* **352 Pt 3**: 763-72.

I

Iaquinta PJ, Lees JA (2007). Life and death decisions by the E2F transcription factors. *Curr Opin Cell Biol* **19**: 649-57.

Imbriano C, Gurtner A, Cocchiarella F, Di Agostino S, Basile V, Gostissa M, et al. (2005). Direct p53 transcriptional repression: in vivo analysis of CCAAT-containing G2/M promoters. *Mol Cell Biol* **25**: 3737-51.

Ivey-Hoyle M, Conroy R, Huber HE, Goodhart PJ, Oliff A, Heimbrook DC (1993). Cloning and characterization of E2F-2, a novel protein with the biochemical properties of transcription factor E2F. *Mol Cell Biol* **13**: 7802-12.

J

Jackson S, Gottlieb T, Hartley K (1993). Phosphorylation of transcription factor Sp1 by the DNA-dependent protein kinase. *Adv Second Messenger Phosphoprotein Res* **28**: 279-86.

Jackson SP, MacDonald JJ, Lees-Miller S, Tjian R (1990). GC box binding induces phosphorylation of Sp1 by a DNA-dependent protein kinase. *Cell* **63**: 155-65.

Jackson SP, Tjian R (1988). O-glycosylation of eukaryotic transcription factors: implications for mechanisms of transcriptional regulation. *Cell* **55**: 125-33.

Jacobson D, Pribnow D, Herson PS, Maylie J, Adelman JP (2003). Determinants contributing to estrogen-regulated expression of SK3. *Biochem Biophys Res Commun* **303**: 660-8.

Jeggo PA, Lobrich M (2006). Contribution of DNA repair and cell cycle checkpoint arrest to the maintenance of genomic stability. *DNA Repair (Amst)* **5**: 1192-8.

Jensen DE, Black AR, Swick AG,Azizkhan JC (1997). Distinct roles for Sp1 and E2F sites in the growth/cell cycle regulation of the DHFR promoter. *J Cell Biochem* **67**: 24-31.

Jin S,Scotto KW (1998). Transcriptional regulation of the MDR1 gene by histone acetyltransferase and deacetylase is mediated by NF-Y. *Mol Cell Biol* **18**: 4377-84.

Johnson DG,Walker CL (1999). Cyclins and cell cycle checkpoints. *Annu Rev Pharmacol Toxicol* **39**: 295-312.

Johnson RS, van Lingen B, Papaioannou VE,Spiegelman BM (1993). A null mutation at the c-jun locus causes embryonic lethality and retarded cell growth in culture. *Genes Dev* **7**: 1309-17.

Jones KA, Kadonaga JT, Rosenfeld PJ, Kelly TJ,Tjian R (1987). A cellular DNA-binding protein that activates eukaryotic transcription and DNA replication. *Cell* **48**: 79-89.

K

Kaczynski J, Cook T,Urrutia R (2003). Sp1- and Kruppel-like transcription factors. *Genome Biol* **4**: 206.

Kadonaga JT, Courey AJ, Ladika J,Tjian R (1988). Distinct regions of Sp1 modulate DNA binding and transcriptional activation. *Science* **242**: 1566-70.

Kaelin WG, Jr., Krek W, Sellers WR, DeCaprio JA, Ajchenbaum F, Fuchs CS, et al. (1992). Expression cloning of a cDNA encoding a retinoblastoma-binding protein with E2F-like properties. *Cell* **70**: 351-64.

Kalff-Suske M, Kunz J, Grzeschik KH,Suske G (1996). Human Sp3 transcriptional regulator gene (SP3) maps to chromosome 2q31. *Genomics* **37**: 410-2.

Kardassis D, Papakosta P, Pardali K,Moustakas A (1999). c-Jun transactivates the promoter of the human p21(WAF1/Cip1) gene by acting as a superactivator of the ubiquitous transcription factor Sp1. *J Biol Chem* **274**: 29572-81.

Karin M (1995). The regulation of AP-1 activity by mitogen-activated protein kinases. *J Biol Chem* **270**: 16483-6.

BIBLIOGRAFIA

- Karin M (1996). The regulation of AP-1 activity by mitogen-activated protein kinases. *Philos Trans R Soc Lond B Biol Sci* **351**: 127-34.
- Karin M (1998). Mitogen-activated protein kinase cascades as regulators of stress responses. *Ann N Y Acad Sci* **851**: 139-46.
- Karlseder J, Rotheneder H,Wintersberger E (1996). Interaction of Sp1 with the growth-and cell cycle-regulated transcription factor E2F. *Mol Cell Biol* **16**: 1659-67.
- Katz JP, Perreault N, Goldstein BG, Actman L, McNally SR, Silberg DG, et al. (2005). Loss of Klf4 in mice causes altered proliferation and differentiation and precancerous changes in the adult stomach. *Gastroenterology* **128**: 935-45.
- Kaufmann J,Smale ST (1994). Direct recognition of initiator elements by a component of the transcription factor IID complex. *Genes Dev* **8**: 821-9.
- Kawakami Y, Esteban CR, Matsui T, Rodriguez-Leon J, Kato S,Belmonte JC (2004). Sp8 and Sp9, two closely related buttonhead-like transcription factors, regulate Fgf8 expression and limb outgrowth in vertebrate embryos. *Development* **131**: 4763-74.
- Kennett SB, Moorefield KS,Horowitz JM (2002). Sp3 represses gene expression via the titration of promoter-specific transcription factors. *J Biol Chem* **277**: 9780-9.
- Kennett SB, Udvadia AJ,Horowitz JM (1997). Sp3 encodes multiple proteins that differ in their capacity to stimulate or repress transcription. *Nucleic Acids Res* **25**: 3110-7.
- Kim CG,Sheffery M (1990). Physical characterization of the purified CCAAT transcription factor, alpha-CP1. *J Biol Chem* **265**: 13362-9.
- Kim K, Thu N, Saville B, Safe S (2003). Domains of estrogen receptor alpha (ERalpha) required for ERalpha/Sp1-mediated activation of GC-rich promoters by estrogens and antiestrogens in breast cancer cells. *Mol Endocrinol* **17**: 804-17.
- Kim TK,Roeder RG (1994). Proline-rich activator CTF1 targets the TFIIB assembly step during transcriptional activation. *Proc Natl Acad Sci U S A* **91**: 4170-4.
- Kingsley C,Winoto A (1992). Cloning of GT box-binding proteins: a novel Sp1 multigene family regulating T-cell receptor gene expression. *Mol Cell Biol* **12**: 4251-61.
- Klevit RE (1991). Recognition of DNA by Cys2,His2 zinc fingers. *Science* **253**: 1367, 1393.

- Knight GB, Gudas JM, Pardee AB (1987). Cell-cycle-specific interaction of nuclear DNA-binding proteins with a CCAAT sequence from the human thymidine kinase gene. *Proc Natl Acad Sci U S A* **84**: 8350-4.
- Kohn KW (1999). Molecular interaction map of the mammalian cell cycle control and DNA repair systems. *Mol Biol Cell* **10**: 2703-34.
- Koutsodontis G, Tentes I, Papakosta P, Moustakas A, Kardassis D (2001). Sp1 plays a critical role in the transcriptional activation of the human cyclin-dependent kinase inhibitor p21(WAF1/Cip1) gene by the p53 tumor suppressor protein. *J Biol Chem* **276**: 29116-25.
- Koutsodontis G, Vasilaki E, Chou WC, Papakosta P, Kardassis D (2005). Physical and functional interactions between members of the tumour suppressor p53 and the Sp families of transcription factors: importance for the regulation of genes involved in cell-cycle arrest and apoptosis. *Biochem J* **389**: 443-55.
- Kouzarides T, Ziff E (1988). The role of the leucine zipper in the fos-jun interaction. *Nature* **336**: 646-51.
- Kovesdi I, Reichel R, Nevins JR (1986). Identification of a cellular transcription factor involved in E1A trans-activation. *Cell* **45**: 219-28.
- Koziczak M, Muller H, Helin K, Nagamine Y (2001). E2F1-mediated transcriptional inhibition of the plasminogen activator inhibitor type 1 gene. *Eur J Biochem* **268**: 4969-78.
- Kriwacki RW, Schultz SC, Steitz TA, Caradonna JP (1992). Sequence-specific recognition of DNA by zinc-finger peptides derived from the transcription factor Sp1. *Proc Natl Acad Sci U S A* **89**: 9759-63.
- Kruger I, Vollmer M, Simmons DG, Elsasser HP, Philipsen S, Suske G (2007). Sp1/Sp3 compound heterozygous mice are not viable: impaired erythropoiesis and severe placental defects. *Dev Dyn* **236**: 2235-44.
- Kuo CT, Veselits ML, Barton KP, Lu MM, Clendenin C, Leiden JM (1997a). The LKLF transcription factor is required for normal tunica media formation and blood vessel stabilization during murine embryogenesis. *Genes Dev* **11**: 2996-3006.

Kuo CT, Veselits ML, Leiden JM (1997b). LKLF: A transcriptional regulator of single-positive T cell quiescence and survival. *Science* **277**: 1986-90.

Kuwahara J, Yonezawa A, Futamura M, Sugiura Y (1993). Binding of transcription factor Sp1 to GC box DNA revealed by footprinting analysis: different contact of three zinc fingers and sequence recognition mode. *Biochemistry* **32**: 5994-6001.

L

La Thangue NB (1994). DP and E2F proteins: components of a heterodimeric transcription factor implicated in cell cycle control. *Curr Opin Cell Biol* **6**: 443-50.

Lacroix I, Lipcey C, Imbert J, Kahn-Perles B (2002). Sp1 transcriptional activity is up-regulated by phosphatase 2A in dividing T lymphocytes. *J Biol Chem* **277**: 9598-605.

Lam EW, Robinson C, Watson RJ (1992). Characterization and cell cycle-regulated expression of mouse B-myb. *Oncogene* **7**: 1885-90.

Laniel MA, Poirier GG, Guerin SL (2001). Nuclear factor 1 interferes with Sp1 binding through a composite element on the rat poly(ADP-ribose) polymerase promoter to modulate its activity in vitro. *J Biol Chem* **276**: 20766-73.

Latchman DS (1990). Eukaryotic transcription factors. *Biochem J* **270**: 281-9.

Laub F, Dragomir C, Ramirez F (2006). Mice without transcription factor KLF7 provide new insight into olfactory bulb development. *Brain Res* **1103**: 108-13.

Laub F, Lei L, Sumiyoshi H, Kajimura D, Dragomir C, Smaldone S, et al. (2005). Transcription factor KLF7 is important for neuronal morphogenesis in selected regions of the nervous system. *Mol Cell Biol* **25**: 5699-711.

Lee HH, Chiang WH, Chiang SH, Liu YC, Hwang J, Ng SY (1995). Regulation of cyclin D1, DNA topoisomerase I, and proliferating cell nuclear antigen promoters during the cell cycle. *Gene Expr* **4**: 95-109.

Lee JA, Suh DC, Kang JE, Kim MH, Park H, Lee MN, et al. (2005). Transcriptional activity of Sp1 is regulated by molecular interactions between the zinc finger DNA binding domain and the inhibitory domain with corepressors, and this interaction is modulated by MEK. *J Biol Chem* **280**: 28061-71.

Lee W, Haslinger A, Karin M,Tjian R (1987a). Activation of transcription by two factors that bind promoter and enhancer sequences of the human metallothionein gene and SV40. *Nature* **325**: 368-72.

Lee W, Mitchell P,Tjian R (1987b). Purified transcription factor AP-1 interacts with TPA-inducible enhancer elements. *Cell* **49**: 741-52.

Lees E, Faha B, Dulic V, Reed SI,Harlow E (1992). Cyclin E/cdk2 and cyclin A/cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev* **6**: 1874-85.

Lees JA, Saito M, Vidal M, Valentine M, Look T, Harlow E, et al. (1993). The retinoblastoma protein binds to a family of E2F transcription factors. *Mol Cell Biol* **13**: 7813-25.

Leggett RW, Armstrong SA, Barry D,Mueller CR (1995). Sp1 is phosphorylated and its DNA binding activity down-regulated upon terminal differentiation of the liver. *J Biol Chem* **270**: 25879-84.

Levrero M, De Laurenzi V, Costanzo A, Gong J, Wang JY,Melino G (2000). The p53/p63/p73 family of transcription factors: overlapping and distinct functions. *J Cell Sci* **113 (Pt 10)**: 1661-70.

Li L, He S, Sun JM,Davie JR (2004). Gene regulation by Sp1 and Sp3. *Biochem Cell Biol* **82**: 460-71.

Li N, Seetharam S,Seetharam B (1998a). Characterization of the human transcobalamin II promoter. A proximal GC/GT box is a dominant negative element. *J Biol Chem* **273**: 16104-11.

Li Q, Herrler M, Landsberger N, Kaludov N, Ogryzko VV, Nakatani Y, et al. (1998b). Xenopus NF-Y pre-sets chromatin to potentiate p300 and acetylation-responsive transcription from the Xenopus hsp70 promoter in vivo. *Embo J* **17**: 6300-15.

Li Q, Imhof A, Collingwood TN, Urnov FD,Wolffe AP (1999). p300 stimulates transcription instigated by ligand-bound thyroid hormone receptor at a step subsequent to chromatin disruption. *Embo J* **18**: 5634-52.

BIBLIOGRAFIA

- Li XY, Hooft van Huijsduijnen R, Mantovani R, Benoist C, Mathis D (1992). Intron-exon organization of the NF-Y genes. Tissue-specific splicing modifies an activation domain. *J Biol Chem* **267**: 8984-90.
- Lin SY, Black AR, Kostic D, Pajovic S, Hoover CN, Azizkhan JC (1996). Cell cycle-regulated association of E2F1 and Sp1 is related to their functional interaction. *Mol Cell Biol* **16**: 1668-75.
- Liptay S, Schmid RM, Nabel EG, Nabel GJ (1994). Transcriptional regulation of NF-kappa B2: evidence for kappa B-mediated positive and negative autoregulation. *Mol Cell Biol* **14**: 7695-703.
- Lisztwan J, Marti A, Sutterluty H, Gstaiger M, Wirbelauer C, Krek W (1998). Association of human CUL-1 and ubiquitin-conjugating enzyme CDC34 with the F-box protein p45(SKP2): evidence for evolutionary conservation in the subunit composition of the CDC34-SCF pathway. *Embo J* **17**: 368-83.
- Liu A, Hoffman PW, Lu W, Bai G (2004). NF-kappaB site interacts with Sp factors and up-regulates the NR1 promoter during neuronal differentiation. *J Biol Chem* **279**: 17449-58.
- Loeffler S, Fayard B, Weis J, Weissenberger J (2005). Interleukin-6 induces transcriptional activation of vascular endothelial growth factor (VEGF) in astrocytes in vivo and regulates VEGF promoter activity in glioblastoma cells via direct interaction between STAT3 and Sp1. *Int J Cancer* **115**: 202-13.
- Look DC, Pelletier MR, Tidwell RM, Roswit WT, Holtzman MJ (1995). Stat1 depends on transcriptional synergy with Sp1. *J Biol Chem* **270**: 30264-7.
- Lou Z, Maher VM, McCormick JJ (2005). Identification of the promoter of human transcription factor Sp3 and evidence of the role of factors Sp1 and Sp3 in the expression of Sp3 protein. *Gene* **351**: 51-9.
- Luscher B, Eisenman RN (1990a). New light on Myc and Myb. Part I. Myc. *Genes Dev* **4**: 2025-35.
- Luscher B, Eisenman RN (1990b). New light on Myc and Myb. Part II. Myb. *Genes Dev* **4**: 2235-41.

M

Macleod KF, Sherry N, Hannon G, Beach D, Tokino T, Kinzler K, et al. (1995). p53-dependent and independent expression of p21 during cell growth, differentiation, and DNA damage. *Genes Dev* **9**: 935-44.

Maity SN, de Crombrugghe B (1992). Biochemical analysis of the B subunit of the heteromeric CCAAT-binding factor. A DNA-binding domain and a subunit interaction domain are specified by two separate segments. *J Biol Chem* **267**: 8286-92.

Maity SN, de Crombrugghe B (1998). Role of the CCAAT-binding protein CBF/NF-Y in transcription. *Trends Biochem Sci* **23**: 174-8.

Majello B, De Luca P, Lania L (1997). Sp3 is a bifunctional transcription regulator with modular independent activation and repression domains. *J Biol Chem* **272**: 4021-6.

Maldonado E, Shiekhattar R, Sheldon M, Cho H, Drapkin R, Rickert P, et al. (1996). A human RNA polymerase II complex associated with SRB and DNA-repair proteins. *Nature* **381**: 86-9.

Malumbres M, Barbacid M (2005). Mammalian cyclin-dependent kinases. *Trends Biochem Sci* **30**: 630-41.

Manni I, Mazzaro G, Gurtner A, Mantovani R, Haugwitz U, Krause K, et al. (2001). NF-Y mediates the transcriptional inhibition of the cyclin B1, cyclin B2, and cdc25C promoters upon induced G2 arrest. *J Biol Chem* **276**: 5570-6.

Mantovani R (1999). The molecular biology of the CCAAT-binding factor NF-Y. *Gene* **239**: 15-27.

Mantovani R, Li XY, Pessara U, Hooft van Huisdijnen R, Benoist C, Mathis D (1994). Dominant negative analogs of NF-YA. *J Biol Chem* **269**: 20340-6.

Mantovani R, Pessara U, Tronche F, Li XY, Knapp AM, Pasquali JL, et al. (1992). Monoclonal antibodies to NF-Y define its function in MHC class II and albumin gene transcription. *Embo J* **11**: 3315-22.

Marco E, Garcia-Nieto R, Gago F (2003). Assessment by molecular dynamics simulations of the structural determinants of DNA-binding specificity for transcription factor Sp1. *J Mol Biol* **328**: 9-32.

BIBLIOGRAFIA

- Marin M, Karis A, Visser P, Grosveld F, Philipsen S (1997). Transcription factor Sp1 is essential for early embryonic development but dispensable for cell growth and differentiation. *Cell* **89**: 619-28.
- Marin MC, Kaelin WG, Jr. (2000). p63 and p73: old members of a new family. *Biochim Biophys Acta* **1470**: M93-M100.
- Marmorstein LY, Ouchi T, Aaronson SA (1998). The BRCA2 gene product functionally interacts with p53 and RAD51. *Proc Natl Acad Sci U S A* **95**: 13869-74.
- Marti A, Wirbelauer C, Scheffner M, Krek W (1999). Interaction between ubiquitin-protein ligase SCFSKP2 and E2F-1 underlies the regulation of E2F-1 degradation. *Nat Cell Biol* **1**: 14-9.
- Martin K, Trouche D, Hagemeier C, Kouzarides T (1995). Regulation of transcription by E2F1/DP1. *J Cell Sci Suppl* **19**: 91-4.
- Matsumoto N, Kubo A, Liu H, Akita K, Laub F, Ramirez F, et al. (2006). Developmental regulation of yolk sac hematopoiesis by Kruppel-like factor 6. *Blood* **107**: 1357-65.
- Matsumoto N, Laub F, Aldabe R, Zhang W, Ramirez F, Yoshida T, et al. (1998). Cloning the cDNA for a new human zinc finger protein defines a group of closely related Kruppel-like transcription factors. *J Biol Chem* **273**: 28229-37.
- Meisterernst M, Gander I, Rogge L, Winnacker EL (1988). A quantitative analysis of nuclear factor I/DNA interactions. *Nucleic Acids Res* **16**: 4419-35.
- Merchant JL, Du M, Todisco A (1999). Sp1 phosphorylation by Erk 2 stimulates DNA binding. *Biochem Biophys Res Commun* **254**: 454-61.
- Meyer TE, Waeber G, Lin J, Beckmann W, Habener JF (1993). The promoter of the gene encoding 3',5'-cyclic adenosine monophosphate (cAMP) response element binding protein contains cAMP response elements: evidence for positive autoregulation of gene transcription. *Endocrinology* **132**: 770-80.
- Michel JJ, Xiong Y (1998). Human CUL-1, but not other cullin family members, selectively interacts with SKP1 to form a complex with SKP2 and cyclin A. *Cell Growth Differ* **9**: 435-49.
- Mighell AJ, Smith NR, Robinson PA, Markham AF (2000). Vertebrate pseudogenes. *FEBS Lett* **468**: 109-14.

Milanini J, Vinals F, Pouyssegur J, Pages G (1998). p42/p44 MAP kinase module plays a key role in the transcriptional regulation of the vascular endothelial growth factor gene in fibroblasts. *J Biol Chem* **273**: 18165-72.

Miller IJ, Bieker JJ (1993). A novel, erythroid cell-specific murine transcription factor that binds to the CACCC element and is related to the Kruppel family of nuclear proteins. *Mol Cell Biol* **13**: 2776-86.

Milner J, Ponder B, Hughes-Davies L, Seltmann M, Kouzarides T (1997). Transcriptional activation functions in BRCA2. *Nature* **386**: 772-3.

Mongay L, Plaza S, Vigorito E, Serra-Pages C, Vives J (2001). Association of the cell cycle regulatory proteins p45(SKP2) and CksHs1. Functional effect on CDK2 complex formation and kinase activity. *J Biol Chem* **276**: 25030-6.

Monteiro AN, August A, Hanafusa H (1996). Evidence for a transcriptional activation function of BRCA1 C-terminal region. *Proc Natl Acad Sci U S A* **93**: 13595-9.

Moran KM, Crusio RH, Chan CH, Grekova MC, Richert JR (2004). Human transcription factor Sp3: genomic structure, identification of a processed pseudogene, and transcript analysis. *Gene* **341**: 235-47.

Morita M, Kobayashi A, Yamashita T, Shimanuki T, Nakajima O, Takahashi S, et al. (2003). Functional analysis of basic transcription element binding protein by gene targeting technology. *Mol Cell Biol* **23**: 2489-500.

Mucenski ML, McLain K, Kier AB, Swerdlow SH, Schreiner CM, Miller TA, et al. (1991). A functional c-myb gene is required for normal murine fetal hepatic hematopoiesis. *Cell* **65**: 677-89.

Murata Y, Kim HG, Rogers KT, Udvadia AJ, Horowitz JM (1994). Negative regulation of Sp1 trans-activation is correlated with the binding of cellular proteins to the amino terminus of the Sp1 trans-activation domain. *J Biol Chem* **269**: 20674-81.

N

Nagata K, Guggenheimer RA, Enomoto T, Lichy JH, Hurwitz J (1982). Adenovirus DNA replication in vitro: identification of a host factor that stimulates synthesis of the preterminal protein-dCMP complex. *Proc Natl Acad Sci U S A* **79**: 6438-42.

BIBLIOGRAFIA

- Nahle Z, Polakoff J, Davuluri RV, McCurrach ME, Jacobson MD, Narita M, et al. (2002). Direct coupling of the cell cycle and cell death machinery by E2F. *Nat Cell Biol* **4**: 859-64.
- Nakashima K, Zhou X, Kunkel G, Zhang Z, Deng JM, Behringer RR, et al. (2002). The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* **108**: 17-29.
- Nehls MC, Rippe RA, Veloz L, Brenner DA (1991). Transcription factors nuclear factor I and Sp1 interact with the murine collagen alpha 1 (I) promoter. *Mol Cell Biol* **11**: 4065-73.
- Ness SA (1999). Myb binding proteins: regulators and cohorts in transformation. *Oncogene* **18**: 3039-46.
- Nguyen-Tran VT, Kubalak SW, Minamisawa S, Fiset C, Wollert KC, Brown AB, et al. (2000). A novel genetic pathway for sudden cardiac death via defects in the transition between ventricular and conduction system cell lineages. *Cell* **102**: 671-82.
- Nicolas M, Noe V, Ciudad CJ (2003). Transcriptional regulation of the human Sp1 gene promoter by the specificity protein (Sp) family members nuclear factor Y (NF-Y) and E2F. *Biochem J* **371**: 265-75.
- Nicolas M, Noe V, Jensen KB, Ciudad CJ (2001). Cloning and characterization of the 5'-flanking region of the human transcription factor Sp1 gene. *J Biol Chem* **276**: 22126-32.
- Nilson DG, Sabatino DE, Bodine DM, Gallagher PG (2006). Major erythrocyte membrane protein genes in EKLF-deficient mice. *Exp Hematol* **34**: 705-12.
- Nishinaka T, Fu YH, Chen LI, Yokoyama K, Chiu R (1997). A unique cathepsin-like protease isolated from CV-1 cells is involved in rapid degradation of retinoblastoma susceptibility gene product, RB, and transcription factor SP1. *Biochim Biophys Acta* **1351**: 274-86.
- Noe V, Alemany C, Chasin LA, Ciudad CJ (1998). Retinoblastoma protein associates with SP1 and activates the hamster dihydrofolate reductase promoter. *Oncogene* **16**: 1931-8.

Noe V, Alemany C, Nicolas M,Ciudad CJ (2001). Sp1 involvement in the 4beta-phorbol 12-myristate 13-acetate (TPA)-mediated increase in resistance to methotrexate in Chinese hamster ovary cells. *Eur J Biochem* **268**: 3163-73.

Noe V, Chen C, Alemany C, Nicolas M, Caragol I, Chasin LA, et al. (1997). Cell-growth regulation of the hamster dihydrofolate reductase gene promoter by transcription factor Sp1. *Eur J Biochem* **249**: 13-20.

Nomura N, Takahashi M, Matsui M, Ishii S, Date T, Sasamoto S, et al. (1988). Isolation of human cDNA clones of myb-related genes, A-myb and B-myb. *Nucleic Acids Res* **16**: 11075-89.

Nuez B, Michalovich D, Bygrave A, Ploemacher R,Grosveld F (1995). Defective haematopoiesis in fetal liver resulting from inactivation of the EKLF gene. *Nature* **375**: 316-8.

O

Oelgeschlager M, Janknecht R, Krieg J, Schreek S,Luscher B (1996). Interaction of the co-activator CBP with Myb proteins: effects on Myb-specific transactivation and on the cooperativity with NF-M. *Embo J* **15**: 2771-80.

Ogra Y, Suzuki K, Gong P, Otsuka F,Koizumi S (2001). Negative regulatory role of Sp1 in metal responsive element-mediated transcriptional activation. *J Biol Chem* **276**: 16534-9.

Oh IH,Reddy EP (1999). The myb gene family in cell growth, differentiation and apoptosis. *Oncogene* **18**: 3017-33.

Oh JE, Han JA,Hwang ES (2007). Downregulation of transcription factor, Sp1, during cellular senescence. *Biochem Biophys Res Commun* **353**: 86-91.

Oleksiak MF,Crawford DL (2002). 5' genomic structure of human Sp3. *Mol Biol Evol* **19**: 2026-9.

Opitz OG,Rustgi AK (2000). Interaction between Sp1 and cell cycle regulatory proteins is important in transactivation of a differentiation-related gene. *Cancer Res* **60**: 2825-30.

Oswald F, Dobner T,Lipp M (1996). The E2F transcription factor activates a replication-dependent human H2A gene in early S phase of the cell cycle. *Mol Cell Biol* **16**: 1889-95.

P

Pages G (2007). Sp3-mediated VEGF regulation is dependent on phosphorylation by extra-cellular signals regulated kinases (Erk). *J Cell Physiol* **213**: 454-63.

Pallas DC, Shahrik LK, Martin BL, Jaspers S, Miller TB, Brautigan DL, et al. (1990). Polyoma small and middle T antigens and SV40 small t antigen form stable complexes with protein phosphatase 2A. *Cell* **60**: 167-76.

Pan MR,Hung WC (2002). Nonsteroidal anti-inflammatory drugs inhibit matrix metalloproteinase-2 via suppression of the ERK/Sp1-mediated transcription. *J Biol Chem* **277**: 32775-80.

Pascal E,Tjian R (1991). Different activation domains of Sp1 govern formation of multimers and mediate transcriptional synergism. *Genes Dev* **5**: 1646-56.

Patton EE, Willems AR,Tyers M (1998). Combinatorial control in ubiquitin-dependent proteolysis: don't Skp the F-box hypothesis. *Trends Genet* **14**: 236-43.

Pavletich NP,Pabo CO (1991). Zinc finger-DNA recognition: crystal structure of a Zif268-DNA complex at 2.1 A. *Science* **252**: 809-17.

Pearson BE, Nasheuer HP,Wang TS (1991). Human DNA polymerase alpha gene: sequences controlling expression in cycling and serum-stimulated cells. *Mol Cell Biol* **11**: 2081-95.

Penuelas S, Alemany C, Noe V,Ciudad CJ (2003). The expression of retinoblastoma and Sp1 is increased by low concentrations of cyclin-dependent kinase inhibitors. *Eur J Biochem* **270**: 4809-22.

Perkins AC, Sharpe AH,Orkin SH (1995). Lethal beta-thalassaemia in mice lacking the erythroid CACCC-transcription factor EKLF. *Nature* **375**: 318-22.

Perkins ND (2004). Regulation of NF-kappaB by atypical activators and tumour suppressors. *Biochem Soc Trans* **32**: 936-9.

Persengiev SP, Saffer JD, Kilpatrick DL (1995). An alternatively spliced form of the transcription factor Sp1 containing only a single glutamine-rich transactivation domain. *Proc Natl Acad Sci U S A* **92**: 9107-11.

Philipsen S, Suske G (1999). A tale of three fingers: the family of mammalian Sp/XKLF transcription factors. *Nucleic Acids Res* **27**: 2991-3000.

Pines J, Hunter T (1990). Human cyclin A is adenovirus E1A-associated protein p60 and behaves differently from cyclin B. *Nature* **346**: 760-3.

Poole JC, Thain A, Perkins ND, Roninson IB (2004). Induction of transcription by p21Waf1/Cip1/Sdi1: role of NFκB and effect of non-steroidal anti-inflammatory drugs. *Cell Cycle* **3**: 931-40.

Pugh BF, Tjian R (1990). Mechanism of transcriptional activation by Sp1: evidence for coactivators. *Cell* **61**: 1187-97.

R

Rafty LA, Khachigian LM (2001). Sp1 phosphorylation regulates inducible expression of platelet-derived growth factor B-chain gene via atypical protein kinase C-zeta. *Nucleic Acids Res* **29**: 1027-33.

Rafty LA, Santiago FS, Khachigian LM (2002). NF1/X represses PDGF A-chain transcription by interacting with Sp1 and antagonizing Sp1 occupancy of the promoter. *Embo J* **21**: 334-43.

Ravasi T, Huber T, Zavolan M, Forrest A, Gaasterland T, Grimmond S, et al. (2003). Systematic characterization of the zinc-finger-containing proteins in the mouse transcriptome. *Genome Res* **13**: 1430-42.

Reisinger K, Kaufmann R, Gille J (2003). Increased Sp1 phosphorylation as a mechanism of hepatocyte growth factor (HGF/SF)-induced vascular endothelial growth factor (VEGF/VPF) transcription. *J Cell Sci* **116**: 225-38.

Rickers A, Peters N, Badock V, Beyaert R, Vandenebeele P, Dorken B, et al. (1999). Cleavage of transcription factor SP1 by caspases during anti-IgM-induced B-cell apoptosis. *Eur J Biochem* **261**: 269-74.

BIBLIOGRAFIA

- Ristiniemi J,Oikarinen J (1989). Histone H1 binds to the putative nuclear factor I recognition sequence in the mouse alpha 2(I) collagen promoter. *J Biol Chem* **264**: 2164-74.
- Robinson C, Light Y, Groves R, Mann D, Marias R,Watson R (1996). Cell-cycle regulation of B-Myb protein expression: specific phosphorylation during the S phase of the cell cycle. *Oncogene* **12**: 1855-64.
- Roder K, Wolf SS, Beck KF,Schweizer M (1997). Cooperative binding of NF-Y and Sp1 at the DNase I-hypersensitive site, fatty acid synthase insulin-responsive element 1, located at -500 in the rat fatty acid synthase promoter. *J Biol Chem* **272**: 21616-24.
- Rohlf C, Ahmad S, Borellini F, Lei J,Glazer RI (1997). Modulation of transcription factor Sp1 by cAMP-dependent protein kinase. *J Biol Chem* **272**: 21137-41.
- Romier C, Cocchiarella F, Mantovani R,Moras D (2003). The NF-YB/NF-YC structure gives insight into DNA binding and transcription regulation by CCAAT factor NF-Y. *J Biol Chem* **278**: 1336-45.
- Roninson IB (2002). Oncogenic functions of tumour suppressor p21(Waf1/Cip1/Sdi1): association with cell senescence and tumour-promoting activities of stromal fibroblasts. *Cancer Lett* **179**: 1-14.
- Ross S, Best JL, Zon LI,Gill G (2002). SUMO-1 modification represses Sp3 transcriptional activation and modulates its subnuclear localization. *Mol Cell* **10**: 831-42.
- Rotheneder H, Gejmayer S,Haidweger E (1999). Transcription factors of the Sp1 family: interaction with E2F and regulation of the murine thymidine kinase promoter. *J Mol Biol* **293**: 1005-15.
- Ryseck RP,Bravo R (1991). c-JUN, JUN B, and JUN D differ in their binding affinities to AP-1 and CRE consensus sequences: effect of FOS proteins. *Oncogene* **6**: 533-42.
- Ryu H, Lee J, Olofsson BA, Mwidau A, Dedeoglu A, Escudero M, et al. (2003a). Histone deacetylase inhibitors prevent oxidative neuronal death independent of expanded polyglutamine repeats via an Sp1-dependent pathway. *Proc Natl Acad Sci U S A* **100**: 4281-6.

Ryu H, Lee J, Zaman K, Kubilis J, Ferrante RJ, Ross BD, et al. (2003b). Sp1 and Sp3 are oxidative stress-inducible, antideath transcription factors in cortical neurons. *J Neurosci* **23**: 3597-606.

Ryu S, Tjian R (1999). Purification of transcription cofactor complex CRSP. *Proc Natl Acad Sci U S A* **96**: 7137-42.

Ryu S, Zhou S, Ladurner AG, Tjian R (1999). The transcriptional cofactor complex CRSP is required for activity of the enhancer-binding protein Sp1. *Nature* **397**: 446-50.

S

Saegusa N, Yokono M, Matsushita K, Sugiura Y (1997). Different contributions of three zinc fingers of transcription factor Sp1 to DNA recognition: novel binding mode of N-terminal finger 1. *Nucleic Acids Symp Ser* 151-2.

Safe S, Abdelrahim M (2005). Sp transcription factor family and its role in cancer. *Eur J Cancer* **41**: 2438-48.

Saffer JD, Jackson SP, Annarella MB (1991). Developmental expression of Sp1 in the mouse. *Mol Cell Biol* **11**: 2189-99.

Sala A (2005). B-MYB, a transcription factor implicated in regulating cell cycle, apoptosis and cancer. *Eur J Cancer* **41**: 2479-84.

Sala A, Kundu M, Casella I, Engelhard A, Calabretta B, Grasso L, et al. (1997). Activation of human B-MYB by cyclins. *Proc Natl Acad Sci U S A* **94**: 532-6.

Sala A, Saitta B, De Luca P, Cervellera MN, Casella I, Lewis RE, et al. (1999). B-MYB transactivates its own promoter through SP1-binding sites. *Oncogene* **18**: 1333-9.

Sala A, Watson R (1999). B-Myb protein in cellular proliferation, transcription control, and cancer: latest developments. *J Cell Physiol* **179**: 245-50.

Salimath B, Marme D, Finkenzeller G (2000). Expression of the vascular endothelial growth factor gene is inhibited by p73. *Oncogene* **19**: 3470-6.

Salomoni P, Perrotti D, Martinez R, Franceschi C, Calabretta B (1997). Resistance to apoptosis in CTLL-2 cells constitutively expressing c-Myb is associated with induction

BIBLIOGRAFIA

- of BCL-2 expression and Myb-dependent regulation of bcl-2 promoter activity. *Proc Natl Acad Sci U S A* **94**: 3296-301.
- Santpere G, Nieto M, Puig B,Ferrer I (2006). Abnormal Sp1 transcription factor expression in Alzheimer disease and tauopathies. *Neurosci Lett* **397**: 30-4.
- Sapetschnig A, Koch F, Rischitor G, Mennenga T,Suske G (2004). Complexity of translationally controlled transcription factor Sp3 isoform expression. *J Biol Chem* **279**: 42095-105.
- Schafer D, Hamm-Kunzelmann B,Brand K (1997). Glucose regulates the promoter activity of aldolase A and pyruvate kinase M2 via dephosphorylation of Sp1. *FEBS Lett* **417**: 325-8.
- Schatt MD, Rusconi S,Schaffner W (1990). A single DNA-binding transcription factor is sufficient for activation from a distant enhancer and/or from a promoter position. *Embo J* **9**: 481-7.
- Schreiber M, Kolbus A, Piu F, Szabowski A, Mohle-Steinlein U, Tian J, et al. (1999). Control of cell cycle progression by c-Jun is p53 dependent. *Genes Dev* **13**: 607-19.
- Schulze A, Zerfass-Thome K, Berges J, Middendorp S, Jansen-Durr P,Henglein B (1996). Anchorage-dependent transcription of the cyclin A gene. *Mol Cell Biol* **16**: 4632-8.
- Schutte J, Viallet J, Nau M, Segal S, Fedorko J,Minna J (1989). jun-B inhibits and c-fos stimulates the transforming and trans-activating activities of c-jun. *Cell* **59**: 987-97.
- Schuur ER, Kruse U, Iacovoni JS,Vogt PK (1995). Nuclear factor I interferes with transformation induced by nuclear oncogenes. *Cell Growth Differ* **6**: 219-27.
- Scohy S, Gabant P, Van Reeth T, Hertveldt V, Dreze PL, Van Vooren P, et al. (2000). Identification of KLF13 and KLF14 (SP6), novel members of the SP/XKLF transcription factor family. *Genomics* **70**: 93-101.
- Scully R, Chen J, Ochs RL, Keegan K, Hoekstra M, Feunteun J, et al. (1997). Dynamic changes of BRCA1 subnuclear location and phosphorylation state are initiated by DNA damage. *Cell* **90**: 425-35.
- Segre JA, Bauer C,Fuchs E (1999). Klf4 is a transcription factor required for establishing the barrier function of the skin. *Nat Genet* **22**: 356-60.

- Selga E, Noe V,Ciudad CJ (2008). Transcriptional regulation of aldo-keto reductase 1C1 in HT29 human colon cancer cells resistant to methotrexate: role in the cell cycle and apoptosis. *Biochem Pharmacol* **75**: 414-26.
- Shan B,Lee WH (1994). Deregulated expression of E2F-1 induces S-phase entry and leads to apoptosis. *Mol Cell Biol* **14**: 8166-73.
- Shaulian E,Karin M (2001). AP-1 in cell proliferation and survival. *Oncogene* **20**: 2390-400.
- Shaulian E,Karin M (2002). AP-1 as a regulator of cell life and death. *Nat Cell Biol* **4**: E131-6.
- Sheikh MS, Li XS, Chen JC, Shao ZM, Ordonez JV,Fontana JA (1994). Mechanisms of regulation of WAF1/Cip1 gene expression in human breast carcinoma: role of p53-dependent and independent signal transduction pathways. *Oncogene* **9**: 3407-15.
- Shi Y,Berg JM (1995). A direct comparison of the properties of natural and designed zinc-finger proteins. *Chem Biol* **2**: 83-9.
- Shields JM,Yang VW (1998). Identification of the DNA sequence that interacts with the gut-enriched Kruppel-like factor. *Nucleic Acids Res* **26**: 796-802.
- Shin DH, Lee HW, Jeon GS, Lee HY, Lee KH,Cho SS (2001). Constitutive expression of c-myb mRNA in the adult rat brain. *Brain Res* **892**: 203-7.
- Shindo T, Manabe I, Fukushima Y, Tobe K, Aizawa K, Miyamoto S, et al. (2002). Kruppel-like zinc-finger transcription factor KLF5/BTEB2 is a target for angiotensin II signaling and an essential regulator of cardiovascular remodeling. *Nat Med* **8**: 856-63.
- Shirodkar S, Ewen M, DeCaprio JA, Morgan J, Livingston DM,Chittenden T (1992). The transcription factor E2F interacts with the retinoblastoma product and a p107-cyclin A complex in a cell cycle-regulated manner. *Cell* **68**: 157-66.
- Shou Y, Baron S,Poncz M (1998). An Sp1-binding silencer element is a critical negative regulator of the megakaryocyte-specific alphallb gene. *J Biol Chem* **273**: 5716-26.
- Simmen RC, Eason RR, McQuown JR, Linz AL, Kang TJ, Chatman L, Jr., et al. (2004). Subfertility, uterine hypoplasia, and partial progesterone resistance in mice lacking the

BIBLIOGRAFIA

- Kruppel-like factor 9/basic transcription element-binding protein-1 (Bteb1) gene. *J Biol Chem* **279**: 29286-94.
- Sinha S, Maity SN, Lu J, de Crombrugghe B (1995). Recombinant rat CBF-C, the third subunit of CBF/NFY, allows formation of a protein-DNA complex with CBF-A and CBF-B and with yeast HAP2 and HAP3. *Proc Natl Acad Sci U S A* **92**: 1624-8.
- Slansky JE, Farnham PJ (1996). Transcriptional regulation of the dihydrofolate reductase gene. *Bioessays* **18**: 55-62.
- Slansky JE, Li Y, Kaelin WG, Farnham PJ (1993). A protein synthesis-dependent increase in E2F1 mRNA correlates with growth regulation of the dihydrofolate reductase promoter. *Mol Cell Biol* **13**: 1610-8.
- Smale ST, Schmidt MC, Berk AJ, Baltimore D (1990). Transcriptional activation by Sp1 as directed through TATA or initiator: specific requirement for mammalian transcription factor IID. *Proc Natl Acad Sci U S A* **87**: 4509-13.
- Song CZ, Gavrilidis G, Asano H, Stamatoyannopoulos G (2005). Functional study of transcription factor KLF11 by targeted gene inactivation. *Blood Cells Mol Dis* **34**: 53-9.
- Song CZ, Keller K, Chen Y, Stamatoyannopoulos G (2003). Functional interplay between CBP and PCAF in acetylation and regulation of transcription factor KLF13 activity. *J Mol Biol* **329**: 207-15.
- Sonnenberg JL, Rauscher FJ, 3rd, Morgan JI, Curran T (1989). Regulation of proenkephalin by Fos and Jun. *Science* **246**: 1622-5.
- Sowa Y, Orita T, Minamikawa-Hiranabe S, Mizuno T, Nomura H, Sakai T (1999). Sp3, but not Sp1, mediates the transcriptional activation of the p21/WAF1/Cip1 gene promoter by histone deacetylase inhibitor. *Cancer Res* **59**: 4266-70.
- Spencer VA, Davie JR (1999). Role of covalent modifications of histones in regulating gene expression. *Gene* **240**: 1-12.
- Spengler ML, Brattain MG (2006). Sumoylation inhibits cleavage of Sp1 N-terminal negative regulatory domain and inhibits Sp1-dependent transcription. *J Biol Chem* **281**: 5567-74.
- Stark LA, Dunlop MG (2005). Nucleolar sequestration of RelA (p65) regulates NF- κ B-driven transcription and apoptosis. *Mol Cell Biol* **25**: 5985-6004.

- Steele-Perkins G, Butz KG, Lyons GE, Zeichner-David M, Kim HJ, Cho MI, et al. (2003). Essential role for NFI-C/CTF transcription-replication factor in tooth root development. *Mol Cell Biol* **23**: 1075-84.
- Su K, Roos MD, Yang X, Han I, Paterson AJ, Kudlow JE (1999). An N-terminal region of Sp1 targets its proteasome-dependent degradation in vitro. *J Biol Chem* **274**: 15194-202.
- Suardet L, Li C, Little JB (1996). Radio-induced modulation of transforming growth factor beta1 sensitivity in a p53 wild-type human colorectal-cancer cell line. *Int J Cancer* **68**: 126-31.
- Subramaniam M, Gorny G, Johnsen SA, Monroe DG, Evans GL, Fraser DG, et al. (2005). TIEG1 null mouse-derived osteoblasts are defective in mineralization and in support of osteoclast differentiation in vitro. *Mol Cell Biol* **25**: 1191-9.
- Sun JM, Chen HY, Moniwa M, Litchfield DW, Seto E, Davie JR (2002). The transcriptional repressor Sp3 is associated with CK2-phosphorylated histone deacetylase 2. *J Biol Chem* **277**: 35783-6.
- Sun P, Dong P, Dai K, Hannon GJ, Beach D (1998). p53-independent role of MDM2 in TGF-beta1 resistance. *Science* **282**: 2270-2.
- Sun X, Wang Y, Qing H, Christensen MA, Liu Y, Zhou W, et al. (2005). Distinct transcriptional regulation and function of the human BACE2 and BACE1 genes. *Faseb J* **19**: 739-49.
- Supp DM, Witte DP, Branford WW, Smith EP, Potter SS (1996). Sp4, a member of the Sp1-family of zinc finger transcription factors, is required for normal murine growth, viability, and male fertility. *Dev Biol* **176**: 284-99.
- Suske G (1999). The Sp-family of transcription factors. *Gene* **238**: 291-300.
- Suske G, Bruford E, Philipsen S (2005). Mammalian SP/KLF transcription factors: bring in the family. *Genomics* **85**: 551-6.
- Sutterluty H, Chatelain E, Marti A, Wirbelauer C, Senften M, Muller U, et al. (1999). p45SKP2 promotes p27Kip1 degradation and induces S phase in quiescent cells. *Nat Cell Biol* **1**: 207-14.

Suzuki T, Muto S, Miyamoto S, Aizawa K, Horikoshi M, Nagai R (2003). Functional interaction of the DNA-binding transcription factor Sp1 through its DNA-binding domain with the histone chaperone TAF-I. *J Biol Chem* **278**: 28758-64.

Suzuki Y, Shimada J, Shudo K, Matsumura M, Crippa MP, Kojima S (1999). Physical interaction between retinoic acid receptor and Sp1: mechanism for induction of urokinase by retinoic acid. *Blood* **93**: 4264-76.

Suzuki Y, Tsunoda T, Sese J, Taira H, Mizushima-Sugano J, Hata H, et al. (2001). Identification and characterization of the potential promoter regions of 1031 kinds of human genes. *Genome Res* **11**: 677-84.

T

Taatjes DJ, Naar AM, Andel F, 3rd, Nogales E, Tjian R (2002). Structure, function, and activator-induced conformations of the CRSP coactivator. *Science* **295**: 1058-62.

Taatjes DJ, Tjian R (2004). Structure and function of CRSP/Med2; a promoter-selective transcriptional coactivator complex. *Mol Cell* **14**: 675-83.

Takahara T, Kanazu SI, Yanagisawa S, Akanuma H (2000). Heterogeneous Sp1 mRNAs in human HepG2 cells include a product of homotypic trans-splicing. *J Biol Chem* **275**: 38067-72.

Takahara T, Kasahara D, Mori D, Yanagisawa S, Akanuma H (2002). The trans-spliced variants of Sp1 mRNA in rat. *Biochem Biophys Res Commun* **298**: 156-62.

Tanaka Y, Patestos NP, Maekawa T, Ishii S (1999). B-myb is required for inner cell mass formation at an early stage of development. *J Biol Chem* **274**: 28067-70.

Tanese N, Pugh BF, Tjian R (1991). Coactivators for a proline-rich activator purified from the multisubunit human TFIID complex. *Genes Dev* **5**: 2212-24.

Tanno B, Negroni A, Vitali R, Pirozzoli MC, Cesi V, Mancini C, et al. (2002). Expression of insulin-like growth factor-binding protein 5 in neuroblastoma cells is regulated at the transcriptional level by c-Myb and B-Myb via direct and indirect mechanisms. *J Biol Chem* **277**: 23172-80.

- Tashiro S, Takemoto Y, Handa H, Ishii S (1995). Cell type-specific trans-activation by the B-myb gene product: requirement of the putative cofactor binding to the C-terminal conserved domain. *Oncogene* **10**: 1699-707.
- Thiesen HJ, Bach C (1990). Target Detection Assay (TDA): a versatile procedure to determine DNA binding sites as demonstrated on SP1 protein. *Nucleic Acids Res* **18**: 3203-9.
- Thomas JE, Smith M, Tonkinson JL, Rubinfeld B, Polakis P (1997). Induction of phosphorylation on BRCA1 during the cell cycle and after DNA damage. *Cell Growth Differ* **8**: 801-9.
- Thomas K, Wu J, Sung DY, Thompson W, Powell M, McCarrey J, et al. (2007). SP1 transcription factors in male germ cell development and differentiation. *Mol Cell Endocrinol* **270**: 1-7.
- Thomas NS, Pizzey AR, Tiwari S, Williams CD, Yang J (1998). p130, p107, and pRb are differentially regulated in proliferating cells and during cell cycle arrest by alpha-interferon. *J Biol Chem* **273**: 23659-67.
- Thoms HC, Dunlop MG, Stark LA (2007a). CDK4 inhibitors and apoptosis: a novel mechanism requiring nucleolar targeting of RelA. *Cell Cycle* **6**: 1293-7.
- Thoms HC, Dunlop MG, Stark LA (2007b). p38-mediated inactivation of cyclin D1/cyclin-dependent kinase 4 stimulates nucleolar translocation of RelA and apoptosis in colorectal cancer cells. *Cancer Res* **67**: 1660-9.
- Tjian R, Maniatis T (1994). Transcriptional activation: a complex puzzle with few easy pieces. *Cell* **77**: 5-8.
- Tommasi S, Pfeifer GP (1995). In vivo structure of the human cdc2 promoter: release of a p130-E2F-4 complex from sequences immediately upstream of the transcription initiation site coincides with induction of cdc2 expression. *Mol Cell Biol* **15**: 6901-13.
- Toscani A, Mettus RV, Coupland R, Simpkins H, Litvin J, Orth J, et al. (1997). Arrest of spermatogenesis and defective breast development in mice lacking A-myb. *Nature* **386**: 713-7.

BIBLIOGRAFIA

Townsend PA, Cragg MS, Davidson SM, McCormick J, Barry S, Lawrence KM, et al. (2005). STAT-1 facilitates the ATM activated checkpoint pathway following DNA damage. *J Cell Sci* **118**: 1629-39.

Trauth K, Mutschler B, Jenkins NA, Gilbert DJ, Copeland NG, Klempnauer KH (1994). Mouse A-myb encodes a trans-activator and is expressed in mitotically active cells of the developing central nervous system, adult testis and B lymphocytes. *Embo J* **13**: 5994-6005.

Treichel D, Schock F, Jackle H, Gruss P, Mansouri A (2003). mBtd is required to maintain signaling during murine limb development. *Genes Dev* **17**: 2630-5.

Tsai LH, Harlow E, Meyerson M (1991). Isolation of the human cdk2 gene that encodes the cyclin A- and adenovirus E1A-associated p33 kinase. *Nature* **353**: 174-7.

Turner J, Crossley M (1999). Mammalian Kruppel-like transcription factors: more than just a pretty finger. *Trends Biochem Sci* **24**: 236-40.

U

Udvadia AJ, Templeton DJ, Horowitz JM (1995). Functional interactions between the retinoblastoma (Rb) protein and Sp-family members: superactivation by Rb requires amino acids necessary for growth suppression. *Proc Natl Acad Sci U S A* **92**: 3953-7.

V

Vallian S, Chin KV, Chang KS (1998). The promyelocytic leukemia protein interacts with Sp1 and inhibits its transactivation of the epidermal growth factor receptor promoter. *Mol Cell Biol* **18**: 7147-56.

Van Loo PF, Bouwman P, Ling KW, Middendorp S, Suske G, Grosveld F, et al. (2003). Impaired hematopoiesis in mice lacking the transcription factor Sp3. *Blood* **102**: 858-66.

van Loo PF, Mahtab EA, Wisse LJ, Hou J, Grosveld F, Suske G, et al. (2007). Transcription factor Sp3 knockout mice display serious cardiac malformations. *Mol Cell Biol* **27**: 8571-82.

van Vliet J, Crofts LA, Quinlan KG, Czolij R, Perkins AC,Crossley M (2006). Human KLF17 is a new member of the Sp/KLF family of transcription factors. *Genomics* **87**: 474-82.

van Wijnen AJ, Massung RF, Stein JL,Stein GS (1988). Human H1 histone gene promoter CCAAT box binding protein HiNF-B is a mosaic factor. *Biochemistry* **27**: 6534-41.

Verger A, Perdomo J,Crossley M (2003). Modification with SUMO. A role in transcriptional regulation. *EMBO Rep* **4**: 137-42.

Vinals F, Fandos C, Santalucia T, Ferre J, Testar X, Palacin M, et al. (1997). Myogenesis and MyoD down-regulate Sp1. A mechanism for the repression of GLUT1 during muscle cell differentiation. *J Biol Chem* **272**: 12913-21.

Vispe S, Cazaux C, Lesca C,Defais M (1998). Overexpression of Rad51 protein stimulates homologous recombination and increases resistance of mammalian cells to ionizing radiation. *Nucleic Acids Res* **26**: 2859-64.

von der Lehr N, Johansson S, Wu S, Bahram F, Castell A, Cetinkaya C, et al. (2003). The F-box protein Skp2 participates in c-Myc proteosomal degradation and acts as a cofactor for c-Myc-regulated transcription. *Mol Cell* **11**: 1189-200.

W

Wang T, Lerer I, Gueta Z, Sagi M, Kadouri L, Peretz T, et al. (2001). A deletion/insertion mutation in the BRCA2 gene in a breast cancer family: a possible role of the Alu-polyA tail in the evolution of the deletion. *Genes Chromosomes Cancer* **31**: 91-5.

Wang W, Dong L, Saville B, Safe S (1999a). Transcriptional activation of E2F1 gene expression by 17beta-estradiol in MCF-7 cells is regulated by NF-Y-Sp1/estrogen receptor interactions. *Mol Endocrinol* **13**: 1373-87.

Wang Z, Zhang Y, Lu J, Sun S, Ravid K (1999b). Mpl ligand enhances the transcription of the cyclin D3 gene: a potential role for Sp1 transcription factor. *Blood* **93**: 4208-21.

Wani MA, Means RT, Jr., Lingrel JB (1998). Loss of LKLF function results in embryonic lethality in mice. *Transgenic Res* **7**: 229-38.

BIBLIOGRAFIA

Wani MA, Wert SE,Lingrel JB (1999). Lung Kruppel-like factor, a zinc finger transcription factor, is essential for normal lung development. *J Biol Chem* **274**: 21180-5.

Webster NJ, Resnik JL, Reichart DB, Strauss B, Haas M,Seely BL (1996). Repression of the insulin receptor promoter by the tumor suppressor gene product p53: a possible mechanism for receptor overexpression in breast cancer. *Cancer Res* **56**: 2781-8.

Westin EH, Wong-Staal F, Gelmann EP, Dalla-Favera R, Papas TS, Lautenberger JA, et al. (1982). Expression of cellular homologues of retroviral onc genes in human hematopoietic cells. *Proc Natl Acad Sci U S A* **79**: 2490-4.

Wimmer EA, Jackle H, Pfeifle C,Cohen SM (1993). A Drosophila homologue of human Sp1 is a head-specific segmentation gene. *Nature* **366**: 690-4.

Wong WK, Chen K,Shih JC (2003). Decreased methylation and transcription repressor Sp3 up-regulated human monoamine oxidase (MAO) B expression during Caco-2 differentiation. *J Biol Chem* **278**: 36227-35.

Wright KL, Moore TL, Vilen BJ, Brown AM,Ting JP (1995). Major histocompatibility complex class II-associated invariant chain gene expression is up-regulated by cooperative interactions of Sp1 and NF-Y. *J Biol Chem* **270**: 20978-86.

X

Xiao W, Hodge DR, Wang L, Yang X, Zhang X,Farrar WL (2004). Co-operative functions between nuclear factors NFkappaB and CCAT/enhancer-binding protein-beta (C/EBP-beta) regulate the IL-6 promoter in autocrine human prostate cancer cells. *Prostate* **61**: 354-70.

Xing Y, Fikes JD,Guarente L (1993). Mutations in yeast HAP2/HAP3 define a hybrid CCAAT box binding domain. *Embo J* **12**: 4647-55.

Xing Y, Zhang S, Olesen JT, Rich A,Guarente L (1994). Subunit interaction in the CCAAT-binding heteromeric complex is mediated by a very short alpha-helix in HAP2. *Proc Natl Acad Sci U S A* **91**: 3009-13.

Y

- Yamabe Y, Shimamoto A, Goto M, Yokota J, Sugawara M, Furuichi Y (1998). Sp1-mediated transcription of the Werner helicase gene is modulated by Rb and p53. *Mol Cell Biol* **18**: 6191-200.
- Yamada K, Tanaka T, Miyamoto K, Noguchi T (2000). Sp family members and nuclear factor-Y cooperatively stimulate transcription from the rat pyruvate kinase M gene distal promoter region via their direct interactions. *J Biol Chem* **275**: 18129-37.
- Yang X, Su K, Roos MD, Chang Q, Paterson AJ, Kudlow JE (2001). O-linkage of N-acetylglucosamine to Sp1 activation domain inhibits its transcriptional capability. *Proc Natl Acad Sci U S A* **98**: 6611-6.
- Ye X, Liu SF (2002). Lipopolysaccharide down-regulates Sp1 binding activity by promoting Sp1 protein dephosphorylation and degradation. *J Biol Chem* **277**: 31863-70.
- Yieh L, Sanchez HB, Osborne TF (1995). Domains of transcription factor Sp1 required for synergistic activation with sterol regulatory element binding protein 1 of low density lipoprotein receptor promoter. *Proc Natl Acad Sci U S A* **92**: 6102-6.
- Yokono M, Saegusa N, Matsushita K, Sugiura Y (1998). Unique DNA binding mode of the N-terminal zinc finger of transcription factor Sp1. *Biochemistry* **37**: 6824-32.
- Yoshida K, Miki Y (2004). Role of BRCA1 and BRCA2 as regulators of DNA repair, transcription, and cell cycle in response to DNA damage. *Cancer Sci* **95**: 866-71.
- Yu B, Datta PK, Bagchi S (2003). Stability of the Sp3-DNA complex is promoter-specific: Sp3 efficiently competes with Sp1 for binding to promoters containing multiple Sp-sites. *Nucleic Acids Res* **31**: 5368-76.
- Yun J, Chae HD, Choy HE, Chung J, Yoo HS, Han MH, et al. (1999). p53 negatively regulates cdc2 transcription via the CCAAT-binding NF-Y transcription factor. *J Biol Chem* **274**: 29677-82.

Z

Zaid A, Hodny Z, Li R,Nelson BD (2001). Sp1 acts as a repressor of the human adenine nucleotide translocase-2 (ANT2) promoter. *Eur J Biochem* **268**: 5497-503.

Zeng YX,el-Deiry WS (1996). Regulation of p21WAF1/CIP1 expression by p53-independent pathways. *Oncogene* **12**: 1557-64.

Zhang H, Kobayashi R, Galaktionov K,Beach D (1995). p19Skp1 and p45Skp2 are essential elements of the cyclin A-CDK2 S phase kinase. *Cell* **82**: 915-25.

Zhang L, Yu D, Hu M, Xiong S, Lang A, Ellis LM, et al. (2000). Wild-type p53 suppresses angiogenesis in human leiomyosarcoma and synovial sarcoma by transcriptional suppression of vascular endothelial growth factor expression. *Cancer Res* **60**: 3655-61.

Zheng XL, Matsubara S, Diao C, Hollenberg MD,Wong NC (2001). Epidermal growth factor induction of apolipoprotein A-I is mediated by the Ras-MAP kinase cascade and Sp1. *J Biol Chem* **276**: 13822-9.

Zhou M, McPherson L, Feng D, Song A, Dong C, Lyu SC, et al. (2007). Kruppel-like transcription factor 13 regulates T lymphocyte survival in vivo. *J Immunol* **178**: 5496-504.

Zhou Q, Lieberman PM, Boyer TG,Berk AJ (1992). Holo-TFIID supports transcriptional stimulation by diverse activators and from a TATA-less promoter. *Genes Dev* **6**: 1964-74.

Zorbas M, Sicurella C, Bertoncello I, Venter D, Ellis S, Mucenski ML, et al. (1999). c-Myb is critical for murine colon development. *Oncogene* **18**: 5821-30.

Zwicker J, Liu N, Engeland K, Lucibello FC,Muller R (1996). Cell cycle regulation of E2F site occupation in vivo. *Science* **271**: 1595-7.