

9. BIBLIOGRAFIA

1. Stanski DR. Monitoring depth of anesthesia. In: R.D. Miller. Anesthesia Quinta edición, Churchill Livingstone: 1087-1116.
2. Jones J G. Perception and memory during general anesthesia. Br. J. Anaesth 1994; 73:31-37.
3. Bergman IJ, Kluger MT, Short TG. Awareness during general anaesthesia: a review of 81 cases from the anaesthetic incident monitoring study. Anaesthesia 2002; 57: 549-556.
4. Domino K B, Posner K L, Caplan R A. Awareness during anesthesia. Anesthesiology 1999; 90:1053-1061.
5. Ruiz V, Koo M, García E, Crespo JM, Mayoral V, Casals P, Cochs J. Incidencia de consciencia intraoperatoria en cirugía urgente y su posible relación con el síndrome de estrés postraumático. A propósito de tres casos. Rev Esp Anestesiol Reanim 1998; 45: 106-109.
6. Pomfrett CJD. Heart rate variability, BIS and depth of anaesthesia. Br J Anaesth 1999; 82: 659-662.
7. Sebel PS, Heneghan CP, Ingram DA. Evoked responses a neurophysiological indicator of depth of anesthesia. Br J Anaesth 1985;57: 841-842.
8. Hudson RJ, Stanski DR, Saidman LJ, Meathe E. A model for studying depth of anesthesia and acute tolerance to thiopental. Anesthesiology 1983; 59: 301-308.
9. Hung O R, Varvel JR, Schafer ST, Stanski DR: Thiopental Pharmacodynamics II Quantification of clinical and electroencephalographic depth of anesthesia. Anesthesiology 1992; 77: 237-244.

10. Levy WJ, Shapiro HM, Maruchak G. Automated EEG processing for intraoperative monitoring. *Anesthesiology* 1980; 53: 223-236.
11. Rampil IJ, Matteo RS Changes in EEG spectral edge frequency correlate with the hemodynamic response to laryngoscopy and intubation. *Anesthesiology* 1987; 67: 139-142.
12. Rampil IJ, Mason P, Singh H. Anesthetic potency (MAC) is independent of forebrain structures in the rat. *Anesthesiology* 1993; 78:707-712.
13. Antognini JF, Schwartz KS. Exaggerated anesthetic requirements in the preferentially anesthetized brain. *Anesthesiology* 1993; 79: 1244-1249.
14. Kissin I. Depth of anesthesia and bispectral index monitoring. *Anesth Analg* 2000; 90: 1114-1117.
15. Kissin I, Stanski DR, Brown PT, Bradley EL,Jr: Pentobarbital-morphine anesthetic interacción in terms of intensity of noxious stimulation required for arousal. *Anesthesiology* 1993;78: 744-749.
16. Menigaux C, Guignard B, Adam F, Sessler DI, Joly V, Chauvin M. Esmolol prevents movement and attenuates the BIS response to orotracheal intubation. *Br J Anaesth* 2002; 89: 857-862.
17. Kock M, Martín N, Scholtes JL. Central effects of epidural and intravenous clonidine in patients anesthetized with enflurane/nitrous oxide. *Anesthesiology* 1992; 77: 457-462
18. Hodgson P, Liu S, Gras TW. Does epidural anesthesia have general anesthetic effects? *Anesthesiology* 1999; 91:1687-1692.
19. Hodgson S, Liu S. Epidural lidocaine decreases sevoflurane requirements for adequate depth of anesthesia as measured by the bispectral index monitor. *Anesthesiology* 2001; 91: 799-803.

20. Willenkin RL, Polk SL. Management of general anesthesia. In: R.D. Miller. Anesthesia Tercera edición, Churchill Livingstone: 1045-1056.
21. White DC. Anaesthesia: a privation of the senses. An historical introduction and some definitions In: Consciousness awareness and pain in general anaesthesia Rosen M, Lunn JN Editors Butterworths 1987.
22. Kissin I. General anesthetic action: an obsolete notion? *Anesth Analg* 1993; 76: 215.
23. Antognini Jf, Carstens E. In vivo characterization of clinical anesthesia and its components. *Br J Anaesth* 2002; 89 : 156-166.
24. Heine W, Schwarzbauer C. In vivo imaging of anaesthetic action in humans: approaches with PET and fMRI. *Br J Anaesth* 2002; 89: 112-122.
25. Urban BW, Bleckwenn M. Concepts and correlations relevant to general anaesthesia. *Br J Anaesth* 2002; 89: 3-16
26. Schwinn DA, Shafer SL. Basic principles of pharmacology related to anesthesia. In R.D. Miller. Anesthesia. Quinta edición, Churchill Livingstone: 15-47.
27. Edmond I, Eger II. Uptake and distribution In R.D. Miller. Anesthesia. Quinta edición, Churchill Livingstone: 74-95.
28. Gold M, Abello M, Herrington C. Minimum alveolar concentration of desflurane in patients older than 65 years. *Anesthesiology* 1993; 79: 710-714.
29. Sebel P, Glass P, Fletcher J E, Murphy MR, Gallager C, Quill T. Reduction of the MAC of desflurane with fentanyl. *Anesthesiology* 1992 ; 76: 52-59.
30. Evans JM, Davies WL. Monitoring anaesthesia. *Clinics in Anesthesiology* 1984; 2: 243-263.

31. Russell IF. Midazolam alfentanil: an anaesthetic? An investigation using the isolated forearm technique. *Br J Anaesth* 1993; 70: 42-46.
32. Jones JG, Aggarwal S. Monitoring the depth of anesthesia. In Ghoneim MM Awareness during anesthesia. Butterworth-Heinemann . Oxford. 2001: 60-92.
33. King FG. Functional neurophysiology. In Russel G B, Rodihock L D. Intraoperative neurophysiologic monitoring. Butterworth-Heinemann. Washington. 1995: 27-40.
34. Simón O. Electroencefalografía. Introducción y atlas. Salvat editores. Barcelona. 1983.
35. Schwentker MC. Technical standards and techniques for basic electroencephalography. In Russel G B, Rodihock L D. Intraoperative neurophysiologic monitoring. Butterworth-Heinemann. Washington. 1995: 51-64.
36. Pichlmayer I, Lehmhuhl P, Lips U. EEG atlas for anesthesiologists. Springer-Verlag editors. Berlin 1987.
37. Rampil IJ. What every neuroanesthesiologist should know about electroencephalograms and computerized monitors. *Anesthesiol Clin North Am* 1992; 10: 683-718.
38. Gambus PL. Empleo del electroencefalograma como medida del nivel de profundidad del efecto de los anestésicos. Tesis doctoral. Barcelona, Septiembre 2002.
39. Billard V, Constant I. Analyse automatique de l'électroencephalogramme: quel intérêt en l'an 2000 dans le monitorage de la profondeur de l'anesthésie? *Ann Fr Anesth Réanim* 2001;20: 763-85.
40. Black S, Mahla M. Cuchiara RF. In Neurologic monitoring. R.D. Miller. Anesthesia. Quinta edición, Churchill Livingstone: 1324-1350.

41. Ropcke H, Rehnerg B, Koenen-Bergman M, Bouillon T, Bruñí J, Hoeft A. Surgical stimulation shifts EEG concentration-response relationship of desflurane. *Anesthesiology* 2001; 94: 390-399.
42. Kochs E, Bischoff P, Pichlmeier U, Schulte J. Surgical stimulation induces changes in brain electrical activity during isoflurane/nitrous oxide anesthesia. A topographical electroencephalographic analysis. *Anesthesiology* 1994; 80: 1026-1034.
43. Kiyama S, Tsuaki K. Processed electroencephalogram during combined extradural and general anaesthesia. *Br J Anaesth* 1997; 78: 751-753.
44. Rampil Y J. A primer for EEG signal processing in anaesthesia. *Anesthesiology* 1998; 89: 980-1002.
45. Warren J, Levy MD, Shapiro HM, Maruchak G, Meathe E. Automated EEG processing for intraoperative monitoring: a comparison of techniques. *Anesthesiology* 1980; 53: 223-236.
46. Bührer M, Maitre PO, Hung O, Ebling WF, Schafer SL, Stanski DR. Thiopental pharmacodynamics. I. Defining the pseudo-steady state serum concentration-EEG effect relationship. *Anesthesiology* 1992; 77: 226-236.
47. Hans P, Lecoq P, Brichant F, Dewandre P, Lamy M. Effect of epidural bupivacaine on the relationship between the bispectral index and end-expiratory concentration of desflurane. *Anesthesia* 1999; 54: 899-908.
48. Katoh T, Suzuki A, Ikeda K. Electroencephalographic derivatives as a tool for predicting depth of sedation and anesthesia induced by sevoflurane. *Anesthesiology* 1998; 88:642-650.
49. Thomsen CE, Prior PF. Quantitative EEG in assessment of anaesthetic depth: comparative study of methodology. *Br J Anaesth*; 1996; 77: 172-178.

50. Rehberg B, Bouillon T, Zinserlin J. Comparative pharmacodynamic modeling of the electroencephalography-slowing effect of isoflurane, sevoflurane and desflurane. *Anesthesiology* 1999; 91: 397-405.
51. Schwender D, Daunderer M, Klasing S. Power spectral analysis of the electroencephalogram during increasing end-expiratory concentration of isoflurane, desflurane and sevoflurane. *Anaesthesia* 1998; 53: 335-342.
52. Doi M, Gajraj RJ, Mantzaridis H. Relationship between calculated blood concentration of propofol and electroencephalographic variables during emergence from anaesthesia: comparison of bispectral index, spectral edge frequency, median frequency and evoked potential index. *Br J Anaesth* 1997; 78: 180-184.
53. Leslie K, Fanzca MBBS, Sessler DI, Schroeder M, Walters K. Propofol blood concentration and the bispectral index predict suppression of learning during propofol/epidural anesthesia in volunteers. *Anesth Analg* 1995; 81: 1269-1274.
54. Sleigh JW, Donovan J. Comparison of bispectral index, 95% spectral edge frequency and approximate entropy of EEG, with changes in heart rate variability during induction of general anaesthesia. *Br J Anaesth* 1999; 82: 666-671.
55. Irwin MG, Hui WC, Milne SE, Kenny GNC. Propofol effective concentration 50 and its relationship to bispectral index. *Anaesthesia* 2002; 57: 242-248.
56. Milne SE, Troy A, Irwin MG, Kenny GNC. Relationship between bispectral index, auditory evoked potential index and effect site EC₅₀ for propofol at two clinical endpoints. *Br J Anaesth* 2002; 90: 127-131.
57. Gajraj RJ, Doi M, Mantzaridis H. Comparison of bispectral EEG analysis and auditory evoked potentials for monitoring depth of anaesthesia during propofol anaesthesia. *Br J Anaesth* 1999; 82: 672-678.
58. Drummond JC, Brann CA, Perkins DE. A comparison of median frequency, spectral edge frequency, a frequency band power ratio, total power and dominance shift in

- the determination of depth of anesthesia. *Acta Anaesthesiol Scand* 1991; 35: 693-699.
59. Kearse LA, Manberg P, DeBros F, Chamoun N, Sinai V. Bispectral análisis of the electroencephalogram during induction of anesthesia may predict hemodynamic responses to laryngoscopy and intubation. *Electroenceph Clin Neurophysiol*. 1994; 90: 194-200.
60. White PF, Boyle WA. Relationship between hemodynamic and electroencephalographic changes during anesthesia. *Anesth Analg* 1989; 68:177-181.
61. Wei Dong Mi, Tetsushiro Sakai. Haemodynamic and electroencephalographic responses to intubation during induction with propofol or propofol/fentanyl. *Can J Anaesth* 1998; 45: 19-22.
62. Sidis A, Halimi P, Shamay C. Estimating anesthetic depth by electroencephalography during induction and intubation in patients undergoing cardiac surgery. *J Clin Anesth*; 2: 101-107.
63. Doi M, Gajraj RJ. Prediction of movement at laryngeal mask airway insertion: comparison of auditory evoked potential index, bispectral index, spectral edge frequency and median frequency. *Br J Anaesth* 1999; 82: 203-207.
64. Liu Jin, Singh H. Electroencephalographic bispectral index correlates with intraoperative recall and depth of propofol-induced sedation. *Anesth Analg* 1997; 84: 185-189.
65. Glass P Bloom M. Bispectral analysis measures sedation and memory effects of propofol, midazolam, isoflurane and remifentanil in healthy volunteers. *Anesthesiology* 1997; 86: 836-847.
66. Sandin H, Enlund G, Samuelson P, Lennmarken C. Awareness during anaesthesia: a prospective case study. *Lancet* 2000; 335: 707-711.

67. Luginbühl M, Schnider TW. Detection of awareness with the bispectral index: two case reports. *Anesthesiology* 2002; 1: 241-243.
68. Mychaskiw G, Horowitz M, Sachdev V, Heath BL. Explicit intraoperative recall at a bispectral index of 47. *Anesth Analg* 2001; 92: 808-809.
69. Lubke G H, Kerssens C, Phaf H. Dependence of explicit and implicit memory on hypnotic state in trauma patients. *Anesthesiology* 1999; 90: 670-609.
70. Kerssens C, Klein J, Woerd A, Bonke B. Auditory information processing during adequate propofol anesthesia monitored by electroencephalogram bispectral index. *Anesth Analg* 2001; 92: 1210-1214.
71. Kerssens C, Lubke G, Klein J, Woerd A, Bonke B. Memory function during propofol and alfentanil anesthesia. *Anesthesiology* 2002; 97: 382-389.
72. O'Connor MF, Daves SM, Tung A, Cock RI, Thisted R, Apfelbaum J. BIS monitoring to prevent awareness during general anesthesia. *Anesthesiology* 2001; 94: 520-522.
73. Sebel PS, Glass P, Fletcher JE, Murphy MR, Gallager CD. Reduction of the MAC of desflurane with fentanyl. *Anesthesiology* 1992; 76: 52-59.
74. Rampil IJ, Lockhart SH, Zwass MS, Peterson N, Yasuda N. Clinical characteristics of desflurane in surgical patients: minimum alveolar concentration. *Anesthesiology* 1991; 74: 429-433.
75. Liu W, Thorpe T, Graham S, Aitkenhead A. Incidence of awareness with recall during general anesthesia. *Anaesthesia* 1991; 46: 435-437.
76. Billard V, Gambus PL, Chamoun N, Stanski DR, Schafer L. A comparison of spectral edge, delta power, and bispectral index as EEG measures of alfentanil, propofol, and midazolam drug effect. *Clin Pharmacol Ther* 1997; 61: 45-58.

77. Scott JC, Ponganis KV, Stanski DR. EEG quantitation of narcotic effect: the comparative pharmacodynamics of fentanyl and alfentanil. *Anesthesiology* 1985; 62: 234-241.
78. Rampil IJ, Lockhart SH, Eger EY, Yasuda N, Weiskopf RB, Cahalan MK. The electroencephalographic effects of desflurane in humans. *Anesthesiology* 1991; 74: 434-439.
79. Schwilden H, Sxhüttler J, Stoeckel H. Closed-loop feedback of methohexitol anesthesia by quantitative EEG analysis in humans. *Anesthesiology* 1987; 67: 341-347.
80. Schwilden H, Stoeckel H, Schüttler J. Closed-loop feedback control of propofol anaesthesia by quantitative EEG analysis in humans. *Br J Anaesth* 1989; 62: 290-296.
81. Detsch O, Schneider G, Kochs E, Hapfelmeier G, Werner C. Increasing isoflurane concentration may cause paradoxical increases in the EEG bispectral index in surgical patients. *Br J Anaesth* 2000; 84: 33-37.
82. Katoh T, Ikeda K: The effects of fentanyl on sevoflurane requirements for loss of consciousness and skin incision. *Anesthesiology* 1998; 88: 18-24.
83. Schwender D, Daunderer M, Mulzer S et al: Spectral edge frequency of the electroencephalogram to monitor "depth" of anaesthesia with isoflurane or propofol. *Br J Anaesth* 1996; 77: 179-184.
84. Kiyama S, Takeda J. Effect of extradural analgesia on the paradoxical arousal response of the electroencephalogram. *Br J Anaesth* 1997; 79: 750-753.
85. Vernon J, Lang E, Sebel P, Manberg P. Prediction of movement using bispectral electroencephalographic analysis during propofol/alfentanil or isoflurane/alfentanil anesthesia. *Anesth Analg* 1995; 80: 780-785.

86. Deyne C, Struys M, Heylen R et al: Influence of intravenous clonidine pretreatment on anesthetic requirements during bispectral EEG-guided sevoflurane anesthesia. *J Clin Anesth* 2000; 12: 52-57.
87. Olofson E, Dahan A. The dynamic relationship between end-tidal sevoflurane and isoflurane concentrations and bispectral index and spectral edge frequency of the electroencephalogram. *Anesthesiology* 1999; 90: 1345-1353.
88. Morley A, Chung D, Wong A, Short T. The sedative and electroencephalographic effects of regional anesthesia. *Anaesthesia* 2000; 55: 864-69.
89. Cousins, Bromage P: Epidural neural blockade. In: Cousins MJ, Bridenbaugh PO (eds): *Neural Blockade in clinical anesthesia and management of pain*, Philadelphia: Lippincot, 1988; 253-360.
90. Webster J, Barnard M, Carli F. Metabolic response to colonic surgery: extradural vs continuous spinal. *Br J Anaesth* 1991; 67: 467-469.
91. Dahl JB, Hansen BL, Hjortso NC, Erichsen CJ, Moiniches S, Kehlet H. Influence of timing on the effect of continuous extradural analgesia with bupivacaine and morphine after major abdominal surgery. *Br J Anaesth* 1992; 69: 4-8.
92. Liu S, Carpenter RL, Mackey DC et al. Effects of perioperative analgesic technique on rate of recovery after colon surgery. *Anesthesiology* 1995; 83: 757-765.
93. Hirabayashi Y, Shimizu R. Effect of age on extradural dose requirement in thoracic extradural anaesthesia. *Br J Anaesth* 1993; 71: 445-446.
94. Veerring B, Burm A, Vletter A, et al. The effect of age on the systemic absorption, disposition and pharmacodynamics of bupivacaine after epidural administration. *Clin Pharmacokinet* 1992; 22: 75-84.
95. Sear J. Recent advances and developments in the clinical use of iv opioids during the peroperative period. *Br J Anesth* 1998; 81: 38-50.

96. Mc Ewan A, Smith C, Dyar O, et al. Isoflurane MAC reduction by fentanyl. *Anesthesiology* 1993; 78: 864-869.
97. Peng P, Sandler A. A review of the use of fentanyl analgesia in the management of acute pain in adults. *Anesthesiology* 1999; 90: 576-599.
98. Glass P, Doherty M, Jacobs J, et al. Plasma concentration of fentanyl, with 70% nitrous oxide, to prevent movement at skin incision. *Anesthesiology* 1993; 78: 842-847.
99. Cousins M, Mather L. Intrathecal and epidural administration of opioids. *Anesthesiology* 1984; 61: 276-310.
100. Chribasik S, Chubrasik J Pfisterer M, et al. Comparison of morphine with and without fentanyl for epidural analgesia after major abdominal surgery. *Reg Anesth* 1996; 21: 175-181.
101. Cousins MJ, Cherry DA, Gourlay GK. Acute and chronic pain: use of spinal opioids In: MJ Cousins, Bridenbaugh PO (eds): Neural Blockade in clinical anesthesia and management of pain. Philadelphia: Lippincot, 1988; 955-1030.
102. Sjöström S, Hartving P, Persson P, Tamsen A. Pharmacokinetics of epidural morphine and meperidine in humans. *Anesthesiology* 1987; 67: 877-888.
103. Struys M, Versichelen L., Byttebier G, Mortier E, Moerman E, Rolly G. Clinical usefulness of the bispectral index for titrating propofol target effect-site concentration. *Anaesthesia* 1998; 53: 4-12.
104. Sebel P, Lang E, Rampil I, White P, Cork R, Jopling M, Smith N, Glass P, Manberg P. A multycenter study of bispectral electroencephalogram analysis for monitoring anesthetic effect. *Anesth Analg* 1997; 84: 891-899.
105. Dwyer RC, Rampil IJ, Eger EI, Bennett HL. The electroencephalogram does not predict depth of isoflurane anesthesia. *Anesthesiology* 1994; 81: 403-409.

106. Liu J, Singh H, White PF. Electroencephalogram bispectral analysis predicts the depth of midazolam-induced sedation. *Anesthesiology* 1996; 84: 64-69.
107. Ibrahim A, Taraday JK, Kharasch ED. Bispectral index monitoring during sedation with sevoflurane, Midazolam and propofol. *Anesthesiology* 2001; 95: 1151-1159.
108. Gurman GM. Assessment of depth of general anesthesia. *Int J Clin Monit Comput* 1994; 11: 185-189.
109. Kearse LA, Manberg P, Chamoun N, deBros F, Zaslavsky A. Bispectral analysis of the electroencephalogram correlates with patient movement to skin incision during propofol/nitrous oxide anesthesia. *Anesthesiology* 1994; 81: 1365-1370.
110. Sakai T, Singh H, Kudo T, Kudo M, Matsuki A. Hipnotic endpoints vs. the bispectral index, 95% spectral edge frequency and median frequency during propofol infusion with or without fentanyl. *Eur J Anaesthesiol* 1999; 16: 47-52.
111. Sharpe RM, Nathwani D, Brunner MD, Thornton C, Dore CJ, Newton DEF. Auditory evoked response, median frequency and 95% spectral edge during anaesthesia with desflurane and nitrous oxide. *Br J Anaesth* 1997; 78: 282-285.
112. Song D, Girish P, White P. Titration of volatile anesthetics using bispectral index facilitates recovery after ambulatory anesthesia. *Anesthesiology* 1997; 87: 842-848.
113. Song D, Vlymen J, White P, Fanzca. Is the bispectral index useful in predicting fast-track eligibility after ambulatory anesthesia with propofol and desflurane? *Anesth Analg* 1998; 87: 1245-1248.
114. Pavlin J, Hong J, Freund PR, Koerschgen ME, Bower JO, Bowdle TA. The effect of bispectral index monitoring on end-tidal gas concentration and recovery after outpatient anesthesia. *Anesth Analg* 2001; 93: 613-619.

