

ESTUDIO DE LA FUNCION RENAL A PARTIR DE SECUENCIAS
DINAMICAS DE IMAGENES GAMMAGRAFICAS

Tesis para aspirar al grado
de doctor presentada por
ANGEL GONZALEZ SISTAL

Laboratori de Biofisica i Bioenginyeria
Departament de Ciències Fisiològiques
Humanes i de la Nutrició
Facultat de Medicina
Universitat de Barcelona

Barcelona, 1990

7. REFERENCIAS

Alderson P O, Douglass K H, Mendenhall K G, Guadiani V A, Watson D C, Links J M, y Wagner Jr. H N. Deconvolution analysis in radionuclide quantitation of left-to-right cardiac shunts, J Nucl Med, 20, 502-506, 1979.

Anger H O. Scintillation camera, Rev Sci Instrum, 29, 27-33, 1958.

Bankir L y Grünfeld J P. en Fisiología Humana. Meyer P. Salvat Editores, 1985.

Bendat J S y Piersol A G. Engineering applications of correlation and spectral analysis, John Wiley and sons, 1980.

Blaufox M D. Compartmental analysis of the radiorenogram and kinetics of ^{131}I -Hippuran, Progress in Nuclear Medicine, 2, 107-124, 1972.

Britton K E, Brown N J G y Slater J D H. Isotope renography, Lancet, 1, 986, 1969.

Britton K E y Brown N J G. Clinical renography. London. Lloyd-Luke ltd, 1971.

Britton K E y Brown N J G. The value in obstructive nephropathy of the Hippuran output curve derived by computer analysis of the renogram, en Dynamic studies with radioisotopes in medicine, International Atomic Energy Agency, 1970.

Britton K E, Brown N J G, Cruz F, Chang H C, Ralphs D y Myers M. Deconvolution analysis of dynamic studies, en Information Processing in Scintigraphy. Proc 4th Int Conf, Raynaud, C., y Todd-Pokropek, A., eds., 244-260, 1976.

Brown N J G y Britton K E. A new system of renography, Biomed Eng, 4, 268-274, 1969.

Brown N J G. The application of interpolative background subtraction to ^{99m}Tc -DTPA gammacamara renography data, en Joekes, A.M., Constable, A.R., Brown, N.J.G., Tauxe, W.N.(eds.), Radionuclides in nephrology, London, Academic Press, 1982.

Burbank M, Tauxe W, Maher F y Hunt J. Evaluation of radioiodinated hippuran for the estimation of renal plasma flow. Proc. Staff Meet. Mayo Clin., 36, 372-386, 1961.

Carlsen J E, Moller M L, Lund J O, y Trap-Jensen J. Comparison of four commercial $\text{Tc}^{99m}(\text{Sn})$ -DTPA preparations used for the measurement of GFR: concise communication. J Nucl Med, 21, 126-129, 1980.

Carlsen O. Absorbed radiation dose in adults from Iodine-131 and Iodine-123 Orthoiodohippurate and Technetium- 99m DTPA renography, J Nucl Med, 29, 400-406, 1988.

Catafau A, Piera C, Pavia J, Herranz A y Setoain J. RPF and Renogram with ^{131}I -OIH and Tc^{99m} -MAG₃ in hypertensive patients. Eur J Nucl Med, 14, 296, 1988.

Coveney J R y Robbins, M. S. Biological characterization of Tc-99m mercaptoacetyltriglycine (MAG₃) kit for renal function, J Nucl Med, 28, 732, 1987.

Diffey B L, Hall F M y Corfield J R. The ^{99m}Tc -DTPA dynamic renal scan with deconvolution analysis, J Nucl Med, 17, 352-355, 1976.

Diffey B L y Corfield J R. The data bounding technique in discrete deconvolution, Med Biol Eng, 14, 478, 1976.

Diffey B L, Hall F M, Piepsz A y Erbsmann F. Renal deconvolution and the poor injection, Eur J Nucl Med, 3, 145-146, 1978.

Dore E K, Taplin G V y DeLores E J. Current interpretation of the sodium iodohippurate ^{131}I renocystogram, J.A.M.A., 185, 925-932, 1963.

Dubovsky E V y Yester M V, en Gelfand M J y Thomas S R. GFR, RTT, ERPF and DRF. Effective use of computers in Nuclear Medicine, McGraw-Hill Book company, 1988.

Eshima D, Taylor A, Fritzberg A, Kasina S, Hansen L y Sorenson F. J Nucl Med, 28, 1180-1186, 1987.

Elliott A T y Britton K E. A review of the physiological parameters in the dosimetry of ^{123}I and ^{131}I -labelled hippuran, Int J Appl Rad Isot, 29, 571-573, 1978.

Fleming J S y Goddard B A. A technique for the deconvolution of the renogram, Phys Med Biol, 19, 546-549, 1974.

Fleming J S y Kenny R W. A comparison of techniques for the filtering of noise in the renogram, Phys Med Biol, 22, 359-366, 1977.

Fritzberg A R, Kasina S, Eshima D y Johnson D L. Syntesis and biological evaluation of Technetium-99m MAG3 as a Hippuran replacement, J Nucl Med, 27, 111-116, 1986.

Gates G F. Glomerular filtration rate: Estimation from fractional renal accumulation of $^{99\text{m}}\text{Tc}$ -DTPA (stannous), Am J Roentgenol, 138: 565, 1982.

González R C y Wintz P. Digital Image Processing, Addison-Wesley publishing company, Massachusetts, 1977.

González A, Ros D, Pavía J y Puchal R. Obtención de tiempo de tránsito intrarrenal mediante deconvolución del renograma isotópico, Proceedings VIII Reunión anual de la Agrupación Española de Bioingeniería, 56, 1988.

González A. Ros D. Pavía J y Puchal R. Comparación de dos métodos de filtrado lineal optimizado utilizados en la deconvolución del renograma, Proceedings IX Reunión anual de la Agrupación Española de Bioingeniería, 1989.

Goris M L, Daspit S G, McLaughlin P y Kriss J P. Interpolative background subtraction, J Nucl Med, 17, 744-747, 1976.

Gullquist R R, Abrahamson H, Brandt R, Philip J y Strandell T. New method to calculate the renal transit time spectrum from conventional threee-probe renography, Med Biol Eng Comput, 21, 599-602, 1983.

Gullquist R R y Fleming J S. Error analysis by simulation studies in renography deconvolution, Phys Med Biol, 32, 383-395, 1987.

Hall F M y Monks G K. The renogram. A method of separating vascular and renal components, Invest Radiol, 220-224, 1966.

Hiramatsu Y, O'Mara R E, McAfee J G y Markarian B. Intrarenal distribution of diagnostic agents. Invest Radiol, 5, 295-310, 1970.

Holroyd A M, Chrisholm G D y Glass H I. The quantitative analysis of renograms using the gamma camera, Phys Med Biol, 15, 483-492, 1970.

Hunermann B, Weissbach L y Winkler C. Significance of extra parenchymal activity correction in individual renal radio Hippurate clearance determination. In Zum Winkel, K., Blaufox, M.D. y Funck-Bretano, J.L. (Eds.). Radionuclides in Nephrology, 298-302. Stuttgart. Georg Thieme, 1975.

Hvid-Jacobsen K, Thomsen HS y Nielsen SL. Diuresis renographic. A simultaneous comparison between ^{131}I -hippuran and ^{99}Tcm -MAG₃. Acta Radiol, 31, 83-86, 1990.

Jafri R A, Britton K E, Nimmon C C, Solanski K, Al-Nahhass-A, Bomanji J, Fettich J y Hawkins L A. Technetium- ^{99}m MAG₃, a comparison with iodine-123 and iodine-131 orthoiodohippurate, in patients with renal disorders, J Nucl Med, 29, 147-158, 1988.

Jewkes R F y Jeyasingh K. Comparison of ^{123}I -Hippuran and ^{99}mTc DTPA, Nucl Med Comm, 278-288, 1981.

Kenny R W, Ackery D M, Fleming J S, Goddard B A y Grant R W. Deconvolution analysis of the scintillation camera renogram, Br J Radiol, 48, 481-486, 1975.

Knesaurek K y Spaventi S. Comparison of three deconvolution techniques in renography, Eur J Nucl Med, 9, 254-256, 1984.

Keske U, Cordes M, Wilfling M, Hunger J, Hosten N y Gahl G. ^{99}mTc -MAG₃ and ^{131}I -OIH in patients with kidney insufficient and with normal kidneys, ROFO, 152, 287-292, 1990.

Krueger R P, Sanders A P, Demaria W y Baylin G J. Analysis of the radiorenogram curve, Amer J Roentgen, 86, 81-89, 1961.

Kuruc A, Caldicott W y Treves S. An improved deconvolution technique for calculating organ retentions, J Nucl Med, 22, 59, 1981.

Kuruc A, Treves S y Parker J A. Accuracy of deconvolution algorithms assessed by simulation studies: concise communication, J Nucl Med, 24, 258-263, 1983.

Lawson C L y Hanson R J. Solving least squares problems, Prentice-Hall inc., Englewood Cliffs. New Jersey, 1974.

Lawson R S, en O'Reilly P H, Shields R A y Testa H J. Nuclear Medicine in Urology and Nephrology, London: Butterworths, 1986.

McAfee J G y Subramanian G. Radioactive agents for imaging, en Freeman and Johnson's clinical radionuclide imaging. Grune and Stratton, 1984.

Maini CL, Antonacci P, Sargiotti A, Castellano G y Podio U. Dynamic renal scanning using ^{99m}Tc -MAG₃ in man, Eur J Nucl Med, 15, 635-640, 1989.

Matthews C M E. The theory of tracer experiments with ^{131}I -labelled plasma proteins, Phys Med Biol, 2, 36-53, 1957.

- Middleton G W, Thomson W H, Davies I H y Morgan A. A multiple regression analysis for accurate background subtraction in 99m Tc-DTPA renography. Nucl Med Comm, 10, 315-324, 1989.
- Muller-Suur R y Muller-Suur C. Renal and extrarenal handling of a new imaging compound (99m Tc-MAG-3) in the rat. Eur J Nucl Med, 12, 438-442, 1986.
- Nakagawa T, Maeda H, Terada N, Ono M, Ohi M, Ito T, Takeda K, Tagami T y Matsushita T. Clinical value of deconvolution analysis in radionuclide renal study. Radiat Med, 7, 236-242, 1989.
- Nimmon C C, McAlister J M y Cattell W R. Kidney position and the management of relative uptake of 131 I-Hippuran in renography. Brit J Radiol, 48, 286-290, 1975.
- Norman N. Effective plasma flow of the individual kidney. Determination on the basis of the 131 I-Hippuran renogram. Scand J Clin Lab Invest, 30, 395-403, 1972.
- Papoulis A. Circuits and systems: A modern approach, 1980, Holt, Rinehart and Winston, Inc, 1980.
- Pavia J, Puchal R y Ros D. Obtención de la función de retención renal por técnicas digitales Proceedings IV Reunión Nacional de Física Médica, 488-493, 1983.
- Piepsz A, Dobbeleir A y Ham H R. Background determination for 99m Tc DTPA renal studies. A comparison between interpolative background subtraction and surface ratio method, Nucl Med Commun, 4, 276, 1983.

Piepsz A, Ham H R, Erbsmann F, Hall M Diffey B L, Goggin M J, Hall F M, Miller J A, Lumbroso J, Di Paola R, Bazin J P, Di Paola M y Fries D. A cooperative study on the clinical value of dynamic renal scanning with deconvolution analysis, Br J Radiol, 55, 419-433, 1982.

Puchal R, Pavía J, González A y Ros D. Optimal filtering values in renogram deconvolution, Phys Med Biol, 33, 831-845, 1988.

Pynsent P B y Hanka R. A simple program for a phaseless recursive digital filter. J Biomed Eng, 4, 252-254, 1982.

Reeve J, Crawley J C W, Hamilton S M y Goldberg A D. Deconvolution analysis of the standard renogram and its clinical application, Proc 3th Int Symp Radionucl Nephrol, 321-325, 1975.

Reeve J y Crawley J C W. Quantitative radioisotope renography: the derivation of physiological data by deconvolution analysis using a single-injection technique, Clin Sci Mol Med, 47, 317-330, 1974.

Ros D, Puchal R, Pavía J y González A. Data filtering optimization in renogram deconvolution, Proceedings III Simp. Int. Ing. Biomed., 457-460, 1987.

Rutland M D. A single injection technique for subtraction of blood background in ^{131}I -hippuran renograms. British Journal of Radiology, 52, 134-137, 1979

Sapirstein L A, Vidt D G, Mandel M J y Hanusek G. Volumes of distribution and clearances of intravenously injected creatinine in the dog. American Journal of Physiology, 181, 330-336, 1955.

Savitsky A y Golay J R. Smoothing and differentiation of data by simplified least squares procedures, Analytical Chemistry, 36, 1627-1639, 1964.

Short M D, Glass H I y Vernon P. Use of the gammacamera for ^{131}I renography: a case study on a nefrectomized patient, J Nucl Med, 11, 758-760, 1970.

Smith H W. The Kidney: Structure and Function in Health and Disease. London. Oxford University Press, 1951.

Stephenson J L. Theory of the measurements of blood flow by the dilution of an indicator, Bull Math Biophys 10, 117-121, 1948.

Stockell M, Bröchner-Mortennen J y Rödbro P. Determination of fractional renal function in renography with and without individually (^{131}I) albumin-determined blood background. Scand J Clin Lab Invest, 35, 511-516, 1975.

Szabó Z, Vosberg H, Sondhaus C A y Feinendegen L E. Model identification and estimation of organ-function parameters using radioactive tracers and the impulse-response function, Eur J Nucl Med, 11, 265-274, 1985.

Szabó Z, Kutkuhn B, Georgescu G, Mecklenbeck W, Suatmadjia A y Vosberg H. Parametric representation of kidney function using $^{99\text{m}}\text{Tc}$ -mercaptoacetyl-triglycine (MAG₃), Nuklearmedizin, 28, 73-83, 1989.

Tamminen T E, Rihimaki E J y Thati E E. A gammacamera method for quantitation of split renal function in children followed for vesicoureteric reflux, Pediatr Radio, 7, 78-84, 1978.

Tauxe W N, Hunt J C y Burbank M K. The radioisotope renogram (orthoiodohippurate I-131). Standardization of technic and expression of data. Am J Clin Path, 37, 567-578, 1962.

Tauxe W N, Tobin M, Dubovsky E V, Bueschen A J y Kontzen F. A macrofunction for computer processing of comprehensive renal function studies, Eur J Nucl Med, 5, 103-108, 1980.

Taplin G V, Meredith O M, Kade H y Winter C C. The radioisotope renogram: an external test for individual kidney function and upper urinary tract patency. J Lab Clin Med, 48, 886-901, 1956.

Taylor A, Eshima D, Fritzberg A R, et al. Comparison of iodine-131-OIH and technetium-99m MAG3 renal imaging in volunteers, J Nucl Med, 27, 795-803, 1986.

Taylor A, Eshima D y Alazraki N. 99mTc-MAG3, a new renal imaging agent: preliminary results in patients, Eur J Nucl Med, 12, 510-514, 1987.

Taylor A, Eshima D, Christian P E, Milton W. Evaluation of Tc-99m mercaptoacetyltriglycine in patients with impaired renal function, Radiology, 162, 365-370, 1987.

Taylor A, Eshima D, Christian P E, Wooten W W, Hansen L y McElvany K. Technetium-99m MAG3 kit formulation: Preliminary results in normal volunteers and patients with renal failure, J Nucl Med, 29, 616-622, 1988.

Timmermans L y Merchie G. The renogram, en Radioisotopes in the diagnosis of diseases of the kidneys and the urinary tract, Proc 1st Int Symp, Timmermans, L., y Merchie, G., eds, Liege, 18-41, 1967.

Todd-Pokropek A y Di Paola R. The use of computers for image processing in Nuclear Medicine, IEEE Trans NS, 29, 1299-1309, 1982.

Tonnesen K H, Muncho O, Hadd T et al. Influence on the renogram of variation in skin-to-kidney distance and the clinical importance there of, en zum Winkel K, Blaufox MD, Funck-Brentano JL eds: Radionuclides in Nephrology, 1974. Stuttgart. Georg Thieme Verlag, 1975.

Valentinuzzi M E y Montaldo Volachec E M. Discrete deconvolution, Med Biol Eng, 13, 123-5, 1975.

Van Huffel S, Vandewalle J, De Roo M C y Willem J L. Reliable and efficient deconvolution technique based on total linear squares for calculating the renal retention function, Med Biol Eng Comput, 25, 26-33, 1987.

Van Stekelenburg L H M, Al, N, Kooman A y Tertoolen J F W. A three compartment model for the transport and distribution of hippuran, Phys Med Biol, 21, 74-84, 1976.

Wedeen R P, Goldstein M H y Levitt M F. The radioisotope renogram in normal subjects, Amer J Med, 34, 765-774, 1963.

Winter C C. A clinical study of new renal function test, the radioactive Diodrast renogram, J Urol, 76, 182-196, 1956.

Winter C C. Radioisotope renography, The Williams and Wilkins company, Baltimore, 1963.

Wooten W W, Eshima D y Taylor A. Radiation absorbed dose from I-131 OIH, and Tc-99m MAG3, J Nucl Med, 898, 1986.

Yester M V. Renal transit time, en Tauxe, W.N., y Dubovysky E V. Nuclear Medicine in clinical urology and nephrology, Norwalk, Conn., Appleton-Century-Crofts, 1985.

Zierler K. A critique of compartmental analysis, Ann Rev Biophys Bioeng, 10, 531-562, 1981.