

TESIS DOCTORAL

EL PRECONDICIONAMIENTO ISQUÉMICO COMO ESTRATEGIA QUIRÚRGICA ÚTIL EN EL TRASPLANTE HEPÁTICO CON INJERTO DE TAMAÑO REDUCIDO



Rosa Franco Gou

Universidad de Barcelona
Departamento de Fisiología Animal
2006

7. BIBLIOGRAFÍA

1. Zakim, Boyer. Hepatology. A textbook of liver disease, 4 edition Saunders Ed. 2002.
2. Sherlock S, Dooley J. Diseases of the liver and biliary system. Blackwell Publishing ED 11 edition. 1988.
3. Arias, Boyer, Fausto, Jakoby Schacter Shafritz. The liver biology and pathobiology, 3 edition. Raven Press Ed. 1994.
4. Ginns LC, Cosimi AB, Morris PJ. Transplantation. Blackwell Science ED. 1999.
5. Bismuth H, Castaing D, Ericzon BG, Otte JB, Rolles K, Ringe B, Sloof M. Hepatic transplantation in Europe. First Report of the European Liver Transplant Registry. Lancet. 1987;2(8560):674-676.
6. Cuervas-Mons V., del Castillo-Olivares J.L.. Introducción al trasplante de órganos y tejidos. 2^a edición Arán Ediciones 1999.
7. Quinta memoria de resultados del Registro Español de Trasplante Hepático, 1984-2001.
8. Bzeizi KI, Jalan R, Plevris JN, Hayes PC. Primary graft dysfunction after liver transplantation: from pathogenesis to prevention. Liver Transpl Surg. 1997;3(2):137-148.
9. Neuberger, J. Recent advances in clinical practice. Developments in liver transplantation. Gut 2004; 53:759-768.
10. Grande L, Matus D, Rimola A, Manyalic M, Cabrer C, Garcia-Valdecasas JC, Visa J. Expanded liver donor age over 60 years for hepatic transplantation. Clin Transpl. 1998:297-301.
11. Grande L, Matus D, Manyalic M, Cabrer C, Rodriguez-Montalvo C, Rimola A, Navasa M, Garcia-Valdecasas JC, Visa J. Effect of donor age on graft outcome after liver transplantation. Transplant Proc. 1999;31(6):2482-2483.
12. Alexander JW, Zola JC. Expanding the donor pool: use of marginal donors for solid organ transplantation. Clin Transplant. 1996;10(1 Pt 1):1-19.
13. Melendez HV, Heaton ND. Understanding "marginal" liver grafts. Transplantation. 1999;68(4):469-471.
14. Inomata Y, Nakamura T, Uemoto S, Tanaka K, Wakabayashi G, Shimazu M. Domino split-liver transplantation from a living donor: case reports of in situ and ex situ splitting. Liver Transpl. 2001;7(2):150-153.
15. Azoulay D, Samuel D, Castaing D, Adam R, Adams D, Said G, Bismuth H. Domino liver transplants for metabolic disorders: experience with familial amyloidotic polyneuropathy. J Am Coll Surg. 1999;189(6):584-593.
16. Schmidt HH, Nashan B, Propsting MJ, Nakazato M, Flemming P, Kubicka S, Boker K, Pichlmayr R, Manns MP. Familial Amyloidotic Polyneuropathy: domino liver transplantation. J Hepatol. 1999;30(2):293-298.
17. Monteiro E, Perdigoto R, Furtado AL. Liver transplantation for familial amyloid polyneuropathy. Hepatogastroenterology. 1998;45(23):1375-1380.
18. D'alessandro AM, Hoffmann RM, Knechtle SJ, Odorico JS, Becker YT, Musat A, Pirsch JD, Sollinger HW, Kalayoglu M. Liver transplantation from controlled non-heart-beating donors. Surgery. 2000;128(4):579-588.

19. Casavilla A, Ramirez C, Shapiro R, Nghiem D, Miracle K, Bronsther O, Randhawa P, Broznick B, Fung JJ, Starzl T. Experience with liver and kidney allografts from non-heart-beating donors. *Transplantation*. 1995;59(2):197-203.
20. D'Alessandro AM, Hoffmann RM, Knechtle SJ, Eckhoff DE, Love RB, Kalayoglu M, Sollinger HW, Belzer FO. Successful extrarenal transplantation from non-heart-beating donors. *Transplantation*. 1995;59(7):977-982.
21. Malago M, Rogiers X, Broelsch CE. Reduced-size hepatic allografts. *Annu Rev Med* 1995;46:507-512. Review.
22. Ploeg RJ, D'Alessandro AM, Knechtle et al. Risk factors for primary dysfunction after liver transplantation, a multivariate analysis. *Transplantation* 1993;55:807-813.
23. Strasberg SM, Howard TK, Molmenti EP, Hertzl M. Selecting the donor liver: risk factors for poor function after orthotopic liver transplantation. *Hepatology* 1994;20:829-838.
24. Starzl TE, Demetris AJ. Liver transplantation: a 31-year experience, part I. *Curr Probl Surg* 1990;17(2): 92-93.
25. Broelsch CE, Neuhaus P, Burdelski M, Bojar H. Orthotope lebertransplantation von lebersegmenten bei kleinkindern mit gallengangsatzesieen. *Langenbecks Arch Chir* 1984;362:105-109 (Suppl).
26. Bismuth H, Houssin D. Reduced-sized orthotopic liver graft in hepatic transplantation in children. *Surgery*. 1984;95(3):367-370.
27. Emond JC, Whitington PF, Thistlethwaite JR, Alonso EM, Broelsch CE. Reduced-size orthotopic liver transplantation: use in the management of children with chronic liver disease. *Hepatology*. 1989;10(5):867-872.
28. De Ville de Goyet J, Hausleithner V, Reding R, Lerut J, Janssen M, Otte JB. Impact of innovative techniques on the waiting list and results in pediatric liver transplantation. *Transplantation*. 1993;56(5):1130-1136.
29. Pichlmayer R, Ringe B, Gubernatis G, Hauss J, Bunzendahl H. Transplantation einer spenderleber auf zwei empfänger (splitting-transplantation) eine neue methode in der weiterentwicklung der lebersegment-transplantation. *Langenbecks Arch Chir* 1988;373:127-130.
30. Emond JC, Whitington PF, Thistlethwaite JR, Cherqui D, Alonso EA, Woodle IS, Vogelbach P, Busse-Henry SM, Zucker AR, Broelsch CE. Transplantation of two patients with one liver. Analysis of a preliminary experience with 'split-liver' grafting. *Ann Surg*. 1990;212(1):14-22.
31. Busuttil RW, Goss JA. Split liver transplantation. *Ann Surg*. 1999;229(3):313-21.
32. Sauer IM, Pascher A, Steinmuller T, Settmacher U, Muller AR, Bechstein WO, Neuhaus P. Split liver and living donation liver transplantation: the Berlin experience. *Transplant Proc*. 2001;33(1-2):1459-1460.
33. Broelsch CE, Whitington PF, Emond JC, Heffron TG, Thistlethwaite JR, Stevens L, Piper J, Whitington SH, Lichtor JL. Liver transplantation in children from living related donors. Surgical techniques and results. *Ann Surg*. 1991;214(4):428-437; discussion 437-439.

34. Broelsch CE, Emond JC, Thistlethwaite JR, Rouch DA, Whitington PF, Lichtor JL. Liver transplantation with reduced-size donor organs. *Transplantation*. 1988;45(3):519-524.
35. Otte JB, de Ville de Goyet J, Reding R, Van Obbergh L, Veyckemans F, Carlier MA et al. Pediatric liver transplantation: from the full-size liver graft to reduced, split, and living related liver transplantation. *Pediatr Surg Int* 1998;13(5-6):308-318. Review.
36. Tanaka K, Uemoto S, Tokunaga Y, Fujita S, Sano K, Nishizawa T, Sawada H, Shirahase I, Kim HJ, Yamaoka Y. Surgical techniques and innovations in living related liver transplantation. *Ann Surg* 1993; 217(1):82-91.
37. Ozawa K, Uemoto S, Tanaka K, Kumada K, Yamaoka Y, Kobayashi N, namoto T, Shimahara Y, Mori K, Honda K. An appraisal of pediatric liver transplantation from living relatives. Initial clinical experiences in 20 pediatric liver transplants from living relatives as donors. *Ann Surg* 1992; 216(5):547-553.
38. Marcos A, Fisher RA, Ham JM, Shiffman ML, Sanyal AJ, Luketic VA, Sterling RK, Posner MP. Right lobe living donor liver transplantation. *Transplantation*. 1999;68(6):798-803.
39. Koniaris LG, McKillop IH, Schwartz SI, Zimmers TA. Liver regeneration. *J Am Coll Surg*. 2003;197(4):634-659. Review
40. Michalopoulos GK, DeFrances MC. Liver regeneration. *Science*. 1997 Apr 4;276(5309):60-66.
41. Fausto N. Liver regeneration. *J Hepatol*. 2000;32(1 Suppl):19-31. Review.
42. Malik R, Selden C, Hodgson H. The role of non-parenchymal cells in liver growth. *Semin Cell Dev Biol*. 2002 Dec;13(6):425-431. Review.
43. Yamada Y, Kirillova I, Peschon JJ, Fausto N. Initiation of liver growth by tumor necrosis factor: deficient liver regeneration in mice lacking type I tumor necrosis factor receptor. *Proc Natl Acad Sci U S A*. 1997;94(4):1441-1446.
44. Uchiyama H, Yanaga K, Nishizaki T, Soejima Y, Yoshizumi T, Sugimachi K. Effects of deletion variant of hepatocyte growth factor on reduced-size liver transplantation in rats. *Transplantation* 1999; 68(1):39-44.
45. Burr AW, Toole K, Chapman C, Hines JE, Burt AD. Anti-hepatocyte growth factor antibody inhibits hepatocyte proliferation during liver regeneration. *J Pathol* 1998; 185(3):298-302.
46. Khalil N. TGF-beta: from latent to active. *Microbes Infect*. 1999;1(15):1255-1263. Review.
47. Russell WE, Coffey RJ Jr, Ouellette AJ, Moses HL. Type beta transforming growth factor reversibly inhibits the early proliferative response to partial hepatectomy in the rat. *Proc Natl Acad Sci U S A*. 1988;85(14):5126-5130.
48. Enami Y, Kato H, Murakami M, Fujioka T, Aoki T, Niiya T, Murai N, Ohtsuka K, Kusano M. Anti-transforming growth factor-beta1 antibody transiently enhances DNA synthesis during liver regeneration after partial hepatectomy in rats. *J Hepatobiliary Pancreat Surg* 2001;8(3):250-258.
49. Harvey PR, Iu S, McKeown CM, Petrunka CN, Ilson RG, Strasberg SM. Adenine nucleotide tissue concentrations and liver allograft viability after cold preservation and warm ischemia. *Transplantation*. 1988 ;45(6):1016-1020.

50. Maruyama H, Harada A, Kurokawa T, Kobayashi H, Nonami T, Nakao A, Takagi H. Duration of liver ischemia and hepatic regeneration after hepatectomy in rats. *J Surg Res* 1995; 58(3):290-294.
51. Portugal V, García-Alonso I, Barceló P, Mendez J. Effect of allopurinol, folinic acid, SOD and cyclosporine A on ischemic liver regeneration. *Eur Surg Res* 1995; 27(2):69-76.
52. Foschi D, Castoldi L, Lesma A, Musazzi M, Benevento A, Trabucchi E. Effects of ischaemia and reperfusion on liver regeneration in rats. *Eur J Surg* 1993; 159(8):393-398.
53. Clavien PA, Harvey PR, Strasberg SM. Preservation and reperfusion injuries in liver allografts. An overview and synthesis of current studies. *Transplantation* 1992; 53(5):957-978. Review.
54. Reilly PM, Schiller HJ, Bulkley GB. Pharmacologic approach to tissue injury mediated by free radicals and other reactive oxygen metabolites. *Am J Surg*. 1991;161(4):488-503.
55. Biasi F, Bosco M, Chiappino I, Chiarpotto E, Lanfranco G, Ottobrelli A, Massano G, Donadio PP, Vaj M, Andorno E. Oxidative damage in human liver transplantation. *Free Radic Biol Med*. 1995;19(3):311-317.
56. Portugal V, García-Alonso I, Bilbao J, Barceló P y Méndez J. Efecto de terapias antioxidantes en la regeneración del hígado isquémico. *Rev esp Enf Digest*, 1993.83(6): 434-438.
57. Harris CM, Massey V. The oxidative half-reaction of xanthine dehydrogenase with NAD; reaction kinetics and steady-state mechanism. *J Biol Chem*. 1997;272(45):28335-28341.
58. Stirpe F, Della Corte E. The regulation of rat liver xanthine oxidase. Conversion in vitro of the enzyme activity from dehydrogenase (type D) to oxidase (type O). *J Biol Chem*. 1969;264:3855-3863.
59. Peralta C, Bulbena O, Xaus C, Prats N, Cutrin JC, Poli G, Gelpi E, Rosello-Catafau J. Ischemic preconditioning: a defense mechanism against the reactive oxygen species generated after hepatic ischemia reperfusion. *Transplantation*. 2002;73(8):1203-1211.
60. Jaeschke H, Smith CW, Clemens MG, Ganey PE, Roth RA. Mechanisms of inflammatory liver injury: adhesion molecules and cytotoxicity of neutrophils. *Toxicol Appl Pharmacol*. 1996;139(2):213-226.
61. Ghezzi P, Dinarello CA, Bianchi M, Rosandich ME, Repine JE, White CW. Hypoxia increases production of interleukin-1 and tumor necrosis factor by human mononuclear cells. *Cytokine*. 1991;3(3):189-194.
62. Zhou W, McCollum MO, Levine BA, Olson MS. Inflammation and platelet-activating factor production during hepatic ischemia/reperfusion. *Hepatology*. 1992;16(5):1236-1240.
63. Cutrin JC, Boveris A, Zingaro B, Corvetti G, Poli G. In situ determination by surface chemiluminescence of temporal relationships between evolving warm ischemia-reperfusion injury in rat liver and phagocyte activation and recruitment. *Hepatology*. 2000;31(3):622-632.

64. Jaeschke H, Bautista AP, Spolarics Z, Spitzer JJ. Superoxide generation by neutrophils and Kupffer cells during in vivo reperfusion after hepatic ischemia in rats. *J Leukoc Biol.* 1992;52(4):377-382.
65. Jaeschke H. Reactive oxygen and ischemia/reperfusion injury of the liver. *Chem Biol Interact.* 1991;79(2):115-136.
66. Jaeschke H, Farhood A, Smith CW. Neutrophils contribute to ischemia/reperfusion injury in rat liver in vivo. *FASEB J.* 1990;4(15):3355-3359.
67. Hisama N, Yamaguchi Y, Ishiko T, Miyanari N, Ichiguchi O, Goto M, Mori K, Watanabe K, Kawamura K, Tsurufuji S, Ogawa M. Kupffer cell production of cytokine-induced neutrophil chemoattractant following ischemia/reperfusion injury in rats. *Hepatology.* 1996;24(5):1193-1198.
68. Jaeschke H, Farhood A, Bautista AP, Spolarics Z, Spitzer JJ, Smith CW. Functional inactivation of neutrophils with a Mac-1 (CD11b/CD18) monoclonal antibody protects against ischemia-reperfusion injury in rat liver. *Hepatology.* 1993;17(5):915-923.
69. Lentsch AB, Kato A, Yoshidome H, McMasters KM, Edwards MJ. Inflammatory mechanisms and therapeutic strategies for warm hepatic ischemia/reperfusion injury. *Hepatology.* 2000;32(2):169-173.
70. Martinez-Mier G, Toledo-Pereyra LH, McDuffie JE, Warner RL, Ward PA. Neutrophil depletion and chemokine response after liver ischemia and reperfusion. *J Invest Surg.* 2001;14(2):99-107.
71. Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med.* 1989;320(6):365-376. Review.
72. Lemasters JJ, Thurman RG. Reperfusion injury after liver preservation for transplantation. *Annu Rev Pharmacol Toxicol.* 1997;37:327-38. Review.
73. Klein AS, Zhadkevich M, Wang D, Margolick JB, Winkelstein JA, Bulkley GB. Discriminant quantitation of posttransplant hepatic reticuloendothelial function. The impact of ischemic preservation. *Transplantation.* 1996;61(8):1156-61.
74. Dinarello CA. Biologic basis for interleukin-1 in disease. *Blood.* 1996;87:2095-2147.
75. Beutler B. TNF, immunity and inflammatory disease: lessons of the past decade. *J Investig Med.* 1995;43:227-235.
76. Cutrin JC, Perrelli MG, Cavalieri B, Peralta C, Rosell Catafau J, Poli G. Microvascular dysfunction induced by reperfusion injury and protective effect of ischemic preconditioning. *Free Radic Biol Med.* 2002;33(9):1200-1208. Review.
77. Colletti LM, Remick DG, Burtch GD, Kunkel SL, Strieter RM, Campbell DA Jr. Role of tumor necrosis factor-alpha in the pathophysiologic alterations after hepatic ischemia/reperfusion injury in the rat. *J Clin Invest.* 1990;85(6):1936-1943.
78. Thornton AJ, Strieter RM, Lindley I, Baggiolini M, Kunkel SL. Cytokine-induced gene expression of a neutrophil chemotactic factor/IL-8 in human hepatocytes. *J Immunol.* 1990;144(7):2609-2613.
79. Pober JS. Warner-Lambert/Parke-Davis award lecture. Cytokine-mediated activation of vascular endothelium. Physiology and pathology. *Am J Pathol.* 1988;133(3):426-433.

80. Colletti LM, Kunkel SL, Walz A, Burdick MD, Kunkel RG, Wilke CA, Strieter RM. The role of cytokine networks in the local liver injury following hepatic ischemia/reperfusion in the rat. *Hepatology*. 1996;23(3):506-514.
81. Shibuya H, Ohkohchi N, Tsukamoto S, Satomi S. Tumor necrosis factor-induced, superoxide-mediated neutrophil accumulation in cold ischemic/reperfused rat liver. *Hepatology*. 1997;26(1):113-120.
82. Shirasugi N, Wakabayashi G, Shimazu M, Oshima A, Shito M, Kawachi S, Karahashi T, Kumamoto Y, Yoshida M, Kitajima M. Up-regulation of oxygen-derived free radicals by interleukin-1 in hepatic ischemia/reperfusion injury. *Transplantation*. 1997;64(10):1398-1403.
83. Shito M, Wakabayashi G, Ueda M, Shimazu M, Shirasugi N, Endo M, Mukai M, Kitajima M. Interleukin 1 receptor blockade reduces tumor necrosis factor production, tissue injury, and mortality after hepatic ischemia-reperfusion in the rat. *Transplantation*. 1997;63(1):143-148.
84. Colletti LM, Burtch GD, Remick DG, Kunkel SL, Strieter RM, Guice KS, Oldham KT, Campbell DA Jr. The production of tumor necrosis factor alpha and the development of a pulmonary capillary injury following hepatic ischemia/reperfusion. *Transplantation*. 1990;49(2):268-272.
85. Hato S, Urakami A, Yamano T, Uemura T, Ota T, Hirai R, Shimizu N. Attenuation of liver and lung injury after hepatic ischemia and reperfusion by a cytokine-suppressive agent, FR167653. *Eur Surg Res* 2001;33(3):202-209.
86. Watanabe M, Yamaguchi K, Chijiwa K, Tanaka M. FR167653 improves survival and pulmonary injury after partial hepatectomy under ischemia/reperfusion in rats. *J Surg Res* 2001;101(2):146-151.
87. Court FG, Wemyss-Holden SA, Dennison AR, Maddern GJ. The mystery of liver regeneration. *Br J Surg*. 2002;89(9):1089-1095. Review.
88. Boulton R, Woodman A, Calnan D, Selden C, Tam F, Hodgson H. Nonparenchymal cells from regenerating rat liver generate interleukin-1 α and -1 β : a mechanism of negative regulation of hepatocyte proliferation. *Hepatology* 1997; 26(1):49-58.
89. Brundage SI, Zautke NA, Holcomb JB, Spain DA, Lam JC, Mastrangelo MA, Macaitis JM, Tweardy DJ. Interleukin-6 infusion blunts proinflammatory cytokine production without causing systematic toxicity in a swine model of uncontrolled hemorrhagic shock. *J Trauma*. 2004;57(5):970-977; discussion 977-978.
90. Jaeschke H. Molecular mechanisms of hepatic ischemia-reperfusion injury and preconditioning. *Am J Physiol Gastrointest Liver Physiol*. 2003;284(1):G15-G26. Review.
91. Hong F, Radaeva S, Pan HN, Tian Z, Veech R, Gao B. Interleukin 6 alleviates hepatic steatosis and ischemia/reperfusion injury in mice with fatty liver disease. *Hepatology*. 2004;40(4):933-941.
92. Tange S, Hofer Y, Welte M, Anthuber M, Jauch KW, Geissler EK, Ertel W. Local secretion of TNF-alpha from the liver does not correlate with endotoxin, IL-6, or organ function in the early phase after orthotopic liver transplantation. *Transpl Int*. 2001;14(2):80-86.
93. Van Hoorn EC, van Middelaar-Voskuilen MC, van Limpt CJ, Lamb KJ, Bouritius , Vriesema AJ, van Leeuwen PA, van Norren K. Preoperative supplementation with

- a carbohydrate mixture decreases organ dysfunction-associated risk factors. *Clin Nutr.* 2005;24(1):114-123.
94. Ledda-Columbano GM, Curto M, Piga R, Zedda AI, Menegazzi M, Sartori C, Shinozuka H, Bluethmann H, Poli V, Ciliberto G, Columbano A. In vivo hepatocyte proliferation is inducible through a TNF and IL-6-independent pathway. *Oncogene* 1998; 17(8):1039-1044.
 95. Liu P, Xu B, Hock CE, Nagele R, Sun FF, Wong PY. NO modulates P-selectin and ICAM-1 mRNA expression and hemodynamic alterations in hepatic I/R. *Am J Physiol.* 1998;275(6 Pt 2):H2191-H2198.
 96. Rockey DC, Chung JJ. Reduced nitric oxide production by endothelial cells in cirrhotic rat liver: endothelial dysfunction in portal hypertension. *Gastroenterology.* 1998;114(2):344-351.
 97. Shah V, Haddad FG, Garcia-Cardena G, Frangos JA, Mennone A, Groszman RJ, Sessa WC. Liver sinusoidal endothelial cells are responsible for nitric oxide modulation of resistance in the hepatic sinusoids. *J Clin Invest.* 1997;100(11):2923-2930.
 98. Liu P, Yin K, Nagele R, Wong PY. Inhibition of nitric oxide synthase attenuates peroxynitrite generation, but augments neutrophil accumulation in hepatic ischemia-reperfusion in rats. *J Pharmacol Exp Ther.* 1998;284(3):1139-1146.
 99. Caplan MS, Hedlund E, Hill N, MacKendrick W. The role of endogenous nitric oxide and platelet-activating factor in hypoxia-induced intestinal injury in rats. *Gastroenterology.* 1994;106(2):346-352.
 100. Yin DP, Sankary HN, Chong AS, Ma LL, Shen J, Foster P, Williams JW. Protective effect of ischemic preconditioning on liver preservation-reperfusion injury in rats. *Transplantation.* 1998;66(2):152-157.
 101. Pannen BH, Al-Adili F, Bauer M, Clemens MG, Geiger KK. Role of endothelins and nitric oxid in hepatic reperfusion injury in the rat. *Hepatology.* 1998;27(3):755-764.
 102. Cottart CH, Do L, Blanc MC, Vaubourdolle M, Descamps G, Durand D, Galen FX, Clot JP. Hepatoprotective effect of endogenous nitric oxide during ischemia-reperfusion in the rat. *Hepatology.* 1999;29(3):809-813.
 103. Jones SM, Thurman RG. L-arginine minimizes reperfusion injury in a low-flow, reflow model of liver perfusion. *Hepatology.* 1996;24(1):163-168.
 104. Uhlmann D, Scommatou S, Witzigmann H, Spiegel HU. Exogenous L-arginine protects liver microcirculation from ischemia reperfusion injury. *Eur Surg Res.* 1998;30(3):175-184.
 105. Calabrese F, Valente M, Pettenazzo E, Ferrarese M, Burra P, Cadrobbi R, Cardin R, Bacelle L, Parnigotto A, Rigotti P. The protective effects of L-arginine after liver ischaemia/reperfusion injury in a pig model. *J Pathol.* 1997;183(4):477-485.
 106. Jaeschke H, Schini VB, Farhood A. Role of nitric oxide in the oxidant stress during ischemia/reperfusion injury of the liver. *Life Sci.* 1992;50(23):1797-1804.
 107. Koeppel TA, Thies JC, Schemmer P, Trauner M, Gebhard MM, Otto G, Post S. Inhibition of nitric oxide synthesis in ischemia/reperfusion of the rat liver is followed by impairment of hepatic microvascular blood flow. *J Hepatol.* 1997;27(1):163-169.

108. Pabla R, Buda AJ, Flynn DM, Blesse SA, Shin AM, Curtis MJ, Lefer DJ. Nitric oxide attenuates neutrophil-mediated myocardial contractile dysfunction after ischemia and reperfusion. *Circ Res.* 1996;78(1):65-72.
109. Naito Y, Yoshikawa T, Matsuyama K, Yagi N, Arai M, Nakamura Y, Kaneko T, Yoshida N, Kondo M. Neutrophils, lipid peroxidation, and nitric oxide in gastric reperfusion injury in rats. *Free Radic Biol Med.* 1998;24(3):494-502.
110. Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxy nitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci U S A.* 1990;87(4):1620-1624.
111. Liu P, Hock CE, Nagele R, Wong PY. Formation of nitric oxide, superoxide, and peroxy nitrite in myocardial ischemia-reperfusion injury in rats. *Am J Physiol.* 1997;272(5 Pt 2):H2327-H2336.
112. Szabo C. The pathophysiological role of peroxy nitrite in shock, inflammation, and ischemia-reperfusion injury. *Shock.* 1996;6(2):79-88.
113. Rai RM, Lee FY, Rosen A, Yang SQ, Lin HZ, Koteish A, Liew FY, Zaragoza C, Lowenstein C, Diehl AM. Impaired liver regeneration in inducible nitric oxide synthase-deficient mice. *Proc Natl Acad Sci U S A.* 1998;95(23):13829-13834.
114. Goto M, Takei Y, Kawano S, Tsuji S, Fukui H, Fushimi H, Nishimura Y, Kashiwagi T, Fusamoto H, Kamada T. Tumor necrosis factor and endotoxin in the pathogenesis of liver and pulmonary injuries after orthotopic liver transplantation in the rat. *Hepatology.* 1992;16(2):487-493.
115. Matuschak GM, Rinaldo GE, Pinsky MR, Gavaler JS, Van Thiel DH. Effect of end stage liver failure on the incidence and resolution of the adult respiratory distress syndrome. *J Crit Care.* 1987;2:162-173.
116. Yost CS, Matthay MA, Gropper MA. Etiology of acute pulmonary edema during liver transplantation: a series of cases with analysis of the edema fluid. *Chest.* 2001;119(1):219-223.
117. Zheng SS, Xu SG, Liang TB, et al. Early expression of pro-inflammatory cytokines in the lung of rat with small-for-size liver transplantation and the significance. *Zhonghua Yi Xue Za Zhi* 2003;83(7):588-592.
118. Mack CL, Millis JM, Whitington PF, et al. Pulmonary complications following liver transplantation in pediatric patients. *Pediatr Transplant.* 2000;4(1):39-44.
119. Whitington PF, Alonso EM, Piper JB. Pediatric liver transplantation. *Semin Liver Dis.* 1994;14(3):303-317. Review.
120. Jaeschke H, Farhood A. Neutrophil and Kupffer cell-induced oxidant stress and ischemia-reperfusion injury in rat liver. *Am J Physiol.* 1991;260(3 Pt 1):G355-G362.
121. Sankary HN, Foster P, Brown E, Williams J. Do splanchnic viscera contribute to liver preservation reperfusion injury? *Transplantation.* 1996;61(8):1142-1147.
122. Sankary HN, Yin DP, Chong AS, Ma LL, Blinder L, Shen JK, Foster P, Liu LP, Li C, Williams JW. The portosystemic shunt protects liver against ischemic reperfusion injury. *Transplantation.* 1999;68(7):958-93.
123. Okuaki Y, Miyazaki H, Zeniya M, Ishikawa T, Ohkawa Y, Tsuno S, Sakaguchi M, Hara M, Takahashi H, Toda G. Splenectomy-reduced hepatic injury induced by ischemia/reperfusion in the rat. *Liver.* 1996;16(3):188-194.

124. Carvalho EM, Massarollo PC, Isern MR, Toledo NS, Kawacama J, Mies S, Raia S. Pulmonary evolution in conventional liver transplantation with venovenous bypass and the piggyback method. *Transplant Proc.* 1999;31(7):3064-3066.
125. Urata K, Brault A, Rocheleau B, Huet PM. Role of Kupffer cells in the survival after rat liver transplantation with long portal vein clamping times. *Transpl Int.* 2000;13(6):420-427.
126. Strieter RM, Kunkel SL, Bone RC. Role of tumor necrosis factor-alpha in disease states and inflammation. *Crit Care Med.* 1993;21(10 Suppl):S447-S463.
127. Shibuya H, Ohkohchi N, Seya K, Satomi S. Kupffer cells generate superoxide anions and modulate reperfusion injury in rat livers after cold preservation. *Hepatology.* 1997;25(2):356-360.
128. Sheron N, Eddleston A. Preservation-reperfusion injury, primary graft non-function and tumour necrosis factor. *J Hepatol.* 1992;16(3):262-265.
129. Nishimura Y, Takei Y, Kawano S, Yoshihara H, Goto M, Hijioka T, Fukui H, Oshita M, Fusamoto H, Kamada T. Long-term storage of liver grafts is associated with a postoperative elevation of tumor necrosis factor: a possible role in the pathogenesis of primary nonfunction of the graft and pulmonary complications. *Transplant Proc.* 1993;25(1 Pt 1):130-131.
130. Suzuki S, Nakamura S, Serizawa A, Sakaguchi T, Konno H, Muro H, Kosugi I, Baba S. Role of Kupffer cells and the spleen in modulation of endotoxin-induced liver injury after partial hepatectomy. *Hepatology.* 1996;24(1):219-225.
131. Liu P, Xu B, Hock CE. Inhibition of nitric oxide synthesis by L-NAME exacerbates acute lung injury induced by hepatic ischemia-reperfusion. *Shock.* 2001;16(3):211-217.
132. Wanner GA, Ertel W, Muller P, et al. Liver ischemia and reperfusion induces a systemic inflammatory response through Kupffer cell activation. *Shock.* 1996;5(1):34-40.
133. Shito M, Wakabayashi G, Ueda M, et al. Interleukin 1 receptor blockade reduces tumor necrosis factor production, tissue injury, and mortality after hepatic ischemia-reperfusion in the rat. *Transplantation.* 1997;63(1):14314-14318.
134. Seekamp A, Warren JS, Remick DG, Till GO, Ward PA. Requirements for tumor necrosis factor-alpha and interleukin-1 in limb ischemia/reperfusion injury and associated lung injury. *Am J Pathol.* 1993;143(2):453-463.
135. Grenz A, Schenk M, Zipfel A, Viebahn R. TNF-alpha and its receptors mediate graft rejection and loss after liver transplantation. *Clin Chem Lab Med.* 2000;38(11):1183-1185.
136. Weil D. What's new about tumor necrosis factors? Report on the fourth TNF International Congress. *Eur Cytokine Netw.* 1992;3(3):347-351.
137. Leist M, Gantner F, Jilg S, Wendel A. Activation of the 55 kDa TNF receptor is necessary and sufficient for TNF-induced liver failure, hepatocyte apoptosis, and nitrite release. *J Immunol.* 1995;154(3):1307-1316.
138. Niederle N, Ganser A, Bergmann L. *Zytokine.* Gustav Fischer verlag 1996; 15-19.
139. Lin SY, Wang YY, Sheu WH. Increased serum leptin concentrations correlate with soluble tumour necrosis factor receptor levels in patients with cirrhosis. *Clin Endocrinol (Oxf).* 2002;57(6):805-811.

140. Majno PE, Morel P, Mentha G. Mini-review: tumor necrosis factor (TNF) and TNF soluble receptors (TNF-sR) in liver disease and liver transplantation. *Swiss Surg* 1995;(4):182-185. Review.
141. on Baehr V, Docke WD, Plauth M, Liebenthal C, Kupferling S, Lochs H, Baumgarten R, Volk HD. Mechanisms of endotoxin tolerance in patients with alcoholic liver cirrhosis: role of interleukin 10, interleukin 1 receptor antagonist, and soluble tumour necrosis factor receptors as well as effector cell desensitisation. *Gut* 2000;47(2):281-287.
142. Fulop AK, Pocsik E, Brozik M, Karabelyos C, Kiss A, Novak I, Szalai C, Dobozy O, Falus A. Hepatic regeneration induces transient acute phase reaction: systemic elevation of acute phase reactants and soluble cytokine receptors. *Cell Biol Int* 2001;25(7):585-592.
143. Carney DE, Lutz CJ, Picone AL, Gatto LA, Schiller HJ, Finck CM, Searles B, Paskanik AM, Snyder KP, Edwards C, Nieman GF. Soluble tumor necrosis factor receptor prevents post-pump syndrome. *J Surg Res* 1999;83(2):113-121.
144. Sorkine P, Setton A, Halpern P, Miller A, Rudick V, Marmor S, Klausner JM, Goldman G. Soluble tumor necrosis factor receptors reduce bowel ischemia-induced lung permeability and neutrophil sequestration. *Crit Care Med* 1995;23(8):1377-1381.
145. Holtmann H, Wallach D. Down regulation of the receptors for tumor necrosis factor by interleukin 1 and 4 beta-phorbol-12-myristate-13-acetate. *J Immunol* 1987;139(4):1161-1167.
146. Winzen R, Wallach D, Kemper O, Resch K, Holtmann H. Selective up-regulation of the 75-kDa tumor necrosis factor (TNF) receptor and its mRNA by TNF and IL-1. *J Immunol* 1993;150(10):4346-4353.
147. Hart PH, Hunt EK, Bonder CS, atson CJ, Finlay-Jones JJ. Regulation of surface and soluble TNF receptor expression on human monocytes and synovial fluid macrophages by IL-4 and IL-10. *J Immunol* 1996;157(8):3672-3680.
148. Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation*. 1986;74(5):1124-1136.
149. Heurteaux C, Lauritzen I, Widmann C, Lazdunski M. Essential role of adenosine, adenosine A1 receptors, and ATP-sensitive K⁺ channels in cerebral ischemic preconditioning. *Proc Natl Acad Sci U S A*. 1995;92(10):4666-4670.
150. Peralta C, Hotter G, Closa D, Prats N, Xaus C, Gelpi E, Rosello-Catafau J. The protective role of adenosine in inducing nitric oxide synthesis in rat liver ischemia preconditioning is mediated by activation of adenosine A2 receptors. *Hepatology*. 1999;29(1):126-132.
151. Nakayama H, Yamamoto Y, Kume M, Yamagami K, Yamamoto H, Kimoto S, Ishikawa Y, Ozaki N, Shimahara Y, Yamaoka Y. Pharmacologic stimulation of adenosine A2 receptor supplants ischemic preconditioning in providing ischemic tolerance in rat livers. *Surgery* 1999;126(5):945.
152. Ishida T, Yarimizu K, Gute DC, Korthuis RJ. Mechanisms of ischemic preconditioning. *Shock*. 1997;8(2):86-94.
153. Ricciardi R, Meyers WC, Schaffer BK, Kim RD, Shah SA, Wheeler SM, Donohue SE, Sheth KR, Callery MP, Chari RS. Protein kinase C inhibition abrogates hepatic ischemic preconditioning responses. *J Surg Res*. 2001;97(2):144-149.

154. Speechley-Dick ME, Mocanu MM, Yellon DM. Protein kinase C. Its role in ischemic preconditioning in the rat. *Circ Res.* 1994;75(3):586-590.
155. Carini R, De Cesaris MG, Splendore R, Vay D, Domenicotti C, Nitti MP, Paola D, Pronzato MA, Albano E. Signal pathway involved in the development of hypoxic preconditioning in rat hepatocytes. *Hepatology.* 2001;33:131-139.
156. Carini R, De Cesaris MG, Splendore R, Domenicotti C, Nitti MP, Pronzato MA, Albano E. Signal pathway responsible for hepatocyte preconditioning by nitric oxide. *Free Radic Biol Med.* 2003;34:1047-1055.
157. Ricciardi R, Schaffer BK, Kim RD, Shah SA, Donohue SE, Wheeler SM, Quarfordt SH, Callery MP, Meyers WC, Chari RS. Protective effects of ischemic preconditioning on the cold-preserved liver are tyrosine kinase dependent. *Transplantation.* 2001;72(3):406-412.
158. Carini R, De Cesaris MG, Splendore R, Albano E. Stimulation of p38 MAP kinase reduces acidosis and Na(+) overload in preconditioned hepatocytes. *FEBS Lett.* 2001;491:180-183.
159. Teoh N, Dela Pena A, Farrell G. Hepatic ischemic preconditioning in mice is associated with activation of NF-kappaB, p38 kinase, and cell cycle entry. *Hepatology.* 2002;36:94-102.
160. Funaki H, Shimizu K, Harada S, Tsuyama H, Fushida S, Tani T, Miwa K. Essential role for nuclear factor kappaB in ischemic preconditioning for ischemia-reperfusion injury of the mouse liver. *Transplantation.* 2002;74(4):551-556.
161. Ricciardi R, Shah SA, Wheeler SM, Quarfordt SH, Callery MP, Meyers WC, Chari RS. Regulation of NFkappaB in hepatic ischemic preconditioning. *J Am Coll Surg.* 2002;195(3):319-326.
162. Gullo CA, Teoh G. Heat shock proteins: to present or not, that is the question. *Immunol Lett.* 2004;94(1-2):1-10. Review.
163. Morimoto RI, Santoro MG. Stress-inducible responses and heat shock proteins: new pharmacologic targets for cytoprotection. *Nat Biotechnol.* 1998;16(9):833-838. Review.
164. Benjamin IJ, McMillan DR. Stress (heat shock) proteins: molecular chaperones in cardiovascular biology and disease. *Circ Res.* 1998;83(2):117-132. Review.
165. LoCicero J 3rd, Xu X, Zhang L. Heat shock protein suppresses the senescent lung cytokine response to acute endotoxemia. *Ann Thorac Surg* 1999;68(4):1150-1153.
166. Hsu BR, Juang JH, Chen ST, Hsu S, Fu SH. Cobalt-Protoporphyrin treatment renders islets tolerant to interleukin-1 beta suppression. *Transplant Proc* 2004;36(4):1181-1182.
167. Ye J, Laychock SG. A protective role for heme oxygenase expression in pancreatic islets exposed to interleukin-1beta. *Endocrinology* 1998;139(10):4155-4163.
168. Ding XZ, Fernandez-Prada CM, Bhattacharjee AK, Hoover DL. Over-expression of hsp-70 inhibits bacterial lipopolysaccharide-induced production of cytokines in human monocyte-derived macrophages. *Cytokine* 2001;16(6):210-219.
169. Yoo CG, Lee S, Lee CT, Kim YW, Han SK, Shim YS. Anti-inflammatory effect of heat shock protein induction is related to stabilization of I kappa B alpha through

- preventing I kappa B kinase activation in respiratory epithelial cells. *J Immunol* 2000;164(10):5416-5423.
170. Man K, Lee TK, Liang TB, Lo CM, Fung PC, Tsui SH, Li XL, Ng KT, Fan ST. FK 409 ameliorates small-for-size liver graft injury by attenuation of portal hypertension and down-regulation of Egr-1 pathway. *Ann Surg* 2004;240(1):159-168.
171. Katori M, Tamaki T, Takahashi T, Tanaka M, Kawamura A, Kakita A. Prior induction of heat shock proteins by a nitric oxide donor attenuates cardiac ischemia/reperfusion injury in the rat. *Transplantation* 2000;69(12):2530-2537.
172. Kalia N, Brown NJ, Hopkinson K, Stephenson TJ, Wood RF, Pockley AG. FK409 inhibits both local and remote organ damage after intestinal ischaemia. *J Pathol* 2002;197(5):595-602.
173. Kume M, Yamamoto Y, Saad S, Gomi T, Kimoto S, Shimabukuro T, Yagi T, Nakagami M, Takada Y, Morimoto T, Yamaoka Y. Ischemic preconditioning of the liver in rats: implications of heat shock protein induction to increase tolerance of ischemia-reperfusion injury. *J Lab Clin Med*. 1996;128:251-258.
174. Redaelli CA, Tian YH, Schaffner T, Ledermann M, Baer HU, Dufour JF. Extended preservation of rat liver graft by induction of heme oxygenase-1. *Hepatology*. 2002;35(5):1082-1092.
175. Bauer M, Bauer I. Heme oxygenase-1: redox regulation and role in the hepatic response to oxidative stress. *Antioxid Redox Signal*. 2002;4(5):749-758.
176. Uchinami H, Yamamoto Y, Kume M, Yonezawa K, Ishikawa Y, Taura K, Nakajima A, Hata K, Yamaoka Y. Effect of heat shock preconditioning on NF-kappaB/I-kappaB pathway during I/R injury of the rat liver. *Am J Physiol Gastrointest Liver Physiol*. 2002;282(6):G962-G967.
177. Arai M, Thurman RG, Lemasters JJ. Ischemic preconditioning of rat livers against cold storage-reperfusion injury: role of nonparenchymal cells and the phenomenon of heterologous preconditioning. *Liver Transpl*. 2011;7(4):292-299.
178. Yonezawa K, Yamamoto Y, Yamamoto H, Ishikawa Y, Uchinami H, Taura K, Nakajima A, Yamaoka Y. Suppression of tumor necrosis factor-alpha production and neutrophil infiltration during ischemia-reperfusion injury of the liver after heat shock preconditioning. *J Hepatol*. 2001;35(5):619-627.
179. Schlossberg H, Zhang Y, Dudus L, Engelhardt JF. Expression of c-fos and c-jun during hepatocellular remodeling following ischemia/reperfusion in mouse liver. *Hepatology*. 1996;23(6):1546-1555.
180. Ishii S, Abe T, Saito T, Tsuchiya T, Kanno H, Miyazawa M, Suzuki M, Motoki R, Gotoh M. Effects of preconditioning on ischemia/reperfusion injury of hepatocytes determined by immediate early gene transcription. *J Hepatobiliary Pancreat Surg*. 2001;8:461-468.
181. Saito T, Ishii S, Abe T, Tsuchiya T, Kanno H, Miyazawa M, Suzuki M, Gotoh M. Effect of preconditioning in the liver against ischemia/reperfusion injury, protection of sinusoidal cells and alterations of gene transcription. *Transplant Proc*. 2001;33:849.
182. Sindram D, Rudiger HA, Upadhyaya AG, Strasberg SM, Clavien PA. Ischemic preconditioning protects against cold ischemic injury through an oxidative stress dependent mechanism. *J Hepatol*. 2002;36:78-84.

183. Xiao JS, Cai FG, Niu Y, Zhang Y, Xu XL, Ye QF. Preconditioning effects of expression of proto-oncogenes c-fos and c-jun after hepatic ischemia/reperfusion in rats. *Hepatobiliary Pancreat Dis Int.* 2005;4(2):197-202.
184. Teoh N, Leclercq I, Pena AD, Farrell G. Low-dose TNF-alpha protects against hepatic ischemia-reperfusion injury in mice: implications for preconditioning. *Hepatology.* 2003;37:118-128.
185. Peralta C, Closa D, Hotter G, Gelpi E, Prats N, Roselló-Catafau J. Liver ischemic preconditioning is mediated by the inhibitory action of nitric oxide on endothelin. *Biochem Biophys Res Commun.* 1996;229(1):264-270.
186. Gong JP, Tu B, Wang W, Peng Y, Li SB, Yan LN. Protective effect of nitric oxide induced by ischemic preconditioning on reperfusion injury of rat liver graft. *World J Gastroenterol.* 2004;10(1):73-76.
187. Peralta C, Hotter G, Closa D, Gelpi E, Bulbena O, Rosello-Catafau J. Protective effect of preconditioning on the injury associated to hepatic ischemia-reperfusion in the rat: role of nitric oxide and adenosine. *Hepatology.* 1997;25(4):934-937.
188. Arai M, Thurman RG, Lemasters JJ. Contribution of adenosine A(2) receptors and cyclic adenosine monophosphate to protective ischemic preconditioning of sinusoidal endothelial cells against Storage/Reperfusion injury in rat livers. *Hepatology.* 2000;32(2):297-302.
189. Peralta C, Rull R, Rimola A, Deulofeu R, Rosello-Catafau J, Gelpi E, Rodes J. Endogenous nitric oxide and exogenous nitric oxide supplementation in hepatic ischemia-reperfusion injury in the rat. *Transplantation.* 2001;71(4):529-536.
190. Peralta C, Closa D, Xaus C, Gelpi E, Rosello-Catafau J, Hotter G. Hepatic preconditioning in rats is defined by a balance of adenosine and xanthine. *Hepatology.* 1998;28(3):768-773.
191. Peralta C, Bartrons R, Riera L, Manzano A, Xaus C, Gelpi E, Rosello-Catafau J. Hepatic preconditioning preserves energy metabolism during sustained ischemia. *Am J Physiol Gastrointest Liver Physiol.* 2000;279(1):G163-G171.
192. Peralta C, Bartrons R, Serafin A, Blazquez C, Guzman M, Prats N, Xaus C, Cutillas B, Gelpi E, Rosello-Catafau J. Adenosine monophosphate-activated protein kinase mediates the protective effects of ischemic preconditioning on hepatic ischemia-reperfusion injury in the rat. *Hepatology.* 2001;34(6):1164-1173.
193. Compagnon P, Wang HB, Southard JH, et al. Ischemic preconditioning in a rodent hepatocyte model of liver hypothermic preservation injury. *Cryobiology.* 2002;44:269-278.
194. Compagnon P, Lindell S, Ametani MS, Gilligan B, Wang HB, D'Alessandro AM, Southard JH, Mangino MJ. Ischemic preconditioning and liver tolerance to warm or cold ischemia: experimental studies in large animals. *Transplantation.* 2005;79:1393-1400.
195. Howell JG, Zibari GB, Brown MF, Burney DL, Sawaya DE, Olinde JG, Granger DN, McDonald JC. Both ischemic and pharmacological preconditioning decrease hepatic leukocyte/endothelial cell interactions. *Transplantation.* 2000;69(2):300-303.
196. Koti RS, Yang W, Dashwood MR, Davidson BR, Seifalian AM. Effect of ischemic preconditioning on hepatic microcirculation and function in a rat model of ischemia reperfusion injury. *Liver Transpl.* 2002;8(12):1182-1191.

197. Iwasaki Y, Tagaya N, Hattori Y, Yamaguchi K, Kubota K. Protective effect of ischemic preconditioning against intermittent warm-ischemia-induced liver injury. *J Surg Res.* 2002;107(1):82-92.
198. Sawaya DE Jr, Brown M, Minardi A, Bilton B, Burney D, Granger DN, McDonald JC, Zibari GB. The role of ischemic preconditioning in the recruitment of rolling and adherent leukocytes in hepatic venules after ischemia/reperfusion. *J Surg Res.* 1999;85(1):163-170.
199. Zapletal C, Maksan SM, Lehmann T, Guenther L, Fallsehr C, Mehrabi A, Weiss G, Golling M, Gebhard MM, Herfarth C, Klar E. Ischemic preconditioning improves liver microcirculation after ischemia/reperfusion. *Transplant Proc.* 1999;31(8):3260-3262.
200. Shinoda M, Shimazu M, Wakabayashi G, Tanabe M, Hoshino K, Kitajima M. Tumor necrosis factor suppression and microcirculatory disturbance amelioration in ischemia/reperfusion injury of rat liver after ischemic preconditioning. *J Gastroenterol Hepatol.* 2002;17(11):1211-1219.
201. Peralta C, Perales JC, Bartrons R, Mitchell C, Gilgenkrantz H, Xaus C, Prats N, Fernandez L, Gelpi E, Panes J, Rosello-Catafau J. The combination of ischemic preconditioning and liver Bcl-2 overexpression is a suitable strategy to prevent liver and lung damage after hepatic ischemia-reperfusion. *Am J Pathol.* 2002 Jun;160(6):2111-2122.
202. Peralta C, Prats N, Xaus C, Gelpi E, Rosello-Catafau J. Protective effect of liver ischemic preconditioning on liver and lung injury induced by hepatic ischemia-reperfusion in the rat. *Hepatology.* 1999;30(6):1481-1419.
203. Serafin A, Rosello-Catafau J, Prats N, Gelpi E, Rodes J, Peralta C. Ischemic preconditioning affects interleukin release in fatty livers of rats undergoing ischemia/reperfusion. *Hepatology.* 2004;39(3):688-698.
204. Peralta C, Fernandez L, Panes J, Prats N, Sans M, Pique JM, Gelpi E, Rosello-Catafau J. Preconditioning protects against systemic disorders associated with hepatic ischemia-reperfusion through blockade of tumor necrosis factor-induced P-selectin up-regulation in the rat. *Hepatology.* 2001;33(1):100-113.
205. Fernandez L, Heredia N, Grande L, Gomez G, Rimola A, Marco A, Gelpi E, Rosello-Catafau J, Peralta C. Preconditioning protects liver and lung damage in rat liver transplantation: role of xanthine/xanthine oxidase. *Hepatology.* 2002;36(3):562-572.
206. Teoh NC, Farrell GC. Hepatic ischemia reperfusion injury: Pathogenic mechanisms and basis for hepatoprotection. *J Gastroenterol Hepatol.* 2003;18(8):891-902.
207. Pasupathy S, Homer-Vanniasinkam S. Ischaemic preconditioning protects against ischaemia/reperfusion injury: emerging concepts. *Eur J Vasc Endovasc Surg.* 2005;29(2):106-115.
208. Terajima H, Enders G, Thiaener A, Hammer C, Kondo T, Thiery J, Yamamoto Y, Yamaoka Y, Messmer K. Impact of hyperthermic preconditioning on postischemic hepatic microcirculatory disturbances in an isolated perfusion model of the rat liver. *Hepatology.* 2000;31(2):407-415.
209. Matsumoto K, Honda K, Kobayashi N. Protective effect of heat preconditioning of rat liver graft resulting in improved transplant survival. *Transplantation.* 2001;71(7):862-868.

210. Yonezawa K, Yamamoto Y, Yamamoto H, Ishikawa Y, Uchinami H, Taura K, Nakajima A, Yamaoka Y. Suppression of tumor necrosis factor-alpha production and neutrophil infiltration during ischemia-reperfusion injury of the liver after heat shock preconditioning. *J Hepatol.* 2001;35(5):619-627.
211. Ito K, Ozasa H, Sanada K, Horikawa S. Doxorubicin preconditioning: a protection against rat hepatic ischemia-reperfusion injury. *Hepatology.* 2000;31:416-419.
212. Chen H, Yu YY, Zhang MJ, Deng XX, Yang WP, Ji J, Peng CH, Li HW. Protective effect of doxorubicin induced heat shock protein 72 on cold preservation injury of rat livers. *World Gastroenterol.* 2004;10(9):1375-1378.
213. Gerbes AL, Vollmar AM, Kiemer AK, Bilzer M. The guanylate cyclase-coupled natriuretic peptide receptor: a new target for prevention of cold ischemia-reperfusion damage of the rat liver. *Hepatology.* 1998;28:1309-1317.
214. Bilzer M, Jaeschke H, Vollmar AM, Paumgartner G, Gerbes AL. Prevention of Kupffer cell-induced oxidant injury in rat liver by atrial natriuretic peptide. *Am J Physiol.* 1999;276:G1137-G1144.
215. Kiemer AK, Gerbes AL, Bilzer M, Vollmar AM. The atrial natriuretic peptide and cGMP: novel activators of the heat shock response in rat livers. *Hepatology.* 2002;35:88-94.
216. Malhi H, Gorla GR, Irani AN, Annamaceni P, Gupta S. Cell transplantation after oxidative hepatic preconditioning with radiation and ischemia-reperfusion leads to extensive liver repopulation. *PNAS.* 2002;99(20):13114-13119.
217. Yellon DM, Alkhulaifi AM, Pugsley WB. Preconditioning the human myocardium. *Lancet.* 1993;342(8866):276-277.
218. Marber MS, Latchman DS, Walker JM, Yellon DM. Cardiac stress protein elevation 24 hours after brief ischemia or heat stress is associated with resistance to myocardial infarction. *Circulation.* 1993;88(3):1264-1272.
219. Jenkins DP, Pugsley WB, Alkhulaifi AM, Kemp M, Hooper J, Yellon DM. Ischaemic preconditioning reduces troponin T release in patients undergoing coronary artery bypass surgery. *Heart.* 1997;77(4):314-318.
220. Chen S, Li G, Long L. Clinical research of ischemic preconditioning on lung protection. *Hunan Yi Ke Da Xue Xue Bao.* 1999;24(4):357-359.
221. Kharbanda RK, Mortensen UM, White PA, Kristiansen SB, Schmidt MR, Hoschitzky JA, Vogel M, Sorensen K, Redington AN, MacAllister R. Transient limb ischemia induces remote ischemic preconditioning in vivo. *Circulation.* 2002;106(23):2881-2883.
222. Yang Y, Chen SX, Zhang WX. Effect of ischemic preconditioning on human lung cell apoptosis in vivo and the expression of regulating gene bcl-2. *Hunan Yi Ke Da Xue Xue Bao.* 2002;27(1):43-45.
223. Sitzer M, Foerch C, Neumann-Haefelin T, Steinmetz H, Misselwitz B, Kugler C, Back T. Transient ischaemic attack preceding anterior circulation infarction is independently associated with favourable outcome. *J Neurol Neurosurg Psychiatry.* 2004;75(4):659-
224. Wegener S, Gottschalk B, Jovanovic V, Knab R, Fiebach JB, Schellinger PD, Kucinski T, Jungehulsing GJ, Brunecker P, Muller B, Banasik A, Amberger N, Wernecke KD, Siebler M, Rother J, Villringer A, Weih M; MRI in Acute Stroke Study Group of the German Competence Network Stroke. Transient ischemic

- attacks before ischemic stroke: preconditioning the human brain? A multicenter magnetic resonance imaging study. *Stroke*. 2004;35(3):616-621.
225. Clavien PA, Yadav S, Sindram D, Bentley RC. Protective effects of ischemic preconditioning for liver resection performed under inflow occlusion in humans. *Ann Surg*. 2000;232(2):155-162.
226. Clavien PA, Selzner M, Rudiger HA, Graf R, Kadry Z, Rousson V, Jochum W. A prospective randomized study in 100 consecutive patients undergoing major liver resection with versus without ischemic preconditioning. *Ann Surg*. 2003;238(6):843-850.
227. Nuzzo G, Giulante F, Vellone M, De Cosmo G, Ardito F, Murazio M, D'Acapito F, Giovannini I. Pedicle clamping with ischemic preconditioning in liver resection. *Liver Transpl*. 2004;10(2 Suppl 1):S53-S57.
228. Chouker A, Schachtner T, Schauer R, Dugas M, Lohe F, Martignoni A, Pollwein B, Niklas M, Rau HG, Jauch KW, Peter K, Thiel M. Effects of Pringle manoeuvre and ischaemic preconditioning on haemodynamic stability in patients undergoing elective hepatectomy: a randomized trial. *Br J Anaesth*. 2004;93(2):204-211.
229. Li SQ, Liang LJ, Huang JF, Li Z. Ischemic preconditioning protects liver from hepatectomy under hepatic inflow occlusion for hepatocellular carcinoma patients with cirrhosis. *World J Gastroenterol*. 2004;10(17):2580-2584.
230. Kamada N, Calne RY. Orthotopic liver transplantation in the rat. Technique using cuff for portal vein anastomosis and biliary drainage. *Transplantation*. 1979;28(1):47-50.
231. Xia R, Emond JC. Orthotopic partial liver transplantation in the rat: a model of 70% hepatectomy and reduced size liver transplantation. *Transplantation* 1993; 56(4):1041-1043.
232. Omura T, Ascher NL, Emond JC. Fifty-percent partial liver transplantation in the rat. *Transplantation* 1996; 62(2):291-302.
233. Takei Y, Gao WS, Hijioka T, Savier E, Lindert KA, Lemasters JJ. Ischemic preconditioning protects liver from hepatectomy under hepatic inflow occlusion for hepatocellular or hepatic microcirculation. *Transplantation* 1991;52(2):225-230.
234. Schauer RJ, Bilzer M, Kalmuk S, Gerbes AL, Leiderer R, Schildberg FW, Messmer K. Microcirculatory failure after rat liver transplantation is related to kupffer cell-derived oxidant stress but not involved in early graft dysfunction. *Transplantation* 2001;72(10):1692-1699.
235. Okamura N, Tsukada K, Sakaguchi T, Otake M, Yoshida K, Muto T. Enhanced liver regeneration by FK 506 can be blocked by interleukin-1 alpha and interleukin-2. *Transplant Proc*. 1992;24(1):413-415.
236. Kaur D, Cruess DF, Potter WZ. Effect of IL-1alpha on the release of norepinephrine in rat hypothalamus. *J Neuroimmunol*. 1998;90(2):122-127.
237. Kelly KJ, Baird NR, Greene AL. Induction of stress response proteins and experimental renal ischemia/reperfusion. *Kidney Int* 2001; 59(5):1798-1802.
238. Shoskes DA. Effect of bioflavonoids quercetin and curcumin on ischemic renal injury: a new class of renoprotective agents. *Transplantation* 1998; 66(2):147-152.

239. Singleton KD, Serkova N, Beckey VE, Wischmeyer PE. Glutamine attenuates lung injury and improves survival after sepsis: role of enhanced heat shock protein expression. *Crit Care Med.* 2005;33(6):1206-1213.
240. Awasthi N, Wagner BJ. Upregulation of heat shock protein expression by proteasome inhibition: an antiapoptotic mechanism in the lens. *Invest Ophthalmol Vis Sci.* 2005;46(6):2082-2091.
241. Amersi F, Buelow R, Kato H, Ke B, Coito AJ, Shen XD, Zhao D, Zaky J, Melinek J, Lassman CR, Kolls JK, Alam J, Ritter T, Volk HD, Farmer DG, Ghobrial RM, Busuttil RW, Kupiec-Weglinski JW. Upregulation of heme oxygenase-1 protects genetically fat Zucker rat livers from ischemia/reperfusion injury. *J Clin Invest* 1999; 104(11):1631-1639.
242. Huang TJ, McCoubrey WK Jr, Maines MD. Heme oxygenase-2 interaction with metalloporphyrins: function of heme regulatory motifs. *Antioxid Redox Signal.* 2001;3(4):685-696.
243. Lai IR, Ma MC, Chen CF, Chang KJ. The protective role of heme oxygenase-1 on the liver after hypoxic preconditioning in rats. *Transplantation.* 2004;77(7):1004-1008.
244. Singh D, Chander V, Chopra K. The effect of quercetin, a bioflavonoid on ischemia/reperfusion induced renal injury in rats. *Arch Med Res.* 2004;35(6):484-494.
245. Muthian G, Bright JJ. Quercetin, a flavonoid phytoestrogen, ameliorates experimental allergic encephalomyelitis by blocking IL-12 signaling through JAK-STAT pathway in T lymphocyte. *J Clin Immunol.* 2004;24(5):542-552.
246. Ozaki M, Fuchinoue S, Teraoka S, Ota K. The in vivo cytoprotection of ascorbic acid against ischemia/reoxygenation injury of rat liver. *Arch Biochem Biophys* 1995; 318(2):439-445.
247. Tong V, Chang TK, Chen J, Abbott FS. The effect of valproic acid on hepatic and plasma levels of 15-F2t-isoprostane in rats. *Free Radical Bio Med.* 2003; 34(11):1435-1446.
248. Cavalieri B, Perrelli MG, Aragno M, Mastrocola R, Corvetti G, Durazzo M. Ischemic preconditioning attenuates the oxidant-dependent mechanism of reperfusion cell damage and death in rat liver. *Liver Transplant.* 2002; 8(11):990-999.
249. Wolf SP. Ferrous ion oxidation in presence of ferric ion indicator xylenol orange for measurement of hydroperoxides. *Methods Enzymol* 1994; 233:182-189.
250. Schmekel B, Karlsson SE, Linden M, Sundstrom C, Tegner H, Venge P. Myeloperoxidase in human lung lavage. I. A marker of local neutrophil activity. *Inflammation.* 1990 ;14(4):447-54.
251. Corte ED, Stirpe F. The regulation of rat liver xanthine oxidase. Involvement of thiol groups in the conversion of the enzyme activity from dehydrogenase (type D) into oxidase (type O) and purification of the enzyme. *Biochem J.* 197;126(3):739-45.
252. Baykal A, Kaynaroglu V, Demirpence E, Kilinc K, Sayek I, Sanac Y. Experimental study of the effect of adrenaline tolerance on intestinal ischaemia-reperfusion. *Br J Surg.* 1998 ;85(7):947-50.

253. Link EM. Enzymic pathways involved in cell response to H₂O₂. *Free Radic Res Commun.* 1990;11(1-3):89-97.
254. Hortelano S, Genaro AM, Bosca L. Phorbol esters induce nitric oxide synthase activity in rat hepatocytes. Antagonism with the induction elicited by lipopolysaccharide. *J Biol Chem.* 1992;267(35):24937-24940.
255. Selzner N, Selzner M, Odermatt B, Tian Y, Van Rooijen N, Clavien PA. ICAM-1 triggers liver regeneration through leukocyte recruitment and Kupffer cell-dependent release of TNF-alpha/IL-6 in mice. *Gastroenterology* 2003; 124(3):692-700.
256. Tomiya T, Ogata I, Fujiwara K. Transforming growth factor alpha levels in liver and blood correlate better than hepatocyte growth factor with hepatocyte proliferation during liver regeneration. *Am J Pathol* 1998; 153(3):955-961.
257. Yamada A, Matsumoto K, Sekiguchi K, Kawata S, Matsuzawa Y, Nakamura T. Rapid and sensitive enzyme-linked-immunosorbent assay for measurement of HGF in rat and human tissues. *Biomed. Res* 1995; 16:105-114.
258. Plebani M, Panozzo MP, Basso D, De Paoli M, Biasin R, Infantolino D. Cytokines and the progression of liver damage in experimental bile duct ligation. *Clin Exp Pharmacol Physiol* 1999; 26(4):358-363.
259. Roth-Eichhorn S, Kuhl K, Gressner AM. Subcellular localization of (latent) transforming growth factor beta and the latent TGF-beta binding protein in rat hepatocytes and hepatic stellate cells. *Hepatology* 1998;28(6):1588-1596.
260. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001;25(4):402-408.
261. Selzner N, Selzner M, Tian Y, Kadry Z, Clavien PA. Cold ischemia decreases liver regeneration after partial liver transplantation in the rat: A TNF-alpha/IL-6-dependent mechanism. *Hepatology* 2002;36(4 Pt 1):812-818.
262. Selzner M, Camargo CA, Clavien PA. Ischemia impairs liver regeneration after major tissue loss in rodents: protective effects of interleukin-6. *Hepatology* 1999; 30(2):469-475.
263. Camargo CA Jr, Madden JF, Gao W, Selvan RS, Clavien PA. Interleukin-6 protects liver against warm ischemia/reperfusion injury and promotes hepatocyte proliferation in the rodent. *Hepatology* 1997;26(6):1513-1520.
264. Serafin A, Rosello-Catafau J, Prats N, Xaus C, Gelpi E, Peralta C. Ischemic preconditioning increases the tolerance of fatty liver to hepatic ischemia-reperfusion injury in the rat. *Am J Pathol* 2002;161(2):587-601.
265. Masson S, Daveau M, Hiron M, Lyoumi S, Lebreton JP, Teniere P, Scotte M. Differential regenerative response and expression of growth factors following hepatectomy of variable extent in rats. *Liver* 1999;19(4):312-317.
266. Tomiya T, Ogata I, Yamaoka M, Yanase M, Inoue Y, Fujiwara K. The mitogenic activity of hepatocyte growth factor on rat hepatocytes is dependent upon endogenous transforming growth factor-alpha. *Am J Pathol* 2000;157(5):1693-1701.
267. Okamoto K, Suzuki S, Kurachi K, Sunayama K, Yokoi Y, Konno H, Baba S, Nakamura S. Beneficial effect of deletion variant of hepatocyte growth factor for

- impaired hepatic regeneration in the ischemically damaged liver. *World J Surg* 2002;26(10):1260-1266.
268. Tashiro H, Fudaba Y, Itoh H, Mizunuma K, Ohdan H, Itamoto T, Asahara T. Hepatocyte growth factor prevents chronic allograft dysfunction in liver-transplanted rats. *Transplantation* 2003;76(5):761-765.
269. Ikegami T, Nishizaki T, Uchiyama H, Kakizoe S, Yanaga K, Sugimachi K. Deletion variant of hepatocyte growth factor prolongs allograft survival after liver transplantation in rats. *Surgery* 1999;125(6):602-607.
270. Pesonen EJ, Linder N, Raivio KO, Sarnesto A, Lapatto R, Hockerstedt K, Makisalo H, Andersson S. Circulating xanthine oxidase and neutrophil activation during human liver transplantation. *Gastroenterology* 1998;114(5):1009-1115.
271. Zhong Z, Connor H, Stachlewitz RF, Frankenberg M, Mason RP, Lemasters JJ, Thurman RG. Role of free radicals in primary nonfunction of marginal fatty grafts from rats treated acutely with ethanol. *Mol Pharmacol*. 1997;52(5):912-919.
272. Nagayama M, Katsuramaki T, Kimura H, Isobe M, Meguro M, Matsuno T, Nui A, Itata K. Prediction of graft viability from non-heart-beating donor pigs using hepatic microdialysate hypoxanthine levels. *J Surg Res*. 2002;107:210-218.
273. Net M, Valero R, Almenara R, Rull R, González FJ, Taurá P, Lopez-Boado MA, Deulofeu R, Elena M, Capdevila L, Cabrer C, Visa J, Garcia-Valdecasas JC. Hepatic xanthine levels as viability predictor of livers procured from non-heart-beating donor pigs. *Transplantation* 2001;71(9):1232-1237.
274. Fernández L, Carrasco-Chaumel E, Serafín A, Xaus C, Grande L, Rimola A, Roselló-Catafau J, Peralta C. Is ischemic preconditioning a useful strategy in steatotic liver transplantation?. *Am J Transplant*. 2004;4(6):888-899.
275. Hassoun PM, Yu FS, Zulueta JJ, White AC, Lanzillo JJ. Effect of nitric oxide and cell redox status on the regulation of endothelial cell xanthine dehydrogenase. *Am J Physiol*. 1995;268:L809-L817.
276. Cote CG, Yu FS, Zulueta JJ, Vosatka RJ, Hassoun PM. Regulation of intracellular xanthine oxidase by endothelial-derived nitric oxide. *Am J Physiol*. 1996; 271:L869-L874.
277. Rinaldo JE, Clark M, Parinello J, Shepherd VL. Nitric oxide inactivates xanthine dehydrogenase and xanthine oxidase in interferon-gamma-stimulated macrophages. *Am J Resp Cell Mol Biol*. 1994;11:625-630.
278. Ichimori K, Fukahori M, Nakazawa H, Okamoto K, Nishino T. Inhibition of xanthine oxidase and xanthine dehydrogenase by nitric oxide. Nitric oxide converts reduced xanthine-oxidizing enzymes into the desulfo-type inactive form. *J Biol Chem*. 1999;274:7763-7768.
279. Cavalieri B, Perrelli MG, Aragno M, Ramadori P, Poli G, Cutrin JC. Ischaemic preconditioning modulates the activity of kupffer cells during in vivo reperfusion injury of rat liver. *Cell Biochem Funct*. 2003;21(4):299-305.
280. Schauer RJ, Gerbes AL, Vonier D, op den Winkel M, Fraunberger P, Bilzer M. Induction of cellular resistance against kupffer cell-derived oxidant stress: a novel concept of hepatoprotection by ischemic preconditioning. *Hepatology* 2003;37(2):286-295.

281. Rai RM, Yang SQ, McClain C, Karp CL, Klein AS, Diehl AM. Kupffer cell depletion by gadolinium chloride enhances liver regeneration after partial hepatectomy in rats. *Am J Physiol* 1996;270:G909-G918.
282. Glanemann M, Vollmar B, Nussler AK, Schaefer T, Neuhaus P, Menger MD. Ischemic preconditioning protects from hepatic ischemia/reperfusion-injury by preservation of microcirculation and mitochondrial redox-state. *J Hepatol*. 2003;38(1):59-66.
283. Fernández L, Heredia N, Peralta C, Xaus C, Roselló-Catafau J, Rimola A, Marco A, Serafin A, Deulofeu R, Gelpí E, Grande L. Role of ischemic preconditioning and the portosystemic shunt in the prevention of liver and lung damage after rat liver transplantation. *Transplantation* 2003;75 (12):282-289.
284. Khan D, Robson SC, Hickman R. The effect of tumor necrosis factor antibody on the regenerative response after partial hepatectomy in rats. *Transpl Int* 1994;7 Suppl 1:S181-S182.
285. Ledda-Columbano GM, Curto M, Piga R, Zedda AI, Menegazzi M, Sartori C et al. In vivo hepatocyte proliferation is inducible through a TNF and IL-6-independent pathway. *Oncogene* 1998;17(8):1039-1044.
286. Starkel P, Lambotte L, Sempoux C, De Saeger C, Saliez A, Maiter D, Horsmans Y. After portal branch ligation in the rat, cellular proliferation is associated with selective induction of c-Ha-ras, p53, cyclin E, and Cdk2. *Gut* 2001;49(1):119-130.
287. Fausto N, Laird AD, Webber EM. Liver regeneration. 2. Role of growth factors and cytokines in hepatic regeneration. *FASEB J* 1995; 9(15):1527-1536. Review.
288. Michalopoulos GK, DeFrances MC. Liver regeneration. *Science* 1997;276(5309):60-66. Review.
289. Nakamura T, Tomita Y, Hirai R, Yamaoka K, Kaji K, Ichihara A. Inhibitory effect of transforming growth factor-beta on DNA synthesis of adult rat hepatocytes in primary culture. *Biochem Biophys Res Commun* 1985;133(3):1042-1050.
290. Nakamura T, Arakaki R, Ichihara A. Interleukin-1 beta is a potent growth inhibitor of adult rat hepatocytes in primary culture. *Exp Cell Res* 1988;179(2):488-497.
291. Liu P, Xu B, Spokas E, Lai PS, Wong PY. Role of endogenous nitric oxide in TNF-alpha and IL-1beta generation in hepatic ischemia-reperfusion. *Shock* 2000;13(3):217-223.
292. Kim YM, Talanian RV, Li J, Billiar TR. Nitric oxide prevents IL-1beta and IFN-gamma-inducing factor (IL-18) release from macrophages by inhibiting caspase-1 (IL-1beta-converting enzyme). *J Immunol* 1998;161(8):4122-4128.
293. Ozaki M, Haga S, Zhang HQ, Irani K, Suzuki S. Inhibition of hypoxia/reoxygenation-induced oxidative stress in HGF-stimulated antiapoptotic signaling: role of PI3-K and Akt kinase upon rac1. *Cell Death Differ* 2003;10(5):508-515.
294. Oe S, Hiros T, Fujii H, Yasuchika K, Nishio T, Iimuro Y, Morimoto T, Nagao M, Yamaoka Y. Continuous intravenous infusion of deleted form of hepatocyte growth factor attenuates hepatic ischemia-reperfusion injury in rats. *J Hepatol* 2001;34(6):832-839.
295. Herrera B, Murillo MM, Alvarez-Barrientos A, Beltran J, Fernandez M, Fabregat I. Source of early reactive oxygen species in the apoptosis induced by transforming growth factor-beta in fetal rat hepatocytes. *Free Radic Biol Med* 2004;36(1):16-26.

296. Herrera B, Fernandez M, Alvarez AM, Roncero C, Benito M, Gil J, Fabregat I. Activation of caspases occurs downstream from radical oxygen species production, Bcl-xL down-regulation, and early cytochrome C release in apoptosis induced by transforming growth factor beta in rat fetal hepatocytes. *Hepatology* 2001;34(3):548-556.
297. Beutler B, Milsark IW, Cerami AC. Passive immunization against cachectin/tumor necrosis factor protects mice from lethal effect of endotoxin. *Science* 1985;229(4716):869-871.
298. Ertel W, Scholl FA, Gallati H, et al. Increased release of soluble tumor necrosis factor receptors into blood during clinical sepsis. *Arch Surg* 1994;129(12):1330-1337.