

Author's Proof

Please check your proof carefully and mark all corrections in the appropriate place.

Annotate your corrections on-screen using the Adobe Reader PDF editing tools, save and upload as Author's Proof Corrections.

Login → Journal Tab → Manage Articles → Author's Proof → Select Article → Press Enter
Production Forum → Manuscript → Upload Files

If there are only a few minor corrections and query replies, these can be listed by referring to the specific line number in the proof and be directly communicated in the Discussion Forum of your article.

Login → Journal Tab → Manage Articles → Author's Proof → Select Article → Press Enter
Production Forum → Interactive Discussion → Enter Discussion Forum

Make sure to also answer all the queries thoroughly before submitting your comments, as failing to do so will cause delays.

Do not make any corrections by submitting a new manuscript file.

To ensure fast publication of your paper please return your corrections as soon as possible.

If you have any questions contact the Science Production Office.

- Ensure to proofread the entire article, including figures and tables, captions, equations, citations, and references.
- Double-check the spelling of all author names, accuracy of affiliations and addresses.
- Verify that all the special characters are displayed correctly.
- Be sure that you have obtained permission for any reprinted material.
- Carefully reply to all of the author queries to avoid any production delays.

Author Queries Form

Query No.	Details required	Author's Response
Q1	Kindly confirm if the first name and surname of all the authors have been identified correctly in the front page and citation text.	
Q2	Please ask the following authors to register with Frontiers (at https://www.frontiersin.org/Registration/Register.aspx) if they would like their names on the article abstract page and PDF to be linked to a Frontiers profile. Please ensure to register the authors before submitting the proof corrections. Non-registered authors will have the default profile image displayed by their name on the article page. "Alexandre Martínez-Castillo."	

Query No.	Details required	Author's Response
Q3	<p>We have changed the following references to their respective ones as per the reference list. Please confirm if this is fine.</p> <p>“Dumke, 2006” to “Dumke et al., 2006” “Rooks, 2012” to “Rooks et al., 2012” “Rooks, 2010” to “Rooks et al., 2010” “Picozzi et al., 2012” to “Picozzi et al., 2011” “Yamamoto, 2003” to “Yamamoto et al., 2003” “Aersten, 2005” to “Aertsen et al., 2005” “Kimmit et al., 2000” to “Kimmitt et al., 2000” “Mauro et al., 2011” to “Mauro and Koudelka, 2011.” “Aersten et al., 2005” to “Aertsen et al., 2005” “Rode, 2011” to “Rode et al., 2011” “Yue, 2012” to “Yue et al., 2012” “Baquero and Tobes, 2013” to “Baquero and Tobes, 2003” “Loukadis et al., 2008” to “Loukiadis et al., 2008” “Mann et al., 1997” to “Mann and Slauch, 1997.” “Van Reis and Zidney, 2001” to “Van Reis and Zydne, 2001” “Johansen” to “Johannessen et al., 2005.”</p>	
Q4	<p>We have changed “Muniesa et al., 2004” to “Muniesa et al., 2004a,b” here and elsewhere inside the text as per the reference list. Please confirm if this is fine.</p>	
Q5	<p>Please add the following references to the reference list. “Muniesa and Jofre, 1998, 1999; Tanji et al., 2002; Muniesa et al., 1991; Imamovic et al., 2011; USDA, 2011.”</p>	
Q6	<p>Please provide the volume number and page range for the following references. “Allué-Guardia et al., 2014; Langsrud et al., 2013; Trevisani et al., 2014.”</p>	
Q7	<p>Please provide doi for the following references. “Anonymous, 2005; Erickson and Doyle, 2007; Gould et al., 2009; Köhler et al.,2000; Pradel et al., 2000.”</p>	
Q8	<p>Please confirm the deletion of second occurrence of “Muniesa and Jofre, 2000” in the reference list as it is appearing twice in the list.</p>	
Q9	<p>Please provide the page range for “Paton and Paton, 2003.”</p>	
Q10	<p>Please cite “Zhu, 2009” inside the text.</p>	
Q11	<p>Please provide the name of the editors and city name for “Jofre, 2007.”</p>	



Implications of free Shiga toxin-converting bacteriophages occurring outside bacteria for the evolution and the detection of Shiga toxin-producing *Escherichia coli*

Alexandre Martínez-Castillo and Maite Muniesa*

Department of Microbiology, Faculty of Biology, University of Barcelona, Barcelona, Spain

Edited by:

Eelco Franz, Centre for Infectious Disease Control, Netherlands

Reviewed by:

Eelco Franz, Centre for Infectious Disease Control, Netherlands
Stefano Morabito, Istituto Superiore di Sanità, Italy

*Correspondence:

Maite Muniesa, Department of Microbiology, Faculty of Biology, University of Barcelona, Av. Diagonal 643, 08028 Barcelona, Spain
e-mail: mmuniesa@ub.edu

In this review we highlight recent work that has increased our understanding of the distribution of Shiga toxin-converting phages that can be detected as free phage particles, independently of Shiga toxin-producing bacteria (STEC). Stx phages are a quite diverse group of temperate phages that can be found in their prophage state inserted within the STEC chromosome, but can also be found as phages released from the cell after activation of their lytic cycle. They have been detected in extraintestinal environments such as water polluted with feces from humans or animals, food samples or even in stool samples of healthy individuals. The high persistence of phages to several inactivation conditions makes them suitable candidates for the successful mobilization of *stx* genes, possibly resulting in the genes reaching a new bacterial genomic background by means of transduction, where ultimately they may be expressed, leading to Stx production. Besides the obvious fact that Stx phages circulating between bacteria can be, and probably are, involved in the emergence of new STEC strains, we review here other possible ways in which free Stx phages could interfere with the detection of STEC in a given sample by current laboratory methods and how to avoid such interference.

Keywords: bacteriophages, Shiga toxin, STEC, environment, transduction

Shiga toxin-producing *Escherichia coli* (STEC) are important food-borne pathogens and represent a challenge for the scientific community. Effort has been devoted to developing methods for its isolation, detection and identification, to guarantee the quality of products and the health of consumers. Many studies have aimed at gaining an understanding of the mobility of the important gallery of virulence factors that compose the virulome of the *E. coli* pathogenic strains. The mobility of these factors among strains is a key factor for the genotypic and phenotypic variability of this bacterium, and hinders the process of determining what should be considered a true pathogen and how to detect it effectively (Erickson and Doyle, 2007; Karch et al., 2012; Melton-Celsa et al., 2012).

Thus, the methods for *E. coli* detection have focused on those factors that are clearly related with virulence and, particularly, those that are closely related with high incidence and severity of human infections. Among these factors, the Shiga toxin (Stx) is considered one of the most significant and, if not the only one determining pathogenicity, the one that leads to the most undesirable complications of the infection, such as HUS (Tarr et al., 2005).

Stx is a good example of a mobile virulence factor, since the genes encoding for this toxin are located in the genome of temperate bacteriophages (Newland et al., 1985; Huang et al., 1987; O'Brien et al., 1989). Many phages simply multiply by infecting bacteria, killing the host by lysis and releasing new phages. Temperate phages, however, adopt a benign relationship with their hosts, called lysogeny, which allows an attachment of phage

DNA to a bacterial chromosome. For Stx phages, once a phage integrates its genome within the bacterial genome, the bacteria acquires the *stx* gene and with it the capacity to express the toxin; thereby becoming a STEC. Temperate phages can revert from the lysogenic state by entering the lytic cycle, mostly due to environmental conditions or exogenous factors. The lytic pathway starts by generating multiple copies of its genome, causing an increase in the Stx produced by the cell during the process (Neely and Friedman, 1998; Wagner et al., 2001; Tyler et al., 2004). By the expression of the phage genome, new phage capsids are generated, assembled with the phage DNA and, once formed, phage particles are released from the cell by lysis. Once released, phage particles remain free outside the bacterial cell, and here is where our story starts.

The major question we address is: what happens to phages outside the cell and to what extent does their occurrence as free particles play a role in the evolution of new STEC strains?

Phages are rather simple particles and Stx phages are no exception. Their persistence outside the cell is guaranteed by their ability to circumvent natural and artificial inactivation processes (as reviewed in the next section), many of which would inactivate their bacterial host. It is the fact that the life cycle of phages switches to the lytic state that allows phages to be released once the cell has been threatened by factors that activate its SOS response (Muhldorfer et al., 1996; Kimmitt et al., 2000; Köhler et al., 2000; Yamamoto et al., 2003; Aertsen et al., 2005; Tushima et al., 2007; Pacheco and Sperandio, 2009).

Stx PHAGES IN FECALLY POLLUTED WATERS

Several reports have shown the occurrence of free Stx phages in water environments with fecal pollution or directly on homogenates of fecal samples (Table 1). Most studies have only focused on the detection of Stx phages in general or Stx2 phages in particular, and only a few have evaluated Stx1 phages (Dumke et al., 2006; Yan et al., 2011; Rooks et al., 2012). Those studies show a clear predominance of Stx2 phages over Stx1 phages and, despite the fact that there is still a lack of information on the abundance of Stx1 phages in the environment, the results correlate with reported data on the predominance of stx2 phages over Stx1 phages in lysates induced from environmental *E. coli* isolates (Muniesa et al., 2004a,b; Garcia-Aljaro et al., 2009; Yan et al., 2011).

Regardless of the type of toxin variant, the high occurrence of free Stx phages detected suggests that Stx phages could do more

than provide intestinal bacteria with a new genetic character. Although a certain level of human fecal contamination is always observed in environments where free Stx phages have been detected (Muniesa and Jofre, 1999; Tanji et al., 2003; Muniesa et al., 2004a,b; Dumke et al., 2006; Imamovic et al., 2010; Rooks et al., 2010, 2012) (Table 1), there is no total correlation between Stx phage occurrence and fecal pollution, as shown by comparison with fecal indicators (Dumke et al., 2006; Imamovic et al., 2010). This lack of correlation suggests that the fecal origin is the main but not necessarily the sole source of Stx phages. Accordingly, the description of Stx2-positive extraintestinal *E. coli* strains (Wester et al., 2013) supports the suspicion that *stx* is not restricted to a fecal origin. Stx phages are not limited to waters containing human contamination and they have also been found to be highly prevalent when analyzing wastewater from animals (Imamovic et al., 2010; Yan et al., 2011). Water with lower levels of fecal pollution, such as river water, also shows

Table 1 | Occurrence and abundance of free Stx bacteriophages in diverse environments.

Sample	Country	Detection of Stx phages			Detection method	References
		Stx1 phages	Stx2 phages	Abundance		
Human wastewater	Spain	–	Positive (I)	10 PFU.ml ⁻¹	MPN+PCR	Muniesa and Jofre, 1998
Human wastewater	Germany,	–	Positive	≥ 1 PFU.ml ⁻¹	PCR	Muniesa and Jofre, 2000
	Austria,		Positive	>0.1PFU.ml ⁻¹		
	France		Positive	≥ 10 PFU.ml ⁻¹		
	Ireland		Positive	≥ 10 PFU.ml ⁻¹		
	South Africa		Positive	≥ 10 PFU.ml ⁻¹		
	New		Positive	≥ 10 PFU.ml ⁻¹		
	Zaeland					
Human wastewater	Japan	–	Positive (I)	–	PCR	Tanji et al., 2002
Human wastewater	Spain	–	Positive (I)	4.24 log ₁₀ PFU.ml ⁻¹	Plaque count+PCR	Muniesa et al., 2004a,b
Cattle wastewater	Spain	–	Positive (I)	4.45 log ₁₀ PFU.ml ⁻¹	Plaque count+PCR	Muniesa et al., 2004a,b
Human wastewater	Germany	Positive (I)	Positive (I)	0.34 PFU.ml ⁻¹ stx ₁ 3.4 PFU.ml ⁻¹ stx ₂	Plaque count PCR estimation	Dumke et al., 2006
Human treated waste water	Germany	Negative	Positive (I)	–	Plaque count+PCR	Dumke et al., 2006
River water	Germany	Positive (I)	Positive (I)	–	Plaque count+PCR	Dumke et al., 2006
Human wastewater	UK	Negative	Positive (I)	9.41 log ₁₀ PFU.ml ⁻¹	Plaque count+PCR	Rooks et al., 2010
Human wastewater	UK	Negative	Positive	2.39 log ₁₀ GC.ml ⁻¹	qPCR	Rooks et al., 2010
Human wastewater	Spain	–	Positive	1.37 log ₁₀ GC.ml ⁻¹	qPCR	Imamovic et al., 2010
Cattle wastewater	Spain	–	Positive	2.77 log ₁₀ GC.ml ⁻¹	qPCR	Imamovic et al., 2010
Pig wastewater	Spain	–	Positive	4.59 log ₁₀ GC.ml ⁻¹	qPCR	Imamovic et al., 2010
Poultry wastewater	Spain	–	Positive	1.11 log ₁₀ GC.ml ⁻¹	qPCR	Imamovic et al., 2010
Cattle feces	Spain	–	Positive	2.32 log ₁₀ GC.g ⁻¹	qPCR	Imamovic et al., 2010
Beef	Spain	–	Positive (I)	4.10 log ₁₀ GC.g ⁻¹	qPCR	Imamovic and Muniesa, 2011
Salad	Spain	–	Positive (I)	3.36 log ₁₀ GC.g ⁻¹	qPCR	Imamovic and Muniesa, 2011
Swine feces	China	Positive (I)	Positive (I)	–	Plaque count+PCR	Yan et al., 2011
Wastewater	UK	–	Positive (I)	–	Lysogen isolation	Rooks et al., 2012
Human feces	Spain	–	Positive (I)	4.41 log ₁₀ GC.g ⁻¹	qPCR	Martinez-Castillo et al., 2013

GC, gene copy; PFU, plaque forming unit; MPN, most probable number. (I) those studies reporting infectious Stx phages.

the presence of Stx phages (Muniesa et al., 1991; Dumke et al., 2006).

Stx PHAGES IN FECES

If fecally polluted waters present Stx phages, and assuming that Stx phages could have been generated in the intestinal gut, the question that remains to be answered is whether the phages are released into wastewaters by induction from STEC strains present in these environments, or whether free Stx phages are directly excreted through feces. Recent reports indicate that free infectious (those able to infect and propagate in a host strain), Stx2 phages are present in 62% of human feces (Martínez-Castillo et al., 2013). Sequencing of the PCR amplimers suggested that subtypes *stx*_{2a}, *stx*_{2c}, and *stx*_{2d} were the most frequently detected (Martínez-Castillo et al., 2013). From the positive samples, 90% showed infectious Stx phages. Two facts are to be highlighted from that study: there is not a high incidence of STEC in the area of this study; and the stool samples were taken from healthy individuals without reported gastrointestinal symptoms, and that STEC were not detected in or isolated from these samples.

Stx PHAGES IN FOOD

Assuming that Stx phages are excreted in feces, they must have been ingested, either as free phages or as STEC bacteria that were later induced, once in the intestinal tract. If they are ingested as free phages, then they must be present in food or drinking water, assuming that these are free of significant levels of bacteria. The occurrence of infectious Stx2 phages in commercial samples of beef and salad (Imamovic et al., 2011), that were acceptable for process hygiene microbiological criteria under EU regulation (Anonymous, 2005), highlights the gap that exists in legislation regarding the presence of phages in food samples.

The fact that phages detected in food were infectious is interesting, considering the likelihood of transduction in food matrices at different pH and temperatures (Imamovic et al., 2009) as well as under dairy processing conditions (Picozzi et al., 2011). This fact is of special relevance considering the observations that many food-processing conditions, such as thermal treatment, high hydrostatic pressures, or UV or other irradiation (Yamamoto et al., 2003; Aertsen et al., 2005; Yue et al., 2012), and the addition of certain compounds during the food production, such as chelating agents (Imamovic et al., 2011), salt (Harris et al., 2012) or antimicrobials (Kimmitt et al., 2000), not only fail to inactivate of the Stx phages, but can enhance Stx phage induction from their STEC hosts.

PERSISTENCE OF Stx PHAGES

Generally speaking, phages could persist in different environments or under disinfection processes or inactivation conditions (Dee and Fogleman, 1992; Durán et al., 2002; Jofre, 2007; Lee and Sobsey, 2011). In habitats in which the host bacteria are alien, as fecal bacteria are in the environment, it is likely that phages persist much better than the bacteria (Ogunseitan et al., 1990; Muniesa et al., 1999; Durán et al., 2002). Because of their structural characteristics, their persistence in the environment is high, and these survival capabilities make bacteriophages especially

suitable for movement and gene transfer between different biomes.

Certain morphologies can be considered more persistent than others, tailed phages being the ones showing higher persistence (Muniesa et al., 1999; Prigent et al., 2005; Lin et al., 2010). Notably, these are the most abundant in different water environments (Prigent et al., 2005; Lin et al., 2010). Stx phages mostly belong to the *Siphoviridae* and *Podoviridae* morphological types (Rietra et al., 1989; Muniesa et al., 2004a,b; Beutin et al., 2012), showing similar persistence to that of other groups of phages (Muniesa et al., 1999; Dumke et al., 2006; Allué-Guardia et al., 2014) and higher persistence than STEC (Muniesa et al., 1999; Mauro and Koudelka, 2011; Rode et al., 2011; Allué-Guardia et al., 2014). Stx phages have been shown to be very stable under food-processing conditions (Yamamoto et al., 2003; Aertsen et al., 2005; Imamovic et al., 2009; Rode et al., 2011; Harris et al., 2012; Yue et al., 2012; Langsrud et al., 2013; Allué-Guardia et al., 2014). In contrast, they do not seem to persist so well under strict treatments such as that linked to the compost model (Johannessen et al., 2005). Nevertheless, their persistence, as for other phages, enhances their mobility between the different environments (feces-food-water) (Mauro and Koudelka).

Another question is whether the STEC hosts could survive the time necessary to allow phage release before being killed by the treatment applied, or if phages, once released, could remain infectious. Infectivity of Stx phages has been demonstrated after exposure to certain conditions, and some Stx phages remain able to generate lysogens, hence to transduce *stx* (Muniesa and Jofre, 1998; Imamovic et al., 2009; Rode et al., 2011; Yue et al., 2012). The capacity to remain infectious after being subjected to a given condition will determine the chances of *stx* transduction, the real threat in the generation of new STEC strains.

INTERFERENCE OF Stx PHAGES IN STEC DETECTION

The identification of STEC by culture methods in food or clinical samples is advisable to confirm the presence of the pathogen and to further characterize it. However, sometimes strain recovery is not possible because of the low number of STEC cells in a specific sample, the fact that cells could be stressed or in a non-culturable state, or because of the interference of accompanying microbiota, particularly other *E. coli* strains that could confound the detection of the pathogen even when using a specific culture medium. Culture detection of STEC is, in addition, a time-consuming method that hinders early diagnosis of STEC.

Early diagnosis of STEC infection in humans is nevertheless critical for the treatment of the disease, particularly because of the contraindication for treating STEC using antimicrobial agents, and the intense supportive care needed if renal failure occurs (Wong et al., 2000). The need for fast detection and the associated difficulties of STEC isolation from stools in patients treated with antibiotics makes the use of fast and robust molecular methods advisable for STEC detection.

Current laboratory methods for STEC detection, some standardized and approved by national legislation, include PCR-based techniques: either end-point or real-time PCR (Paton and Paton, 2003; Perelle et al., 2004; Gould et al., 2009; Kagkli et al., 2011; Bibbal et al., 2014). Recently, ISO 13136:2012 (Anonymous,

2012), which uses real-time PCR as the reference technology for the detection of the virulence and serogroup-associated genes, has been included in the amended EU regulation for microbiological criteria for sprouts and the sampling rules for poultry carcasses and fresh poultry meat (European commission regulation No. 209/2013). With the same aim in mind, molecular techniques, many based on the information obtained by the use of next-generation sequencing (NGS) methods, are being explored and advised (Baquero and Tobes, 2003; Underwood et al., 2013).

As indicated above, the genomic plasticity of STEC represents a challenge for discriminating pathogenic strains from other non-pathogenic *E. coli* strains present in a given sample. Thus, PCR-based molecular methods are mainly focused on genes related with virulence and on genes that identify a specific serotype, such as *rfb* genes encoding different O antigens (Maurer et al., 1999). Identification of the serotype could be useful for epidemiological purposes and, generally, virulent strains belong to one of the serogroups that account for the non-O157 serotype, including O26, O45, O103, O111, O121, and O145, plus O157:H7 (USDA, 2011). Lessons learnt from the last outbreak in Germany (Karch et al., 2012) forced researchers to include O104:H4 on the list, and showed that identification of serotype, albeit very useful when treating known pathotypes or for epidemiological purposes, is of limited value for dealing with a newly emerging strain. If STEC

isolation is not possible, identifying virulence genes, alone or preferably in combination, appears to be the best approach for assessing the presence of STEC in a sample.

However, when dealing with molecular methods applied to a sample, the presence of free Stx phages, and also other phages that could harbor virulence genes, represents a challenge for the use of PCR-based and NGS methodologies. Moreover, when the protocols include a previous step of selective enrichment of the target bacteria, this step could also maintain or even propagate bacteriophages. This is even more significant if we assume that the methods used for bacterial DNA extraction in these complex matrices will also extract phage DNA (Paton and Paton, 2003; Monday et al., 2007; Grys et al., 2009; ISO 13136:2012). Therefore, any virulence genes present in phage DNA, notably *stx* genes, but also other virulence genes reported in prophages (e.g., *cdt*, *cif*, etc.) (Asakura et al., 2007; Loukiadis et al., 2008) would generate amplicons (PCR) or reads (NGS) that will be interpreted as belonging to STEC, while it could be that they originated in phages. Since in STEC these genes belong to prophages, and therefore are flanked by phage sequences, there is no easy way to distinguish whether the target detected is a bacteria or a phage. If it is a phage, the threat of virulence in humans must be discussed, but it obviously has reduced potential for virulence and would not be enough to raise the alarm.

Table 2 | Frequency of STEC isolation by various methods in samples showing positive result for *stx* (either *stx*₁ or *stx*₂) by PCR methods.

	No of <i>stx</i> -positive samples by PCR	No of <i>stx</i> -positive samples by PCR with STEC isolation	Percentage of PCR <i>stx</i> -positive samples with STEC isolation (%)	References
Human stools (healthy slaughterhouse workers)	90	8	8.9	Hong et al., 2009
Humans stools (asymptomatic)	196	47	24	Stephan et al., 2000
Human stools (volunteers)	21	1	4.8	Urdahl et al., 2012
Human stools (hospital)	150	1	0.67	Urdahl et al., 2012
Human stools (hospital)	20*	10*	50	Buchan et al., 2013
Children stools (hospital)	21	5	24	Vallièrès et al., 2013
Children stools (hospital)	19	10	52.6	Pradel et al., 2000
Cattle feces	145	80	55.2	Fremaux et al., 2006
Cattle feces	154	67	43.5	Rogerie et al., 2001
Cattle feces	417	18	4.3	Hofer et al., 2012
Cattle feces	330	162	49.0	Pradel et al., 2000
Bovine hides	301	25	8.3	Monaghan et al., 2012
Bovine carcasses	122	5	4.1	Monaghan et al., 2012
Bovine carcasses	77	16	20.8	Breum and Boel, 2010
Bovine carcasses	91	16	17.6	Rogerie et al., 2001
Cattle environment	179	38	21.2	Fremaux et al., 2006
Beef meat	47	16	34.0	Pradel et al., 2000
Dairy buffalo (feces and milk)	56	20	35.7	Beraldo et al., 2014
Milk (bulk)	32	1	3.1	Trevisani et al., 2014
Milk (filters)	68	7	10.3	Trevisani et al., 2014
Cheese	60	5	8.3	Pradel et al., 2000
Healthy pigs feces	255	62	24.3	Meng et al., 2014
Swine feces	484	196	40.5	Fratamico et al., 2004
Meat products	36	8	22.2	Díaz-Sánchez et al., 2012

* Calculated from the % of positive samples.

In addition to those well-characterized phages encoding virulence genes in STEC, many lytic phages could mobilize bacterial genes by generalized transduction. In transduction, after random packaging of bacterial DNA fragments, the genetic material can be mobilized by a phage from a donor bacterium and inserted into a recipient bacterium when it becomes infected by the phage particle. Despite a lack of clear evidence of generalized transduction relating to STEC, any bacterial gene can be mobilized via generalized transduction, including chromosome fragments, though plasmids, transposons and insertion elements, and examples in enterobacteria can also be found (Mann and Schlauch, 1997; Schmieger and Schicklmaier, 1999).

Generalized transduction is considered a rare event (Bushman, 2002); however, many approaches for STEC detection based on genomic techniques envisage preliminary steps for the selective or non-selective enrichment of the target microorganism (Hussein and Bollinger, 2008). Enrichment to propagate bacteria can cause the propagation of any sort of phages present in a sample too, a fact that, firstly, would cause a bias in the population of enriched bacteria (Muniesa et al., 2005), and secondly, by increasing the number of phages, would also theoretically cause an increase in the frequency of generalized transduction. As an example to illustrate this point, the protocols for P1 generalized transduction include a propagation step between the donor and the phage to increase the number of generalized transducing particles (Thomason et al., 2007).

Since Shiga toxin is the main virulence threat in STEC, there are many examples of *stx* positive PCR results from samples showing negative detection of culturable STEC. The percentage of samples showing STEC isolates among PCR *stx*-positive samples (either stools of food samples) is in many reports no higher than 50% (Table 2). In clinical stools samples this could be attributable to previous treatment with antimicrobial agents or to a disease being diagnosed late in its course. In food samples it could also be attributable to the presence of non-culturable microorganisms or the presence of bacteria in very low numbers. However, considering the presence of phages as described above, their role in these positive PCR detections or their involvement in sequences generated by NGS cannot be excluded.

AVOIDING PHAGE INTERFERENCE

The issue that arises is how to avoid phage interference. In theory, several methods could be applied to eliminate phages from the equation. These are mostly based on dismissing the phage population from the sample without interfering with the bacterial population and avoiding complicated steps in the methodology. Some protocols use a centrifugation step after the enrichment culture, and only the pellet containing bacteria is used for DNA extraction (Feng et al., 2011). This method would reduce the number of phages present in the supernatant of the enrichment; though to what extent, has not been tested. Still, phages present in the original samples would have propagated and would still be present in the bacterial pellet.

Filtration has also been proposed to separate phages from the sample and has been widely used to purify phage stocks or phages from a sample before counting (Brock, 1983; EPA, 2000). Microfiltration is mostly applied to retain viruses using

the filter and clarify the sample (Van Reis and Zydney, 2001; Saxena et al., 2008), while here the aim is to eliminate phages by passing them through the filter but retaining the bacteria. The selection of the most efficient membrane, in terms of pore size and chemical structure, would be critical with this purpose in mind. The use of 0.22 or 0.45 μm low protein binding membranes that do not retain phages would be advisable. Among these, polyvinylidene fluoride (PVDF), polyethersulfone (PES) or cellulose ester membranes saturated with beef extract have been proposed (Anonymous, 2000; EPA, 2000; Papageorgiou et al., 2000; Mocé-Llivina et al., 2003), and PVDF membranes have been shown to reduce by 2–3 \log_{10} the phages present in a bacterial enrichment culture (Muniesa et al., 2005). Complete elimination of all phages present in the samples will probably not be accomplished only by adding a single filtration step, but optimization of this approach would reduce the phage population enough to avoid interference in the molecular detection methods without adding complicated steps to the already-established protocols and without reducing the bacterial population. However, it must be borne in mind that filtration would turn out to be more useful when treating aqueous samples or clear homogenates of solid samples and with a variable efficiency depending on the complexity of the matrix, with lower recoveries expected when it is applied to solid or mixed samples that could clog the membranes.

ACKNOWLEDGMENTS

This study was supported by the Spanish Ministry of Education and Science (AGL2012-30880) and the Generalitat de Catalunya (2009SGR1043) and by the Spanish Reference Network of Biotechnology (XeRBA). Alex Martínez-Castillo has a grant from the Spanish Ministry.

REFERENCES

- Aertsen, A., Faster, D., and Michiels, C. W. (2005). Induction of Shiga toxin-converting prophage in *Escherichia coli* by high hydrostatic pressure. *Appl. Environ. Microbiol.* 71, 1155–1162. doi: 10.1128/AEM.71.3.1155-1162.2005
- Allué-Guardia, A., Martínez-Castillo, A., and Muniesa, M. (2014). Persistence of infectious Stx bacteriophages after disinfection treatments. *Appl. Environ. Microbiol.* doi: 10.1128/AEM.04006-13
- Anonymous. (2000). *ISO 10705-2: Water Quality. Detection and Enumeration of Bacteriophages -Part 2: Enumeration of Somatic Coliphages*. Geneva: International Organisation for Standardisation.
- Anonymous. (2012). *ISO 13136:2012. Microbiology of Food and Animal Feed – Real-time Polymerase Chain Reaction (PCR)-Based Method for the Detection of Food-borne Pathogens – Horizontal Method for the Detection of Shiga Toxin-producing *Escherichia coli* (STEC) and the Determination of O157, O111, O26, O103 and O145 Serogroups*. Geneva: International Organisation for Standardisation.
- Anonymous. (2005). Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs. *Official J. Eur. Commun.* 338, 11–25.
- Asakura, M., A. Hinenoya, M. S. Alam, K. Shima, S. H., Zahid, L. Shi, N., et al. (2007). An inducible lambdoid prophage encoding cytolethal distending toxin (Cdt-I) and a type III effector protein in enteropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 14483–14488. doi: 10.1073/pnas.0706695104
- Baquero, F., and Tobes, R. (2003). Bloody coli: a gene cocktail in *Escherichia coli* O104:H4. *MBio* 4, e00066–13. doi: 10.1128/mBio.00066-13
- Beraldo, L. G., Borges, C. A., Maluta, R. P., Cardozo, M. V., Rigobelo, E. C., and de Ávila, F. A. (2014). Detection of Shiga toxicogenic (STEC) and enteropathogenic (EPEC) *Escherichia coli* in dairy buffalo. *Vet. Microbiol.* 170, 162–166. doi: 10.1016/j.vetmic.2014.01.023
- Beutin, L., Hammerl, J. A., Strauch, E., Reetz, J., Dieckmann, R., Kelner-Burgos, Y., et al. (2012). Spread of a distinct Stx2-encoding phage prototype among

- 571 *Escherichia coli* O104:H4 strains from outbreaks in Germany, Norway, and
572 Georgia. *J. Virol.* 86, 10444–10455. doi: 10.1128/JVI.00986-12
- 573 Bibbal, D., Loukiadis, E., Kérourédan, M., Peytavin de Garam, C., Ferré, F., Cartier,
574 P., et al. (2014). Intimin gene (eae) subtype-based real-time PCR strategy for
575 specific detection of shiga toxin-producing *Escherichia coli* serotypes O157:H7,
576 O26:H11, O103:H2, O111:H8, and O145:H28 in cattle feces. *Appl. Environ.
577 Microbiol.* 80, 1177–1184. doi: 10.1128/AEM.03161-13
- 578 Breum, S. O., and Boel, J. (2010). Prevalence of *Escherichia coli* O157 and verocytotoxin
579 producing *E. coli* (VTEC) on Danish beef carcasses. *Int. J. Food Microbiol.*
580 141, 90–96. doi: 10.1016/j.ijfoodmicro.2010.03.009
- 581 Brock, T. D. (1983). *Membrane Filtration: A User's Guide and Reference Manual.*
582 Madison, WI: Science Technology Springer-Verlag.
- 583 Buchan, B. W., Olson, W. J., Pezewski, M., Marcon, M. J., Novicki, T., Uphoff,
584 T. S., et al. (2013). Clinical evaluation of a real-time PCR assay for identification
585 of *Salmonella*, *Shigella*, *Campylobacter* (*Campylobacter jejuni* and *C. coli*),
586 and shiga toxin-producing *Escherichia coli* isolates in stool specimens. *J. Clin.
587 Microbiol.* 51, 4001–4007. doi: 10.1128/JCM.02056-13
- 588 Bushman, F. (2002). *Lateral DNA Transfer.* New York, NY: Cold Spring Harbor
589 Laboratory.
- 590 Dee, S. W., and Fogleman, J. C. (1992). Rates of inactivation of waterborne
591 coliphages by monochloramine. *Appl. Environ. Microbiol.* 58, 3136–3141.
- 592 Díaz-Sánchez, S., Sánchez, S., Sánchez, M., Herrera-León, S., Hanning, I., and Vidal,
593 D. (2012). Detection and characterization of Shiga toxin-producing *Escherichia coli*
594 in game meat and ready-to-eat meat products. *Int. J. Food Microbiol.* 160,
595 179–182. doi: 10.1016/j.ijfoodmicro.2012.09.016
- 596 Dumke, R., Schroter-Bobsin, U., Jacobs, E., and Roske, I. (2006). Detection
597 of phages carrying the Shiga toxin 1 and 2 genes in waste water and
598 river water samples. *Lett. Appl. Microbiol.* 42, 48–53. doi: 10.1111/j.1472-
599 765X.2005.01809.x
- 600 Durán, A. E., Muniesa, M., Méndez, X., Valero, F., Lucena, F., and Jofre, J. (2002).
601 Removal and inactivation of indicator bacteriophages in fresh waters. *J. Appl.
602 Microbiol.* 92, 338–347. doi: 10.1046/j.1365-2672.2002.01536.x
- 603 Environmental Protection Agency (EPA). (2000). *Method 1602: Male-specific
604 (F+) and Somatic Coliphages in Water by Single Agar Layer (SAL) Procedure.*
605 Washington DC: Office of Water. United States Environmental Protection
606 Agency.
- 607 Erickson, M. C., and Doyle, M. P. (2007). Food as a vehicle for transmission of
608 Shiga toxin-producing *Escherichia coli*. *J. Food Prot.* 70, 2426–2449.
- 609 Feng, P., Weagant, S. D., and Jinneman, K. (2011). “Diarrheagenic
610 *Escherichia coli*,” in *Inited States Food and Drug Administration. Bacteriological
611 Analytical Manual.* Available online at: [http://
612 www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm070080.htm](http://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm070080.htm)
- 613 Fratamico, P. M., Bagi, L. K., Bush, E. J., and Solow, B. T. (2004). Prevalence and
614 characterization of shiga toxin-producing *Escherichia coli* in swine feces recovered
615 in the National Animal Health Monitoring System's Swine 2000 study. *Appl. Environ. Microbiol.* 70,
616 7173–7178. doi: 10.1128/AEM.70.12.7173-71
617 78.2004
- 618 Fremaux, B., Raynaud, S., Beutin, L., and Rozand, C.V. (2006). Dissemination and
619 persistence of Shiga toxin-producing *Escherichia coli* (STEC) strains on French
620 dairy farms. *Vet. Microbiol.* 117, 180–191. doi: 10.1016/j.vetmic.2006.04.030
- 621 Garcia-Aljaro, C., Muniesa, M., Jofre, J., and Blanch, A. R. (2009). Genotypic and
622 phenotypic diversity among induced, stx2-carrying bacteriophages from environ-
623 mental *Escherichia coli* strains. *Appl. Environ. Microbiol.* 75, 329–336. doi:
624 10.1128/AEM.01367-08
- 625 Gould, L. H., Bopp, C., Strockbine, N., Atkinson, R., Baselski, V., Body, B., et al.
626 (2009). Recommendations for diagnosis of shiga toxin-producing *Escherichia coli*
627 infections by clinical laboratories. *MMWR Recomm. Rep.* 58, 1–14.
- 628 Grys, T. E., Sloan, L. M., Rosenblatt, J. E., and Patel, R. (2009). Rapid and sensitive
629 detection of Shiga toxin-producing *Escherichia coli* from non-enriched stool
630 specimens by real-time PCR in comparison to enzyme immunoassay and culture. *J. Clin. Microbiol.* 47,
631 2008–2012. doi: 10.1128/JCM.02013-08
- 632 Harris, S. M., Yue, W. F., Olsen, S. A., Hu, J., Means W. J., McCormick, R. J., et al.
633 (2012). Salt at concentrations relevant to meat processing enhances Shiga toxin
634 2 production in *Escherichia coli* O157:H7. *Int. J. Food Microbiol.* 159, 186–192.
635 doi: 10.1016/j.ijfoodmicro.2012.09.007
- 636 Hofer, E., Stephan, R., Reist, M., and Zweifel, C. (2012). Application of a real-time
637 PCR-based system for monitoring of O26, O103, O111, O145 and O157 Shiga
638 toxin-producing *Escherichia coli* in cattle at slaughter. *Zoonoses Public Health* 59,
639 408–415. doi: 10.1111/j.1863-2378.2012.01468.x
- 640 Hong, S., Oh, K. H., Cho, S. H., Kim, J. C., Park, M. S., Lim, H. S., et al. (2009).
641 Asymptomatic healthy slaughterhouse workers in South Korea carrying Shiga
642 toxin-producing *Escherichia coli*. *FEMS Immunol. Med. Microbiol.* 56, 41–47.
643 doi: 10.1111/j.1574-695X.2009.00545.x
- 644 Huang, A., Friesen, J., and Brunton, J. L. (1987). Characterization of a bacterio-
645 phage that carries the genes for production of Shiga-like toxin 1 in *Escherichia coli*.
646 *J. Bacteriol.* 169, 4308–4312.
- 647 Hussein, H. S., and Bollinger, L. M. (2008). Influence of selective media
648 on successful detection of Shiga toxin-producing *Escherichia coli* in food,
649 fecal, and environmental samples. *Foodborne Pathog. Dis.* 5, 227–244. doi:
650 10.1089/fpd.2008.0081
- 651 Imamovic, L., Ballesté, E., Jofre, J., and Muniesa, M. (2010). Quantification
652 of Shiga toxin-converting bacteriophages in wastewater and in fecal samples
653 by real-time quantitative PCR. *Appl. Environ. Microbiol.* 76, 5693–5701. doi:
654 10.1128/AEM.00107-10
- 655 Imamovic, L., Jofre, J., Schmidt, H., Serra-Moreno, R., and Muniesa, M. (2009).
656 Phage-mediated Shiga toxin 2 gene transfer in food and water. *Appl. Environ.
657 Microbiol.* 75, 1764–1768. doi: 10.1128/AEM.02273-08
- 658 Imamovic, L., and Muniesa, M. (2011). Quantification and evaluation of infec-
659 tivity of shiga toxin-encoding bacteriophages in beef and salad. *Appl. Environ.
660 Microbiol.* 77, 3536–3540. doi: 10.1128/AEM.02703-10
- 661 Jofre, J. (2007). “Indicators of waterborne enteric viruses in human viruses in water”,
662 in *Series Perspectives in Medical Virology*, Vol. 17. (Elsevier), 227–249. doi:
663 10.1016/S0168-7069(07)17011-7
- 664 Johannessen, G. S., James, C. E., Allison, H. E., Smith, D. L., Saunders, J.
665 R., and McCarthy, A. J. (2005). Survival of a Shiga toxin-encoding bacteriophage
666 in a compost model. *FEMS Microbiol. Lett.* 245, 369–375. doi:
667 10.1016/j.femsle.2005.03.031
- 668 Kagkli, D. M., Weber, T. P., Van den Bulcke, M., Folloni, S., Tozzoli, R., Morabito,
669 S., et al. (2011). Application of the modular approach to an in-house validation
670 study of real-time PCR methods for the detection and serogroup determination
671 of verocytotoxicigenic *Escherichia coli*. *Appl. Environ. Microbiol.* 77, 6954–6963.
672 doi: 10.1128/AEM.05357-11
- 673 Karch, H., Denamur, E., Dobrindt, U., Finlay, B. B., Hengge, R., Johannes,
674 L., et al. (2012). The enemy within us: lessons from the 2011 European
675 *Escherichia coli* O104:H4 outbreak. *EMBO Mol. Med.* 4, 841–848. doi:
676 10.1002/emmm.201201662
- 677 Kimmitt, P. T., Harwood, C. R., and Barer, M. R. (2000). Toxin gene expression by
678 Shiga toxin-producing *Escherichia coli*: the role of antibiotics and the bacterial
679 SOS response. *Emerg. Infect. Dis.* 6, 458–465. doi: 10.3201/eid0605.000503
- 680 Köhler, B., Karch, H., and Schmidt, H. (2000). Antibacterials that are used as
681 growth promoters in animal husbandry can affect the release of Shiga-toxin-
682 2-converting bacteriophages and Shiga toxin 2 from *Escherichia coli* strains.
683 *Microbiology* 146, 1085–1090.
- 684 Langsrud, S., Heir, E., and Rode, T. M. (2013). Survival of Shiga toxin-producing
685 *Escherichia coli* and Stx bacteriophages in moisture enhanced beef. *Meat Sci.* doi:
686 10.1016/j.meatsci.2013.09.022
- 687 Lee, H. S., and Sobsey M. D. (2011). Survival of prototype strains of somatic
688 coliphage families in environmental waters and when exposed to UV low-
689 pressure monochromatic radiation or heat. *Water Res.* 45, 3723–3734. doi:
690 10.1016/j.watres.2011.04.024
- 691 Lin, L., Honh, W., Ji, X., Han, J., Huang, L., and Wei, Y. (2010). Isolation
692 and characterization of an extremely long tail Thermus bacteriophage
693 from Tegchong hot springs in China. *J. Basic Microbiol.* 50, 452–456. doi:
694 10.1002/jobm.201000116
- 695 Loukiadis, E., Nobe, R., Herold, S., Tramuta, C., Ogura, Y., Ooka, T., et al.
696 (2008). Distribution, functional expression, and genetic organization of Cif, a
697 phage-encoded type III-secreted effector from enteropathogenic and enterohe-
698 morrhagic *Escherichia coli*. *J. Bacteriol.* 190, 275–285. doi: 10.1128/JB.00844-07
- 699 Mann, B. A., and Schlauch, J. M. (1997). Transduction of low copy numbers of
700 plasmids by bacteriophages by bacteriophage P22. *Genetics* 146, 447–456.
- 701 Martínez-Castillo, A., Quirós, P., Navarro, F., Miró, E., and Muniesa, M. (2013).
702 Shiga toxin 2-encoding bacteriophages in human fecal samples from healthy
703 individuals. *Appl. Environ. Microbiol.* 79, 4862–4868. doi: 10.1128/AEM.01
704 158-13
- 705 Maurer, J. J., Schmidt, D., Petrosko, P., Sanchez, S., Bolton, L., and Lee, M.
706 D. (1999). Development of primers to O-antigen biosynthesis genes for specific
707 detection of *Escherichia coli* O157 by PCR. *Appl. Environ. Microbiol.* 65,
708 2954–2960.

- 685 Mauro, S. A., and Koudelka, G. B. (2011). Shiga toxin: expression, distribution, and
686 its role in the environment. *Toxins* 3, 608–625. doi: 10.3390/toxins3060608
- 687 Melton-Celsa, A., Mohawk, K., Teel, L., and O'Brien, A. (2012). Pathogenesis
688 of Shiga-toxin producing *Escherichia coli*. *Curr. Top Microbiol. Immunol.* 357, 67–103. doi: 10.1007/82_2011_176
- 689 Meng, Q., Bai, X., Zhao, A., Lan, R., Du, H., Wang, T., et al. (2014).
690 Characterization of Shiga toxin-producing *Escherichia coli* isolated from healthy
691 pigs in China. *BMC Microbiol.* 14:5. doi: 10.1186/1471-2180-14-5
- 692 Mocé-Llivina, L., Jofre, J., and Muniesa, M. (2003). Comparison of polyvinylidene
693 fluoride and polyether sulfone membranes in filtering viral suspensions. *J. Virol.*
694 *Methods.* 109, 99–101. doi: 10.1016/S0166-0934(03)00046-6
- 695 Monaghan, A., Byrne, B., Fanning, S., Sweeney, T., McDowell, D., and Bolton,
696 D. J. (2012). Serotypes and virulotypes of non-O157 shiga-toxin producing
697 *Escherichia coli* (STEC) on bovine hides and carcasses. *Food Microbiol.* 32,
698 223–229. doi: 10.1016/j.fm.2012.06.002
- 699 Monday, S. R., Beisaw, A., and Feng, P. C. (2007). Identification of Shiga toxigenic
700 *Escherichia coli* seropathotypes A and B by multiplex PCR. *Mol. Cell Probes* 21,
701 308–311. doi: 10.1016/j.mcp.2007.02.002
- 702 Muhldorfer, I., Hacker, J., Kusch, G. T., Acheson, D. W. K., Tschape, H., Kane, A.
703 V., et al. (1996). Regulation of the Shiga-like toxin II operon in *Escherichia coli*.
704 *Infect. Immun.* 64, 495–502.
- 705 Muniesa, M., Blanch, A. R., Lucena, F., and Jofre, J. (2005). Bacteriophages may
706 bias outcome of bacterial enrichment cultures. *Appl. Environ. Microbiol.* 71,
707 4269–4275. doi: 10.1128/AEM.71.8.4269-4275.2005
- 708 Muniesa, M., Blanco, J. E., De Simón, M., Serra-Moreno, R., Blanch, A. R., and
709 Jofre, J. (2004a). Diversity of stx2 converting bacteriophages induced from
710 Shiga-toxin-producing *Escherichia coli* strains isolated from cattle. *Microbiology*
711 150, 2959–2971. doi: 10.1099/mic.0.27188-0
- 712 Muniesa, M., and Jofre, J. (2000). Occurrence of phages infecting *Escherichia coli*
713 O157H7 carrying the stx 2 gene in sewage from different countries. *FEMS*
714 *Microbiol. Lett.* 183, 197–200. doi: 10.1111/j.1574-6968.2000.tb08957.x
- 715 Muniesa, M., Lucena, F., and Jofre, J. (1999). Comparative survival of free shiga
716 toxin 2-encoding phages and *Escherichia coli* strains outside the gut. *Appl.*
717 *Environ. Microbiol.* 65, 5615–5618.
- 718 Muniesa, M., Serra-Moreno, R., and Jofre, J. (2004b). Free Shiga toxin bac-
719 teriophages isolated from sewage showed diversity although the stx genes
720 appeared conserved. *Environ. Microbiol.* 6, 716–725. doi: 10.1111/j.1462-
721 2920.2004.00604.x
- 722 Neely, M. N., and Friedman, D. I. (1998). Functional and genetic analysis of
723 regulatory regions of coliphage H-19B: location of shiga-like toxin and lysis
724 genes suggest a role for phage functions in toxin release. *Mol. Microbiol.* 28,
725 1255–1267. doi: 10.1046/j.1365-2958.1998.00890.x
- 726 Newland, J. W., Strockbine, N. A., Miller, S. F., O'Brien, A. D., and Holmes, R.
727 K. (1985). Cloning of Shiga-like toxin structural genes from a toxin converting
728 phage of *Escherichia coli*. *Science* 230, 179–181. doi: 10.1126/science.2994228
- 729 O'Brien, A. D., Marques, L. R., Kerry, C. F., Newland, J. W., and Holmes, R. K.
730 (1989). Shiga-like toxin converting phage of enterohemorrhagic *Escherichia coli*
731 strain 933. *Microb. Pathog.* 6, 381–390. doi: 10.1016/0882-4010(89)90080-6
- 732 Ogunseitan, O. A., Saylor, G. S., and Miller, R. V. (1990). Dynamic interactions
733 of *Pseudomonas aeruginosa* and bacteriophages in lake water. *Microb. Ecol.* 19,
734 171–185.
- 735 Pacheco, A. R., and Sperandio, V. (2009). Inter-kingdom signaling: chemical lan-
736 guage between bacteria and host. *Curr. Opin. Microbiol.* 12, 192–198. doi:
737 10.1016/j.mib.2009.01.006
- 738 Papageorgiou, G. T., Mocé-Llivina, L., Christodoulou, C. G., Lucena, F., Akkelidou,
739 D., Ioannou, E., et al. (2000). A simple methodological approach for count-
740 ing and identifying culturable viruses adsorbed to cellulose nitrate membrane
741 filters. *Appl. Environ. Microbiol.* 66, 194–198. doi: 10.1128/AEM.66.1.194-
742 198.2000
- 743 Paton, J. C., and Paton, A. W. (2003). “Methods for detection of STEC in humans:
744 an overview,” in *Escherichia coli: Shiga Toxin Methods and Protocols. Methods in*
745 *Molecular Medicine*. Vol. 73, eds D. Philpott and F. Ebel (Totowa, NJ: Humana
746 Press).
- 747 Perelle, S., Dilasser, F., Grout, J., and Fach, P. (2004). Detection by 5'-nuclease PCR
748 of Shiga-toxin producing *Escherichia coli* O26, O55, O91, O103, O111, O113,
749 O145 and O157:H7, associated with the world's most frequent clinical cases.
750 *Mol. Cell. Probe.* 18, 185–192. doi: 10.1016/j.mcp.2003.12.004
- 751 Picozzi, C., Volponi, G., Vigentini, I., Grassi, S., and Foschino, R. (2011).
752 Assessment of transduction of *Escherichia coli* Stx2-encoding phage in
753 dairy process conditions. *Int. J. Food Microbiol.* 153, 388–394. doi:
754 10.1016/j.ijfoodmicro.2011.11.031
- 755 Pradel, N., Livrelli, V., De Champs, C., Palcoux, J. B., Reynaud, A., Scheutz, F., et al.
756 (2000). Prevalence and characterization of Shiga toxin-producing *Escherichia*
757 *coli* isolated from cattle, food, and children during a one-year prospective study
758 in France. *J. Clin. Microbiol.* 38, 1023–1031.
- 759 Prigent, M., Leroy, M., Confalonieri, F., Dutertre, M., and DuBow, M. S. (2005). A
760 diversity of bacteriophages forms and genomes can be isolated from the surface
761 sands of Sahara Desert. *Extremophiles* 9, 289–296. doi: 10.1007/s00792-005-
762 0444-5
- 763 Rietra, P. J., Willshaw, G. A., Smith, H. R., Field, A. M., Scotland, S. M., and Rowe,
764 B. (1989). Comparison of Vero-cytotoxin-encoding phages from *Escherichia*
765 *coli* of human and bovine origin. *J. Gen. Microbiol.* 135, 2307–2318. doi:
766 10.1099/00221287-135-8-2307
- 767 Rode, T. M., Axelsson, L., Granum, P. E., Heir, E., Holck, A., and L'abée-Lund,
768 T. M. (2011). High stability of Stx2 phage in food and under food-processing
769 conditions. *Appl. Environ. Microbiol.* 77, 5336–5341. doi: 10.1128/AEM.00
770 180-11
- 771 Rogerie, F., Marecata, A., Gambadeb, S., Duponda, F., Beaubois, P., and Langea,
772 M. (2001). Characterization of Shiga toxin producing *E. coli* and O157 serotype
773 *E. coli* isolated in France from healthy domestic cattle. *Int. J. Food Microbiol.*
774 63, 217–223. doi: 10.1016/S0168-1605(00)00422-0
- 775 Rooks, D. J., Libberton, B., Woodward, M. J., Allison, H. E., and McCarthy, A.
776 J. (2012). Development and application of a method for the purification of
777 free shigatoxigenic bacteriophage from environmental samples. *J. Microbiol.*
778 *Methods* 91, 240–245. doi: 10.1016/j.mimet.2012.08.017
- 779 Rooks, D. J., Yan, Y., McDonald, J. E., Woodward, M. J., McCarthy, A. J., and
780 Allison, H. E. (2010). Development and validation of a qPCR-based method for
781 quantifying Shiga toxin-encoding and other lambdoid bacteriophages. *Environ.*
782 *Microbiol.* 12, 1194–1204. doi: 10.1111/j.1462-2920.2010.02162.x
- 783 Saxena, A., Tripathi, B. P., Kumar, M., and Shahi, V. K. (2008). Membrane-based
784 techniques for the separation and purification of proteins: an overview. *Adv.*
785 *Colloid Interface* 145, 1–22. doi: 10.1016/j.cis.2008.07.004
- 786 Schmieger, H., and Schicklmaier, P. (1999). Transduction of multiple drug resis-
787 tance of *Salmonella enterica* serovar Typhimurium DT104. *FEMS Microbiol.*
788 *Lett.* 170, 251–256. doi: 10.1111/j.1574-6968.1999.tb13381.x
- 789 Stephan, R., Ragettli, S., and Untermann, F. (2000). Prevalence and character-
790 istics of verotoxin-producing *Escherichia coli* (VTEC) in stool samples from
791 asymptomatic human carriers working in the meat processing industry in
792 Switzerland. *J. Appl. Microbiol.* 88, 335–341. doi: 10.1046/j.1365-2672.2000.
793 00965.x
- 794 Tanji, Y., Mizoguchi, K., Yoichi, M., Morita, M., Kijima, N., Kator, H., et al. (2003).
795 Seasonal change and fate of coliphages infected to *Escherichia coli* O157:H7 in
796 a waste water treatment plant. *Water Res.* 37, 1136–1142. doi: 10.1016/S0043-
797 1354(02)00464-5
- 798 Tarr, P. I., Gordon, C. A., and Chandler, W. L. (2005). Shiga-toxin-producing
799 *Escherichia coli* and haemolytic uraemic syndrome. *Lancet* 365, 1073–1086. doi:
800 10.1016/S0140-6736(05)71144-2
- 801 Thomason, L. C., Costantino, N., and Court, D. L. (2007). *E. coli* genome manip-
802 ulation by P1 transduction. *Curr. Protoc. Mol. Biol.* Chapter 1:Unit 1.17. doi:
803 10.1002/0471142727.mb0117s79
- 804 Toshima, H., Yoshimura, A., Arikawa, K., Hidaka, A., Ogasawara, J., Hase, A.,
805 et al. (2007). Enhancement of Shiga toxin production in enterohemorrhagic
806 *Escherichia coli* serotype O157:H7 by DNase colicins. *Appl. Environ. Microbiol.*
807 73, 7582–7588. doi: 10.1128/AEM.01326-07
- 808 Trevisani, M., Mancusi, R., Delle, Donne, G., Bacci, C., Bassi, L., et al. (2014).
809 Detection of Shiga toxin (Stx)-producing *Escherichia coli* (STEC) in bovine
810 dairy herds in Northern Italy. *Int. J. Food Microbiol.* doi: 10.1016/j.ijfoodmicro.
811 789
- 812 Tyler, J. S., Miller, M. J., and Friedman, D. I. (2004). The operator and early pro-
813 moter region of the Shiga toxin type 2-encoding bacteriophage 933W and control
814 of toxin expression. *J. Bacteriol.* 186, 7670–7679. doi: 10.1128/JB.186.22.7670-
815 7679.2004
- 816 Underwood, A. P., Dallman, T., Thomson, N. R., Williams, M., Harker, K., Perry,
817 N., et al. (2013). Public health value of next-generation DNA sequencing of
818 enterohemorrhagic *Escherichia coli* isolates from an outbreak. *J. Clin. Microbiol.*
819 51, 232–237. doi: 10.1128/JCM.01696-12
- 820 Urdahl, A. M., Solheim, H. T., Vold, L., Hasseltvedt, V., and Wasteson, Y. (2012).
821 Shiga toxin-encoding genes (stx genes) in human fecal samples. *APMIS* 121,
822 202–210. doi: 10.1111/j.1600-0463.2012.02957.x

- 799 Vallières, E., Saint-Jean, M., and Rallu, F. (2013). Comparison of three different
800 methods for detection of Shiga toxin-producing *Escherichia coli* in a ter-
801 tiary pediatric care center. *J. Clin. Microbiol.* 51, 481–486. doi: 10.1128/JCM.
02219-12
- 802 Van Reis, R., and Zydney, A. (2001). Membrane separations in biotechnology. *Curr.*
803 *Opin. Biotechnol.* 12, 208–211. doi: 10.1016/S0958-1669(00)00201-9
- 804 Wagner, P. L., Neely, M. N., Zhang, X., Acheson, D. W., Waldor, M. K., and
805 Friedman, D. I. (2001). Role for a phage promoter in Shiga toxin 2 expres-
806 sion from a pathogenic *Escherichia coli* strain. *J. Bacteriol.* 183, 2081–2085. doi:
10.1128/JB.183.6.2081-2085.2001
- 807 Wester, A. L., Brandal, L. T., and Dahle, U. R. (2013). Extraintestinal pathogenic
808 *Escherichia coli* carrying the Shiga Toxin gene *stx*₂. *J. Clin. Microbiol.* 51,
809 4279–4280. doi: 10.1128/JCM.01349-13
- 810 Wong, C. S., Jelacic, S., Habeeb, R. L., Watkins, S. L., and Tarr, P. I. (2000).
811 The risk of the hemolytic-uremic syndrome after antibiotic treatment of
812 *Escherichia coli* O157:H7 infections. *N. Engl. J. Med.* 342, 1930–1936. doi:
10.1056/NEJM200006293422601
- 813 Yamamoto, T., Kojo, S., Taneike, I., Nakagawa, S., Iwakura, N., and Wakisaka-
814 Saito, N. (2003). ⁶⁰Co irradiation of Shiga toxin (Stx)-producing *Escherichia*
815 *coli* induces Stx phage. *FEMS Microbiol. Lett.* 222, 115–121. doi: 10.1016/S0378-
1097(03)00259-3
- 816 Yan, Y., Shi, Y., Cao, D., Meng, X., Xia, L., and Sun, J. (2011). Prevalence of Stx
817 phages in environments of a pig farm and lysogenic infection of the field *E.*
818 *coli* O157 isolates with a recombinant converting Phage. *Curr. Microbiol.* 62,
819 458–464. doi: 10.1007/s00284-010-9729-8
- 820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
- Yue, W. F., Du, M., and Zhu, M. J. (2012). High temperature in combination with
UV irradiation enhances horizontal transfer of *stx*₂ gene from *E. coli* O157:H7 to
non-pathogenic *E. coli*. *PLoS ONE* 7:e31308. doi: 10.1371/journal.pone.0031308
- Zhu, B. (2009). Degradation of plasmid and plant DNA in water microcosms
monitored by natural transformation and real-time polymerase chain reaction
(PCR). *Water Res.* 40, 3231–3238. doi: 10.1016/j.watres.2006.06.040
- Conflict of Interest Statement:** The authors declare that the research was con-
ducted in the absence of any commercial or financial relationships that could be
construed as a potential conflict of interest.
- Received: 06 March 2014; paper pending published: 22 March 2014; accepted: 27
March 2014; published online: xx April 2014.
- Citation: Martínez-Castillo A and Muniesa M (2014) Implications of free Shiga
toxin-converting bacteriophages occurring outside bacteria for the evolution and the
detection of Shiga toxin-producing *Escherichia coli*. *Front. Cell. Infect. Microbiol.* 4:46.
doi: 10.3389/fcimb.2014.00046
- This article was submitted to the journal *Frontiers in Cellular and Infection
Microbiology*.
- Copyright © 2014 Martínez-Castillo and Muniesa. This is an open-access article
distributed under the terms of the Creative Commons Attribution License (CC BY).
The use, distribution or reproduction in other forums is permitted, provided the
original author(s) or licensor are credited and that the original publication in this
journal is cited, in accordance with accepted academic practice. No use, distribution or
reproduction is permitted which does not comply with these terms.
- 856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912